

Dual curvature measures and the dual Minkowski problem

Yiming Zhao

Department of Mathematics
Tandon School of Engineering, New York University

Analysis & Probability seminar, Case Western Reserve University

- Introduce *dual curvature measures*.
 - background, what are they, why are they called “dual” curvature measures

- Introduce *dual curvature measures*.
 - background, what are they, why are they called “dual” curvature measures
- Introduce *the dual Minkowski problem*.

- Introduce *dual curvature measures*.
 - background, what are they, why are they called “dual” curvature measures
- Introduce *the dual Minkowski problem*.
- Solve *the dual Minkowski problem* — in some cases.

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.
- support function $h_K : S^{n-1} \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.
- support function $h_K : S^{n-1} \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.
- support function $h_K : S^{n-1} \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$

can extend h_K to \mathbb{R}^n by making it homogeneous of degree 1.

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.
- support function $h_K : S^{n-1} \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$

can extend h_K to \mathbb{R}^n by making it homogeneous of degree 1.

- For K with $o \in \text{int } K$, radial function $\rho_K : S^{n-1} \rightarrow \mathbb{R}$

$$\rho_K(u) = \max\{r > 0 : ru \in K\}.$$

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.
- support function $h_K : S^{n-1} \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$

can extend h_K to \mathbb{R}^n by making it homogeneous of degree 1.

- For K with $o \in \text{int } K$, radial function $\rho_K : S^{n-1} \rightarrow \mathbb{R}$

$$\rho_K(u) = \max\{r > 0 : ru \in K\}.$$

Basic concepts and notations

- convex bodies K — compact convex subset of \mathbb{R}^n with non-empty interior.
- support function $h_K : S^{n-1} \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$

can extend h_K to \mathbb{R}^n by making it homogeneous of degree 1.

- For K with $o \in \text{int } K$, radial function $\rho_K : S^{n-1} \rightarrow \mathbb{R}$

$$\rho_K(u) = \max\{r > 0 : ru \in K\}.$$

can extend ρ_K to $\mathbb{R}^n \setminus o$ by making it homogeneous of degree -1.

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

- *volume* and *Minkowski combination*

$$V(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

- *volume and Minkowski combination*

$$V(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

- *volume and Minkowski combination*

$$V(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n - i)$ -th quermassintegral $W_{n-i}(K)$

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

- *volume and Minkowski combination*

$$V(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n - i)$ -th quermassintegral $W_{n-i}(K)$

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

- *volume and Minkowski combination*

$$V(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n-i)$ -th quermassintegral $W_{n-i}(K)$
in integral geometry, it is the average of projection areas:

$$W_{n-i}(K) = c \int_{G(n,i)} \mathcal{H}^i(P_\xi K) d\xi$$

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, measures associated with convex bodies.

- *volume and Minkowski combination*

$$V(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n-i)$ -th quermassintegral $W_{n-i}(K)$
in integral geometry, it is the average of projection areas:

$$W_{n-i}(K) = c \int_{G(n,i)} \mathcal{H}^i(P_\xi K) d\xi$$

in differential geometry, integral of mean curvatures:

$$W_{n-i}(K) = c \int_{\partial K} H_{n-i-1}(K, x) dS(x)$$

The Brunn-Minkowski theory

- localized Steiner formula

The Brunn-Minkowski theory

- localized Steiner formula

The Brunn-Minkowski theory

- localized Steiner formula

For $x \notin K$

The Brunn-Minkowski theory

- localized Steiner formula

For $x \notin K$

$p_K(x)$ — closest point to x in K

The Brunn-Minkowski theory

- localized Steiner formula

For $x \notin K$

$p_K(x)$ — closest point to x in K

$d(K, x)$ — distance from x to K

The Brunn-Minkowski theory

- localized Steiner formula

For $x \notin K$

$p_K(x)$ — closest point to x in K

$d(K, x)$ — distance from x to K

$\nu_K(x) = \frac{x - p_K(x)}{|x - p_K(x)|}$ — a unit normal of K at $p_K(x)$

The Brunn-Minkowski theory

- localized Steiner formula

For $x \notin K$

$p_K(x)$ — closest point to x in K

$d(K, x)$ — distance from x to K

$\nu_K(x) = \frac{x - p_K(x)}{|x - p_K(x)|}$ — a unit normal of K at $p_K(x)$

ω, η Borel subsets of S^{n-1} . Define

$$A_t(K, \omega) = \{x \in \mathbb{R}^n : 0 < d(K, x) \leq t \text{ and } \frac{p_K(x)}{|p_K(x)|} \in \omega\}$$

$$B_t(K, \eta) = \{x \in \mathbb{R}^n : 0 < d(K, x) \leq t \text{ and } \nu_K(x) \in \eta\}$$

The Brunn-Minkowski theory

- localized Steiner formula

For $x \notin K$

$p_K(x)$ — closest point to x in K

$d(K, x)$ — distance from x to K

$\nu_K(x) = \frac{x - p_K(x)}{|x - p_K(x)|}$ — a unit normal of K at $p_K(x)$

ω, η Borel subsets of S^{n-1} . Define

$$A_t(K, \omega) = \{x \in \mathbb{R}^n : 0 < d(K, x) \leq t \text{ and } \frac{p_K(x)}{|p_K(x)|} \in \omega\}$$

$$B_t(K, \eta) = \{x \in \mathbb{R}^n : 0 < d(K, x) \leq t \text{ and } \nu_K(x) \in \eta\}$$

Localized Steiner Formula [Schneider, 2014]:

$$V(A_t(K, \omega)) = \frac{1}{n} \sum_{i=0}^{n-1} \binom{n}{i} t^{n-i} G_i(K, \omega) \quad \text{— curvature measures}$$

$$V(B_t(K, \eta)) = \frac{1}{n} \sum_{i=0}^{n-1} \binom{n}{i} t^{n-i} S_i(K, \eta) \quad \text{— area measures}$$

The Brunn-Minkowski theory

In particular,

$C_0(K, \cdot)$ — Aleksandrov's integral curvature

$S_{n-1}(K, \cdot)$ — surface area measure

The Brunn-Minkowski theory

In particular,

$C_0(K, \cdot)$ — Aleksandrov's integral curvature

$S_{n-1}(K, \cdot)$ — surface area measure

Problem

The Minkowski problem. Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that

$$\mu = S_{n-1}(K, \cdot)$$

The Brunn-Minkowski theory

In particular,

$C_0(K, \cdot)$ — Aleksandrov's integral curvature

$S_{n-1}(K, \cdot)$ — surface area measure

Problem

The Minkowski problem. *Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that*

$$\mu = S_{n-1}(K, \cdot)$$

Ans: μ is not concentrated in any closed hemisphere and

$$\int_{S^{n-1}} v d\mu(v) = o.$$

The Brunn-Minkowski theory

In particular,

$C_0(K, \cdot)$ — Aleksandrov's integral curvature

$S_{n-1}(K, \cdot)$ — surface area measure

Problem

The Minkowski problem. *Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that*

$$\mu = S_{n-1}(K, \cdot)$$

Ans: μ is not concentrated in any closed hemisphere and

$$\int_{S^{n-1}} v d\mu(v) = o.$$

Minkowski, Aleksandrov, Fenchel–Jessen, Cheng–Yau

The Brunn-Minkowski theory

Problem

The Minkowski problem. Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that $\mu = S_{n-1}(K, \cdot)$.

The Brunn-Minkowski theory

Problem

The Minkowski problem. Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that $\mu = S_{n-1}(K, \cdot)$.

Remark

- when K is smooth, the density of $S_{n-1}(K, \cdot)$ is the reciporical Gauss curvature. In this case, the Minkowski problem is the problem of prescribing Gauss curvature.

The Brunn-Minkowski theory

Problem

The Minkowski problem. Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that $\mu = S_{n-1}(K, \cdot)$.

Remark

- when K is smooth, the density of $S_{n-1}(K, \cdot)$ is the reciporical Gauss curvature. In this case, the Minkowski problem is the problem of prescribing Gauss curvature.
- also in the smooth case, when μ has a density f , the MP becomes the Monge-Ampère equation

$$\det(h_{ij} + h\delta_{ij}) = f.$$

The Brunn-Minkowski theory

Problem

The Minkowski problem. Given a non-zero finite Borel measure μ on S^{n-1} . Under what condition is there a K such that $\mu = S_{n-1}(K, \cdot)$.

Remark

- when K is smooth, the density of $S_{n-1}(K, \cdot)$ is the reciporical Gauss curvature. In this case, the Minkowski problem is the problem of prescribing Gauss curvature.
- also in the smooth case, when μ has a density f , the MP becomes the Monge-Ampère equation

$$\det(h_{ij} + h\delta_{ij}) = f.$$

- other Minkowski problems: Aleksandrov problem, L_p Minkowski problem — in particular, the logarithmic Minkowski problem [Böröczky-LYZ, JAMS '12]

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s.

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s.

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.
- Busemann-Petty problem. K, L o-symmetric,
 $\mathcal{H}^{n-1}(K \cap H) \leq \mathcal{H}^{n-1}(L \cap H)$ for every hyperplane H that crosses the origin
Is it true that $V(K) \leq V(L)$?

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.
- Busemann-Petty problem. K, L o-symmetric,
 $\mathcal{H}^{n-1}(K \cap H) \leq \mathcal{H}^{n-1}(L \cap H)$ for every hyperplane H that crosses the origin
Is it true that $V(K) \leq V(L)$?

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.
- Busemann-Petty problem. K, L o-symmetric,
 $\mathcal{H}^{n-1}(K \cap H) \leq \mathcal{H}^{n-1}(L \cap H)$ for every hyperplane H that crosses the origin
Is it true that $V(K) \leq V(L)$? — true when $n = 3, 4$, false when $n \geq 5$.

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.
- Busemann-Petty problem. K, L o-symmetric,
 $\mathcal{H}^{n-1}(K \cap H) \leq \mathcal{H}^{n-1}(L \cap H)$ for every hyperplane H that crosses the origin
Is it true that $V(K) \leq V(L)$? — true when $n = 3, 4$, false when $n \geq 5$.
- no measure, no PDE

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.
- Busemann-Petty problem. K, L o-symmetric,
 $\mathcal{H}^{n-1}(K \cap H) \leq \mathcal{H}^{n-1}(L \cap H)$ for every hyperplane H that crosses the origin
Is it true that $V(K) \leq V(L)$? — true when $n = 3, 4$, false when $n \geq 5$.
- no measure, no PDE

The dual Brunn-Minkowski theory

- introduced by Lutwak in 1970s. [Schneider, '14], [Gardner, '06].
- many “dual” concepts and results — radial summation, dual mixed volume, intersection body, etc.
- Busemann-Petty problem. K, L o-symmetric,
 $\mathcal{H}^{n-1}(K \cap H) \leq \mathcal{H}^{n-1}(L \cap H)$ for every hyperplane H that crosses the origin
Is it true that $V(K) \leq V(L)$? — true when $n = 3, 4$, false when $n \geq 5$.
- no measure, no PDE — until [Huang-LYZ, ACTA 2016]

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} t \cdot L$

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} t \cdot L$

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+}_t L$

$$\rho_{K \tilde{+}_t L} = \rho_K + t \rho_L$$

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} t \cdot L$

$$\rho_{K \tilde{+} t \cdot L} = \rho_K + t \rho_L$$

- dual Steiner formula

$$V(K \tilde{+} t \cdot B) = \sum_{i=0}^n t^{n-i} \binom{n}{i} \tilde{W}_{n-i}(K)$$

— dual quermassintegral

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} t \cdot L$

$$\rho_{K \tilde{+} t \cdot L} = \rho_K + t \rho_L$$

- dual Steiner formula

$$V(K \tilde{+} t \cdot B) = \sum_{i=0}^n t^{n-i} \binom{n}{i} \tilde{W}_{n-i}(K) \quad \text{— dual quermassintegral}$$

- dual quermassintegral $\tilde{W}_{n-i}(K)$

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} t \cdot L$

$$\rho_{K \tilde{+} t \cdot L} = \rho_K + t \rho_L$$

- dual Steiner formula

$$V(K \tilde{+} t \cdot B) = \sum_{i=0}^n t^{n-i} \binom{n}{i} \tilde{W}_{n-i}(K) \quad \text{— dual quermassintegral}$$

- dual quermassintegral $\tilde{W}_{n-i}(K)$

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} t \cdot L$

$$\rho_{K \tilde{+} t \cdot L} = \rho_K + t \rho_L$$

- dual Steiner formula

$$V(K \tilde{+} t \cdot B) = \sum_{i=0}^n t^{n-i} \binom{n}{i} \tilde{W}_{n-i}(K) \quad \text{— dual quermassintegral}$$

- dual quermassintegral $\tilde{W}_{n-i}(K)$
in integral geometry, mean of intersection areas:

$$\tilde{W}_{n-i}(K) = c \int_{G(n,i)} \mathcal{H}^i(K \cap \xi) d\xi.$$

The dual Brunn-Minkowski theory

- radial combination $K \tilde{+} tL$

$$\rho_{K \tilde{+} tL} = \rho_K + t\rho_L$$

- dual Steiner formula

$$V(K \tilde{+} tB) = \sum_{i=0}^n t^{n-i} \binom{n}{i} \tilde{W}_{n-i}(K) \quad \text{— dual quermassintegral}$$

- dual quermassintegral $\tilde{W}_{n-i}(K)$

in integral geometry, mean of intersection areas:

$$\tilde{W}_{n-i}(K) = c \int_{G(n,i)} \mathcal{H}^i(K \cap \xi) d\xi.$$

has integral form: $\tilde{W}_{n-i}(K) = \frac{1}{n} \int_{S^{n-1}} \rho_K^i(u) du = \frac{i}{n} \int_K |x|^{i-n} dx.$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

For $x \in \mathbb{R}^n \setminus o$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

For $x \in \mathbb{R}^n \setminus o$

$\tilde{\rho}_K(x) = \rho_K(x)x \in \partial K$ — the point on ∂K that shares the same direction as x .

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

For $x \in \mathbb{R}^n \setminus o$

$\tilde{p}_K(x) = \rho_K(x)x \in \partial K$ — the point on ∂K that shares the same direction as x .

$$\tilde{d}(K, x) = |x - \tilde{p}_K(x)|$$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

For $x \in \mathbb{R}^n \setminus o$

$\tilde{p}_K(x) = \rho_K(x)x \in \partial K$ — the point on ∂K that shares the same direction as x .

$$\tilde{d}(K, x) = |x - \tilde{p}_K(x)|$$

$$\tilde{d}(K, x) = 0 \text{ if } x \in K$$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

For $x \in \mathbb{R}^n \setminus o$

$\tilde{p}_K(x) = \rho_K(x)x \in \partial K$ — the point on ∂K that shares the same direction as x .

$$\tilde{d}(K, x) = |x - \tilde{p}_K(x)|$$

$$\tilde{d}(K, x) = 0 \text{ if } x \in K$$

ω, η Borel subsets of S^{n-1} . Define

$$\tilde{A}_t(K, \eta) = \{x \in \mathbb{R}^n : 0 \leq \tilde{d}(K, x) \leq t$$

and K at $\tilde{p}_K(x)$ has outer normal in $\eta\}$

$$\tilde{B}_t(K, \omega) = \{x \in \mathbb{R}^n : 0 \leq \tilde{d}(K, x) \leq t$$

and $\frac{x}{|x|} \in \omega\}$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

$$V(\tilde{A}_t(K, \eta)) = \sum_{i=0}^n \binom{n}{i} t^{n-i} \tilde{C}_i(K, \eta) \quad \text{— dual curvature measure}$$

$$V(\tilde{B}_t(K, \eta)) = \sum_{i=0}^n \binom{n}{i} t^{n-i} \tilde{S}_i(K, \omega) \quad \text{— dual area measure}$$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

$$V(\tilde{A}_t(K, \eta)) = \sum_{i=0}^n \binom{n}{i} t^{n-i} \tilde{C}_i(K, \eta) \quad \text{— dual curvature measure}$$

$$V(\tilde{B}_t(K, \eta)) = \sum_{i=0}^n \binom{n}{i} t^{n-i} \tilde{S}_i(K, \omega) \quad \text{— dual area measure}$$

- $\tilde{C}_0(K, \cdot)$ is “the same as” $C_0(K^*, \cdot)$

The dual Brunn-Minkowski theory

- localized dual Steiner formula [Huang-LYZ, ACTA 2016]

$$V(\tilde{A}_t(K, \eta)) = \sum_{i=0}^n \binom{n}{i} t^{n-i} \tilde{C}_i(K, \eta) \quad \text{— dual curvature measure}$$

$$V(\tilde{B}_t(K, \eta)) = \sum_{i=0}^n \binom{n}{i} t^{n-i} \tilde{S}_i(K, \omega) \quad \text{— dual area measure}$$

- $\tilde{C}_0(K, \cdot)$ is “the same as” $C_0(K^*, \cdot)$
- $\tilde{C}_n(K, \cdot)$ is “the same as” cone volume measure (L_p surface area measure when $p = 0$)

- integral representation:

$$\tilde{C}_i(K, \eta) = \frac{1}{n} \int_{\alpha_K^*(\eta)} \rho_K^i(u) du,$$

where α_K^* is very similar to inverse Gauss map.

- integral representation:

$$\tilde{C}_i(K, \eta) = \frac{1}{n} \int_{\alpha_K^*(\eta)} \rho_K^i(u) du,$$

where α_K^* is very similar to inverse Gauss map.

- easily extends to all real numbers. (will use $\tilde{C}_q(K, \cdot)$)

The dual Minkowski problem

Problem

The dual Minkowski problem: Given a nonzero finite Borel measure μ on S^{n-1} and $q \in \mathbb{R}$, under what condition(s) on μ does there exist a K such that $\mu = \tilde{C}_q(K, \cdot)$?

The dual Minkowski problem

Problem

The dual Minkowski problem: Given a nonzero finite Borel measure μ on S^{n-1} and $q \in \mathbb{R}$, under what condition(s) on μ does there exist a K such that $\mu = \tilde{C}_q(K, \cdot)$?

$$\det(h_{ij}(v) + h(v)\delta_{ij}) = nh^{-1}(v)|\nabla_{S^{n-1}}h(v) + h(v)v|^{n-q}f(v)$$

The dual Minkowski problem

Problem

The dual Minkowski problem: Given a nonzero finite Borel measure μ on S^{n-1} and $q \in \mathbb{R}$, under what condition(s) on μ does there exist a K such that $\mu = \tilde{C}_q(K, \cdot)$?

$$\det(h_{ij}(v) + h(v)\delta_{ij}) = nh^{-1}(v)|\nabla_{S^{n-1}}h(v) + h(v)v|^{n-q}f(v)$$

Problem

The even dual Minkowski problem: Given a nonzero even finite Borel measure μ on S^{n-1} and $q \in \mathbb{R}$, under what condition(s) on μ does there exist an o -symm K such that $\mu = \tilde{C}_q(K, \cdot)$?

The dual Minkowski problem

Problem

The dual Minkowski problem: Given a nonzero finite Borel measure μ on S^{n-1} and $q \in \mathbb{R}$, under what condition(s) on μ does there exist a K such that $\mu = \tilde{C}_q(K, \cdot)$?

$$\det(h_{ij}(v) + h(v)\delta_{ij}) = nh^{-1}(v)|\nabla_{S^{n-1}}h(v) + h(v)v|^{n-q}f(v)$$

Problem

The even dual Minkowski problem: Given a nonzero even finite Borel measure μ on S^{n-1} and $q \in \mathbb{R}$, under what condition(s) on μ does there exist an o -symm K such that $\mu = \tilde{C}_q(K, \cdot)$?

$q = 0$ Aleksandrov problem, $q = n$ logarithmic Minkowski problem

When $q < 0$, complete solution to the DMP

Theorem (Z., CVPDEs 2017)

Suppose μ is a non-zero finite Borel measure on S^{n-1} and $q < 0$. There exists a K such that $\mu = \tilde{C}_q(K, \cdot)$ iff μ is not concentrated in any closed hemisphere. Moreover, K is unique (if exists).

Results

When $0 < q < n$, even case

Results

When $0 < q < n$, even case
 q -th subspace mass inequality (I):

We say that μ satisfies the q -th subspace mass inequality, if
when $1 < q < n$

$$\frac{\mu(S^{n-1} \cap \xi_i)}{|\mu|} < 1 - \frac{(n-i)(q-1)}{(n-1)q},$$

for each $i = 1, \dots, n-1$ and each i -dim subspace ξ_i ;

Results

When $0 < q < n$, even case
 q -th subspace mass inequality (I):

We say that μ satisfies the q -th subspace mass inequality, if
when $1 < q < n$

$$\frac{\mu(S^{n-1} \cap \xi_i)}{|\mu|} < 1 - \frac{(n-i)(q-1)}{(n-1)q},$$

for each $i = 1, \dots, n-1$ and each i -dim subspace ξ_i ;
when $0 < q \leq 1$

μ is not concentrated in any subspheres.

Theorem (Huang-LYZ, ACTA 2016)

Suppose μ is a nonzero even finite Borel measure on S^{n-1} and $0 < q < n$. If μ satisfies the q -th subspace mass inequality (I), then there exists an o-symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Results

Theorem (Huang-LYZ, ACTA 2016)

Suppose μ is a nonzero even finite Borel measure on S^{n-1} and $0 < q < n$. If μ satisfies the q -th subspace mass inequality (I), then there exists an o -symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Remark

- $0 < q \leq 1$, μ not concentrated in any subsphere, obviously optimal.

Theorem (Huang-LYZ, ACTA 2016)

Suppose μ is a nonzero even finite Borel measure on S^{n-1} and $0 < q < n$. If μ satisfies the q -th subspace mass inequality (I), then there exists an o -symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Remark

- $0 < q \leq 1$, μ not concentrated in any subsphere, obviously optimal.
- $1 < q < n$, the condition is not necessary.

Theorem (Huang-LYZ, ACTA 2016)

Suppose μ is a nonzero even finite Borel measure on S^{n-1} and $0 < q < n$. If μ satisfies the q -th subspace mass inequality (I), then there exists an o -symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Remark

- $0 < q \leq 1$, μ not concentrated in any subsphere, obviously optimal.
- $1 < q < n$, the condition is not necessary.

Theorem (Huang-LYZ, ACTA 2016)

Suppose μ is a nonzero even finite Borel measure on S^{n-1} and $0 < q < n$. If μ satisfies the q -th subspace mass inequality (I), then there exists an o -symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Remark

- $0 < q \leq 1$, μ not concentrated in any subsphere, obviously optimal.
- $1 < q < n$, the condition is not necessary.

$$T_a = \left\{ x \in \mathbb{R}^n : x_1^2 + \cdots + x_i^2 \leq a^2 \text{ and } x_{i+1}^2 + \cdots + x_n^2 \leq 1 \right\}$$

Results

when $q \in (1, n)$

Results

when $q \in (1, n)$

q -th subspace mass inequality (II):

Results

when $q \in (1, n)$

q -th subspace mass inequality (II):

We say that μ satisfies **the q -th subspace mass inequality**, if

$$\frac{\mu(S^{n-1} \cap \xi_i)}{|\mu|} < \begin{cases} \frac{i}{q}, & \text{if } i < q, \\ 1, & \text{if } i \geq q, \end{cases}$$

for each $i = 1, \dots, n-1$ and each i -dim subspace ξ_i .

Results

when $q \in (1, n)$

q -th subspace mass inequality (II):

We say that μ satisfies **the q -th subspace mass inequality**, if

$$\frac{\mu(S^{n-1} \cap \xi_i)}{|\mu|} < \begin{cases} \frac{i}{q}, & \text{if } i < q, \\ 1, & \text{if } i \geq q, \end{cases}$$

for each $i = 1, \dots, n-1$ and each i -dim subspace ξ_i .

Theorem (Z., JDG, to appear)

Suppose μ is a non-zero even finite Borel measure on S^{n-1} and q is an integer in $(1, n)$. If μ satisfies the q -th subspace mass inequality (II), then there exists o-symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Results

when $q \in (1, n)$

q -th subspace mass inequality (II):

We say that μ satisfies **the q -th subspace mass inequality**, if

$$\frac{\mu(S^{n-1} \cap \xi_i)}{|\mu|} < \begin{cases} \frac{i}{q}, & \text{if } i < q, \\ 1, & \text{if } i \geq q, \end{cases}$$

for each $i = 1, \dots, n-1$ and each i -dim subspace ξ_i .

Theorem (Z., JDG, to appear)

Suppose μ is a non-zero even finite Borel measure on S^{n-1} and q is an integer in $(1, n)$. If μ satisfies the q -th subspace mass inequality (II), then there exists o-symm K such that $\mu = \tilde{C}_q(K, \cdot)$.

Theorem (Böröczky & Henk & Pollehn, JDG, to appear)

q -th subspace mass inequality (II) is also necessary for $q \in (1, n)$.

- 1 find a maximization problem whose Euler-Lagrange equation implies that the maximizer is the solution to the DMP.

- 1 find a maximization problem whose Euler-Lagrange equation implies that the maximizer is the solution to the DMP.
- 2 show the maximum can be obtained.

The maximization problem

Given μ even.

The maximization problem

Given μ even. Define $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h]),$$

where $[h] = \{x \in \mathbb{R}^n : x \cdot v \leq h(v), \forall v \in S^{n-1}\}$ is *the Wulff shape*.

The maximization problem

Given μ even. Define $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h]),$$

where $[h] = \{x \in \mathbb{R}^n : x \cdot v \leq h(v), \forall v \in S^{n-1}\}$ is *the Wulff shape*. Obviously Φ is homogeneous of degree 0.

The maximization problem

Given μ even. Define $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h]),$$

where $[h] = \{x \in \mathbb{R}^n : x \cdot v \leq h(v), \forall v \in S^{n-1}\}$ is *the Wulff shape*. Obviously Φ is homogeneous of degree 0.

Problem (The maximization problem)

$$\sup\{\Phi(h) : h \in C_e^+(S^{n-1})\}$$

The maximization problem

Given μ even. Define $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h]),$$

where $[h] = \{x \in \mathbb{R}^n : x \cdot v \leq h(v), \forall v \in S^{n-1}\}$ is *the Wulff shape*. Obviously Φ is homogeneous of degree 0.

Problem (The maximization problem)

$$\sup\{\Phi(h) : h \in C_e^+(S^{n-1})\}$$

Lemma

If a maximizer exists, then it must be the support function of some o-symm convex body.

The maximization problem

Given μ even. Define $\Phi : C_e^+(S^{n-1}) \rightarrow \mathbb{R}$ by

$$\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h]),$$

where $[h] = \{x \in \mathbb{R}^n : x \cdot v \leq h(v), \forall v \in S^{n-1}\}$ is the Wulff shape. Obviously Φ is homogeneous of degree 0.

Problem (The maximization problem)

$$\sup\{\Phi(h) : h \in C_e^+(S^{n-1})\}$$

Lemma

If a maximizer exists, then it must be the support function of some o-symm convex body. Moreover, the maximizer (up to a dilation) is a solution to the DMP.

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

Recall $\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h]).$

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

Recall $\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h])$.

Proof to the lemma.

Suppose h_K is the maximizer.

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

Recall $\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h])$.

Proof to the lemma.

Suppose h_K is the maximizer. Because Φ is homogeneous of degree 0, we may dilate K so that $\tilde{W}_{n-q}(K) = |\mu|$.

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

Recall $\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h])$.

Proof to the lemma.

Suppose h_K is the maximizer. Because Φ is homogeneous of degree 0, we may dilate K so that $\tilde{W}_{n-q}(K) = |\mu|$.

For arbitrary even cont. f , consider $h_t = h_K e^{tf}$.

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

Recall $\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h])$.

Proof to the lemma.

Suppose h_K is the maximizer. Because Φ is homogeneous of degree 0, we may dilate K so that $\tilde{W}_{n-q}(K) = |\mu|$.

For arbitrary even cont. f , consider $h_t = h_K e^{tf}$.

$$0 = \frac{\partial}{\partial t} \bigg|_{t=0} \Phi(h_t)$$

The maximization problem

Variational formula [Huang-LYZ, ACTA 2016]: convex body K , continuous function f on the sphere

$$\frac{\partial}{\partial t} \bigg|_{t=0} \log \tilde{W}_{n-q}([h_K e^{tf}]) = \frac{q}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v).$$

Recall $\Phi(h) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}([h])$.

Proof to the lemma.

Suppose h_K is the maximizer. Because Φ is homogeneous of degree 0, we may dilate K so that $\tilde{W}_{n-q}(K) = |\mu|$.

For arbitrary even cont. f , consider $h_t = h_K e^{tf}$.

$$\begin{aligned} 0 &= \frac{\partial}{\partial t} \bigg|_{t=0} \Phi(h_t) \\ &= -\frac{1}{|\mu|} \int_{S^{n-1}} f(v) d\mu(v) + \frac{1}{\tilde{W}_{n-q}(K)} \int_{S^{n-1}} f(v) d\tilde{C}_q(K, v). \end{aligned}$$

Existence of maximizer

Problem (The maximization problem)

$$\sup\{\Phi(h_K) : K \in \mathcal{K}_e^n\},$$

$$\text{where } \Phi(h_K) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_K d\mu + \frac{1}{q} \log \tilde{W}_{n-q}(K).$$

Existence of maximizer

Problem (The maximization problem)

$$\sup\{\Phi(h_K) : K \in \mathcal{K}_e^n\},$$

$$\text{where } \Phi(h_K) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_K d\mu + \frac{1}{q} \log \tilde{W}_{n-q}(K).$$

sketch of the proof.

Suppose o-symm Q_j is a maximizing sequence.

Existence of maximizer

Problem (The maximization problem)

$$\sup\{\Phi(h_K) : K \in \mathcal{K}_e^n\},$$

$$\text{where } \Phi(h_K) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_K d\mu + \frac{1}{q} \log \tilde{W}_{n-q}(K).$$

sketch of the proof.

Suppose o-symm Q_l is a maximizing sequence. Since Φ is homogeneous of degree 0, assume diameter of each Q_l is 1.

Existence of maximizer

Problem (The maximization problem)

$$\sup\{\Phi(h_K) : K \in \mathcal{K}_e^n\},$$

$$\text{where } \Phi(h_K) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_K d\mu + \frac{1}{q} \log \tilde{W}_{n-q}(K).$$

sketch of the proof.

Suppose o-symm Q_l is a maximizing sequence. Since Φ is homogeneous of degree 0, assume diameter of each Q_l is 1. By Blaschke's selection theorem, (taking a subsequence) we may assume $Q_l \rightarrow Q_0$.

Existence of maximizer

Problem (The maximization problem)

$$\sup\{\Phi(h_K) : K \in \mathcal{K}_e^n\},$$

$$\text{where } \Phi(h_K) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_K d\mu + \frac{1}{q} \log \tilde{W}_{n-q}(K).$$

sketch of the proof.

Suppose o-symm Q_l is a maximizing sequence. Since Φ is homogeneous of degree 0, assume diameter of each Q_l is 1. By Blaschke's selection theorem, (taking a subsequence) we may assume $Q_l \rightarrow Q_0$. In general Q_0 is compact, convex, o-symm.

Existence of maximizer

Problem (The maximization problem)

$$\sup\{\Phi(h_K) : K \in \mathcal{K}_e^n\},$$

$$\text{where } \Phi(h_K) = -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_K d\mu + \frac{1}{q} \log \tilde{W}_{n-q}(K).$$

sketch of the proof.

Suppose o -symm Q_l is a maximizing sequence. Since Φ is homogeneous of degree 0, assume diameter of each Q_l is 1. By Blaschke's selection theorem, (taking a subsequence) we may assume $Q_l \rightarrow Q_0$.

In general Q_0 is compact, convex, o -symm.

If we can further show $o \in \text{int } Q_0$, or $Q_0 \in \mathcal{K}_e^n$, then done (by continuity of Φ). □

Existence of maximizer

- Show $o \in \text{int } Q_0$

Existence of maximizer

- Show $o \in \text{int } Q_0$
Assume otherwise.

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace.

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace. **At the end of the day, show $\Phi(h_{Q_l}) \rightarrow -\infty$.**

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace. **At the end of the day, show $\Phi(h_{Q_l}) \rightarrow -\infty$.**

John ellipsoid E_l of Q_l , $E_l \subset Q_l \subset \sqrt{n}E_l$.

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace. **At the end of the day, show $\Phi(h_{Q_l}) \rightarrow -\infty$.**

John ellipsoid E_l of Q_l , $E_l \subset Q_l \subset \sqrt{n}E_l$.

Say

$$E_l = \{x \in \mathbb{R}^n : \frac{|x \cdot e_{1l}|^2}{a_{1l}^2} + \dots + \frac{|x \cdot e_{nl}|^2}{a_{nl}^2} \leq 1\},$$

where $0 < a_{1l} \leq a_{2l} \leq \dots \leq a_{nl} \leq 1$.

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace. **At the end of the day, show $\Phi(h_{Q_l}) \rightarrow -\infty$.**

John ellipsoid E_l of Q_l , $E_l \subset Q_l \subset \sqrt{n}E_l$.

Say

$$E_l = \{x \in \mathbb{R}^n : \frac{|x \cdot e_{1l}|^2}{a_{1l}^2} + \dots + \frac{|x \cdot e_{nl}|^2}{a_{nl}^2} \leq 1\},$$

where $0 < a_{1l} \leq a_{2l} \leq \dots \leq a_{nl} \leq 1$.

Assume $a_{1l}, \dots, a_{kl} \rightarrow 0$ and $a_{k+1,l}, \dots, a_{nl} \rightarrow > 0$

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace. **At the end of the day, show $\Phi(h_{Q_l}) \rightarrow -\infty$.**

John ellipsoid E_l of Q_l , $E_l \subset Q_l \subset \sqrt{n}E_l$.

Say

$$E_l = \{x \in \mathbb{R}^n : \frac{|x \cdot e_{1l}|^2}{a_{1l}^2} + \dots + \frac{|x \cdot e_{nl}|^2}{a_{nl}^2} \leq 1\},$$

where $0 < a_{1l} \leq a_{2l} \leq \dots \leq a_{nl} \leq 1$.

Assume $a_{1l}, \dots, a_{kl} \rightarrow 0$ and $a_{k+1,l}, \dots, a_{nl} \rightarrow > 0$

Let

$$T_l = \{x \in \mathbb{R}^n : \frac{|x \cdot e_{1l}|^2}{a_{1l}^2} + \dots + \frac{|x \cdot e_{kl}|^2}{a_{kl}^2} \leq 1$$
$$\text{and } |x \cdot e_{k+1,l}|^2 + \dots + |x \cdot e_{nl}|^2 \leq 1\}.$$

Existence of maximizer

- Show $o \in \text{int } Q_0$

Assume otherwise. Then Q_0 is contained in some lower dimensional subspace. **At the end of the day, show $\Phi(h_{Q_l}) \rightarrow -\infty$.**

John ellipsoid E_l of Q_l , $E_l \subset Q_l \subset \sqrt{n}E_l$.

Say

$$E_l = \{x \in \mathbb{R}^n : \frac{|x \cdot e_{1l}|^2}{a_{1l}^2} + \dots + \frac{|x \cdot e_{nl}|^2}{a_{nl}^2} \leq 1\},$$

where $0 < a_{1l} \leq a_{2l} \leq \dots \leq a_{nl} \leq 1$.

Assume $a_{1l}, \dots, a_{kl} \rightarrow 0$ and $a_{k+1,l}, \dots, a_{nl} \rightarrow > 0$

Let

$$T_l = \{x \in \mathbb{R}^n : \frac{|x \cdot e_{1l}|^2}{a_{1l}^2} + \dots + \frac{|x \cdot e_{kl}|^2}{a_{kl}^2} \leq 1$$
$$\text{and } |x \cdot e_{k+1,l}|^2 + \dots + |x \cdot e_{nl}|^2 \leq 1\}.$$

Obviously $E_l \subset T_l$.

Hence $E_I \subset Q_I \subset \sqrt{n}T_I$.

Hence $E_I \subset Q_I \subset \sqrt{n}T_I$. This implies

$$\Phi(h_{Q_I}) \leq -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_{E_I}(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}(T_I) + c.$$

Hence $E_I \subset Q_I \subset \sqrt{n}T_I$. This implies

$$\Phi(h_{Q_I}) \leq -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_{E_I}(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}(T_I) + c.$$

Estimate the first term by “partition of the sphere” and “subspace mass inequality”

Hence $E_I \subset Q_I \subset \sqrt{n}T_I$. This implies

$$\Phi(h_{Q_I}) \leq -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_{E_I}(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}(T_I) + c.$$

Estimate the first term by “partition of the sphere” and “subspace mass inequality”

$$\text{first term} \leq -\left(\frac{1}{q} - \varepsilon_0\right) \log(a_{1I} \cdots a_{kI}) + c$$

Hence $E_I \subset Q_I \subset \sqrt{n}T_I$. This implies

$$\Phi(h_{Q_I}) \leq -\frac{1}{|\mu|} \int_{S^{n-1}} \log h_{E_I}(v) d\mu(v) + \frac{1}{q} \log \tilde{W}_{n-q}(T_I) + c.$$

Estimate the first term by “partition of the sphere” and “subspace mass inequality”

$$\text{first term} \leq -\left(\frac{1}{q} - \varepsilon_0\right) \log(a_{1I} \cdots a_{kI}) + c$$

Estimate the second term by using “spherical coordinates”

$$\text{second term} \leq \frac{1}{q} \log(a_{1I} \cdots a_{kI}) + c$$

Remaining problems

- What about $q > n$?

Remaining problems

- What about $q > n$?
- What about non-even case?

Remaining problems

- What about $q > n$?
- What about non-even case?
- Uniqueness when $q > 0$?

Remaining problems

- What about $q > n$?
- What about non-even case?
- Uniqueness when $q > 0$?
- Regularity?

Remaining problems

- What about $q > n$?
- What about non-even case?
- Uniqueness when $q > 0$?
- Regularity?
- Everything that can be done with surface area measure.