

Geometric quantities and their Minkowski problems

Yiming Zhao

Department of Mathematics
Tandon School of Engineering, New York University

University of Minnesota Morris

Basics and Notations

- The space: Euclidean space \mathbb{R}^2

Basics and Notations

- The space: Euclidean space \mathbb{R}^2

Basics and Notations

- The space: Euclidean space \mathbb{R}^2
— but everything extends to \mathbb{R}^3 or \mathbb{R}^n

Basics and Notations

- The space: Euclidean space \mathbb{R}^2
— but everything extends to \mathbb{R}^3 or \mathbb{R}^n
- The objects to study: *convex bodies* K, L

Basics and Notations

- The space: Euclidean space \mathbb{R}^2
— but everything extends to \mathbb{R}^3 or \mathbb{R}^n
- The objects to study: *convex bodies* K, L

Basics and Notations



- The space: Euclidean space \mathbb{R}^2
— but everything extends to \mathbb{R}^3 or \mathbb{R}^n
- The objects to study: *convex bodies* K, L
— compact (closed and bounded) convex subset of \mathbb{R}^2 with positive area

Basics and Notations

- The space: Euclidean space \mathbb{R}^2
— but everything extends to \mathbb{R}^3 or \mathbb{R}^n
- The objects to study: *convex bodies* K, L
— compact (closed and bounded) convex subset of \mathbb{R}^2 with positive area
e.g., disk, square, or, any convex polygon.

Basics and Notations

- The space: Euclidean space \mathbb{R}^2
— but everything extends to \mathbb{R}^3 or \mathbb{R}^n
- The objects to study: *convex bodies* K, L
— compact (closed and bounded) convex subset of \mathbb{R}^2 with positive area
e.g., disk, square, or, any convex polygon.




Not a convex body!

Basics and Notations

- support function $h_K : S^1 \rightarrow \mathbb{R}$

$$h_K(v) = \max\{x \cdot v : x \in K\}$$



The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- *Area and Minkowski combination*

$$A(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- *Area and Minkowski combination*

$$A(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$A(K + t \cdot D) = \binom{2}{0} w_0(K) + \binom{2}{1} w_1(K)t + \binom{2}{2} w_2(K)t^2,$$

where D is the unit disk in \mathbb{R}^2 .

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- *Area and Minkowski combination*

$$A(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$A(K + t \cdot D) = \binom{2}{0} w_0(K) + \binom{2}{1} w_1(K)t + \binom{2}{2} w_2(K)t^2,$$

where D is the unit disk in \mathbb{R}^2 .

- $w_i(K)$ *quermassintegrals* of K

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- Area and *Minkowski combination*

$$A(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$A(K + t \cdot D) = \binom{2}{0} w_0(K) + \binom{2}{1} w_1(K)t + \binom{2}{2} w_2(K)t^2,$$

where D is the unit disk in \mathbb{R}^2 .

- $w_i(K)$ *quermassintegrals* of K
- $w_0(K)$ — Area of K

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- Area and *Minkowski combination*

$$A(K) \quad K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$A(K + t \cdot D) = \binom{2}{0} W_0(K) + \binom{2}{1} W_1(K)t + \binom{2}{2} W_2(K)t^2,$$

where D is the unit disk in \mathbb{R}^2 .

- $W_i(K)$ *quermassintegrals* of K
- $W_0(K)$ — Area of K
- $W_1(K)$ — (up to a constant independent on K) Perimeter of K

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- Area and *Minkowski combination*

$$A(K + t \cdot L) = \int_{\mathbb{R}^2} \chi_{K + t \cdot L}(x) dx = \int_{\mathbb{R}^2} \chi_K(x) + t \chi_L(x) dx = A(K) + t A(L)$$
$$K + t \cdot L = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$A(K + t \cdot D) = \binom{2}{0} W_0(K) + \binom{2}{1} W_1(K)t + \binom{2}{2} W_2(K)t^2,$$

where D is the unit disk in \mathbb{R}^2 .

- $W_i(K)$ *quermassintegrals* of K
- $W_0(K)$ — Area of K
- $W_1(K)$ — (up to a constant independent on K) Perimeter of K
- $W_2(K)$ — doesn't depend on K at all, in fact *Area of the unit disk*

The Brunn-Minkowski theory

The BM theory studies convex bodies and geometric invariants, *measures* associated with convex bodies.

- Area and *Minkowski combination*

$$A(K + t \cdot L) = \{x + ty : x \in K, y \in L\}$$

- Steiner formula

$$A(K + t \cdot D) = \binom{2}{0} W_0(K) + \binom{2}{1} W_1(K)t + \binom{2}{2} W_2(K)t^2,$$

where D is the unit disk in \mathbb{R}^2 .

- $W_i(K)$ *quermassintegrals* of K
- $W_0(K)$ — Area of K
- $W_1(K)$ — (up to a constant independent on K) Perimeter of K
- $W_2(K)$ — doesn't depend on K at all, in fact *Area of the unit disk*
- Things get more interesting for Euclidean spaces of dimensions beyond 2. In dimension n , you get $n + 1$ quermassintegrals!

The Brunn-Minkowski theory

Let's deviate to dimension n *only* for a minute!

- Steiner formula in \mathbb{R}^n

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

The Brunn-Minkowski theory

Let's deviate to dimension n *only* for a minute!

- Steiner formula in \mathbb{R}^n

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n - i)$ -th quermassintegral $W_{n-i}(K)$

The Brunn-Minkowski theory

Let's deviate to dimension n *only* for a minute!

- Steiner formula in \mathbb{R}^n

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n - i)$ -th quermassintegral $W_{n-i}(K)$

The Brunn-Minkowski theory

Let's deviate to dimension n *only* for a minute!

- Steiner formula in \mathbb{R}^n

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n - i)$ -th quermassintegral $W_{n-i}(K)$
in integral geometry, it is the average of projection areas:

$$W_{n-i}(K) = c \int_{G(n,i)} \mathcal{H}^i(P_\xi K) d\xi$$

The Brunn-Minkowski theory

Let's deviate to dimension n *only* for a minute!

- Steiner formula in \mathbb{R}^n

$$V(K + tB) = \sum_{i=0}^n t^i \binom{n}{i} W_i(K) \quad W_i(K) \text{ quermassintegral}$$

- $(n-i)$ -th quermassintegral $W_{n-i}(K)$
in integral geometry, it is the average of projection areas:

$$W_{n-i}(K) = c \int_{G(n,i)} \mathcal{H}^i(P_\xi K) d\xi$$

in differential geometry, integral of mean curvatures:

$$W_{n-i}(K) = c \int_{\partial K} H_{n-i-1}(K, x) dS(x)$$

The Brunn-Minkowski theory

Quermassintegrals are *global* geometric quantities that describe convex bodies.

The Brunn-Minkowski theory

Quermassintegrals are *global* geometric quantities that describe convex bodies.

Can we do something to *localize* these global concepts and get geometric quantities that capture local properties of K ?

The Brunn-Minkowski theory

Quermassintegrals are *global* geometric quantities that describe convex bodies.

Can we do something to *localize* these global concepts and get geometric quantities that capture local properties of K ?

Yes!

The Brunn-Minkowski theory

Quermassintegrals are *global* geometric quantities that describe convex bodies.

Can we do something to *localize* these global concepts and get geometric quantities that capture local properties of K ?

Yes! There is a localized Steiner formula.

The Brunn-Minkowski theory

Quermassintegrals are *global* geometric quantities that describe convex bodies.

Can we do something to *localize* these global concepts and get geometric quantities that capture local properties of K ?

Yes! There is a localized Steiner formula.

Generates two families of geometric measures: *area measures*, *curvature measures*

The Brunn-Minkowski theory

Quermassintegrals are *global* geometric quantities that describe convex bodies.

Can we do something to *localize* these global concepts and get geometric quantities that capture local properties of K ?


Yes! There is a localized Steiner formula.

Generates two families of geometric measures: *area measures*, *curvature measures*

They replace curvatures that are only defined for surfaces that are smooth enough.

Surface area measure for polygons

Suppose P is a polygon in \mathbb{R}^2 . S_P is the surface area measure of P .



S_P encodes the set of all outer unit normals v_1, v_2, \dots, v_N of P and attach to each v_i a positive number a_i which is equal to the length of the edge with outer normal v_i .

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N ,

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges?

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges? Moreover, if P exists, is it unique?

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges? Moreover, if P exists, is it unique?

Remark

- *The discrete Minkowski problem was solved by Minkowski in 1897.*

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges? Moreover, if P exists, is it unique?

Remark

- The discrete Minkowski problem was solved by Minkowski in 1897.
- Ans: As long as $\text{span}\{v_1, \dots, v_N\} = \mathbb{R}^2$ and

$$a_1 v_1 + \dots + a_N v_N = 0,$$

P will exist.

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges? Moreover, if P exists, is it unique?

Remark

- The discrete Minkowski problem was solved by Minkowski in 1897.
- Ans: As long as $\text{span}\{v_1, \dots, v_N\} = \mathbb{R}^2$ and

$$a_1 v_1 + \dots + a_N v_N = 0,$$

P will exist.

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges? Moreover, if P exists, is it unique?

Remark

- The discrete Minkowski problem was solved by Minkowski in 1897.
- Ans: As long as $\text{span}\{v_1, \dots, v_N\} = \mathbb{R}^2$ and

$$a_1 v_1 + \dots + a_N v_N = 0,$$

P will exist. If P_1, P_2 are both solutions, then they differ by a translation.

The Minkowski problem

Problem

The discrete Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as lengths of corresponding edges? Moreover, if P exists, is it unique?

Remark

- The discrete Minkowski problem was solved by Minkowski in 1897.
- Ans: As long as $\text{span}\{v_1, \dots, v_N\} = \mathbb{R}^2$ and

$$a_1 v_1 + \dots + a_N v_N = 0,$$

P will exist. If P_1, P_2 are both solutions, then they differ by a translation.

- Proof is mostly just Calculus.

The Minkowski problem

Problem

The Minkowski problem. Given a Borel measure μ on S^1 ,

The Minkowski problem

Problem

The Minkowski problem. Given a Borel measure μ on S^1 , under what conditions is there a convex body K such that the surface area measure of K is exactly μ ?

The Minkowski problem

Problem

The Minkowski problem. Given a Borel measure μ on S^1 , under what conditions is there a convex body K such that the surface area measure of K is exactly μ ? Moreover, if K exists, is it unique?

The Minkowski problem

Problem

The Minkowski problem. Given a Borel measure μ on S^1 , under what conditions is there a convex body K such that the surface area measure of K is exactly μ ? Moreover, if K exists, is it unique?

Remark

- *The Minkowski problem, in this general form, was solved by Aleksandrov, Fenchel-Jessen in 1938.*

The Minkowski problem

Problem

The Minkowski problem. Given a Borel measure μ on S^1 , under what conditions is there a convex body K such that the surface area measure of K is exactly μ ? Moreover, if K exists, is it unique?

Remark

- *The Minkowski problem, in this general form, was solved by Aleksandrov, Fenchel-Jessen in 1938.*
- *Regularity results are due to Lewy, Nirenberg, Pogorelov, Cheng-Yau, Caffarelli between 1938 and 1990.*

The Minkowski problem

Problem

The Minkowski problem. Given a Borel measure μ on S^1 , under what conditions is there a convex body K such that the surface area measure of K is exactly μ ? Moreover, if K exists, is it unique?


Remark

- *The Minkowski problem, in this general form, was solved by Aleksandrov, Fenchel-Jessen in 1938.*
- *Regularity results are due to Lewy, Nirenberg, Pogorelov, Cheng-Yau, Caffarelli between 1938 and 1990.*
- *Well known problem both in PDE and differential geometry.*

The Minkowski problem

Remark

E. Calabi



The importance of the Minkowski problem and its solution is to be felt both in differential geometry and elliptic partial differential equations, on either count going far beyond the impact that the literal statement may have. From the geometric point of view it is the Rosetta Stone, from which several other related problems can be solved.

The Minkowski problem

Remark

- *In differential geometry, the problem of prescribing Gauss curvature.*

The Minkowski problem

Remark

- *In differential geometry, the problem of prescribing Gauss curvature.*
- *In PDE, the following Monge-Ampère equation:*

$$\det(h_{ij} + h\delta_{ij}) = f,$$

where unknown is h . Here (h_{ij}) is the Hessian matrix on the unit sphere, δ_{ij} is the Kronecker delta, and f is a given function.

Modern development of the BMt

The L_p Brunn-Minkowski theory

Firey, Lutwak

Modern development of the BMt

The L_p Brunn-Minkowski theory

$\rightsquigarrow L_p$ surface area measure

Firey, Lutwak

Modern development of the BMt

The L_p Brunn-Minkowski theory

$\rightsquigarrow L_p$ surface area measure

$p = 1$, same as surface area measure

Firey, Lutwak

Modern development of the BMt

The L_p Brunn-Minkowski theory

$\rightsquigarrow L_p$ surface area measure

$p = 1$, same as surface area measure

$p = 0$, cone volume measure V_K

Firey, Lutwak

Modern development of the BMt

The L_p Brunn-Minkowski theory


Firey, Lutwak

$\rightsquigarrow L_p$ surface area measure

$p = 1$, same as surface area measure

$p = 0$, cone volume measure V_K

$$V_K = \frac{1}{2} h_K \cdot S_K$$



For polygons, $V_P \rightsquigarrow v_1, \dots, v_N$ and a_1, \dots, a_N .

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N ,

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as areas of corresponding triangles?

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as areas of corresponding triangles? Moreover, if P exists, is it unique?

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as areas of corresponding triangles? Moreover, if P exists, is it unique?

Remark

- *Still unsolved! Including dimension 2!*

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as areas of corresponding triangles? Moreover, if P exists, is it unique?

Remark

- *Still unsolved! Including dimension 2!*
- *When the given data is even, or in another word, when $\pm v_1, \dots, \pm v_N$, solved in [Böröczky-Lutwak-Yang-Zhang; JAMS 2012] for arbitrary dimension.*

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as areas of corresponding triangles? Moreover, if P exists, is it unique?

Remark

- *Still unsolved! Including dimension 2!*
- *When the given data is even, or in another word, when $\pm v_1, \dots, \pm v_N$, solved in [Böröczky-Lutwak-Yang-Zhang; JAMS 2012] for arbitrary dimension.*

The logarithmic Minkowski problem

Problem

The discrete logarithmic Minkowski problem. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as areas of corresponding triangles? Moreover, if P exists, is it unique?

Remark

- *Still unsolved! Including dimension 2!*
- *When the given data is even, or in another word, when $\pm v_1, \dots, \pm v_N$, solved in [Böröczky-Lutwak-Yang-Zhang; JAMS 2012] for arbitrary dimension. measure concentration phenomenon.*

The logarithmic Minkowski problem

Remark

- *Uniqueness unsolved!*

The logarithmic Minkowski problem

Remark

- *Uniqueness unsolved!*
- *When dimension is 2, data is even, uniqueness in [Böröczky-Lutwak-Yang-Zhang; Adv Math 2014]*

The logarithmic Minkowski problem

Remark

- *Uniqueness unsolved!*
- *When dimension is 2, data is even, uniqueness in [Böröczky-Lutwak-Yang-Zhang; Adv Math 2014]*
- *PDE:*

$$\det(h_{ij} + h\delta_{ij}) = \frac{f}{h}.$$

Modern development of the BMt

The dual Brunn-Minkowski theory

Lutwak

Modern development of the BMt

The dual Brunn-Minkowski theory

Lutwak


- many “dual” concepts and results — radial summation, dual quermassintegral, **intersection body**, etc.

Modern development of the BMt

The dual Brunn-Minkowski theory

Lutwak

- many “dual” concepts and results — radial summation, dual quermassintegral, **intersection body**, etc.
- Busemann-Petty problem. K, L o-symmetric, if you slice both of them using the same line through the origin



length of K' is always larger than length of L' , no matter how you slice them. Is it true K has greater area?

- Solution to Busemann-Petty problem:

- Solution to Busemann-Petty problem:

- Solution to Busemann-Petty problem: $n = 2$ true, easier case;

Modern development of the BMt

- Solution to Busemann-Petty problem: $n = 2$ true, easier case; $n = 3, 4$ true, took three Annals papers;

Modern development of the BMt

- Solution to Busemann-Petty problem: $n = 2$ true, easier case;
 $n = 3, 4$ true, took three Annals papers;
 $n \geq 5$, **false**!

Modern development of the BMt

- Solution to Busemann-Petty problem: $n = 2$ true, easier case;
 $n = 3, 4$ true, took three Annals papers;
 $n \geq 5$, **false**!
- no measure, no PDE

Modern development of the BMt

- Solution to Busemann-Petty problem: $n = 2$ true, easier case;
 $n = 3, 4$ true, took three Annals papers;
 $n \geq 5$, **false**!
- no measure, no PDE

Modern development of the BMt


- Solution to Busemann-Petty problem: $n = 2$ true, easier case; $n = 3, 4$ true, took three Annals papers; $n \geq 5$, **false**!
- no measure, no PDE — until [Huang-Lutwak-Yang-Zhang, ACTA 2016]

Modern development of the BMt

- Solution to Busemann-Petty problem: $n = 2$ true, easier case; $n = 3, 4$ true, took three Annals papers; $n \geq 5$, **false**!
- no measure, no PDE — until [Huang-Lutwak-Yang-Zhang, ACTA 2016]
 \rightsquigarrow **dual curvature measures** and **dual area measures**

Dual curvature measures for polygons

For each real number q , the q -th dual curvature measure $\tilde{C}_q(P, \cdot)$
 $\rightsquigarrow v_1, \dots, v_N$ the set of outer unit normals, and a_1, \dots, a_N .



Here $a_i = \frac{q}{2} \int_{\Delta_i} |x|^{q-2} dx$

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N ,

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as the quantity defined on the previous slide?

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as the quantity defined on the previous slide? Moreover, if P exists, is it unique?

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as the quantity defined on the previous slide? Moreover, if P exists, is it unique?

Remark

- *The dual Minkowski problem, in general, deals with measures and convex bodies.*

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as the quantity defined on the previous slide? Moreover, if P exists, is it unique?

Remark

- The dual Minkowski problem, in general, deals with measures and convex bodies.
- The dual Minkowski problem was posed in [Huang-Lutwak-Yang-Zhang; ACTA, 2016].

The dual Minkowski problem

Problem

The discrete dual Minkowski problem. Let $q \in \mathbb{R}$. Given unit vectors v_1, \dots, v_N and positive numbers a_1, \dots, a_N , under what conditions is there a polygon P such that P has v_1, \dots, v_N as outer unit normals and a_1, \dots, a_N as the quantity defined on the previous slide? Moreover, if P exists, is it unique?

Remark

- The dual Minkowski problem, in general, deals with measures and convex bodies.
- The dual Minkowski problem was posed in [Huang-Lutwak-Yang-Zhang; ACTA, 2016].
- PDE:

$$\det(h_{ij}(v) + h(v)\delta_{ij}) = nh^{-1}(v)|\nabla_{S^{n-1}}h(v) + h(v)v|^{n-q}f(v).$$

The dual Minkowski problem

Remark

- *Unsolved when $q > 2$! Including dimension 2!*

The dual Minkowski problem

Remark

- *Unsolved when $q > 2$! Including dimension 2!*
- *Unsolved when $q > 0$ when the given data is not even. Including dimension 2!*

The dual Minkowski problem

Remark

- *Unsolved when $q > 2$! Including dimension 2!*
- *Unsolved when $q > 0$ when the given data is not even. Including dimension 2!*
- *Uniqueness unsolved when $q > 0$. Including dimension 2!*

Results on the dMP

When $q < 0$, both existence and uniqueness of the solution were solved.

Results on the dMP

When $q < 0$, both existence and uniqueness of the solution were solved.

Theorem (Z., CVPDEs 2017)


Suppose $q < 0$. There is a solution P to the discrete dual Minkowski problem if and only if v_1, \dots, v_n are not contained in any half plane. Moreover, P is unique (if exists).

Results on the dMP

When $q < 0$, both existence and uniqueness of the solution were solved.

Theorem (Z., CVPDEs 2017)

Suppose $q < 0$. There is a solution P to the discrete dual Minkowski problem if and only if v_1, \dots, v_n are not contained in any half plane. Moreover, P is unique (if exists).



The theorem was proved for measures and in any dimension.

Results on the dMP

When $0 < q < 2$ (2 is the dimension of the space), only existence results when the given data is even.

Results on the dMP

When $0 < q < 2$ (2 is the dimension of the space), only existence results when the given data is even.

Theorem

Suppose $0 < q < 2$ and the vectors come in pairs $\pm v_1, \dots, \pm v_N$. There is a solution P to the discrete dual Minkowski problem if and only if the ratio

$$\frac{a_i}{\sum_{j=1}^N a_j} < \min \left\{ \frac{1}{q}, 1 \right\}.$$

Results on the dMP

When $0 < q < 2$ (2 is the dimension of the space), only existence results when the given data is even.

Theorem

Suppose $0 < q < 2$ and the vectors come in pairs $\pm v_1, \dots, \pm v_N$. There is a solution P to the discrete dual Minkowski problem if and only if the ratio

$$\frac{a_i}{\sum_{j=1}^N a_j} < \min \left\{ \frac{1}{q}, 1 \right\}.$$

The theorem was proved for measures and in any dimension.

Results on the dMP

When $0 < q < 2$ (2 is the dimension of the space), only existence results when the given data is even.

Theorem

Suppose $0 < q < 2$ and the vectors come in pairs $\pm v_1, \dots, \pm v_N$. There is a solution P to the discrete dual Minkowski problem if and only if the ratio

$$\frac{a_i}{\sum_{j=1}^N a_j} < \min \left\{ \frac{1}{q}, 1 \right\}.$$

The theorem was proved for measures and in any dimension. The condition will generalize to a subspace mass concentration inequality.

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].

Results on the dMP

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:

Results on the dMP

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:

Results on the dMP

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:
[Z., JDG, in press] (sufficiency)

Results on the dMP

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:
 - [Z., JDG, in press] (sufficiency)
 - [Böröczky-Henk-Pollehn, JDG, in press] (necessity)

Results on the dMP

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:
 - [Z., JDG, in press] (sufficiency)
 - [Böröczky-Henk-Pollehn, JDG, in press] (necessity)
- $1 < q < n$ is not an integer:

Results on the dMP

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:
 - [Z., JDG, in press] (sufficiency)
 - [Böröczky-Henk-Pollehn, JDG, in press] (necessity)
- $1 < q < n$ is not an integer:

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:
 - [Z., JDG, in press] (sufficiency)
 - [Böröczky-Henk-Pollehn, JDG, in press] (necessity)
- $1 < q < n$ is not an integer:
 - [Böröczky-Lutwak-Yang-Zhang-Z., preprint] (sufficiency)

- $0 < q \leq 1$: [Huang-Lutwak-Yang-Zhang, Acta, 2016].
- $q = 2, \dots, n - 1$:
 - [Z., JDG, in press] (sufficiency)
 - [Böröczky-Henk-Pollehn, JDG, in press] (necessity)
- $1 < q < n$ is not an integer:
 - [Böröczky-Lutwak-Yang-Zhang-Z., preprint] (sufficiency)
 - [Böröczky-Henk-Pollehn, JDG, in press] (necessity).