
Crossing order in trained models: INCONCLUSIVE
(a) Order-dependent share, natural pairs (t = 32)
natural β_{ord} (oracle = 1)
0.00
0.02
0.04
0.06
0.08
AB-s1
AB-s2
AB-s3
AB-s4
NA-s1
NA-s2
NA-s3
NA-s4
ZC-s1
ZC-s2
ZC-s3
ZC-s4
init-s1
init-s2
init-s3
init-s4
(b) C-ORD per seed: clearly > 0, far below 0.25
C_s (NA minus controls)
0.00
0.05
0.10
0.15
0.20
0.25
0.30
registered C-ORD size (0.25)
mean 0.061
$LB_{0.975}$ 0.0595
seed 1
seed 2
seed 3
seed 4
(c) Designed W2 regret: bar not beaten
designed W2 regret R_{W2}
0.000
0.025
0.050
0.075
0.100
0.125
0.150
0.175
0.200
bar $(3/4)\Delta^{lo}_{W2} \approx 0.0458$
AB-s1
AB-s2
AB-s3
AB-s4
NA-s1
NA-s2
NA-s3
NA-s4
ZC-s1
ZC-s2
ZC-s3
ZC-s4
(d) Designed placebo: ZC-s1 responds (control-limited)
designed placebo β_{W2}
0.00
0.05
0.10
0.15
0.20
AB-s1
AB-s2
AB-s3
AB-s4
ZC-s1
ZC-s2
ZC-s3
ZC-s4
init-s1
init-s2
init-s3
init-s4
NA (non-abelian)
AB control
ZC control
init