Transcript: The 1.0–1.2 TeV Thermal Higgsino Against Planck, LZ and XENONnT: Toy-Model Relic and Inelastic-Recoil Calculations with a Literature Survey

HEP-Genesis-Agent session record
Project: ramp-combo-higgsino-dd

2026-10-22

The conversation verbatim, generated mechanically from the session record.

A Conversation transcript

The exchange verbatim (assistant text as shown to the user; tool calls summarized by name — see the tool-call log for arguments and results).

User (2026-10-16 14:31:17)

Survey recent arXiv work on the higgsino thermal relic target (1.0-1.2 TeV). Use the inelastic-DM tools to compute the relic density at several masses; compare with LZ/XENONnT sensitivities and the Planck abundance. Where does the higgsino sit relative to current limits?

Assistant (research mode, 2026-10-16 14:31:17)

Plan:

1. (closed) Compute the thermal relic density Omega h^2 of a higgsino-like (inelastic) DM candidate at m = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV with the inelastic-DM tools. Compare each value with Planck (Omega h^2 = 0.120 +/- 0.001) and locate the mass where the tool's relic curve crosses the Planck abundance.
2. (closed) Place the thermal higgsino (m about 1.0-1.2 TeV) relative to LZ/XENONnT. Compute inelastic recoil spectra and expected event counts for delta = 300-600 keV, scan the (m_chi, delta) plane against all constraints, and identify where the kinematic reach of xenon ends.
3. (blocked) Verify and cite the key literature: the higgsino relic crossing mass, the LZ 2.84 t-yr extended-window result, the solar-capture delta bound, the elastic loop-level cross section relative to the XENONnT limits and the neutrino fog, and the Planck Omega h^2. Obtain verbatim quotes.
4. (blocked) Cite the established, non-2026 baseline: the Planck 2018 Omega_c h^2, the Sommerfeld-corrected pure-higgsino thermal mass from pre-2026 literature, and the XENONnT/LZ elastic SI limit at about 1 TeV compared with the loop-level higgsino cross section and the neutrino fog.

5. (closed) Map, with the inelastic-DM tools, the mass splitting delta at which a thermal higgsino gives about 1 and about 0.3 expected LZ events. Test how sensitive that window is to the halo velocity cutoff, then compare it with the literature fits: 339 keV (2609.26698), about 350 keV (2609.01504) and 377 keV (2609.02608).

6. (closed) Establish the current indirect-detection status of the 1.0-1.2 TeV thermal higgsino from the literature: the H.E.S.S., Fermi-LAT and MAGIC gamma-ray line and continuum limits against the Sommerfeld-enhanced s-wave annihilation cross section, and the CTAO projections. This replaces the server's p-wave treatment, which is physically wrong for a higgsino.

Tool calls: `list_sterile_nu_skills`, `list_observables`, `list_agent_files`, `list_input_files`, `list_datasets`, `list_files`, `get_dispatch`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv`, `web_search`, `search_arxiv`, `search_arxiv`, `search_arxiv`, `search_arxiv`, `search_arxiv`, `search_arxiv`, `describe_tools`, `describe_tools`, `predict_recoil`, `compute_relic_density`, `predict_recoil`, `compute_relic_density`, `predict_recoil`, `compute_relic_dens` `predict_recoil`, `compute_relic_density`, `predict_recoil`, `compute_relic_density`, `check_all_constrai` `check_all_constraints`, `check_all_constraints`, `check_all_constraints`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `scan_mass_splitting`, `plot_recoil_spectrum`, `predict_annual_modula` `post_finding`, `post_finding`, `post_question`, `post_finding`, `post_finding`, `post_finding`, `plot_constraint_summary`, `post_finding`, `get_arxiv_metadata`, `get_arxiv_metadata`, `download_arxiv_fu` `download_arxiv_full_text`, `download_full_arxiv_paper`, `search_arxiv`, `cite`, `search_arxiv`, `search_arxiv`, `post_finding`, `post_finding`, `post_dead_end`, `search_arxiv`, `post_question`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `search_arxiv`, `search_arxiv`, `get_arxiv_metadata`, `search_arxiv`, `post_finding`, `post_finding`, `get_arxiv_metadata`, `download_arxiv_full_text`, `post_finding`, `post_finding`, `get_arxiv_metadata`, `get_arxiv_metadata`, `download_full_arxiv_paper`, `download_arxiv_full_text`, `search_arxiv`, `cite`, `search_arxiv`, `search_arxiv`, `post_finding`, `post_finding`, `post_dead_end`

Survey — 52 tools on 5 servers, 7 read-only probes, 18 papers found.

Note first: I did not compute any relic densities. No inelastic-DM or relic-density tools were available to me, only literature search tools, so everything below comes from paper abstracts. Second note: the 'recent arXiv work' shown here was found through the OpenAlex-backed search_arxiv tool, which returned papers dated June to October 2026. These IDs should be checked on arxiv.org before anyone cites them. The direct arXiv API search_arxiv_abstracts tool returned HTTP 503/429 errors. On the physics: when the Sommerfeld effect and coannihilation are included, a pure higgsino gives the full Planck abundance at m about 1.0-1.2 TeV. The cited values are about 1.08-1.1 TeV (2609.02775, 2609.02608), about 1.2 TeV (2410.15733) and an upper end near 1150 GeV (2403.19598). Lighter thermal higgsinos are under-abundant, typically by a factor of 5-20 in natural SUSY (1803.11210). For direct detection, the pure higgsino is inelastic dark matter: its Z-mediated scattering requires a neutral-state mass splitting delta of up to about a few hundred keV. If delta is larger than the kinematic reach of xenon, elastic loop-level scattering sits near or below the neutrino fog (2609.01504, 2410.02723). Higgsino-gaugino mixing raises the elastic rate and is what lets direct detection probe the thermal target (2606.31013). In 2026, LZ released a 2.84 tonne-year search with an extended nuclear-recoil window (2609.02823). It found one event at 248 keV, with 2.6 sigma global and 3.4 sigma local significance. A roughly 1.1 TeV higgsino with delta of about 340-380 keV under the Standard Halo Model, or about 480-520 keV if the Large Magellanic Cloud's effect on the halo velocity tail is included, can produce this event. However, IceCube solar-capture neutrino limits require delta $>$ 566 keV (2609.02775, confirmed by 2609.11833 and 2609.07807; 2609.06760 quotes 0.51-0.56 MeV). Two further problems: the empty sideband above 350 keV is in tension with the thermal-mass fit (2609.09385, 2609.26698), and halo uncertainties are not enough to close the gap (2609.15321).

Bottom line: the full-abundance thermal higgsino at about 1.1 TeV is not excluded as a model. Its sub-MeV-splitting interpretation of the LZ event is disfavoured by solar capture. Splittings of order an MeV or more remain allowed (2610.00226) and leave it effectively out of reach of current xenon searches. Indirect gamma-ray line searches (CTA/CTAO, 2008.00692) are the main future test.

Quantities worth checking: Thermal relic Omega h^2 versus higgsino mass at about 0.9, 1.0, 1.1, 1.2 and 1.3 TeV, with and without the Sommerfeld effect. Compare with the Planck value Omega h^2 of about 0.120 +/- 0.001 (standard Planck 2018 value; the Planck paper was not retrieved here). The literature crossing point is about 1.08-1.2 TeV.; Kinematic maximum splitting delta_max = mu v^2/2 for xenon (and argon) at m = 1.1 TeV, for v_esc of about 544 km/s in the Earth frame (about 544+250 km/s).; Endothermic recoil spectrum dR/dE_R for m = 1.0-1.2 TeV and delta = 300-600 keV, with Helm form factor. Count events in [14,225), [225,271] and (271,800] keV for 2.84 tonne-years.; Z-mediated inelastic cross section per nucleon for a pure higgsino (vector coupling, roughly 1e-39 cm^2 scale) versus the rate needed to give 1 event.; Elastic loop-level higgsino-nucleon SI cross section (about 1e-48 to 1e-49 cm^2) versus the LZ/XENONnT elastic limits at 1.1 TeV and the neutrino fog.; Solar-capture threshold delta of about 566 keV versus the LZ-preferred delta (about 340-380 keV for the Standard Halo Model, about 480-520 keV with the LMC)..

Plan — 4 lines of investigation:

1. Compute the thermal relic density Omega h^2 of a higgsino-like (inelastic) DM candidate at m = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV with the inelastic-DM tools. Compare each value with Planck (Omega h^2 = 0.120 +/- 0.001) and locate the mass where the tool's relic curve crosses the Planck abundance. — This answers the core quantitative part of the task directly. The survey computed no relic densities. Its literature crossing point of about 1.08-1.2 TeV needs a tool-based cross-check, and we need to know which physics the tool includes (Sommerfeld effect, coannihilation, higgsino couplings).

2. Place the thermal higgsino (m about 1.0-1.2 TeV) relative to LZ/XENONnT. Compute inelastic recoil spectra and expected event counts for delta = 300-600 keV, scan the (m_chi, delta) plane against all constraints, and identify where the kinematic reach of xenon ends. — This answers 'where does the higgsino sit relative to current limits'. The survey's key claims need tool-based checks: the LZ 248 keV event is fit by delta of about 340-380 keV, solar capture requires delta > 566 keV, and large-delta higgsinos fall out of xenon reach.

3. Verify and cite the key literature: the higgsino relic crossing mass, the LZ 2.84 t-yr extended-window result, the solar-capture delta bound, the elastic loop-level cross section relative to the XENONnT limits and the neutrino fog, and the Planck Omega h^2. Obtain verbatim quotes. — The survey's 2026-dated IDs came from an OpenAlex-backed search. They must be confirmed on arXiv before anything is cited, and the Planck value was never retrieved. The tool results from d-01 and d-02 need a validated literature baseline to compare against.

4. Cite the established, non-2026 baseline: the Planck 2018 Omega_c h^2, the Sommerfeld-corrected pure-higgsino thermal mass from pre-2026 literature, and the XENONnT/LZ elastic SI limit at about 1 TeV compared with the loop-level higgsino cross section and the neutrino fog. — The original d-03 cannot cover these within 13 calls. These anchors are independent of whether the 2026 IDs resolve, so they protect the final answer if the 2026 papers prove unverifiable. (added by the critic) Critic's view: revise — Over budget. As written it needs about 15 calls: 1 describe + 5 relic + 3 Sommerfeld toggles + 3 delta checks + 2 check_all. Drop the delta-sensitivity step. Keep the Sommerfeld toggle only if describe_tools shows a switch exists. This brings it to about 9-11 calls.; The

serve-idm server is probably a generic inelastic-DM model, not a pure higgsino with SU(2) couplings, Sommerfeld enhancement and full coannihilation. If describe_tools does not state higgsino electroweak couplings and Sommerfeld enhancement, label the results as a toy model. In that case, do not report a 'higgsino Planck-crossing mass'. Report only the tool's crossing mass and how far it sits from the 1.0-1.2 TeV literature target. Quote the tool's model description verbatim.; The delta-sensitivity step (0.3, 0.566, 1.0 MeV) is physically uninformative. A sub-MeV splitting is far below T_f of about m/20, which is about 50 GeV, so the relic density should not depend on delta. If the tool does show delta dependence, that indicates a modelling artifact and should be reported as one. Check whether delta is entered in keV or MeV before calling..

Wave 1 — now exploring, in parallel: Compute the thermal relic density Omega h^2 of a higgsino-like (inelastic) DM candidate at m = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV with the inelastic-DM tools. Compare each value with Planck (Omega h^2 = 0.120 +/- 0.001) and locate the mass where the tool's relic curve crosses the Planck abundance.; Place the thermal higgsino (m about 1.0-1.2 TeV) relative to LZ/XENONnT. Compute inelastic recoil spectra and expected event counts for delta = 300-600 keV, scan the (m_chi, delta) plane against all constraints, and identify where the kinematic reach of xenon ends.; Cite the established, non-2026 baseline: the Planck 2018 Omega_c h^2, the Sommerfeld-corrected pure-higgsino thermal mass from pre-2026 literature, and the XENONnT/LZ elastic SI limit at about 1 TeV compared with the loop-level higgsino cross section and the neutrino fog..

Finding: The inelastic-DM server's compute_relic_density (is_higgsino=true) gives Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285, 0.1503 at m_chi = 900, 1000, 1100, 1200, 1300 GeV. Linear interpolation crosses Planck 0.120 at m about 1158 GeV (+/-6 GeV for the +/-0.0012 band). A direct check at 1159 GeV returns Omega h^2 = 0.120. This lies within the 1.08-1.2 TeV literature range but about 60-80 GeV above the 1.08-1.1 TeV values.

Finding: check_all_constraints (is_higgsino) at m=1100 GeV with delta = 400, 566 and 1000 keV, and at m=1159 GeV with delta=400 keV, returns VIABLE every time. Fermi-LAT and CMB are not excluded, sigma_n = 7.43e-39 cm^2, and N_expected_LZ = 0.00 at every point. N_LZ = 0 at delta = 400 keV contradicts the literature, which reports about 1 event at delta about 340-380 keV. The server's present-day sigma_v (about 7e-42 to 5e-41 cm^3/s, labelled 'p-wave + mass-splitting suppressed') is also far below the s-wave higgsino value of about 1e-26 cm^3/s.

Completed — Compute the thermal relic density Omega h^2 of a higgsino-like (inelastic) DM candidate at m = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV with the inelastic-DM tools. Compare each value with Planck (Omega h^2 = 0.120 +/- 0.001) and locate the mass where the tool's relic curve crosses the Planck abundance.. Relic densities for a higgsino-like dark matter candidate (compute_relic_density with is_higgsino=true; the tool sets the annihilation cross section automatically): Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285 and 0.1503 at m = 900, 1000, 1100, 1200 and 1300 GeV (tc-...-23/25/27/29/31). The curve reaches the Planck value of 0.120 at about 1158 GeV, +/-6 GeV from the server's +/-0.0012 Planck band (the brief's +/-0.001 gives about +/-5 GeV). A direct call at 1159 GeV returns 0.120 (tc-...-33). This is inside the 1.08-1.2 TeV literature range. It is about 60-80 GeV above the 1.08-1.1 TeV values (2609.02775, 2609.02608) and close to the about 1.2 TeV value (2410.15733), so agreement is only partial.

Physics in the tool: the higgsino setting is a simple scaling of the cross section, sigma_v = 2.5e-26 cm^3/s x (1100 GeV/m)^2, fed into a Kolb-Turner freeze-out estimate (x_f about 24.5, g* = 86.25). There is no Sommerfeld or coannihilation switch; the only inputs are m_chi_GeV, is_higgsino and sigma_v_cm3_s. As a result, the with/without-Sommerfeld comparison (step 3) and the splitting-sensitivity runs (step 4) could not be done with compute_relic_density. Its

closeness to the literature reflects how it was calibrated, not a full calculation.

Constraint checks (check_all_constraints): at 1100 GeV with delta = 400, 566 and 1000 keV, and at 1159 GeV with delta = 400 keV, every point is rated VIABLE. In each case Fermi-LAT and Planck CMB show no exclusion, sigma_n = 7.43e-39 cm^2 and N_expected_LZ = 0.00 (tc-...-32/33/34/35). The relic verdict passes at 1100 GeV despite the tool's own 9.8 sigma tension.

Caveats:

- The pass/consistent flags are True even at 7-25 sigma tension, which looks like a bug.
- N_LZ = 0 at delta = 400 keV conflicts with the literature, where splittings of about 340-380 keV give about one event (posted as q-06).
- The indirect-detection treatment assumes p-wave annihilation suppressed by the mass splitting, giving sigma_v of about 1e-41 cm^3/s today. That is inappropriate for an s-wave, Sommerfeld-enhanced higgsino.

Posted as findings f-001 and f-002 and question q-06. No figures or files were produced.

Finding: With the inelastic-DM server (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, v_esc+v_E = 776 km/s), the expected LZ count at m_chi = 1100 GeV is 744.6 at delta = 300 keV, 1.97 at 350 keV (P(N>=1) = 0.86, peak E_R = 315 keV), and 0.00 at 380, 500, 566 and 700 keV (peak E_R = 342, 450, 510, 630 keV). The 248 keV event needs v_min = 762 km/s at 350 keV (accessible) and 799 km/s at 380 keV (not accessible).

Finding: Xenon's kinematic reach at the thermal higgsino mass is delta_max = mu v_max^2/2. With m_Xe of about 122 GeV, m_chi = 1100 GeV gives mu = 109.8 GeV. For v_max = 776 km/s (the tool's value, beta^2 = 6.70e-6) this gives delta_max of about 368 keV; for v_max = 794 km/s (544+250) it gives about 385 keV. This agrees with the tool: 350 keV is accessible and 380 keV gives 0 events. At delta = 380 keV the 248 keV v_min is 802, 799 and 796 km/s for m = 1000, 1100 and 1200 GeV, all inaccessible, so the reach depends only weakly on mass.

Finding: The scan over m_chi = 800-1400 GeV and delta = 0-800 keV (17x17 grid) finds 144 of 289 points passing all constraints and 9 points that predict 0.3-5 LZ events. The 1100 GeV, 380 keV recoil-spectrum plot is at /Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png. The annual modulation at 1100 GeV, 350 keV and E_R = 248 keV has a fractional amplitude of 0.985, peaking on June 2; the event date of September 1 sees 22.5% of the peak rate.

Finding: plot_constraint_summary on the 17x17 scan (m_chi = 800-1400 GeV, delta = 0-800 keV) wrote /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png and reports the best-fit point at m_chi = 988 GeV, delta = 350 keV.

Completed — Place the thermal higgsino (m about 1.0-1.2 TeV) relative to LZ/XENONnT. Compute inelastic recoil spectra and expected event counts for delta = 300-600 keV, scan the (m_chi, delta) plane against all constraints, and identify where the kinematic reach of xenon ends.. (Open items: (1) The tool gives only total counts, with no breakdown into the [14,225), [225,271] and (271,800] keV bins. (2) The scan and plot files went to the server's default directory, not the run directory, because the tools have no output_dir parameter. (3) The 566 keV and literature 340-380 keV bands were not drawn on the plot; the plotting tool has no overlay option. (4) Halo-parameter sensitivity (q-04) was not tested, because the tool exposes no v0 or v_esc parameters.) The inelastic-DM server models the higgsino with Z exchange (sigma_n = 7.43e-39 cm^2), the LZ 2.84 t-yr extended nuclear-recoil window, and a halo cutoff of v_esc + v_E = 776 km/s (tc-...-20, -21). At m_chi = 1100 GeV,

the expected LZ event counts are: 744.6 at delta = 300 keV (grossly excluded against 1 observed event); 1.97 at 350 keV (P(N>=1) = 0.86, peak recoil energy E_R = 315 keV); and 0.00 at 380, 500, 566 and 700 keV, where the spectrum would peak at 342, 450, 510 and 630 keV (finding f-003). At delta = 380 keV the result does not depend much on mass: the minimum speed needed for the 248 keV event is v_min = 802, 799 and 796 km/s at m = 1000, 1100 and 1200 GeV, all above 776 km/s and so out of reach. Xenon's kinematic reach, worked by hand: delta_max = mu v_max^2/2, with mu = 109.8 GeV. That gives about 368 keV for v_max = 776 km/s and about 385 keV for 544+250 km/s, consistent with the tool's cutoff between 350 and 380 keV (f-004). The scan over m = 800-1400 GeV and delta = 0-800 keV (17x17 points) found 144 of 289 points viable and 9 points predicting 0.3-5 LZ events. Its best fit is m = 988 GeV, delta = 350 keV. Files: /Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz and constraint_summary.png (f-005, f-006). The recoil spectrum plot for 1100 GeV, 380 keV is at /Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png. Annual modulation at 1100 GeV, 350 keV and E_R = 248 keV: 98.5% amplitude, peaking on June 2; the event date of September 1 sees 22.5% of the peak rate. Regions at 1.1 TeV: delta below about 340 keV is excluded by xenon (hundreds of events predicted); delta of about 345-368 keV is allowed and can explain the 248 keV event (tool value); delta above about 370 keV is out of xenon's reach. The literature solar-capture bound (delta > 566 keV, 2609.02775) therefore puts every solar-capture-allowed thermal higgsino out of LZ inelastic reach under the Standard Halo Model.

Finding: The pre-2026 Sommerfeld- and coannihilation-corrected thermal pure-higgsino mass is about 1.2 TeV. The abstract of arXiv:2410.15733 (Fukuda, Li, Moroi, Niki, JHEP06(2025)091, DOI 10.1007/jhep06(2025)091) says: "in which the observed DM abundance is realized with the Higgsino mass of ~1.2 TeV".

Finding: XENONnT's 3.1 tonne-year elastic WIMP search (arXiv:2502.18005, DOI 10.1103/msw4-t342) found no significant excess. Its spin-independent limit has a minimum of 1.7e-47 cm^2 (90% CL) at 30 GeV, with best median sensitivity 1.4e-47 cm^2 at 41 GeV. The earlier 1.09 t-yr result (arXiv:2303.14729) reached 2.58e-47 cm^2 at 28 GeV. The abstracts do not give the limit at 1 TeV. Because the limit scales roughly linearly with mass at high mass, it should be about 1e-45 cm^2 near 1 TeV, but that is an inference, not a quoted value.

Blocked — Cite the established, non-2026 baseline: the Planck 2018 Omega_c h^2, the Sommerfeld-corrected pure-higgsino thermal mass from pre-2026 literature, and the XENONnT/LZ elastic SI limit at about 1 TeV compared with the loop-level higgsino cross section and the neutrino fog.. (The arXiv-API-backed tools (get_arxiv_metadata, download_arxiv_full_text, download_full_arxiv_paper and cite's metadata check) returned 'not found', 429 or timeout errors even for 1807.06209 and 2410.15733 (tc-...-19, -50, -51, -52, -53, -55). So cite() cannot validate any quote and no full text can be read. Only OpenAlex abstracts were available.) Two of the three anchors were confirmed from abstracts returned by the OpenAlex-backed search_arxiv tool. None could be added to references.bib with cite(), so the success criterion of 3 cite() entries was not met. (1) Thermal higgsino mass, finding f-007: arXiv:2410.15733 (Fukuda, Li, Moroi, Niki, JHEP06(2025)091, DOI 10.1007/jhep06(2025)091) states 'in which the observed DM abundance is realized with the Higgsino mass of ~1.2 TeV', with the Sommerfeld effect and coannihilation included (tc-261016_1-w1-d-04-54). (2) XENONnT elastic limits, finding f-008: arXiv:2502.18005 (3.1 t-yr exposure, DOI 10.1103/msw4-t342) gives a minimum spin-independent limit of 1.7e-47 cm^2 at 30 GeV (90% CL) and a best median sensitivity of 1.4e-47 cm^2 at 41 GeV. arXiv:2303.14729 (1.09 t-yr) gives 2.58e-47 cm^2 at 28 GeV (tc-261016_1-w1-d-04-57). The limit at 1 TeV is not quoted in either abstract. The figure of about 1e-45 cm^2 is only an extrapolation from linear high-mass scaling. (3) Planck Omega_c h^2 = 0.120 +/- 0.001 (1807.06209) was NOT retrieved: get_arxiv_metadata and both download tools returned 'not found' or failed, and a targeted

search did not surface the Planck VI paper. The loop-level higgsino cross section (about 1e-48 to 1e-49 cm^2) and the neutrino-fog comparison were not sourced. Dead end x-01 and question q-07 are posted.

Critic, wave 1 — continue: 5 findings stand up to the evidence, 3 questioned ('check_all_constraints (is_higgsino) at m=1100 GeV with delta = 400, 566 and 1000 keV, and ': The excerpts for tc-32/33/34/35 are cut off before the overall verdict and the CMB block, so 'VIABLE every time' and 'CMB not excluded' are not shown. The excerpts also show consistent_with_1_event = False next to N_LZ = 0, which works against a clean VIABLE reading. The claim that N_LZ = 0 at delta = 400 keV contradicts the literature is overstated: 400 keV lies above the quoted 340-380 keV band. The real tension is at 380 keV (tool gives 0 events, 2609.02608 fits about 377 keV), and that point belongs to f-003 and f-004, not here. The remaining numbers (sigma_n, present-day sigma_v of 6.7e-42 to 5.2e-41, Fermi not excluded) do match.; 'The pre-2026 Sommerfeld- and coannihilation-corrected thermal pure-higgsino mass is about ': The quoted excerpt of tc-261016_1-w1-d-04-54 shows only Paper 1, arXiv:1404.0682 (Low and Wang). It does not contain 2410.15733 or the quoted sentence 'Higgsino mass of ~1.2 TeV'. The evidence as posted does not contain the claimed value.; 'XENONnT's 3.1 tonne-year elastic WIMP search (arXiv:2502.18005, DOI 10.1103/msw4-t342) fou': The excerpt of tc-261016_1-w1-d-04-57 shows only the title and author list of 2303.14729. None of the claimed limits (1.7e-47 at 30 GeV, 1.4e-47 at 41 GeV, 2.58e-47 at 28 GeV) appear, and 2502.18005 is not visible. The claimed values are not in the evidence.).

Issues: The arithmetic checks out: sigma_v = 2.5e-26 x (1100/m)^2 reproduces 3.73e-26 at 900 GeV, and interpolating between 1100 and 1200 GeV gives about 1158 GeV. However, the tool is a single-channel Kolb-Turner scaling calibrated to 1.1 TeV. The 1158 GeV crossing is a calibration artefact, not a higgsino prediction. The final report must label it as a toy-model crossing, as critique c-01 required.; The scan counts 144 of 289 points as viable. That count uses the same relic 'passes' flag that returns True at 7-25 sigma tension, so it probably includes points that are far from the Planck abundance. It also relies on the p-wave-suppressed indirect-detection model, which is wrong for a higgsino. The viable fraction should not be quoted as physics until it is recomputed with a correct relic criterion.; The best fit at m = 988 GeV, delta = 350 keV fits the LZ event, not the thermal relic. With the tool's own relic curve, 988 GeV gives Omega h^2 of about 0.087, which is under-abundant. The best fit is therefore not a full-abundance thermal higgsino, and this should be stated..

Next steps after wave 1 — One wave remains, with 19 calls, so the plan is three directions sized at about 6, 7 and 5 calls. d-03, which is open and has not run, carries over first. It already covers the critic's literature-repair suggestion: re-capturing the abstracts behind f-007 and f-008 so the excerpts show the quoted numbers, sourcing the elastic loop-level cross section and the neutrino fog at 1 TeV, retrying the Planck value, and resolving the 2026 IDs (q-03). For that reason I add no duplicate literature direction. d-03 should cap itself at about 6 calls and drop cite() attempts if the arXiv API fails again. The first new direction addresses the critic's highest-severity issue: the delta window for one LZ event and its dependence on the velocity cutoff. That window sets where the thermal higgsino sits relative to xenon reach, between the kinematic edge near 370-385 keV and the solar-capture bound at 566 keV. The second new direction closes the unaddressed indirect-detection leg with sourced H.E.S.S./Fermi/CTAO status, replacing the tool's invalid p-wave treatment. The critic's suggested full rescan with a corrected relic criterion is not run, because it would not fit the budget. Instead, the final report should note that along the thermal line only delta is free: f-005's 144/289 viable count must not be quoted, and f-006's best fit at 988 GeV is under-abundant, so it is not a thermal higgsino. The 1158 GeV crossing must be labelled a calibrated toy-model value. The task does not ask for a facility (HPC) run, so none is scheduled.

New lines of investigation: Map, with the inelastic-DM tools, the mass splitting delta at

which a thermal higgsino gives about 1 and about 0.3 expected LZ events. Test how sensitive that window is to the halo velocity cutoff, then compare it with the literature fits: 339 keV (2609.26698), about 350 keV (2609.01504) and 377 keV (2609.02608).; Establish the current indirect-detection status of the 1.0-1.2 TeV thermal higgsino from the literature: the H.E.S.S., Fermi-LAT and MAGIC gamma-ray line and continuum limits against the Sommerfeld-enhanced s-wave annihilation cross section, and the CTAO projections. This replaces the server's p-wave treatment, which is physically wrong for a higgsino..

Wave 2 — now exploring, in parallel: Verify and cite the key literature: the higgsino relic crossing mass, the LZ 2.84 t-yr extended-window result, the solar-capture delta bound, the elastic loop-level cross section relative to the XENONnT limits and the neutrino fog, and the Planck Omega h^2. Obtain verbatim quotes.; Map, with the inelastic-DM tools, the mass splitting delta at which a thermal higgsino gives about 1 and about 0.3 expected LZ events. Test how sensitive that window is to the halo velocity cutoff, then compare it with the literature fits: 339 keV (2609.26698), about 350 keV (2609.01504) and 377 keV (2609.02608).; Establish the current indirect-detection status of the 1.0-1.2 TeV thermal higgsino from the literature: the H.E.S.S., Fermi-LAT and MAGIC gamma-ray line and continuum limits against the Sommerfeld-enhanced s-wave annihilation cross section, and the CTAO projections. This replaces the server's p-wave treatment, which is physically wrong for a higgsino..

Finding: With predict_recoil (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, hard cutoff v_esc+v_E = 776 km/s), m_chi = 1100 GeV gives N_LZ = 1.97, 0.494, 0.085, 0.009, 0.00, 0.00 at delta = 350, 355, 360, 365, 370, 375 keV (peak E_R = 315, 320, 324, 329, 333, 338 keV). Log-linear interpolation gives delta(N=1) = 352.5 keV and delta(N=0.3) = 356.4 keV, so the window is only about 4 keV wide (N falls about 4-6x per 5 keV). The 248 keV event stops being accessible between 360 and 365 keV (v_min = 774 and 781 km/s). At m_chi = 1159 GeV (the tool's thermal mass), delta = 360 keV gives N = 0.135 against 0.085 at 1100 GeV. That moves delta(N=1) up by only about 1-2 keV, to about 354 keV.

Finding: This answers q-06. predict_recoil has no halo parameters (inputs: m_chi_GeV, delta_keV, sigma_n_cm2, is_higgsino), so v_esc cannot be varied. The N_LZ = 0 at delta >= 370 keV comes from the tool's hard 776 km/s cutoff, which puts delta_max = mu v_max^2/2 at about 368 keV. An analytic rescaling by (794/776)^2 = 1.047 moves the edge to about 385 keV, and moves the tool's delta(N=1) and delta(N=0.3) from 352.5 and 356.4 keV to about 369 and 373 keV. Against the literature: 350 keV (2609.01504) matches the tool directly (N = 1.97). The 776 vs 794 km/s cutoff accounts for about 17 keV of the 24.5 keV offset from 377 keV (2609.02608). The remaining roughly 5-8 keV has not been explained; plausible causes, none tested, are a smooth truncated-Maxwellian tail, a different v_E at the event date, and differences in cross-section and efficiency normalization. The 339 keV value at 1 TeV (2609.26698) is a normalization-free sideband-ratio optimum. The tool would predict far more than 1 event there (744.6 at 300 keV), so 339 keV is not a direct comparison for the absolute count.

Completed — Map, with the inelastic-DM tools, the mass splitting delta at which a thermal higgsino gives about 1 and about 0.3 expected LZ events. Test how sensitive that window is to the halo velocity cutoff, then compare it with the literature fits: 339 keV (2609.26698), about 350 keV (2609.01504) and 377 keV (2609.02608).. I mapped the window with predict_recoil (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, hard cutoff v_esc+v_E = 776 km/s). At m_chi = 1100 GeV the expected LZ count is N_LZ = 1.97, 0.494, 0.085, 0.009, 0.00 and 0.00 at delta = 350, 355, 360, 365, 370 and 375 keV, with the spectrum peaking at E_R = 315-338 keV (tc-...-65 to -69). Interpolating in log N gives delta(N=1) = 352.5 keV and delta(N=0.3) = 356.4 keV at 1100 GeV, so the window is only about 4 keV wide. The 248 keV event becomes inaccessible between 360 and 365 keV, where its v_min rises from 774 to 781 km/s. Mass dependence along the thermal line is weak: at 1159 GeV, delta = 360 keV gives N = 0.135 against

0.085 at 1100 GeV (tc-...-70), which moves delta(N=1) up by about 1-2 keV. The halo cutoff could not be tested with the tool, because predict_recoil takes no halo parameters. Scaling delta_max with v_max^2 by hand, (794/776)^2 = 1.047, moves the kinematic edge from about 368 to about 385 keV and puts delta(N=1) near 369 keV and delta(N=0.3) near 373 keV. This answers q-06: N = 0 at delta >= 370 keV is caused by the hard 776 km/s cutoff, not by the energy window. Against the literature: 350 keV (2609.01504) agrees with the tool directly. The 776 vs 794 km/s cutoff accounts for about 17 keV of the 24.5 keV offset from 377 keV (2609.02608). The remaining roughly 5-8 keV is unexplained; possible causes such as the shape of the halo tail, the value of v_E and the normalization were not tested. The 339 keV value at 1 TeV (2609.26698) comes from a sideband-ratio optimum that does not depend on the overall rate. With the tool's normalization, splittings at or below 340 keV give hundreds of events (744.6 at 300 keV), so 339 keV cannot be compared directly with an absolute event count. Findings f-009 and f-010 are posted. No figures or files were produced. I used 6 compute calls.

Finding: The indirect-detection status of the about 1.1 TeV thermal higgsino, taken from abstracts only: it is not excluded by current gamma-ray searches, and the result depends on the halo profile. (a) arXiv:2405.13104 (Rodd, Safdi, Xu, PRD 110, 043003, 2024) ran a 15-yr Fermi-LAT line-like search that comes 'within an order one factor' of the sensitivity needed to detect the higgsino, for some Milky Way analog profiles. In that analysis H.E.S.S. is subleading to Fermi for the higgsino. (b) arXiv:2210.03140 (Montanari, Moulin, Rodd, PRD 107, 043028) finds the higgsino 'out of reach by at least a factor of a few' even at the ultimate sensitivity of current H.E.S.S.-like IACTs. (c) arXiv:2608.07234 (H.E.S.S. Inner Galaxy Survey, 546 h, DOI 10.1103/d8tj-55kc, dated 2026 and unverified) quotes a line limit of 2.3e-28 cm^3/s at 1 TeV for an Einasto profile and states that thermal higgsino DM is 'probed for DM Milky Way models' for the first time. The abstract does not say whether it excludes the higgsino, or for which profiles.

Finding: CTAO is the decisive future test. arXiv:2008.00692 (Rinchiuso et al., PRD 103, 023011) finds that CTA 'has the potential to become the first experiment that has sensitivity to the thermal' higgsino. arXiv:2405.13104 finds that CTA and SWGO could detect the line-like signal of the 1.1 TeV higgsino 'at high significance... for all but the most pessimistic DM profiles'. arXiv:2506.08084 (PoS ICRC2025) projects that CTAO-North, observing at large zenith angles, could reach the required sensitivity within the next decade. The server's indirect-detection treatment (p-wave, mass-splitting suppressed, sigma_v of about 1e-41 cm^3/s, f-002) is SUPERSEDED. A higgsino annihilates through an s-wave, Sommerfeld-enhanced channel, and the literature treats the thermal target as near the current Fermi/IACT reach rather than 15 orders of magnitude below it.

Completed — Establish the current indirect-detection status of the 1.0-1.2 TeV thermal higgsino from the literature: the H.E.S.S., Fermi-LAT and MAGIC gamma-ray line and continuum limits against the Sommerfeld-enhanced s-wave annihilation cross section, and the CTAO projections. This replaces the server's p-wave treatment, which is physically wrong for a higgsino.. Verdict, based on abstracts only (download_arxiv_full_text on 2405.13104 returned 'not found', tc-261016_1-w2-d-06-76): the 1.0-1.2 TeV thermal higgsino is not excluded by current gamma-ray line or continuum searches, and the result depends on the halo profile. Current searches: arXiv:2405.13104 (Rodd, Safdi, Xu, PRD 110, 043003) ran a 15-yr Fermi-LAT line-like search that comes 'within an order one factor' of the sensitivity needed to detect the higgsino, for some Milky Way analog density profiles from hydrodynamic simulations. The same paper finds H.E.S.S. subleading to Fermi for the higgsino. arXiv:2210.03140 (Montanari, Moulin, Rodd, PRD 107, 043028) finds the higgsino 'out of reach by at least a factor of a few' even at the ultimate sensitivity of current H.E.S.S.-like IACTs; the dominant systematic is the dark matter distribution in the inner Galaxy. The 2026-dated H.E.S.S. Inner Galaxy Survey, arXiv:2608.07234 (546 h, DOI 10.1103/d8tj-55kc, still to be checked on arxiv.org), quotes a line limit of 2.3e-28 cm^3/s at 1 TeV for an Einasto profile and says thermal

higgsino DM is 'probed for DM Milky Way models' for the first time. Its abstract does not say whether it excludes the higgsino, or for which profiles. No MAGIC-specific higgsino limit came up in the searches. Future: arXiv:2008.00692 says CTA could be the first experiment sensitive to the thermal higgsino; 2405.13104 says CTA and SWGO could detect it 'at high significance' for all but the most pessimistic profiles; 2506.08084 projects CTAO-North could reach the required sensitivity within a decade. The server's indirect-detection treatment (p-wave, mass-splitting suppressed, sigma_v of about 1e-41 cm^3/s, f-002) is explicitly flagged as superseded, because higgsino annihilation is s-wave and Sommerfeld-enhanced. The 'no Fermi-LAT exclusion' flags in the f-005 viable count are therefore not physically meaningful for indirect detection. The predicted line cross section at 1.1 TeV, and its ratio to the limit, could not be extracted without a full text. Findings f-011 and f-012 are posted (evidence tc-261016_1-w2-d-06-63, -71, -72). No files or figures were produced.

Finding: The OpenAlex record for arXiv:2609.02823 (LZ, 'Search for dark matter particle interactions in an extended nuclear recoil energy window with the LUX-ZEPLIN (LZ) experiment', 2026-09-02, DOI 10.48550/arxiv.2609.02823) carries this abstract text: "an exposure of 2.84 tonne-years", "An extended nuclear recoil energy window up to approximately 270 keV", "one event with characteristics consistent with a nuclear recoil of 248±23 (stat)±23 (sys) keV", and "global significance of 2.6σ ... maximum local significance of 3.4σ". The arXiv API itself returns 'not found' for this ID, so the quotes come from abstract text only and are not cite()-validated.

Finding: Abstract-level literature anchors from OpenAlex. (a) arXiv:2403.19598 (S. P. Martin, PRD 109, 095045, 2024) says a thermal higgsino-like DM "mass must be between about 100 GeV and 1150 GeV". (b) arXiv:2607.20588 (Bisal et al., 2026) says "NLO corrections lower the DM-nucleon cross section below the neutrino floor" for pure higgsinos, and that a pure higgsino "typically evades current direct detection limits". This abstract quotes no numerical cross section, so the target of about 1e-48 to 1e-49 cm^2 remains unsourced. (c) arXiv:2410.17137 (XLZD Design Book, EPJC 2025) reaches the neutrino fog, with 3 sigma evidence potential down to 3e-49 cm^2 at 40 GeV. (d) Planck 2018 papers 1807.06205 (overview) and 1907.12875 (likelihoods) were found, but the abstract of 1807.06209 (parameters) was not retrieved, so Omega_c h^2 = 0.120 +/- 0.001 is still unquoted.

Blocked — Verify and cite the key literature: the higgsino relic crossing mass, the LZ 2.84 t-yr extended-window result, the solar-capture delta bound, the elastic loop-level cross section relative to the XENONnT limits and the neutrino fog, and the Planck Omega h^2. Obtain verbatim quotes.. (The arXiv API behind get_arxiv_metadata, both download tools and cite's metadata check returns 'not found', HTTP 429 or timeouts even for 1807.06209, so no full text could be read and no quote could be validated with cite(). Only OpenAlex abstracts are reachable.) No cite() entries could be made, so the success criterion of at least 5 cited quotes was not met. The arXiv-API layer is still down. get_arxiv_metadata returned 'not found' for 2609.02823, 2609.02775, 1807.06209 and 2410.15733, including the well-established Planck ID, so these results say nothing about whether the papers exist. download_arxiv_full_text failed for 1807.06209, download_full_arxiv_paper crashed with "'Search' object has no attribute 'results"', and cite() failed its metadata check for 2410.15733 (tc-...-64, -73, -77, -80, -81, -82, -84; dead end x-02). Only the OpenAlex-backed search_arxiv returned anything, and it gives abstracts only.

Reference table from OpenAlex abstracts. None of these quotes are validated by cite().

- arXiv:2609.02823 (LZ, 2026-09-02): the OpenAlex record exists and its abstract text matches the board. Exposure 2.84 tonne-years, recoil window up to about 270 keV, one event at 248 +/- 23 (stat) +/- 23 (sys) keV, 2.6 sigma global and 3.4 sigma local significance (f-013, tc-...-85). The elastic SI limit at 1 TeV is not in the abstract.
- arXiv:2403.19598 (Martin, PRD 109, 095045): a thermal higgsino-like DM mass must be

between about 100 and 1150 GeV (f-014, tc-...-86).

- arXiv:2410.15733: the abstract quote 'Higgsino mass of ~1.2 TeV' was already recorded by d-04 (f-007). cite() failed on it.
- arXiv:2607.20588 (2026): next-to-leading-order corrections push the pure-higgsino nucleon cross section below the neutrino floor. The abstract gives no number, so the value of about 1e-48 to 1e-49 cm^2 is still unsourced.
- arXiv:2410.17137 (XLZD): neutrino-fog reach, with 3 sigma evidence potential at 3e-49 cm^2 for a 40 GeV WIMP.
- Planck 2018: search_arxiv found 1807.06205 and 1907.12875, but 1807.06209 did not appear in the results and its metadata lookup returned 'not found'. The value Omega_c h^2 = 0.120 +/- 0.001 remains unquoted (tc-...-83).

Not checked, for lack of tool budget: 2609.02775 (its metadata lookup returned 'not found'), 2609.02608, 2609.26698, 2609.15321 and 2610.00226. The solar-capture bound delta > 566 keV is therefore still unquoted.

Critic, wave 2 — continue: 2 findings stand up to the evidence, 4 questioned ('The indirect-detection status of the about 1.1 TeV thermal higgsino, taken from abstracts ': The quoted excerpts do not contain the claimed values. tc-63 is cut off at the 2608.07234 author list, so the 546 h exposure, the 2.3e-28 cm^3/s Einasto line limit at 1 TeV and the 'probed for DM Milky Way models' quote are not visible. tc-71 shows only 2609.01590. Neither 2405.13104 ('within an order one factor', H.E.S.S. subleading) nor 2210.03140 ('out of reach by at least a factor of a few') appears in any cited excerpt. The 'not excluded' conclusion is plausible, but it needs to be re-posted with the abstract text that actually holds these values.; 'CTAO is the decisive future test. arXiv:2008.00692 (Rinchiuso et al., PRD 103, 023011) fin': The excerpts show only the title and authors of 2008.00692 and the start of the 2609.01590 entry. The quoted CTA sentence, the 2405.13104 'at high significance... for all but the most pessimistic DM profiles' quote and the 2506.08084 CTAO-North projection are not in the evidence. The physics point that the server's p-wave sigma_v of about 1e-41 cm^3/s is wrong for an s-wave, Sommerfeld-enhanced higgsino is sound. Even so, confidence 'high' is not justified for abstract-level quotes that cannot be seen in the excerpts.; 'The OpenAlex record for arXiv:2609.02823 (LZ, 'Search for dark matter particle interaction': tc-85 is cut off after the LZ author list. The excerpt contains the ID and title but not the date, the DOI, '2.84 tonne-years', '270 keV', '248+/-23 (stat)+/-23 (sys) keV' or the 2.6/3.4 sigma significances. tc-64 and tc-81 only document API failures. The quoted abstract sentences need to be in the excerpt itself.; 'Abstract-level literature anchors from OpenAlex. (a) arXiv:2403.19598 (S. P. Martin, PRD 1': Only part (d) is supported: tc-77 and tc-82 show 1807.06209 'not found', and tc-83 shows 1907.12875. Parts (a) to (c) are not. The visible part of tc-86 shows 2203.02309, not 2403.19598, 2607.20588 or 2410.17137. The Martin 'between about 100 GeV and 1150 GeV' quote, the NLO neutrino-floor statement and the XLZD 3e-49 cm^2 figure are not in any quoted excerpt.).

Issues: The (794/776)^2 rescaling of delta(N=1) and delta(N=0.3) from 352.5/356.4 keV to about 369/373 keV is a heuristic. It assumes the whole N(delta) curve stretches with v_max^2, but near the kinematic edge the count depends on how the halo tail is populated between v_min and v_max, not only on where the edge sits. The arithmetic checks out (mu = 109.8 GeV gives delta_max = 367.8 and 385 keV), but the shifted window has not been computed. The claim that 2609.02608 uses v_max = 794 km/s is also unsourced, so attributing 17 keV of the 24.5 keV offset to the cutoff is an assumption.; The 350 keV point (N = 1.97, peak E_R = 315 keV) and the 300 keV point (744.6) come from wave-1 calls under d-02 and are not among this finding's evidence ids. The interpolations check out: ln-linear between 1.97 and 0.494 gives

352.45 keV, and between 0.494 and 0.085 gives 356.4 keV. The finding should still cite the d-02 call ids for the 350 keV anchor.; The board brief still says 'I did not compute any relic densities... No inelastic-DM or relic-density tools were available'. This contradicts d-01, which computed Omega h^2 at five masses (crossing at about 1158 GeV), and d-02 and d-05, which computed LZ event counts. The brief must be rewritten before synthesis. It should also downgrade the 'no Fermi-LAT exclusion' parts of the f-005 viable count, given the conclusion that f-002 is superseded..

Stopping after wave 2: budget reached (2/2 waves used). Writing the report from the board.

Research report:

The 1.0–1.2 TeV thermal higgsino against LZ, XENONnT, Planck and gamma-ray limits: toy-model calculations and an abstract-level literature survey

Abstract

This investigation asked where the pure higgsino thermal relic, at about 1.0–1.2 TeV, sits relative to current limits.

Tool calculations. An inelastic-dark-matter (inelastic-DM) server was used for relic and recoil calculations.

- **Relic density.** The server's relic model is a single-channel freeze-out scaling calibrated near 1.1 TeV. It reaches the Planck abundance at about 1158 GeV. This lies inside the 1.08–1.2 TeV literature range, but it is a calibration result, not a Sommerfeld/coannihilation prediction.
- **LZ inelastic events.** The server's Z-exchange higgsino predicts these LZ counts at 1.1 TeV for the 2.84 tonne-year extended nuclear-recoil window:
- **Kinematic edge and one-event window.** The cutoff sits at about 368 keV for the tool's 776 km/s halo cutoff. It would sit at about 385 keV for 794 km/s. The "about one event" window is only about 4 keV wide, at 352–356 keV.

Mass splitting delta (keV)	Expected LZ events
300	744.6 (excluded)
350	about 2
380 and above	0

Literature, from abstracts only. None of the following could be verified against full texts.

- Solar-capture limits from IceCube require delta > 566 keV.
- That puts every solar-capture-allowed thermal higgsino beyond the inelastic reach of xenon.
- The pure-higgsino elastic cross section is described as lying below the neutrino floor.
- The higgsino is not excluded by current gamma-ray line searches, and the Cherenkov Telescope Array Observatory (CTAO) is identified as the decisive future test.

Bottom line. The full-abundance thermal higgsino at about 1.1 TeV is not excluded. Its interpretation as the source of the sub-MeV-splitting LZ 248 keV event is disfavoured. With splittings of order an MeV it is effectively invisible to current xenon detectors.

Background

Thermal mass. With Sommerfeld enhancement and coannihilation included, a pure higgsino gives the full Planck abundance at roughly 1.0–1.2 TeV. Quoted values are:

- about 1.08–1.1 TeV (arXiv:2609.02775, arXiv:2609.02608);
- "~1.2 TeV" (arXiv:2410.15733);
- an upper end of about 1150 GeV, with the thermal higgsino-like mass required to lie "between about 100 GeV and 1150 GeV" (arXiv:2403.19598).

Lighter thermal higgsinos are under-abundant by a factor of 5–20 in natural SUSY (arXiv:1803.11210).

Direct detection. The pure higgsino is inelastic dark matter. Z-mediated scattering requires the neutral-state splitting delta to be no more than a few hundred keV. If delta lies beyond the kinematic reach of xenon, only loop-level elastic scattering remains, and that rate sits near or below the neutrino fog (arXiv:2609.01504, arXiv:2410.02723). One 2026 abstract states that next-to-leading-order corrections push the pure-higgsino nucleon cross section "below the neutrino floor" (arXiv:2607.20588). Higgsino–gaugino mixing raises the elastic rate (arXiv:2606.31013).

The 2026 LZ result. LZ reported a 2.84 tonne-year search with a recoil window extended to about 270 keV (arXiv:2609.02823). It found one event at 248 ± 23 (stat) ± 23 (sys) keV, with 2.6σ global and 3.4σ local significance.

Interpretations of that event as a higgsino give these preferred splittings:

Assumption or method	Preferred delta	Source
Standard Halo Model	about 350 keV	arXiv:2609.01504
Standard Halo Model	377 keV	arXiv:2609.02608
Sideband optimum at 1 TeV	339 keV, about 1.9σ tension	arXiv:2609.26698
Including the Large Magellanic Cloud (LMC)	about 480–520 keV	arXiv:2609.01504; 481–519 keV in arXiv:2609.01590

Tensions with the event interpretation.

- **Solar capture.** IceCube non-observation requires delta > 566 keV (arXiv:2609.02775). This is supported by arXiv:2609.11833 and arXiv:2609.07807. arXiv:2609.06760 quotes 0.51–0.56 MeV.
- **Empty sideband.** The absence of events above 350 keV is in tension with the thermal-mass fit (arXiv:2609.09385, arXiv:2609.26698).
- **Halo uncertainties.** These cannot close the gap (arXiv:2609.15321). Under the Standard Halo Model, delta > 566 keV would need v0 > 240 km/s. With the LMC included, the largest workable splitting is 514 keV.

Surviving branch. A Planck-compatible thermal branch survives at delta of about 0.85–1.34 MeV (arXiv:2610.00226). arXiv:2609.09136 proposes a non-higgsino alternative.

Caveat on sources. All 2026-dated identifiers came from an OpenAlex-backed search and could not be resolved through the arXiv API. They should be checked on arxiv.org before anyone cites them.

Results

Relic density from the inelastic-DM server (toy model) The relic tool was run with the higgsino flag set. This flag sets σv = 2.5×10^{-2}? cm^3/s × (1100 GeV/m)^2 and feeds it into a Kolb–Turner freeze-out estimate, with x_f about 24.5 and g* = 86.25. The tool has no Sommerfeld or coannihilation switch.

- **Planck crossing.** Linear interpolation reaches Ω h^2 = 0.120 at about 1158 GeV. The server's ±0.0012 band gives ±6 GeV; the ±0.001 band gives about ±5 GeV. A direct call at 1159 GeV returns 0.120.
- **Comparison with the literature.** This crossing is inside the 1.08–1.2 TeV range. It is about 60–80 GeV above the 1.08–1.1 TeV values and close to the ~1.2 TeV value of arXiv:2410.15733.
- **Interpretation.** The agreement reflects how the tool's cross section is normalized. It should be read as a toy-model crossing, not a higgsino prediction.
- **Tests not run.** The with/without-Sommerfeld comparison and splitting-dependence tests could not be done.
- **Pass flag.** The tool marks results as "consistent" with Planck even at 7–25σ tension; for example, 1100 GeV carries 9.8σ tension. This looks like a bug.

m_χ (GeV)	**Ω h^2**
900	0.0731
1000	0.0898
1100	0.1083
1200	0.1285
1300	0.1503

LZ inelastic event counts and the kinematic reach of xenon The recoil tool uses:

- Z exchange, with σ_n = 7.43×10^{-3}? cm^2 per nucleon;
- the LZ 2.84 tonne-year extended nuclear-recoil window;
- a hard halo cutoff at v_esc + v_E = 776 km/s.

Expected LZ counts at m_χ = 1100 GeV:

delta (keV)	**Expected LZ events**	**Spectrum peak (keV)**	**Notes**
300	744.6	not recorded	Grossly excluded against 1 observed event
350	1.97	315	P(N $\geq$ 1) = 0.86; the 248 keV event needs v_min = 762 km/s, which is accessible

delta (keV)	Expected LZ events	Spectrum peak (keV)	Notes
380	0.00	342	The 248 keV event needs v_min = 799 km/s, which is inaccessible
500	0.00	450	
566	0.00	510	
700	0.00	630	

Hand check of the edge. The maximum splitting is δ_max = μ v_max^2/2. With m_Xe of about 122 GeV, the reduced mass is μ = 109.8 GeV.

- For v_max = 776 km/s, δ_max is about 368 keV.
- For v_max = 794 km/s (544 + 250 km/s), δ_max is about 385 keV.

This matches the tool, which gives events at 350 keV and none at 380 keV.

Mass dependence. The edge depends only weakly on mass. At delta = 380 keV, v_min for the 248 keV event is 802, 799 and 796 km/s at 1000, 1100 and 1200 GeV, all above 776 km/s.

Consequence. The solar-capture bound of delta > 566 keV from the literature therefore puts every solar-capture-allowed thermal higgsino beyond LZ's inelastic reach under the Standard Halo Model.

Annual modulation. At 1100 GeV, 350 keV and E_R = 248 keV, the tool gives:

- a fractional amplitude of 98.5%, peaking on June 2;
- 22.5% of the peak rate on the event date of September 1.

A recoil-spectrum plot for 1100 GeV and 380 keV was written to `/Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png`. No numbered figure was produced for this report.

The one-event splitting window A finer scan at 1100 GeV gives:

- **Window location.** Log-linear interpolation gives delta(N=1) = 352.5 keV and delta(N=0.3) = 356.4 keV. The window is therefore about 4 keV wide, and N falls by about 4–6 times per 5 keV.
- **Where the event becomes inaccessible.** The 248 keV event drops out between 360 and 365 keV, where v_min rises from 774 to 781 km/s.
- **Thermal mass.** At the tool's thermal mass of 1159 GeV, delta = 360 keV gives N = 0.135, against 0.085 at 1100 GeV. This moves delta(N=1) up by only about 1–2 keV.
- **Cause of the zero counts.** The zero counts at delta $\geq$ 370 keV come from the hard 776 km/s cutoff, not the energy window.
- **Halo dependence (not computed).** The recoil tool has no halo inputs, so halo dependence could not be computed. An analytic rescaling by $(794/776)^2 = 1.047$ would move the window to about 369–373 keV. This is a heuristic: near the edge, the count depends on how the velocity tail is populated, not only on where the edge sits.

delta (keV)	Expected LZ events	Spectrum peak (keV)
350	1.97	315
355	0.494	320
360	0.085	324
365	0.009	329
370	0.00	333
375	0.00	338

Comparison with literature fits:

- **350 keV (arXiv:2609.01504).** Agrees with the tool directly.
- **377 keV (arXiv:2609.02608).** The offset is 24.5 keV. About 17 keV of it would be explained if that paper used a 794 km/s cutoff, but this is an assumption, not a sourced fact. The remaining roughly 5–8 keV is unexplained. Possible causes are the halo tail shape, v_E at the event date, and cross-section or efficiency normalization; none was tested.
- **339 keV (arXiv:2609.26698).** This is a normalization-free sideband-ratio optimum. With the tool's normalization, splittings this low give hundreds of events, so it is not a like-for-like comparison.

The tool therefore does not by itself account for the spread among literature fits.

Parameter scan and constraint checker A 17 × 17 scan covered m_χ = 800–1400 GeV and delta = 0–800 keV. It reported 144 of 289 points "viable" and 9 points predicting 0.3–5 LZ events. The best-fit point is m_χ = 988 GeV, delta = 350 keV. Output files are:

- `/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz`
- `/Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png`

These numbers should not be taken as physics, for three reasons:

- The viable count uses the relic pass flag that stays true at 7–25σ.
- It uses an indirect-detection model that is wrong for a higgsino (see below).
- The best fit fits the LZ event, not the relic. At 988 GeV the tool's own relic curve gives Ω h^2 of about 0.087, so this point is under-abundant.

The constraint checker rated four points VIABLE: 1100 GeV with delta = 400, 566 and 1000 keV, and 1159 GeV with 400 keV. All four have N_LZ = 0.00 and no Fermi-LAT or CMB exclusion. Its present-day σv of about $7\times10^{-}?^{2}$ to $5\times10^{-}?^{1}$ cm^3/s is labelled "p-wave + mass-splitting suppressed". That is inappropriate for an s-wave, Sommerfeld-enhanced higgsino, whose cross section is about $10^{-2}?$ cm^3/s.

Elastic scattering (literature only) The XENONnT 3.1 tonne-year search (arXiv:2502.18005) reports:

- a minimum spin-independent limit of $1.7\times10^{-??}$ cm^2 at 30 GeV (90% CL);
- a best median sensitivity of $1.4\times10^{-??}$ cm^2 at 41 GeV.

The earlier 1.09 tonne-year result (arXiv:2303.14729) reached $2.58\times10^{-??}$ cm^2 at 28 GeV.

Neither abstract gives the limit at 1 TeV. The figure of about $10^{-??}$ cm^2 is only an extrapolation using linear high-mass scaling.

XLZD (arXiv:2410.17137) reaches the neutrino fog, with 3σ evidence potential down to $3\times10^{-??}$ cm^2 at 40 GeV.

The pure-higgsino loop-level elastic cross section of about $10^{-??}$–$10^{-??}$ cm^2 was not sourced. arXiv:2607.20588 places it qualitatively below the neutrino floor without giving a number.

Indirect detection (literature only) These findings are based on abstracts only. The full text of arXiv:2405.13104 could not be downloaded.

- **Fermi-LAT.** A 15-year line-like search comes "within an order one factor" of the sensitivity needed to detect the higgsino, for some simulated Milky Way profiles. In the same analysis, H.E.S.S. is subleading to Fermi (arXiv:2405.13104).
- **Current imaging Cherenkov telescopes.** At their ultimate sensitivity, H.E.S.S.-like instruments leave the higgsino "out of reach by at least a factor of a few". The dominant systematic is the dark-matter density in the inner Galaxy (arXiv:2210.03140).
- **H.E.S.S. Inner Galaxy Survey.** This 2026-dated, unverified paper used 546 h of data (arXiv:2608.07234). It quotes a line limit of 2.3×10^{-2}? cm^3/s at 1 TeV for an Einasto profile and says the thermal higgsino is "probed" for the first time. Its abstract does not say whether, or for which profiles, the higgsino is excluded.
- **MAGIC.** No MAGIC-specific limit surfaced.
- **Future instruments.**
- CTA could be the first experiment with sensitivity to the thermal higgsino (arXiv:2008.00692).
- CTA and SWGO could detect it "at high significance" for all but the most pessimistic profiles (arXiv:2405.13104).
- CTAO-North could reach the required sensitivity within a decade (arXiv:2506.08084).

The predicted line cross section at 1.1 TeV, and its ratio to current limits, could not be extracted.

Literature verification status Every arXiv-API-backed tool failed: metadata lookup, both download tools, and the metadata check inside the citation tool. They returned "not found", HTTP 429 or timeouts, even for the Planck 2018 parameters paper (1807.06209). No quote could be validated and no bibliography entry was made.

Abstract-level text was obtained through OpenAlex for:

- the LZ result (arXiv:2609.02823);

- arXiv:2403.19598;
- arXiv:2410.15733;
- arXiv:2607.20588;
- arXiv:2410.17137.

Planck overview and likelihood papers (1807.06205, 1907.12875) were found. The parameters abstract was not, so $\Omega_c\ h^2 = 0.120 \pm 0.001$ is used here as the standard value but remains unquoted.

Discussion

Answering "where does the higgsino sit?" requires assembling three pieces.

First, the relic target. This is set by the literature at about 1.1 TeV, give or take about 0.1 TeV. The tool's crossing at about 1158 GeV is consistent with that only because the tool is calibrated there.

Second, inelastic direct detection. The xenon kinematics are robust and do not depend on the toy model. At the thermal mass, δ_max is about 368–385 keV depending on the velocity cutoff, nearly independent of mass. The sub-MeV LZ interpretation therefore requires a splitting in a narrow band just below this edge. In the tool, that band is 352–356 keV for one to 0.3 events, and lower splittings are grossly excluded.

The solar-capture bound reported in the literature (delta > 566 keV, or 0.51–0.56 MeV) lies far above this edge under the Standard Halo Model. It also lies above the roughly 480–520 keV LMC-tail values. This is why the LZ interpretation is disfavoured while the higgsino itself survives at delta of order an MeV (arXiv:2610.00226). In that regime, inelastic scattering is closed in xenon. The elastic rate is described as below the neutrino floor, although no number was sourced here.

Third, indirect detection. The literature consensus from abstracts is that the higgsino is within an order-one factor of current Fermi-LAT line sensitivity and is not excluded, with strong dependence on the halo profile. CTAO is the decisive future test.

The results agree with the literature on:

- the location of the xenon edge;
- the roughly 350 keV preferred splitting under a Standard-Halo-like cutoff.

They are in tension, or at least unreconciled, with:

- the 377 keV fit of arXiv:2609.02608, where 5–8 keV is unexplained even after the assumed cutoff correction;
- the 339 keV sideband optimum, which is not a comparable quantity.

Dead ends:

- **Verbatim citations.** Verbatim citation of Planck 2018 VI, arXiv:2410.15733, arXiv:2609.02823 and arXiv:2609.02775 was tried twice. Each time, the arXiv API layer returned "not found", 429 errors or internal errors, even for long-established papers. A "not found" result therefore says nothing about whether a paper exists.

- **Tool-based Sommerfeld, halo and indirect-detection studies.** These were impossible because the server has no Sommerfeld switch and no halo parameters, and its indirect-detection model is p-wave.

Limitations

- **Relic model.** The relic calculation is a toy single-channel scaling. No Sommerfeld or coannihilation physics was computed, and no tool for it was available.
- **Halo model.** Recoil predictions use a single hard 776 km/s cutoff. Halo sensitivity, including v0, v_esc, a smooth tail and the LMC, was only estimated by an analytic rescaling. The Standard Halo Model v0 scan of arXiv:2609.15321 and the sideband analysis of arXiv:2609.26698 were not reproduced.
- **Binned counts.** Counts in the [14,225), [225,271] and (271,800] keV bins were not obtained. The tool returns integrated rates only.
- **Constraint checker.** Its relic pass flag and its indirect-detection model are unreliable. Its viability counts and "no Fermi-LAT exclusion" verdicts should be disregarded.
- **

System note (2026-10-16 15:03:49)

Report generated: /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/reports/261016_150203/report.pdf Title: The 1.0–1.2 TeV Thermal Higgsino Against Planck, LZ, XENONnT and Gamma-Ray Limits: A Toy-Model Study Supplement (implementation, tool-call log, board, extra figures, code index): /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/reports/261016_150203/supplement.pdf Transcript: /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/reports/261016_150203/transcript
Folder: /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/reports/261016_150203
(report.tex, supplement.tex, transcript.tex, figures/, bundle.json)

User (2026-10-22 14:31:35)

Survey recent arXiv work on the higgsino thermal relic target (1.0-1.2 TeV). Use the inelastic-DM tools to compute the relic density at several masses; compare with LZ/XENONnT sensitivities and the Planck abundance. Where does the higgsino sit relative to current limits?

Assistant (research mode, 2026-10-22 14:31:35)

Plan:

1. (closed) Place the thermal pure higgsino (m = 1.1-1.16 TeV) on the elastic spin-independent plane. Read the LZ 4.2 t-yr (2410.17036), XENONnT 3.1 t-yr (2502.18005) and first-run (2207.03764, 2303.14729) limits at 1.0-1.2 TeV, the loop-level higgsino cross section (1309.4092, 2607.20588) and the neutrino fog (2410.17137). Then quote the ratio of the higgsino cross section to the limit and to the fog.
2. (closed) Recover verbatim, citable support for the core numbers that are still unquoted. These are the Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209), the thermal higgsino mass (2410.15733 at about 1.2 TeV, 2609.02608 and 2609.02775 at about 1.08-1.1

TeV, 2403.19598 at 1150 GeV), and the solar-capture bound delta > 566 keV (2609.02775). Resolve why the literature mass values disagree.

3. (closed) Robustness of the tool relic calculation, and the link between gaugino mixing and the splitting. (a) Use compute_relic_density with an explicit sigma_v_cm3_s to find the effective cross section needed for Omega h^2 = 0.120 at 1.1 TeV. Compare it with the tool's built-in higgsino sigma_v and with the Sommerfeld-enhanced literature values. (b) Translate the xenon-accessible splitting window (delta of about 345-385 keV) and the solar bound (566 keV) into gaugino masses M1 and M2, and estimate the mixing at which Higgs-exchange elastic scattering reaches the LZ limit.

4. (closed) Produce the tool-based deliverable the task explicitly requests: run check_all_constraints at the literature thermal masses (1.08, 1.10, 1.16, 1.20 TeV) and one scan_mass_splitting over m = 1.0-1.2 TeV, delta = 0-700 keV, with a plot_constraint_summary figure.

5. (blocked) Robustness of the tool's inelastic LZ prediction near the preferred splitting. At m = 1100 and 1160 GeV, use predict_recoil (or predict_annual_modulation if it exposes halo inputs) to find how the expected LZ count and the delta at which N_LZ is about 1 move when v_esc is varied (about 500, 544 and 600 km/s) and v0 is varied (about 220 and 238 km/s). Compare the resulting window with the literature preferred delta (about 377 keV, 2609.02608) and the solar bound (above 566 keV, 2609.02775).

Tool calls: `list_sterile_nu_skills`, `list_observables`, `list_agent_files`, `list_input_files`, `list_datasets`, `list_files`, `get_dispatch`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `get_arxiv_metadata`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `search_arxiv_abstracts`, `describe_tools`, `get_arxiv_metadata`, `list_files`, `get_paper_figures`, `get_paper_figures`, `check_all_constraints`, `check_all_constraints`, `check_all_constraints`, `check_all_constraints`, `check_all_constraints`, `check_all_constraints`, `cite`, `get_arxiv_metadata`, `get_arxiv_metadata`, `get_arxiv_metadata`, `get_arxiv_metadata`, `cite`, `cite`, `cite`, `cite`, `search_arxiv_abstracts`, `post_finding`, `post_finding`, `scan_mass_splitting`, `plot_constraint_summary`, `post_finding`, `post_finding`, `post_question`, `arxiv_agent`, `arxiv_agent`, `arxiv_agent`, `predict_recoil`, `arxiv_agent`, `arxiv_agent`, `post_finding`, `post_finding`, `post_finding`, `post_finding`, `post_finding`, `post_dead_end`, `cite`, `post_dead_end`, `describe_tools`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `compute_relic_den` `compute_relic_density`, `compute_relic_density`, `compute_relic_density`, `compute_relic_density`, `compute_relic_density`, `compute_relic_density`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `predict_recoil`, `predict_annual_modulation`, `post_finding`, `post_dead_end`, `post_finding`, `arxiv_agent`, `post_finding`, `post_finding`, `post_finding`

Survey — 52 tools on 5 servers, 7 read-only probes, 11 papers found.

Warning: two limitations up front. First, my tools were limited to arXiv/web search and metadata, so I did not compute any relic densities. Neither the inelastic-DM tools named in the task nor any relic-density code were available to me. Second, my arXiv searches did not turn up the main recent paper deriving the higgsino thermal mass target (about 1.0-1.2 TeV with Sommerfeld effects). From general knowledge, a pure higgsino (an electroweak doublet) gives the Planck abundance of Omega h^2 of about 0.12 at about 1.1 TeV. Planck 2018 (1807.06209) is the reference for this value; its abstract was cut off, so the exact number should be taken from the paper. Searches did return a pMSSM study of heavy higgsino and wino relic abundance with Sommerfeld enhancement (1411.6930) and general methods for including the Sommerfeld effect in relic calculations (1008.2905, 0910.5221). On direct detection: at tree level a pure higgsino's spin-independent scattering is suppressed. It only reaches xenon detectors through electroweak loops, which give an elastic cross-section well below 1e-46 cm^2 (heavy-WIMP effective theory,

1309.4092). Gaugino mixing adds Higgs exchange; if the mass splitting between the neutral states drops below about 100-200 keV, Z-mediated inelastic scattering switches on and is strongly constrained. The 4.2 tonne-year LZ result (2410.17036), the 3.1 tonne-year XENONnT result (2502.18005) and earlier first runs are the current limits. A 2026 paper (2609.09830) works out loop-corrected elastic and inelastic higgsino scattering rates in LZ. The expected conclusion, still to be confirmed against the published limit curves, is that a thermal pure higgsino near 1.1 TeV sits orders of magnitude below current LZ/XENONnT spin-independent limits, near or under the neutrino fog. Xenon detectors can probe it only if there is enough gaugino mixing or a mass splitting small enough for inelastic scattering.

Quantities worth checking: Omega h^2 against higgsino mass over 0.9-1.3 TeV, with and without the Sommerfeld effect and coannihilation with the charged and second neutral state; find the mass where it crosses about 0.120.; Exact Planck 2018 Omega_c h^2 and its uncertainty (from 1807.06209), and how much the higgsino mass band shifts across that uncertainty.; Charged-neutral mass splitting (about 340 MeV from electroweak loops) and the neutral-neutral splitting as a function of gaugino masses M1 and M2.; Inelastic-scattering cutoff: the largest neutral splitting that xenon can still access (about 100-200 keV for a TeV-mass WIMP at the galactic escape velocity).; Pure-higgsino loop-level elastic cross-section (expected roughly 1e-49 to 1e-48 cm^2; confirm against 1309.4092) compared with LZ 2024 and XENONnT 2025 limits at 1.1 TeV.; How far below the 1.1 TeV LZ limit the higgsino sits (ratio), and its position relative to the neutrino fog..

Plan — 4 lines of investigation:

1. Place the thermal pure higgsino (m = 1.1-1.16 TeV) on the elastic spin-independent plane. Read the LZ 4.2 t-yr (2410.17036), XENONnT 3.1 t-yr (2502.18005) and first-run (2207.03764, 2303.14729) limits at 1.0-1.2 TeV, the loop-level higgsino cross section (1309.4092, 2607.20588) and the neutrino fog (2410.17137). Then quote the ratio of the higgsino cross section to the limit and to the fog. — This is the most direct remaining answer to 'where does the higgsino sit relative to current limits'. The inelastic part is settled (f-003 to f-010). The elastic limit at 1 TeV, the loop cross section and the fog level are still unsourced, because the arXiv API layer failed. get_paper_figures uses arXiv HTML and arxiv_agent is a separate pipeline, so both are untried routes.

2. Recover verbatim, citable support for the core numbers that are still unquoted. These are the Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209), the thermal higgsino mass (2410.15733 at about 1.2 TeV, 2609.02608 and 2609.02775 at about 1.08-1.1 TeV, 2403.19598 at 1150 GeV), and the solar-capture bound delta > 566 keV (2609.02775). Resolve why the literature mass values disagree. — Directions d-03 and d-04 were blocked by the arXiv API. The final answer rests on the Planck value and the relic mass target, so these need sources that can be quoted. The 1.08 vs 1.2 TeV spread (f-001, f-007 contested) should be explained, for example by different Sommerfeld, bound-state formation or electroweak-correction treatments.

3. Robustness of the tool relic calculation, and the link between gaugino mixing and the splitting. (a) Use compute_relic_density with an explicit sigma_v_cm3_s to find the effective cross section needed for Omega h^2 = 0.120 at 1.1 TeV. Compare it with the tool's built-in higgsino sigma_v and with the Sommerfeld-enhanced literature values. (b) Translate the xenon-accessible splitting window (delta of about 345-385 keV) and the solar bound (566 keV) into gaugino masses M1 and M2, and estimate the mixing at which Higgs-exchange elastic scattering reaches the LZ limit. — The tool's relic curve is a calibrated scaling with no Sommerfeld or coannihilation physics (f-001). Varying sigma_v directly quantifies how sensitive the 1158 GeV crossing is to that missing physics. The brief's 'gaugino mixing needed' question is still open. With the tree-level relation delta0 of

about m_Z^2 (sin^2 theta_W/M1 + cos^2 theta_W/M2) and published Higgs-exchange formulas, the inelastic window can be mapped onto decoupled gauginos at roughly 10-30 TeV.

4. Produce the tool-based deliverable the task explicitly requests: run check_all_constraints at the literature thermal masses (1.08, 1.10, 1.16, 1.20 TeV) and one scan_mass_splitting over m = 1.0-1.2 TeV, delta = 0-700 keV, with a plot_constraint_summary figure. — No proposed direction uses check_all_constraints, scan_mass_splitting or plot_constraint_summary. The final answer needs one reproducible table and figure that combine relic density, LZ, Fermi-LAT and CMB from the idm server. That output can then be checked against the literature numbers from d-07 and d-08. (added by the critic) Critic's view: revise — The gaugino-scale estimate in the rationale is off by about three orders of magnitude. Using delta0 ~ m_Z^2/M, a splitting of 350-570 keV needs M ~ 1e7 GeV (about 1e4 TeV), not 10-30 TeV. Gauginos at 10-30 TeV give splittings of order 0.3-1 GeV. The formula also depends on tan(beta) and the relative signs of M1, M2 and mu, through factors of (1 +/- sin 2beta). Redo step 2 with the full expression and state which sign convention is used. Higgs-exchange sigma_SI at M ~ 1e7 GeV is negligible, so step 3 should explicitly test whether the inelastic window and LZ-reachable Higgs exchange are mutually exclusive. They very likely are, and that is a substantive conclusion.; Step 4 ('Optionally call predict_sterile_nu_grid ... no') is not a step; delete it. Before spending about 10 calls on compute_relic_density, confirm with describe_tools that it accepts a sigma_v_cm3_s argument. Also add the Planck-uncertainty propagation the brief asks for. With Omega proportional to m^2 at fixed scaling, dm/m = 0.5 dOmega/Omega, so Omega_c h^2 = 0.1200 +/- 0.0012 shifts the mass by roughly +/- 6 GeV. This is negligible next to the 1.08-1.2 TeV literature spread, and the report should say so.; The direction calls get_paper_figures on 7 papers, plus search, predict_recoil and cite. That is about 11-12 calls with no room for retries. Order the papers by priority: LZ 2024, then the higgsino loop sigma_SI source, then the fog, then XENONnT 2025, then the first runs. The first-run comparison may be dropped if the budget runs out. Extrapolate high-mass limits as sigma proportional to m (the standard large-m scaling), not 'linearly in m' on a log plot. Verify that 2607.20588 and 2410.17137 exist with get_arxiv_metadata or search before relying on them. For 1309.4092 (Hill-Solon), record the large cancellation uncertainty of the pure-higgsino result, which spans roughly 1e-49 to 1e-48 cm^2 with sign-sensitive cancellations, rather than a single central value..

Wave 1 — now exploring, in parallel: Place the thermal pure higgsino (m = 1.1-1.16 TeV) on the elastic spin-independent plane. Read the LZ 4.2 t-yr (2410.17036), XENONnT 3.1 t-yr (2502.18005) and first-run (2207.03764, 2303.14729) limits at 1.0-1.2 TeV, the loop-level higgsino cross section (1309.4092, 2607.20588) and the neutrino fog (2410.17137). Then quote the ratio of the higgsino cross section to the limit and to the fog.; Recover verbatim, citable support for the core numbers that are still unquoted. These are the Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209), the thermal higgsino mass (2410.15733 at about 1.2 TeV, 2609.02608 and 2609.02775 at about 1.08-1.1 TeV, 2403.19598 at 1150 GeV), and the solar-capture bound delta > 566 keV (2609.02775). Resolve why the literature mass values disagree.; Produce the tool-based deliverable the task explicitly requests: run check_all_constraints at the literature thermal masses (1.08, 1.10, 1.16, 1.20 TeV) and one scan_mass_splitting over m = 1.0-1.2 TeV, delta = 0-700 keV, with a plot_constraint_summary figure..

Finding: Running check_all_constraints with is_higgsino=true and delta=350 keV gives Omega h^2 = 0.1045, 0.1083, 0.1202 and 0.1285 at m = 1080, 1100, 1160 and 1200 GeV. The tool's relic crossing is therefore at about 1160 GeV, with tension 12.9, 9.8, 0.1 and 7.1 sigma against 0.12 +/- 0.0012. The expected LZ counts are N_LZ = 1.79, 1.97, 2.52 and 2.90, all

consistent with the 1 observed event. The Fermi-LAT and CMB checks pass because the inelastic annihilation rate today is only about 5e-42 cm^3/s.

Finding: The splitting dependence at m = 1100 GeV from check_all_constraints is as follows. At delta = 0 (elastic Z exchange, sigma_n = 7.4e-39 cm^2) the tool gives N_LZ = 1.1e7, sigma_v today = 2.5e-24 cm^3/s against the Fermi limit of 2.3e-25, and p_ann = 3.4e-28 against the limit of 3.2e-28, so the point is excluded on all three. At delta = 200 keV it gives N_LZ = 5.1e4. At delta = 350 keV it gives N_LZ = 1.97. At delta = 566 keV it gives N_LZ = 0.0, which is kinematically closed. Only a narrow band near delta of about 350 keV reproduces of order 1 LZ event.

Finding: Verbatim-cited (cite() succeeded, references.bib in run dir): Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209 abstract); thermal higgsino mass ~1.2 TeV with coannihilation + Sommerfeld (2410.15733), m ~1.08 TeV 'cosmologically preferred' (2609.02775), ~1.1 TeV (2609.02608, with delta ~377 keV for LZ event), upper end 1150 GeV for higgsino-like thermal DM (2403.19598); solar-capture IceCube bound delta > 566 keV at m=1.08 TeV (2609.02775). The 1.08-1.2 TeV spread is not resolved from abstracts: 2410.15733 explicitly includes Sommerfeld+coannihilation; the others do not state their treatment in the abstract, so attributing the spread to bound-state formation/NLO/Planck input remains unverified (likely ~10% theory systematics; consensus band 1.08-1.2 TeV, central ~1.1 TeV).

Finding: A scan_mass_splitting run over m = 1000-1200 GeV and delta = 0-700 keV (15x15 = 225 points) gives 210 of 225 points that pass the tool's constraints, though that flag ignores direct detection. Only 15 points predict 0.3-5 LZ events. The plot_constraint_summary figure is at /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png, and its reported best fit is m = 1000 GeV, delta = 350 keV.

Completed — Recover verbatim, citable support for the core numbers that are still unquoted. These are the Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209), the thermal higgsino mass (2410.15733 at about 1.2 TeV, 2609.02608 and 2609.02775 at about 1.08-1.1 TeV, 2403.19598 at 1150 GeV), and the solar-capture bound delta > 566 keV (2609.02775). Resolve why the literature mass values disagree.. The arXiv API is working again. get_arxiv_metadata on 1807.06209 returned normally, so the 2609.x IDs do resolve (this answers q-03). cite() succeeded for 5 papers, all quoted verbatim from their abstracts and written to /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib.

Cited values:

- Planck 2018 (1807.06209): 'A combined analysis gives dark matter density Omega_c h^2 = 0.120 +/- 0.001'.
- 2410.15733: thermal higgsino mass about 1.2 TeV. The abstract states it includes coannihilation and the Sommerfeld effect.
- 2609.02775 (Pospelov and Ramani): 'cosmologically preferred mass' of about 1.08 TeV, and the solar-capture bound delta > 566 keV.
- 2609.02608 (Di Mauro): relic density fixes the mass near 1.1 TeV; a splitting delta of about 377 keV reproduces the 248 keV LZ event.
- 2403.19598 (Martin): thermal higgsino-like mass between about 100 and 1150 GeV.

Solar-capture bound: its source is cited. The abstract gives these assumptions: solar-core speeds up to about 1400 km/s, inelastic scattering on heavy elements, annihilation to WW and ZZ, and IceCube's non-observation of high-energy neutrinos from the Sun. I did not extract the spin-dependent cross section or the halo parameters because the full text was not read.

Mass spread: not resolved from the sources. Only 2410.15733 states its treatment in the abstract (Sommerfeld plus coannihilation, about 1.2 TeV). None of the abstracts mention bound-state formation, NLO corrections or the Planck input, so q-09 stays open. The recommended band, 1.08-1.2 TeV with a central value near 1.1 TeV, is my inference from the spread of the cited values, not a published result.

Result recorded as finding f-017. No figures were produced.

Completed — Produce the tool-based deliverable the task explicitly requests: run check_all_constraints at the literature thermal masses (1.08, 1.10, 1.16, 1.20 TeV) and one scan_mass_splitting over m = 1.0-1.2 TeV, delta = 0-700 keV, with a plot_constraint_summary figure.. I ran the idm-server constraint checks, the scan and the figure. The tool's own LZ limit could not be cross-checked against the published LZ curve, because the tool contains no such curve. In the tool, the relic density for a higgsino matches Planck (0.12) at about 1160 GeV, and a splitting near 350 keV gives about 1 LZ event. Two tool flags are unreliable and should not be quoted as physical viability.

Results of check_all_constraints (higgsino, delta = 350 keV), tc-25/28/29/30:

- m = 1080 GeV: Omega h^2 = 0.1045 (12.9 sigma from 0.12 +/- 0.0012), N_LZ = 1.79
- m = 1100 GeV: Omega h^2 = 0.1083 (9.8 sigma), N_LZ = 1.97
- m = 1160 GeV: Omega h^2 = 0.1202 (0.1 sigma), N_LZ = 2.52
- m = 1200 GeV: Omega h^2 = 0.1285 (7.1 sigma), N_LZ = 2.90
- At all four masses Fermi-LAT and CMB pass: sigma_v today is about 5e-42 cm^3/s against a limit of about 2.3e-25.

Splitting dependence at m = 1100 GeV, tc-31/32/33:

- delta = 0: N_LZ = 1.1e7, sigma_v = 2.5e-24 cm^3/s (Fermi limit 2.3e-25), p_ann = 3.4e-28 (CMB limit 3.2e-28). Excluded by LZ, Fermi-LAT and CMB.
- delta = 200 keV: N_LZ = 5.1e4, so LZ excludes it.
- delta = 566 keV: N_LZ = 0, so the scattering is kinematically closed.

Scan and figure, tc-45/46:

- scan_mass_splitting over 1000-1200 GeV x 0-700 keV (225 points): 210 points pass the tool's flag; 15 predict 0.3-5 LZ events, apparently one splitting row near 350 keV.
- Figure: /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png. The reported best fit is m = 1000 GeV, delta = 350 keV, which sits on the scan's mass edge.
- The scan tools have no output_dir argument, so the files went to the server's own directory, not the run directory.

Tool limitations (findings f-015, f-016, f-018; question q-13):

- The overall 'all pass' verdict ignores the LZ result. It reports VIABLE even at delta = 200 keV with N_LZ = 51158.

- The relic flag passes even at 12.9 sigma tension.
- Per describe_tools (tc-20), the LZ check only compares expected counts with the single 248 keV event in a 2.84 t-yr exposure. The tool has no published LZ cross-section limit curve and no XENONnT data, so the LZ limit at 1.1 TeV has to come from d-07's literature reading.

Finding: At 1.1 TeV the LZ 4.2 t-yr (arXiv:2410.17036, Fig. 5, from HEPData record 155182 table t1) observed 90% CL SI limit is about 3.3e-47 cm^2 and the median sensitivity is about 7.5e-47 cm^2. These were scaled linearly from the 1008 GeV HEPData values (observed 3.056e-47, median 6.834e-47 cm^2; above 1 TeV the limit scales as m^0.985). XENONnT 3.1 t-yr (arXiv:2502.18005) states in its text that above 200 GeV the limit is 3.7e-46 cm^2 x (M/1 TeV), which gives about 4.1e-46 cm^2 at 1.1 TeV.

Finding: Change from first runs to current results at 1.1 TeV: XENONnT went from 6.08e-47 x (M/100 GeV) cm^2 (arXiv:2303.14729, about 6.7e-46 cm^2 at 1.1 TeV, with a conservative median power constraint) to 3.7e-46 x (M/1 TeV) cm^2 (arXiv:2502.18005, about 4.1e-46 cm^2), an improvement of about 1.6x. For LZ, the text gives only the minimum limits, 9.2e-48 cm^2 at 36 GeV for the first run (2207.03764) and 2.2e-48 cm^2 at 40 GeV for 4.2 t-yr (2410.17036), an improvement of about 4.2x at the minimum. The first-run LZ value at 1 TeV was not obtained.

Finding: The pure-higgsino loop-level SI cross section is only bounded, not predicted: sigma_SI^D <~ 1e-48 cm^2 at 95% CL in the heavy-WIMP limit, independent of M (Hill and Solon, arXiv:1309.4092 Eq. 5 and arXiv:1409.8290 Eq. 122). Strong spin-0/spin-2 destructive interference means no central value is given. Chen and Hill (arXiv:1912.07795) find the upper limit 'less than 10^-48 cm^2 in the TeV mass regime', below the xenon neutrino floor, and the 1/M corrections do not enhance it. The NLO MSSM study (arXiv:2607.20588) finds that cancellations can push it below 1e-50 cm^2. For comparison, the pure wino is 8(+6/-3)(+3/-3)e-48 cm^2.

Finding: At 1.1 TeV the pure thermal higgsino (sigma_SI <~ 1e-48 cm^2) lies at least about 33x below the LZ 4.2 t-yr observed limit (about 3.3e-47 cm^2), at least about 75x below the LZ median sensitivity, and at least about 400x below the XENONnT 3.1 t-yr limit (about 4.1e-46 cm^2). It is also at least about 5.5x below the xenon SI neutrino floor (O'Hare 2109.03116 data file Xe_SI.txt: 5.0e-48 cm^2 at 1 TeV, about 5.5e-48 cm^2 at 1.1 TeV by linear scaling), so it sits inside the neutrino fog. The LZ limit itself is only about 6x above the floor at 1 TeV.

Finding: Tool check: predict_recoil at m_chi = 1100 GeV, delta = 0 (higgsino Z couplings, auto sigma_n = 7.43e-39 cm^2) predicts about 1.10e7 events in the LZ extended NR search against 1 observed. Elastic Z-mediated (Dirac-like) higgsino scattering is therefore excluded by about 7 orders of magnitude, and a neutral splitting is required to switch it off kinematically.

Completed — Place the thermal pure higgsino (m = 1.1-1.16 TeV) on the elastic spin-independent plane. Read the LZ 4.2 t-yr (2410.17036), XENONnT 3.1 t-yr (2502.18005) and first-run (2207.03764, 2303.14729) limits at 1.0-1.2 TeV, the loop-level higgsino cross section (1309.4092, 2607.20588) and the neutrino fog (2410.17137). Then quote the ratio of the higgsino cross section to the limit and to the fog.. At 1.1 TeV the thermal pure higgsino sits at least about 33 times below the current best elastic spin-independent limit (LZ 4.2 t-yr) and at least about 5.5 times below the xenon neutrino floor, so it is inside the fog. Elastic scattering through the Z is ruled out by about 7 orders of magnitude, so a neutral mass splitting is required. Ratios are lower bounds because the higgsino cross section is itself an upper bound.

(1) Current limits at 1.1 TeV (f-019). LZ 4.2 t-yr (arXiv:2410.17036): the paper text gives no 1 TeV value. HEPData record 155182, table t1 (the data behind the paper's Fig. 5) gives an observed 90% CL limit of 3.056e-47 cm^2 and a median sensitivity of 6.834e-47 cm^2 at 1008 GeV, scaling as m^0.985. Scaled linearly, that is about 3.3e-47 cm^2 (observed) and about

7.5e-47 cm^2 (median) at 1.1 TeV. XENONnT 3.1 t-yr (arXiv:2502.18005): the text states the limit is 3.7e-46 cm^2 x (M/1 TeV) above 200 GeV, about 4.1e-46 cm^2 at 1.1 TeV.

(2) First runs versus now (f-020). XENONnT first run (arXiv:2303.14729): 6.08e-47 cm^2 x (M/100 GeV), about 6.7e-46 cm^2 at 1.1 TeV. That run used a more conservative median power constraint. The improvement to the 3.1 t-yr result is about 1.6x. LZ: the minimum limit went from 9.2e-48 cm^2 at 36 GeV (arXiv:2207.03764) to 2.2e-48 cm^2 at 40 GeV, about 4.2x. The first-run LZ value at 1 TeV was not obtained.

(3) Higgsino cross section (f-021). The literature gives only an upper bound, sigma_SI <~ 1e-48 cm^2 at 95% CL, independent of mass in the heavy-WIMP limit (Hill and Solon, arXiv:1309.4092 Eq. 5 and arXiv:1409.8290 Eq. 122). No central value is quoted because of strong spin-0/spin-2 destructive interference. Chen and Hill (arXiv:1912.07795) state it is below 1e-48 cm^2 at TeV masses, below the neutrino floor. An NLO MSSM study (arXiv:2607.20588) finds regions where cancellations take it below 1e-50 cm^2. For comparison, the pure wino is 8(+6/-3)(+3/-3)e-48 cm^2.

(4) Ratios at 1.1 TeV (f-022). The higgsino lies at least 33x below the LZ observed limit, at least 75x below the LZ median sensitivity and at least 400x below the XENONnT limit. The xenon SI floor is 5.0e-48 cm^2 at 1 TeV, about 5.5e-48 cm^2 at 1.1 TeV (O'Hare, arXiv:2109.03116, public data file Xe_SI.txt). This is the same fog shown in the XLZD design book (arXiv:2410.17137) Fig. 2. The higgsino is at least 5.5x below that floor, and the LZ limit is only about 6x above it.

(5) Tool check (f-023). predict_recoil at m_chi = 1100 GeV, delta = 0 (sigma_n = 7.43e-39 cm^2) predicts 1.1e7 events in the LZ extended nuclear-recoil search against 1 observed.

Caveats: get_paper_figures returned only non-limit figures for both experiments (LZ Figs. 1 and S4, XENONnT Figs. 2 and S1), so nothing was read off a figure; values come from HEPData and the scaling formulas in the papers (dead end x-03). The HEPData units have not been checked against Fig. 5. cite() on 2410.17036 failed its quote check (dead end x-04), so provenance is given as arXiv IDs, HEPData record 155182 and the arXiv HTML pages; references.bib was not written.

The critic's model failed (critic model:); this step continues without a review.

Critic, wave 1 — continue: 9 findings stand up to the evidence.

Next steps after wave 1 — This is the last wave, with 20 tool calls. The critic found no contested results. The elastic-plane comparison (d-07), citations (d-08) and tool constraint deliverable (d-10) are done and stay closed. d-09 is still open and runs first, because it supplies the remaining required pieces. Part (a) checks how robust the tool's relic crossing is (about 1160 GeV against the literature's 1.08-1.2 TeV), via the effective sigma_v needed for Omega h^2 = 0.120 and how far the mass band shifts across the Planck plus or minus 0.001 uncertainty. Part (b) links the splitting window to the gaugino scale. For part (b), d-09 must write out the tree-level neutral splitting formula explicitly, including its M1, M2 and tan beta dependence, because critique c-05 flagged an order-of-magnitude error in the earlier gaugino-scale estimate. I am adding one small robustness direction: how the tool's preferred inelastic splitting changes with halo escape velocity and v0 (q-04), checked against the literature value of about 377 keV. No facility run is needed because the task does not ask for HPC compute. No literature direction is added, since the cross-checks against LZ, XENONnT, the neutrino fog and the thermal-mass papers are already done and cited.

New lines of investigation: Robustness of the tool's inelastic LZ prediction near the preferred splitting. At m = 1100 and 1160 GeV, use predict_recoil (or predict_annual_modulation if it exposes halo inputs) to find how the expected LZ count and the delta at which N_LZ is about 1 move when v_esc is varied (about 500, 544 and 600 km/s) and v0 is varied (about 220 and 238 km/s). Compare the resulting window with the literature preferred delta (about 377 keV, 2609.02608) and the solar bound (above 566 keV, 2609.02775)..

Wave 2 — now exploring, in parallel: Robustness of the tool relic calculation, and the link between gaugino mixing and the splitting. (a) Use compute_relic_density with an explicit sigma_v_cm3_s to find the effective cross section needed for Omega h^2 = 0.120 at 1.1 TeV. Compare it with the tool's built-in higgsino sigma_v and with the Sommerfeld-enhanced literature values. (b) Translate the xenon-accessible splitting window (delta of about 345-385 keV) and the solar bound (566 keV) into gaugino masses M1 and M2, and estimate the mixing at which Higgs-exchange elastic scattering reaches the LZ limit.; Robustness of the tool's inelastic LZ prediction near the preferred splitting. At m = 1100 and 1160 GeV, use predict_recoil (or predict_annual_modulation if it exposes halo inputs) to find how the expected LZ count and the delta at which N_LZ is about 1 move when v_esc is varied (about 500, 544 and 600 km/s) and v0 is varied (about 220 and 238 km/s). Compare the resulting window with the literature preferred delta (about 377 keV, 2609.02608) and the solar bound (above 566 keV, 2609.02775)..

Finding: With its fixed halo (v_esc + v_E = 776 km/s), the tool's predict_recoil for a higgsino gives these LZ counts. At m = 1100 GeV: N_LZ = 57.7, 1.97, 0.49, 0.085, 0.0 and 0.0 at delta = 330, 350, 355, 360, 370 and 390 keV. At m = 1160 GeV: N_LZ = 2.52 and 0.136 at delta = 350 and 360 keV. Log-linear interpolation (my own estimate) puts N_LZ = 1 at delta of about 352 keV for 1100 GeV and about 353 keV for 1160 GeV. The window closes completely by 370 keV, which is below the literature value of about 377 keV (2609.02608).

Finding: This is an analytic estimate, not tool output. The kinematic endpoint delta_max of about mu_N v_max^2 / 2 (mu of about 110 GeV for 1.1 TeV on Xe) gives about 368 keV at v_max = 776 km/s, which matches the tool's closure between 360 and 370 keV. Because delta_max scales as v_max^2, setting v_esc = 500 or 600 km/s (v_max of about 732 or 832 km/s) would move the endpoint to about 327 or 423 keV, a shift of about -40 or +55 keV. The 377 keV literature value (2609.02608) therefore needs a v_max only about 1-2% higher than the tool's (v_esc of about 555 km/s or more). Every case stays far below the solar bound of 566 keV (2609.02775).

Blocked — Robustness of the tool's inelastic LZ prediction near the preferred splitting. At m = 1100 and 1160 GeV, use predict_recoil (or predict_annual_modulation if it exposes halo inputs) to find how the expected LZ count and the delta at which N_LZ is about 1 move when v_esc is varied (about 500, 544 and 600 km/s) and v0 is varied (about 220 and 238 km/s). Compare the resulting window with the literature preferred delta (about 377 keV, 2609.02608) and the solar bound (above 566 keV, 2609.02775).. (The inelastic-DM server hard-codes the halo (v_esc + v_E = 776 km/s), and none of its tools take v_esc or v0, so the requested halo variations cannot be run until the server exposes these parameters.) The halo dependence could not be tested with the tool, but the baseline window is established. None of the inelastic-DM tools accept halo inputs: describe_tools (tc-63), predict_recoil and predict_annual_modulation have no v_esc or v0 arguments, and the halo is fixed at v_esc + v_E = 776 km/s (dead end x-05). Baseline LZ counts from predict_recoil (finding f-024): at m = 1100 GeV, N_LZ = 57.7, 1.97, 0.49, 0.085 and 0 at delta = 330, 350, 355, 360 and 370 keV, and 0 at 390 keV. At m = 1160 GeV, N_LZ = 2.52 at 350 keV and 0.136 at 360 keV. Interpolating, N_LZ = 1 falls at delta of about 352 keV for 1100 GeV and about 353 keV for 1160 GeV. The window is about 15 keV wide and closes by 370 keV, below the literature value of about 377 keV (2609.02608). The tool marks the 248 keV event inaccessible at 370 keV (v_min = 787 km/s). Analytic estimate, not tool output (finding f-025): the endpoint scales as v_max^2, so v_esc = 500 or 600 km/s would shift it by about -40 or +55 keV (to about 327 or 423 keV). Reaching 377 keV needs v_esc of about 555 km/s or more, only 1-2% above the tool's halo. All halo choices stay well below the 566 keV solar bound (2609.02775). The tool's 350 keV result versus the literature's 377 keV is therefore within the range that halo uncertainty could plausibly cover, which answers q-04 in part. The v0 effect was not quantified. The annual modulation at 1100 GeV and 350 keV has a 98.5% fractional

amplitude, peaking on June 2 (tc-79). No figures were produced.

Finding: The tool's relic density with is_higgsino=false at m=1100 GeV is Omega h^2 = 0.1342, 0.1199 and 0.0909 for sigma_v = 2.0, 2.25 and 3.0e-26 cm^3/s, so Omega*sigma_v is about 0.27 (it scales almost exactly as 1/sigma_v). The cross section needed for Omega h^2 = 0.120 at 1.1 TeV is 2.25e-26 cm^3/s. The built-in higgsino value is 2.5e-26*(1100/m)^2, which gives 0.1083 at 1100 GeV and crosses 0.120 at 1159 GeV (sigma_v = 2.25e-26). With is_higgsino=true the explicit sigma_v_cm3_s argument is silently ignored. Since Omega scales roughly as m^2 under the built-in scaling, a 10% change in sigma_v moves the crossing mass by about 5%, roughly 55-60 GeV. To reproduce the literature masses of 1.08, 1.10 and 1.20 TeV, the built-in rate would need effective factors of 0.87, 0.90 and 1.07, respectively. This factor is 0.9*(m/1100)^2 and is relative to the tool's own normalization, not a physical Sommerfeld factor.

Finding: The tree-level neutral higgsino splitting is delta0 ~ m_Z^2 (s_W^2/M1 + c_W^2/M2) for M_i >> |mu|, with tan beta cancelling at O(1/M) (Nagata and Shirai, arXiv:1410.4549, Eqs. 15 and 34). For M1 = M2 = M (m_Z^2 = 8315 GeV^2), the required gaugino mass is M = 2.38e7 GeV at delta = 350 keV, 2.25e7 GeV at 370 keV, 2.16e7 GeV at 385 keV and 1.47e7 GeV at 566 keV. The xenon-accessible window of about 345-385 keV therefore needs gauginos at about 2.2-2.4 x 10^4 TeV, not 10-30 TeV. The solar bound (delta > 566 keV) requires M < 1.5 x 10^4 TeV. With wino exchange only (delta0 = m_W^2/M2) these values are about 22% lower. RG running of the dimension-5 operators (1410.4549, Eq. 35, Fig. 6) and on-shell one-loop mass corrections (2609.09830, Eqs. 19-20) change the splitting multiplicatively by tens of percent. The exact percentage was not extracted.

Finding: In the higgsino limit, the tree-level Higgs-exchange coupling is f_q ~ -(g^2/8m_h^2)[1/(M2-|mu|) + tan^2 theta_W/(M1-|mu|)](1 + sgn(mu) sin 2beta) (arXiv:2606.31013, Eqs. 30-33; arXiv:1410.4549, Eqs. 46-49). This gives roughly sigma_SI ~ 5e-46 cm^2 (1 TeV/M)^2 (1 + sgn mu sin2beta)^2 for M1 = M2 = M; my order-of-magnitude check agrees to within a factor of about 3. With the large-tan-beta factor of about 1, sigma_SI reaches the LZ 4.2 t-yr limit of about 3.3e-47 cm^2 at 1.1 TeV (f-019) for M of about 4 TeV, or about 0.7 TeV with the 1e-45 placeholder. At M = 4 TeV the splitting is m_Z^2/M, about 2 GeV, so inelastic scattering is closed. In the inelastic window (M ~ 2.3e4 TeV) sigma_SI is about 1e-54 cm^2, 7-8 orders below LZ and far below the neutrino floor. The Higgs-exchange elastic signal and the inelastic xenon signal are therefore mutually exclusive, which answers q-11.

Completed — Robustness of the tool relic calculation, and the link between gaugino mixing and the splitting. (a) Use compute_relic_density with an explicit sigma_v_cm3_s to find the effective cross section needed for Omega h^2 = 0.120 at 1.1 TeV. Compare it with the tool's built-in higgsino sigma_v and with the Sommerfeld-enhanced literature values. (b) Translate the xenon-accessible splitting window (delta of about 345-385 keV) and the solar bound (566 keV) into gaugino masses M1 and M2, and estimate the mixing at which Higgs-exchange elastic scattering reaches the LZ limit.. (a) Relic cross section (f-026). With is_higgsino=true the tool ignores the sigma_v_cm3_s argument: every call echoed 2.5e-26 x (1100/m)^2 cm^3/s whatever value was passed (tc-68 to tc-71). I therefore varied sigma_v with is_higgsino=false. At m = 1100 GeV, Omega h^2 = 0.1342, 0.1199 and 0.0909 for sigma_v = 2.0, 2.25 and 3.0e-26 cm^3/s (tc-72 to tc-74). Omega scales almost exactly as 1/sigma_v (Omega x sigma_v is about 0.27e-26), so Omega h^2 = 0.120 needs sigma_v = 2.25e-26 cm^3/s at 1.1 TeV. The built-in higgsino value of 2.5e-26 gives 0.1083 at 1100 GeV and reaches 0.120 at 1159 GeV. Under the built-in (1100/m)^2 scaling Omega goes as about m^2, so a 10% change in sigma_v moves the crossing mass by about 5% (about 55-60 GeV). To match the literature masses of 1.08, 1.10 and 1.20 TeV, the built-in rate would need effective factors of about 0.87, 0.90 and 1.07. These are relative to the tool's own normalization, not physical Sommerfeld or coannihilation factors. (b)

Gaugino scale for the splitting (f-027). The tree-level formula is delta0 ~ m_Z^2 (s_W^2/M1 + c_W^2/M2) (Nagata and Shirai, arXiv:1410.4549, Eqs. 15 and 34). For M1 = M2 = M it gives M = 2.38e7, 2.25e7, 2.16e7 and 1.47e7 GeV for delta = 350, 370, 385 and 566 keV. The xenon window therefore needs gauginos at about 2.2-2.4 x 10^4 TeV, not the 10-30 TeV suggested in the direction. The solar bound requires M below about 1.5 x 10^4 TeV. With wino exchange only (delta0 = m_W^2/M2) these values are about 22% lower. RG running of the operators and on-shell loop corrections (1410.4549 Eq. 35 and Fig. 6; 2609.09830 Eqs. 19-20) change these by tens of percent, but no exact percentage was extracted. (c) Higgs-exchange elastic scattering (f-028). In the higgsino limit sigma_SI is roughly 5e-46 cm^2 x (1 TeV/M)^2 x (1 + sgn(mu) sin 2beta)^2 (2606.31013 Eqs. 30-33; 1410.4549 Eqs. 46-49). This normalization is my estimate from those formulas, not a quoted value. It reaches the LZ limit of about 3.3e-47 cm^2 at 1.1 TeV at M of about 4 TeV, or about 0.7 TeV with the 1e-45 placeholder. At M of about 4 TeV the splitting is about 2 GeV, so inelastic scattering is closed. In the inelastic window, sigma_SI is about 1e-54 cm^2. The elastic Higgs-exchange signal and the inelastic signal therefore cannot both appear, which answers q-11. Caveats: the literature came through arxiv_agent reading the arXiv HTML pages; cite() was not run, no figures were extracted and no files were written.

Critic, wave 2 — continue: 4 findings stand up to the evidence, 1 questioned ('In the higgsino limit, the tree-level Higgs-exchange coupling is f_q ~ -(g^2/8m_h^2)[1/(M2': The only evidence (tc-83) states that the Higgs-exchange elastic cross section is 'around 1e-53 cm^2' at gaugino scales of 2-8 x 10^4 TeV. The claim says about 1e-54 cm^2 at 2.3 x 10^4 TeV, an order of magnitude lower. The key quantitative inputs are not in the quoted evidence: the normalization sigma_SI ~ 5e-46 cm^2 (1 TeV/M)^2 and the M of about 4 TeV at which LZ is reached. The claim itself calls the normalization the agent's own estimate, accurate to about a factor of 3. The qualitative conclusion, that elastic Higgs-exchange and inelastic signals are mutually exclusive, is robust by many orders of magnitude. The specific numbers, however, are unsupported by the cited excerpt. Separately, the 'M of about 0.7 TeV with the 1e-45 placeholder' case puts the gauginos below the 1.1 TeV higgsino. That breaks the higgsino-limit formula being used, so the case should be dropped.).

Issues: The 1-2% (v_esc about 555 km/s) requirement uses the overall kinematic endpoint, mu v_max^2/2 = 368 keV. That endpoint applies at the optimal recoil energy E_R = mu*delta/m_N, not at 248 keV. The relevant quantity is v_min at the 248 keV event, which is (m_N E_R/mu + delta)/sqrt(2 m_N E_R). This formula reproduces the tool's 787 km/s at 370 keV. At 377 keV it gives v_min of about 795 km/s, so v_max must be at least about 795 km/s, about 2.5% above 776. That corresponds to v_esc of about 563 km/s for v_E of about 232 km/s, an assumed value that is not stated. Reproducing about one event, rather than a merely nonzero rate, needs more margin still. The qualitative conclusion holds, but the threshold should be restated.; The v0 dependence is not quantified. Near the endpoint, the rate depends exponentially on the tail of the velocity distribution, so v0, v_esc truncation and the choice of SHM versus non-Maxwellian halo can shift the N_LZ = 1 splitting by several keV even at fixed v_max. Without an independent rate calculation, the statement that halo uncertainty covers 352 versus 377 keV is plausible but not demonstrated.; Defining the window by N_LZ about 1 is a heuristic. With 1 observed event, a proper statement would use the Poisson likelihood plus the event's energy-reconstruction uncertainty: an approximate band in delta of N_LZ between about 0.05 and 4.7, not a single crossing point. The window width of about 15 keV is likely an underestimate once the 248 keV energy resolution is included..

Stopping after wave 2: budget reached (2/2 waves used). Writing the report from the board.

Research report:

The thermal higgsino (1.0-1.2 TeV) relative to Planck, LZ and XENONnT: relic target, elastic limits and the inelastic window

Abstract

This study places a pure higgsino at its thermal-relic mass against current xenon direct-detection limits. It combines a literature survey with calculations from an inelastic-dark-matter tool server.

Relic mass. The cited literature puts the thermal mass between about 1.08 and 1.2 TeV, so a central value near 1.1 TeV is reasonable. The Planck 2018 value used is $\Omega_c\ h^2 = 0.120 \pm 0.001$. In the tool's own higgsino normalization, $\Omega\ h^2$ reaches 0.120 at about 1159-1160 GeV.

Elastic scattering. The pure higgsino's loop-level spin-independent (SI) cross section is only bounded, at σ_SI ? 1e-48 cm^2. At 1.1 TeV that is at least about 33 times below the LZ 4.2 tonne-year observed limit (about 3.3e-47 cm^2). It is at least about 400 times below the XENONnT 3.1 tonne-year limit (about 4.1e-46 cm^2). It is also at least about 5.5 times below the xenon neutrino floor, which places it inside the neutrino fog.

Inelastic scattering. Z-mediated elastic scattering at zero splitting is excluded by about 7 orders of magnitude. A neutral mass splitting is therefore required. With its fixed halo, the tool gives about one LZ event only in a narrow window near $\delta \approx 352$ keV, and the rate closes completely by 370 keV.

Gaugino scale. Tree-level splittings in that window need gauginos at roughly 2×10? TeV. At that scale, Higgs-exchange elastic scattering is negligible. On an estimate the reviewer contested, the inelastic and Higgs-exchange elastic signals appear mutually exclusive.

Background

Relic abundance. A pure higgsino is an electroweak doublet. Its thermal relic abundance includes coannihilation with the charged state and the second neutral state, and Sommerfeld enhancement of the annihilation rate. The relevant literature includes:

- a pMSSM study of heavy higgsino and wino relic abundance with Sommerfeld effects (arXiv:1411.6930);
- analytic and numerical treatments of Sommerfeld-enhanced freeze-out (arXiv:1008.2905, arXiv:0910.5221).

Cosmological reference. The abundance is taken from Planck 2018 (arXiv:1807.06209), whose abstract gives $\Omega_c\ h^2 = 0.120 \pm 0.001$. The planning review recommended quoting the more precise form 0.1200 ± 0.0012 (TT,TE,EE+lowE+lensing). The inelastic-DM tool uses 0.12 ± 0.0012.

Recent thermal-mass estimates (cited verbatim from their abstracts):

Source	Thermal mass	Notes
arXiv:2410.15733	about 1.2 TeV	Abstract states that coannihilation and Sommerfeld effects are included
arXiv:2609.02775 (Pospelov and Ramani)	about 1.08 TeV ("cosmologically preferred")	Also derives a solar-capture bound $\delta > 566$ keV
arXiv:2609.02608 (Di Mauro)	near 1.1 TeV	$\delta \approx 377$ keV reproduces the 248 keV LZ nuclear-recoil candidate

Source	Thermal mass	Notes
arXiv:2403.19598 (Martin)	about 100-1150 GeV	Range for higgsino-like thermal dark matter

The solar-capture bound in arXiv:2609.02775 rests on these assumptions, as stated in its abstract:

- solar-core speeds up to about 1400 km/s;
- inelastic scattering on heavy elements;
- annihilation to WW and ZZ;
- IceCube's non-observation of high-energy neutrinos from the Sun.

A model-building context in which a roughly 1 TeV higgsino lightest supersymmetric particle saturates the relic density is given in arXiv:2302.02905.

Direct detection: pure higgsino. At tree level a pure higgsino's elastic SI scattering is suppressed, and it reaches xenon only through electroweak loops. Hill and Solon's heavy-WIMP effective theory (arXiv:1309.4092, arXiv:1409.8290) gives an upper bound rather than a central value. Further results:

- Chen and Hill (arXiv:1912.07795) place it below 1e-48 cm^2 at TeV masses.
- An NLO MSSM study (arXiv:2607.20588) finds regions where cancellations take it below 1e-50 cm^2.

Direct detection: mixing and splittings. Gaugino mixing adds Higgs exchange. The neutral splitting scales inversely with the gaugino masses (Nagata and Shirai, arXiv:1410.4549). A 2026 paper (arXiv:2609.09830) computes loop-corrected splittings and the resulting elastic and inelastic LZ rates.

Current limits and floor. The current xenon limits are LZ 4.2 tonne-years (arXiv:2410.17036) and XENONnT 3.1 tonne-years (arXiv:2502.18005), preceded by the first runs (arXiv:2207.03764, arXiv:2303.14729). The xenon neutrino floor is taken from O'Hare (arXiv:2109.03116), consistent with the XLZD design book (arXiv:2410.17137).

Results

No figure in the registered list shows a computed result or a limit curve. The four extracted figures are LZ and XENONnT event-distribution and calibration plots, which add nothing to this analysis. The tool's scan figure was written to the server's output directory and was not registered.

Relic density from the tool, and the mass target With the higgsino option and $\delta = 350$ keV, the tool's constraint check gives these relic densities (tension computed against 0.12 ± 0.0012):

m (GeV)	$\Omega\ h^2$	Tension
1080	0.1045	12.9σ

m (GeV)	Ω h^2	Tension
1100	0.1083	9.8σ
1160	0.1202	0.1σ
1200	0.1285	7.1σ

The tool's built-in higgsino cross section is ?σv? = 2.5e-26 $\times$ (1100/m)^2 cm^3/s, and it crosses Ω h^2 = 0.120 at 1159 GeV.

With the higgsino option on, the tool ignored an explicitly supplied ?σv? and echoed its built-in value every time. The reviewer noted that this rests partly on the agent's account of what was passed, though the echoed value supports it. The cross section was therefore varied with the generic-WIMP option at m = 1100 GeV:

?σv? (cm^3/s)	Ω h^2
2.0e-26	0.1342
2.25e-26	0.1199
3.0e-26	0.0909

Ω scales almost exactly as 1/?σv? (Ω $\times$?σv? $\approx$ 0.27e-26), so Ω h^2 = 0.120 at 1.1 TeV needs ?σv? = 2.25e-26 cm^3/s. Under the built-in scaling Ω grows roughly as m^2. Consequences:

- A 10% change in ?σv? moves the crossing mass by about 5%, or 55-60 GeV.
- Reproducing the literature masses of 1.08, 1.10 and 1.20 TeV would need effective factors of about 0.87, 0.90 and 1.07 on the tool's rate. These are relative to the tool's own normalization and are not physical Sommerfeld or coannihilation factors.
- The planning review estimated that the Planck uncertainty shifts the mass by only about $\pm$6 GeV. That is negligible next to the 1.08-1.2 TeV spread in the literature.

Only arXiv:2410.15733 states its treatment (Sommerfeld plus coannihilation) in its abstract. The cause of the 1.08 versus 1.2 TeV spread was therefore not established. The 1.08-1.2 TeV band with a central value near 1.1 TeV is an inference from the spread of cited values, not a published result.

Current elastic SI limits at 1.1 TeV LZ 4.2 tonne-years. The paper's text gives no value at 1 TeV. The HEPData record 155182, table t1 (the data behind its Fig. 5), gives at 1008 GeV:

- an observed 90% CL limit of 3.056e-47 cm^2;
- a median sensitivity of 6.834e-47 cm^2;
- a scaling of roughly m^0.985 above 1 TeV.

Scaled to 1.1 TeV, these become about 3.3e-47 cm^2 (observed) and about 7.5e-47 cm^2 (median). The HEPData units were not checked against the figure.

XENONnT 3.1 tonne-years. The text states 3.7e-46 cm^2 $\times$ (M/1 TeV) above 200 GeV, which gives about 4.1e-46 cm^2 at 1.1 TeV.

Time evolution.

- XENONnT: the first run gave 6.08e-47 cm^2 × (M/100 GeV), about 6.7e-46 cm^2 at 1.1 TeV, under a more conservative median power constraint. The improvement to the current result is about 1.6 times.
- LZ: the minimum limit improved from 9.2e-48 cm^2 at 36 GeV to 2.2e-48 cm^2 at 40 GeV, about 4.2 times. The first-run LZ value at 1 TeV was not obtained.

Higgsino cross section relative to limits and to the neutrino fog The pure higgsino is bounded at σ_SI ? 1e-48 cm^2 at 95% CL. This bound is independent of mass in the heavy-WIMP limit (arXiv:1309.4092 Eq. 5; arXiv:1409.8290 Eq. 122). No central value exists because of strong destructive interference between the spin-0 and spin-2 amplitudes. For comparison, the pure wino is 8(+6/-3)(+3/-3)e-48 cm^2.

Taking the xenon SI floor as 5.0e-48 cm^2 at 1 TeV (about 5.5e-48 cm^2 at 1.1 TeV, from the O'Hare data file Xe_SI.txt), the higgsino lies at 1.1 TeV:

- at least about 33 times below the LZ observed limit;
- at least about 75 times below the LZ median sensitivity;
- at least about 400 times below the XENONnT limit;
- at least about 5.5 times below the neutrino floor.

The LZ limit itself is only about 6 times above the floor at 1 TeV. All of these ratios are lower bounds, because the higgsino cross section is itself an upper bound.

Elastic Z exchange and the inelastic window (tool) At $\delta = 0$ the tool uses the higgsino Z coupling (σ_n = 7.43e-39 cm^2) and predicts about 1.1e7 events in the LZ extended nuclear-recoil search, against 1 observed. The same point also fails indirect detection:

- ?σv? today is 2.5e-24 cm^3/s, against a Fermi-LAT limit of 2.3e-25 cm^3/s;
- p_ann is 3.4e-28, against a CMB limit of 3.2e-28.

Expected LZ counts at m = 1100 GeV from the recoil predictor, which uses a fixed halo with v_esc + v_E = 776 km/s:

δ (keV)	N_LZ
200	5.1e4
330	57.7
350	1.97
355	0.49
360	0.085
370	0
390	0
566	0

At 370 keV the tool marks the 248 keV event as inaccessible, because v_min = 787 km/s exceeds 776 km/s. At m = 1160 GeV, N_LZ = 2.52 at 350 keV and 0.136 at 360 keV.

Log-linear interpolation puts N_LZ = 1 at $\delta \approx 352$ keV for 1100 GeV and about 353 keV for

1160 GeV. The window is roughly 15 keV wide and sits below the literature's 377 keV.

At $\delta = 350$ keV the four masses give N_LZ = 1.79, 1.97, 2.52 and 2.90, all compatible with the single observed event. At those points the present-day annihilation rate is about 5e-42 cm^3/s, so Fermi-LAT and CMB checks pass. The annual modulation at 1100 GeV and 350 keV has a 98.5% fractional amplitude and peaks on June 2.

A 15 × 15 scan over 1000-1200 GeV and 0-700 keV finds 210 of 225 points passing the tool's overall flag, which ignores direct detection. Fifteen points predict 0.3-5 LZ events, apparently one splitting row near 350 keV. The reported best fit (1000 GeV, 350 keV) sits on the mass edge of the scan. The scan file and figure are at /Users/nramachandra/github-integration/inelastic-dm-server/output/.

Halo sensitivity of the window (analytic estimate, not tool output) The kinematic endpoint is δ_max ≈ μ_N v_max^2/2, with $\mu \approx 110$ GeV for 1.1 TeV on xenon. This gives about 368 keV at 776 km/s, consistent with the tool's closure between 360 and 370 keV. Because the endpoint scales as v_max^2, setting v_esc to 500 or 600 km/s would move it to about 327 or 423 keV.

The reviewer pointed out that the relevant quantity is v_min at the 248 keV event, not the overall endpoint. On that basis, 377 keV needs v_max ? 795 km/s, about 2.5% above the tool's value. That corresponds to v_esc ≈ 563 km/s for an assumed v_E ≈ 232 km/s, rather than the 1-2% and 555 km/s first estimated. Producing about one event, rather than a merely nonzero rate, needs more margin still. The v0 dependence was not quantified.

The conclusion that halo uncertainty can plausibly bridge 352 versus 377 keV is therefore plausible but not demonstrated. Every halo choice considered stays well below the 566 keV solar bound.

Gaugino scale implied by the splitting The tree-level relation is $\delta_0 \approx$ m_Z^2(s_W^2/M1 + c_W^2/M2) (arXiv:1410.4549 Eqs. 15 and 34). Setting M1 = M2 = M gives:

δ (keV)	Required M (GeV)
350	2.38e7
370	2.25e7
385	2.16e7
566	1.47e7

The xenon window therefore needs gauginos at about 2.2-2.4 × 10? TeV, about three orders of magnitude above the 10-30 TeV initially assumed in planning. The solar bound requires M below about 1.5 × 10? TeV. With wino exchange only, these values are about 22% lower. Renormalization-group running and on-shell loop corrections (arXiv:1410.4549 Eq. 35; arXiv:2609.09830 Eqs. 19-20) change them by tens of percent; the exact amount was not extracted.

Higgs-exchange elastic scattering versus the inelastic window Warning: the reviewer contested this result; it relies on an unverified normalization.

From the higgsino-limit coupling (arXiv:2606.31013 Eqs. 30-33; arXiv:1410.4549 Eqs. 46-49), the estimate is σ_SI ≈ 5e-46 cm^2 × (1 TeV/M)^2 × (1 + sgn μ sin 2β)^2. This normalization is an estimate made in this study, not a quoted value. On it:

- σ_SI reaches the LZ limit of about 3.3e-47 cm^2 at M ≈ 4 TeV. At that scale the splitting

is about 2 GeV, so inelastic scattering is closed.

- In the inelastic window (M $\approx$ 2.3 $\times$ 10? TeV), σ_SI is about 1e-54 cm^2, 7-8 orders below LZ.

The reviewer raised two objections:

- Keeping 1/(M - $|\mu|$) rather than 1/M changes σ_SI by about a factor of 2 at M = 4 TeV.
- The tan β factor ranges from 0 to 4, which moves the LZ-reach scale anywhere from 0 to about 8 TeV.

The normalization has not been checked against a published σ_SI-versus-M curve. The qualitative conclusion is robust to these factors, because the gap is many orders of magnitude: the two signals cannot both appear.

Tool reliability

- The tool's overall pass verdict ignores the LZ result. It reports a viable point at δ = 200 keV with N_LZ = 51158.
- The relic flag reads "consistent with Planck" even at 12.9σ and 24.2σ tension.
- The LZ check only compares expected counts with the single 248 keV event in 2.84 tonne-years.
- There is no published LZ σ_SI limit curve and no XENONnT data, so the tool's LZ treatment could not be cross-checked against arXiv:2410.17036.

None of the tool's pass flags should be read as physical viability.

Discussion

The answer to the task is as follows. A thermal pure higgsino at about 1.1 TeV is out of reach of elastic SI searches. It sits at least an order of magnitude below the best LZ limit and inside the xenon neutrino fog. Elastic scattering will not become accessible to xenon detectors simply through more exposure. This agrees with the expectation in the survey brief and with Chen and Hill (arXiv:1912.07795).

Xenon detectors can see a higgsino in two ways:

- **Inelastic scattering**, if the neutral splitting falls within a narrow window just below the kinematic endpoint. The window is about 350-370 keV in the tool's halo, and the literature prefers about 377 keV.
- **Higgs exchange**, if gauginos are light, around a few TeV. This option is contested and normalization-dependent.

The splitting relation makes these two routes mutually exclusive.

The tool's inelastic window sits about 25 keV below the 377 keV of arXiv:2609.02608. The analytic estimate suggests that a few-percent increase in the halo's maximum speed could account for this. The reviewer's refined estimate puts the required v_max about 2.5% above the tool's value, and v0 and the shape of the distribution's tail could shift the window by several keV.

The tension is therefore plausibly a halo-modelling difference, but this was not shown with a calculation.

The brief's assumed 100-200 keV cutoff for inelastic scattering is inconsistent with the computed endpoint near 370-400 keV. In the tool, $\delta = 200$ keV gives about 5e4 events and is strongly excluded. The tool's 1160 GeV relic crossing lies within the literature band, but toward its upper end. Because the tool's relic calculation is a simple freeze-out with a fixed normalization, its crossing mass should be treated as model-dependent, not as a check of the Sommerfeld literature.

Several approaches did not work:

- **Reading the limit curves from figures.** Figure extraction returned only two non-limit figures per experiment, so the limits came from HEPData and from scaling formulas in the papers' text.
- **Citing the LZ 4.2 tonne-year paper.** The quote check failed, so its provenance rests on the arXiv ID and HEPData record 155182.
- **Testing halo variations with the tool.** The server exposes no halo parameters, so this direction was blocked.

Limitations

- **No independent relic calculation.** No calculation with Sommerfeld effects or coannihilation was run. The tool's internal physics, such as whether it implements chargino coannihilation, is not documented, and its explicit ?σv? input is ignored in higgsino mode.
- **Unresolved mass spread.** The 1.08-1.2 TeV spread could not be explained, because the full texts of the mass papers were not read.
- **Indirect LZ limit.** The LZ limit at 1.1 TeV comes from a HEPData point at 1008 GeV with linear scaling, and its units were not cross-checked against the pa