Supplementary material: The 1.0–1.2 TeV Thermal Higgsino Against Planck, LZ and XENONnT: Toy-Model Relic and Inelastic-Recoil Calculations with a Literature Survey

HEP-Genesis-Agent session record
Project: ramp-combo-higgsino-dd

2026-10-22

Supplementary material, generated mechanically from the session record: it lists exactly what ran, in order, without interpretation.

A Implementation details

This appendix is generated mechanically from the session record; it lists exactly what ran, in order, without interpretation.

A.1 Session

- Source: cli client; conversation `b128dd5f`
- Started: 2026-10-16 14:31:17; report generated: 2026-10-22 14:54:44
- Project: ramp-combo-higgsino-dd (`/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd`)
- Compute execution site(s) selected in the session: local
- 2 user turn(s), 175 tool call(s) (0 returned errors), 4 figure(s), 0 data file(s)

A.2 Models

Stage	Model	Endpoint
quick	`claude-opus-5-5`	`https://mag.genesis.americansciencecloud.org/v1`
report	`claude-opus-5-5`	`https://mag.genesis.americansciencecloud.org/v1`
coder	`claude-opus-5-5`	`https://mag.genesis.americansciencecloud.org/v1`

A.3 MCP servers

Server	Transport	Location
hep-research-aid	stdio	`python -m hep_genesis.domain.mcp_server`
alcf-tools	stdio	`python -m hep_genesis.iri.alcf.mcp_server`
nersc-tools	stdio	`python -m hep_genesis.iri.nersc.mcp_server`

A.4 Tool-call log

Chronological; arguments and results are truncated for space (the full record is in the project's `provenance/` folder).

#	Tool	Arguments	Outcome
1	`serve-sbbn.sh list_sterile_nu_skills`	`{}`	`{ "status": "ok", "files": [], "message": "2 skills. Load one with load_sterile_nu_skill.", "metadata": { "skills": { "kev-sterile-dm-xray": "keV sterile-neutrino dark matter \u2014 ...`
2	`serve-hydro.sh list_observables`	`{}`	`{ "status": "success", "message": "14 observables available: GSMF, HMF, fGas, Pk-ratio, CSFR, CGD, CGED, CPP, CTP, CEP, CEEP, CMP, CYP, Pk_GO.", "files": [], "metadata": { "observables": {...`
3	`hep-research-aid list_agent_files`	`{}`	`[]`
4	`hep-research-aid list_input_files`	`{}`	`{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }`
5	`hep-research-aid list_datasets`	`{}`	`{'count': 0, 'datasets': []}`
6	`hep-research-aid list_files`	`{}`	`Files in .: - .git - .gitignore - ISSUES.md - SCOPE.md - docs - downloads - logs - projects - reports - research-results - runs - scripts - venvs - workspace`
7	`hep-research-aid get_dispatch`	`{}`	`Dispatch: local execution.`

#	Tool	Arguments	Outcome
8	local search_arxiv_abstracts	{"query": "higgsino dark matter thermal relic 1.1 TeV direct detection", "max_results": 8}	Search failed: Page request resulted in HTTP 503 (https://export.arxiv.org/api/query?search_query=higgsino+dark+matter+thermal+relic+1 list=&sortBy=relevance&sortOrder=descen...
9	local search_arxiv_abstracts	{"query": "pure higgsino inelastic scattering mass splitting xenon detectors", "max_results": 8}	Search failed: Page request resulted in HTTP 429 (https://export.arxiv.org/api/query?search_query=pure+higgsino+inelastic+scattering+ma list=&sortBy=relevance&sortOrder=...
10	local search_arxiv_abstracts	{"query": "higgsino Sommerfeld enhancement relic abundance electroweak", "max_results": 6}	Search failed: Page request resulted in HTTP 429 (https://export.arxiv.org/api/query?search_query=higgsino+Sommerfeld+enhancement+reli list=&sortBy=relevance&sortOrder=descen...
11	hep-research-aid search_arxiv	{"query": "higgsino dark matter thermal relic direct detection", "max_results": 10}	Found 10 papers matching query: 'higgsino dark matter thermal relic direct detection' (source: OpenAlex) Paper 1: arXiv ID: 1611.00804 Title: The last refuge of mixed wino-Higgsino dark matter Author...
12	local web_search	{"query": "higgsino 1.1 TeV thermal relic inelastic dark matter LZ XENONnT arXiv"}	Sep 1, 2026 · The Higgsino dark matter of the minimal supersymmetric standard model is one of the most minimal and canonical WIMP candidates. Sep 2, 2026 · This Supplemental Material details the compu...

#	Tool	Arguments	Outcome
13	hep-research-aid search_arxiv	{"max_results": 10, "query": "inelastic higgsino LZ nuclear recoil excess mass splitting keV"}	Found 10 papers matching query: 'inelastic higgsino LZ nuclear recoil excess mass splitting keV' (source: OpenAlex) Paper 1: arXiv ID: 2609.09136 Title: A Warped Extra Dimensional Candidate for the L...
14	hep-research-aid search_arxiv	{"max_results": 8, "query": "wino higgsino thermal mass Sommerfeld bound state relic 1.1 TeV"}	Found 8 papers matching query: 'wino higgsino thermal mass Sommerfeld bound state relic 1.1 TeV' (source: OpenAlex) Paper 1: arXiv ID: 1010.2172 Title: Relic densities including Sommerfeld enhancemen...
15	hep-research-aid search_arxiv	{"max_results": 8, "query": "LZ high-energy nuclear recoil search inelastic dark matter 2026"}	Found 8 papers matching query: 'LZ high-energy nuclear recoil search inelastic dark matter 2026' (source: OpenAlex) Paper 1: arXiv ID: 2609.10491 Title: Inelastic Dark Matter and High-Energy Recoil S...
16	hep-research-aid search_arxiv	{"max_results": 6, "query": "inelastic higgsino solar capture IceCube neutrino bound"}	Found 6 papers matching query: 'inelastic higgsino solar capture IceCube neutrino bound' (source: OpenAlex) Paper 1: arXiv ID: 2609.02775 Title: Strong Constraints on Higgsino Dark Matter from Solar ...
17	hep-research-aid search_arxiv	{"max_results": 5, "query": "higgsino nucleon elastic cross section loop neutrino fog"}	Found 5 papers matching query: 'higgsino nucleon elastic cross section loop neutrino fog' (source: OpenAlex) Paper 1: arXiv ID: 2609.01504 Title: Higgsino Above the Sea of Fog Authors: JiJi Fan, Matt...

#	Tool	Arguments	Outcome
18	hep-research-aid search_arxiv	{"max_results": 5, "query": "LZ collaboration search high energy nuclear recoils 2.84 tonne-year"}	Found 5 papers matching query: 'LZ collaboration search high energy nuclear recoils 2.84 tonne-year' (source: OpenAlex) Paper 1: arXiv ID: 2609.02608 Title: Dark Matter at the Kinematic Edge: Interpr...
19	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
20	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
21	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 300}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 300.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 270.0, "v_min_at_p...
22	serve-idm.sh compute_relic_density	{"m_chi_GeV": 900}	{ "parameters": { "m_chi_GeV": 900.0, "is_higgsino": true, "sigma_v_cm3_s": 3.7345679012345686e-26 }, "relic_density": { "omega_h2": 0.0731, "planck_value": 0.12, "planck...

#	Tool	Arguments	Outcome
23	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 380}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 380.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 342.1, "v_min_at_p...
24	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000}	{ "parameters": { "m_chi_GeV": 1000.0, "is_higgsino": true, "sigma_v_cm3_s": 3.0250000000000006e-26 }, "relic_density": { "omega_h2": 0.0898, "planck_value": 0.12, "planc...
25	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 500}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 500.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 450.1, "v_min_at_p...
26	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
27	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 566}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 566.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 509.5, "v_min_at_p...

#	Tool	Arguments	Outcome
28	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200}	{ "parameters": { "m_chi_GeV": 1200.0, "is_higgsino": true, "sigma_v_cm3_s": 2.1006944444444443e-26 }, "relic_density": { "omega_h2": 0.1285, "planck_value": 0.12, "planc...
29	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 700}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 700.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 630.1, "v_min_at_p...
30	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1300}	{ "parameters": { "m_chi_GeV": 1300.0, "is_higgsino": true, "sigma_v_cm3_s": 1.789940828402367e-26 }, "relic_density": { "omega_h2": 0.1503, "planck_value": 0.12, "planck...
31	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1100, "delta_keV": 400}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 400.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_si...
32	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1159, "delta_keV": 400}	{ "parameters": { "m_chi_GeV": 1159.0, "delta_keV": 400.0, "sigma_n_cm2": 7.430576349889478e-39, "is_higgsino": true }, "relic_density": { "omega_h2": 0.12, "tension_sigm...

#	Tool	Arguments	Outcome
33	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1100, "delta_ keV": 566}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 566.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_si...
34	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1100, "delta_ keV": 1000}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 1000.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_s...
35	serve-idm.sh predict_recoil	{"delta_keV": 380, "m_chi_ GeV": 1000}	{ "parameters": { "m_chi_GeV": 1000.0, "delta_keV": 380.0, "sigma_n_cm2": 7.428662722453626e-39, "is_higgsino": true }, "kinematics": { "E_R_ peak_keV": 338.7, "v_min_ at_p...
36	serve-idm.sh predict_recoil	{"delta_keV": 380, "m_chi_ GeV": 1200}	{ "parameters": { "m_chi_GeV": 1200.0, "delta_keV": 380.0, "sigma_n_cm2": 7.430987655656208e-39, "is_higgsino": true }, "kinematics": { "E_R_ peak_keV": 344.9, "v_min_ at_p...
37	serve-idm.sh predict_recoil	{"delta_keV": 350, "m_chi_ GeV": 1100}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_ peak_keV": 315.1, "v_min_ at_p...

#	Tool	Arguments	Outcome
38	serve-idm.sh scan_mass_splitting	{"m_chi_min": 800, "m_chi_max": 1400, "delta_min": 0, "delta_max": 800, "n_points": 17}	{ "scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz", "grid": "17 x 17 = 289 points", "ranges": { "m_chi_GeV": [800.0, 1400.0], ...
39	serve-idm.sh plot_recoil_spectrum	{"m_chi_GeV": 1100, "delta_keV": 380}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png", "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 380.0 } }
40	serve-idm.sh predict_annual_modulation	{"m_chi_GeV": 1100, "delta_keV": 350, "E_R_keV": 248}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "E_R_keV": 248.0 }, "modulation": { "peak_day": 153, "peak_date": "June 02", "trough_day": 336, "trough_dat...
41	board post_finding	{"claim": "The inelastic-DM server's compute_relic_density (is_higgsino=true) gives Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285, 0.1503 at m_chi = 900, 1000, 1100, 1200, 1300 GeV. Linear interpolation crosses Planck 0.120...	Recorded finding f-001.
42	board post_finding	{"claim": "check_all_constraints (is_higgsino) at m=1100 GeV with delta = 400, 566 and 1000 keV, and at m=1159 GeV with delta=400 keV, returns VIABLE every time. Fermi-LAT and CMB are not excluded, sigma_n = 7.43e-39 cm^...	Recorded finding f-002.

#	Tool	Arguments	Outcome
43	board post_question	{"text": "Why does check_all_constraints return N_expected_LZ = 0.00 for a 1.1 TeV higgsino at delta = 400 keV (sigma_n = 7.4e-39 cm^2) when the literature finds about 1 event at 340-380 keV? Is the velocity cutoff or en...	Recorded question q-06.
44	board post_finding	{"claim": "With the inelastic-DM server (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, v_esc+v_E = 776 km/s), the expected LZ count at m_chi = 1100 GeV is 744.6 at delta = 300 keV, 1.97 a...	Recorded finding f-003.
45	board post_finding	{"claim": "Xenon's kinematic reach at the thermal higgsino mass is delta_max = mu v_max^2/2. With m_Xe of about 122 GeV, m_chi = 1100 GeV gives mu = 109.8 GeV. For v_max = 776 km/s (the tool's value, beta^2 = 6.70e-6) th...	Recorded finding f-004.
46	board post_finding	{"claim": "The scan over m_chi = 800-1400 GeV and delta = 0-800 keV (17x17 grid) finds 144 of 289 points passing all constraints and 9 points that predict 0.3-5 LZ events. The 1100 GeV, 380 keV recoil-spectrum plot is at...	Recorded finding f-005.
47	serve-idm.sh plot_constraint_summary	{"scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz"}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png", "best_fit": { "m_chi_GeV": 988.0, "delta_keV": 350.0 } }

#	Tool	Arguments	Outcome
48	board post_finding	{"claim": "plot_constraint_ summary on the 17x17 scan (m_chi = 800-1400 GeV, delta = 0-800 keV) wrote /Users/nramachandra/ github-integration/ inelastic-dm-server/output/ constraint_summary.png and reports the best-fit point...	Recorded finding f-006.
49	local get_arxiv_ metadata	{"arxiv_id": "1807.06209"}	Paper 1807.06209 not found
50	local get_arxiv_ metadata	{"arxiv_id": "2410.15733"}	Paper 2410.15733 not found
51	local download_arxiv_ full_text	{"arxiv_id": "1807.06209", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261016_143117"}	Download failed: Paper 1807.06209 not found
52	local download_arxiv_ full_text	{"arxiv_id": "2410.15733", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261016_143117"}	Download failed: Paper 2410.15733 not found
53	hep-research-aid download_full_ arxiv_paper	{"paper_id": "1807.06209", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261016_143117"}	Error: Error downloading/extracting paper: 'Search' object has no attribute 'results'
54	hep-research-aid search_arxiv	{"query": "higgsino dark matter thermal relic Sommerfeld 1.1 TeV", "max_ results": 5}	Found 5 papers matching query: 'higgsino dark matter thermal relic Sommerfeld 1.1 TeV' (source: OpenAlex) Paper 1: arXiv ID: 1404.0682 Title: Neutralino dark matter at 14 TeV and 100 TeV Authors: Mat...

#	Tool	Arguments	Outcome
55	local cite	{"arxiv_id": "2410.15733", "note": "[\"in which the observed DM abundance is realized with the Higgsino mass of ~1.2 TeV\"]", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/2610...	METADATA FAILED: arxiv_id 2410.15733 not found
56	hep-research-aid search_arxiv	{"max_results": 5, "query": "Planck 2018 results VI cosmological parameters"}	Found 5 papers matching query: 'Planck 2018 results VI cosmological parameters' (source: OpenAlex) Paper 1: arXiv ID: 1907.12875 Title: Planck 2018 results Authors: N. Aghanim, Y. Akrami, M. Ashdown,...
57	hep-research-aid search_arxiv	{"max_results": 5, "query": "XENONnT first WIMP search nuclear recoil spin-independent limit"}	Found 5 papers matching query: 'XENONnT first WIMP search nuclear recoil spin-independent limit' (source: OpenAlex) Paper 1: arXiv ID: 2303.14729 Title: First Dark Matter Search with Nuclear Recoils ...
58	board post_finding	{"claim": "The pre-2026 Sommerfeld- and coannihilation-corrected thermal pure-higgsino mass is about 1.2 TeV. The abstract of arXiv:2410.15733 (Fukuda, Li, Moroi, Niki, JHEP06(2025)091, DOI 10.1007/jhep06(2025)091) says:...	Recorded finding f-007.
59	board post_finding	{"claim": "XENONnT's 3.1 tonne-year elastic WIMP search (arXiv:2502.18005, DOI 10.1103/msw4-t342) found no significant excess. Its spin-independent limit has a minimum of 1.7e-47 cm^2 (90% CL) at 30 GeV, with best median...	Recorded finding f-008.

#	Tool	Arguments	Outcome
60	board post_dead_end	{"what": "Verbatim citation of 1807.06209 (Planck 2018 VI) and 2410.15733 using get_arxiv_metadata, download_arxiv_full_text, download_full_arxiv_ paper and cite.", "why": "During this run, every arXiv-API-backed tool ret...	Recorded dead end x-01.
61	hep-research-aid search_arxiv	{"query": "Planck 2018 results cosmological parameters dark matter density 0.120 Hubble constant 67.4", "max_ results": 3}	Found 3 papers matching query: 'Planck 2018 results cosmological parameters dark matter density 0.120 Hubble constant 67.4' (source: OpenAlex) Paper 1: arXiv ID: 2403.14247 Title: Hint of Dark Matter...
62	board post_question	{"text": "Can a later run, once the arXiv API is back, retrieve and cite the verbatim Planck 2018 VI value Omega_c h^2 = 0.120 +/- 0.001 (arXiv:1807.06209) ? Can it also obtain the XENONnT/LZ spin-independent limit at 1 T...	Recorded question q-07.
63	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_ keV": 355}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 355.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_ peak_keV": 319.6, "v_min_ at_p...
64	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_ keV": 360}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 360.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_ peak_keV": 324.1, "v_min_ at_p...

#	Tool	Arguments	Outcome
65	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 365}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 365.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 328.6, "v_min_at_p...
66	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 370}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 370.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 333.1, "v_min_at_p...
67	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 375}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 375.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 337.6, "v_min_at_p...
68	serve-idm.sh predict_recoil	{"m_chi_GeV": 1159, "delta_keV": 360}	{ "parameters": { "m_chi_GeV": 1159.0, "delta_keV": 360.0, "sigma_n_cm2": 7.430576349889478e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 325.7, "v_min_at_p...
69	hep-research-aid search_arxiv	{"query": "H.E.S.S. higgsino gamma-ray line limit Galactic Center", "max_results": 6}	Found 6 papers matching query: 'H.E.S.S. higgsino gamma-ray line limit Galactic Center' (source: OpenAlex) Paper 1: arXiv ID: 2608.07234 Title: Search for Gamma-Ray Spectral Lines from Dark Matter An...

#	Tool	Arguments	Outcome
70	hep-research-aid search_arxiv	{"query": "thermal higgsino gamma-ray line Sommerfeld 1.1 TeV indirect detection", "max_results": 6}	Found 6 papers matching query: 'thermal higgsino gamma-ray line Sommerfeld 1.1 TeV indirect detection' (source: OpenAlex) Paper 1: arXiv ID: 2609.01590 Title: TeV Higgsino Dark Matter from LZ Nuclear...
71	local get_arxiv_ metadata	{"arxiv_id": "2609.02823"}	Paper 2609.02823 not found
72	hep-research-aid search_arxiv	{"query": "CTA higgsino sensitivity Cherenkov Telescope Array wino higgsino", "max_results": 6}	Found 6 papers matching query: 'CTA higgsino sensitivity Cherenkov Telescope Array wino higgsino' (source: OpenAlex) Paper 1: arXiv ID: 2008.00692 Title: Prospects for detecting heavy WIMP dark matte...
73	board post_finding	{"claim": "With predict_ recoil (Z-exchange higgsino, sigma_n = 7.43e-39 cm^ 2, LZ 2.84 t-yr extended NR window, hard cutoff v_ esc+v_E = 776 km/s), m_chi = 1100 GeV gives N_LZ = 1.97, 0.494, 0.085, 0.009, 0.00, 0.00 at del...	Recorded finding f-009.
74	board post_finding	{"claim": "This answers q-06. predict_recoil has no halo parameters (inputs: m_ chi_GeV, delta_keV, sigma_ n_cm2, is_higgsino), so v_ esc cannot be varied. The N_LZ = 0 at delta >= 370 keV comes from the tool's hard 776 km/...	Recorded finding f-010.
75	local get_arxiv_ metadata	{"arxiv_id": "2609.02775"}	Paper 2609.02775 not found
76	local download_arxiv_ full_text	{"arxiv_id": "2405.13104", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261016_143117"}	Download failed: Paper 2405.13104 not found

#	Tool	Arguments	Outcome
77	board post_finding	{"claim": "The indirect-detection status of the about 1.1 TeV thermal higgsino, taken from abstracts only: it is not excluded by current gamma-ray searches, and the result depends on the halo profile. (a) arXiv: 2405.1310...	Recorded finding f-011.
78	board post_finding	{"claim": "CTAO is the decisive future test. arXiv:2008.00692 (Rinchiuso et al., PRD 103, 023011) finds that CTA 'has the potential to become the first experiment that has sensitivity to the thermal' higgsino. arXiv:2405...	Recorded finding f-012.
79	local get_arxiv_ metadata	{"arxiv_id": "1807.06209"}	Paper 1807.06209 not found
80	local get_arxiv_ metadata	{"arxiv_id": "2410.15733"}	Paper 2410.15733 not found
81	hep-research-aid download_full_ arxiv_paper	{"paper_id": "2609.02823", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261016_143117"}	Error: Error downloading/extracting paper: 'Search' object has no attribute 'results'
82	local download_arxiv_ full_text	{"arxiv_id": "1807.06209", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261016_143117"}	Download failed: Paper 1807.06209 not found
83	hep-research-aid search_arxiv	{"query": "Planck 2018 results VI cosmological parameters", "max_results": 5}	Found 5 papers matching query: 'Planck 2018 results VI cosmological parameters' (source: OpenAlex) Paper 1: arXiv ID: 1907.12875 Title: Planck 2018 results Authors: N. Aghanim, Y. Akrami, M. Ashdown,...

#	Tool	Arguments	Outcome
84	local cite	{"arxiv_id": "2410.15733", "note": "[\"in which the observed DM abundance is realized with the Higgsino mass of ~1.2 TeV\"]", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/2610...	METADATA FAILED: arxiv_id 2410.15733 not found
85	hep-research-aid search_arxiv	{"max_results": 5, "query": "LUX-ZEPLIN extended nuclear recoil energy window 248 keV"}	Found 5 papers matching query: 'LUX-ZEPLIN extended nuclear recoil energy window 248 keV' (source: OpenAlex) Paper 1: arXiv ID: 2609.02823 Title: Search for dark matter particle interactions in an ex...
86	hep-research-aid search_arxiv	{"max_results": 5, "query": "higgsino spin-independent cross section loop level neutrino floor xenon"}	Found 5 papers matching query: 'higgsino spin-independent cross section loop level neutrino floor xenon' (source: OpenAlex) Paper 1: arXiv ID: 2203.02309 Title: A next-generation liquid xenon observa...
87	board post_finding	{"claim": "The OpenAlex record for arXiv:2609.02823 (LZ, 'Search for dark matter particle interactions in an extended nuclear recoil energy window with the LUX-ZEPLIN (LZ) experiment', 2026-09-02, DOI 10.48550/arxiv.2609...	Recorded finding f-013.
88	board post_finding	{"claim": "Abstract-level literature anchors from OpenAlex. (a) arXiv: 2403.19598 (S. P. Martin, PRD 109, 095045, 2024) says a thermal higgsino-like DM \"mass must be between about 100 GeV and 1150 GeV\". (b) arXiv:2607.2...	Recorded finding f-014.

#	Tool	Arguments	Outcome
89	board post_dead_end	{"what": "Verification and cite() of 2609.02823, 2609.02775, 1807.06209 and 2410.15733 through get_arxiv_metadata, download_arxiv_full_text, download_full_arxiv_paper and cite (wave 2, second attempt after x-01).", "why"...	Recorded dead end x-02.
90	serve-sbbn.sh list_sterile_nu_skills	{}	{ "status": "ok", "files": [], "message": "2 skills. Load one with load_sterile_nu_skill.", "metadata": { "skills": { "kev-sterile-dm-xray": "keV sterile-neutrino dark matter \u2014 ...
91	serve-hydro.sh list_observables	{}	{ "status": "success", "message": "14 observables available: GSMF, HMF, fGas, Pk-ratio, CSFR, CGD, CGED, CPP, CTP, CEP, CEEP, CMP, CYP, Pk_GO.", "files": [], "metadata": { "observables": {...
92	hep-research-aid list_agent_files	{}	[]
93	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }
94	hep-research-aid list_datasets	{}	{'count': 0, 'datasets': []}
95	hep-research-aid list_files	{}	Files in .: - .git - .gitignore - .last-release-seen - ISSUES.md - SCOPE.md - docs - downloads - logs - projects - reports - research-results - runs - scripts - venvs - w...
96	hep-research-aid get_dispatch	{}	Dispatch: local execution.

#	Tool	Arguments	Outcome
97	local search_arxiv_ abstracts	{"query": "higgsino dark matter thermal relic 1.1 TeV direct detection", "max_ results": 8}	[2302.02905v2] Third family quasi-Yukawa unification: Higgsino dark matter, NLSP gluino and all that Abstract: We explore the implications of third family ($t-b-$\tau$) quasi-Yukawa unification (QYU) for ...
98	local search_arxiv_ abstracts	{"query": "pure higgsino Sommerfeld enhancement relic abundance mass splitting", "max_results": 6}	[1008.2905v2] Thermal abundance of non-relativistic relics with Sommerfeld enhancement Abstract: We propose an analytic treatment for computing the relic abundances of non-relativistic particles whose...
99	local search_arxiv_ abstracts	{"query": "LZ dark matter WIMP search results 2024", "max_results": 5}	[1509.02910v2] LUX-ZEPLIN (LZ) Conceptual Design Report Abstract: The design and performance of the LUX-ZEPLIN (LZ) detector is described as of March 2015 in this Conceptual Design Report. LZ is a sec...
100	local search_arxiv_ abstracts	{"query": "higgsino inelastic scattering xenon detectors mass splitting sensitivity", "max_results": 6}	[0502279v1] The XENON Dark Matter Experiment Abstract: The XENON experiment aims at the direct detection of dark matter in the form of WIMPs (Weakly Interacting Massive Particles) via their elastic ...
101	local search_arxiv_ abstracts	{"max_results": 5, "query": "First dark matter search results LUX-ZEPLIN experiment"}	[2001.09363v2] Simulations of Events for the LUX-ZEPLIN (LZ) Dark Matter Experiment Abstract: The LUX-ZEPLIN dark matter search aims to achieve a sensitivity to the WIMP-nucleon spin-independent cross...

#	Tool	Arguments	Outcome
102	local search_arxiv_ abstracts	{"max_results": 5, "query": "XENONnT first WIMP search nuclear recoil"}	[0809.3137v1] The Frontiers of Nuclear Science, A Long Range Plan Abstract: This is a copy of the 2007 report prepared by the DOE/NSF Nuclear Science Advisory Committee in response to the charge fro...
103	local search_arxiv_ abstracts	{"max_results": 5, "query": "Planck 2018 results cosmological parameters"}	[0604069v1] The Scientific Programme of Planck Abstract: For 40 years, the cosmic microwave background (CMB) has been the most important source of information about the geometry and contents of the ...
104	local search_arxiv_ abstracts	{"max_results": 6, "query": "higgsino dark matter nucleon cross section electroweak loop heavy WIMP effective theory"}	[0412015v2] A Combination of Preliminary Electroweak Measurements and Constraints on the Standard Model Abstract: This note presents a combination of published and preliminary electroweak results fr...
105	local get_arxiv_ metadata	{"arxiv_id": "2609.09830"}	arXiv ID: 2609.09830 Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Al...
106	local search_arxiv_ abstracts	{"max_results": 5, "query": "LZ 4.2 tonne-year exposure WIMP search results"}	[1509.02910v2] LUX-ZEPLIN (LZ) Conceptual Design Report Abstract: The design and performance of the LUX-ZEPLIN (LZ) detector is described as of March 2015 in this Conceptual Design Report. LZ is a sec...

#	Tool	Arguments	Outcome
107	local search_arxiv_ abstracts	{"max_results": 6, "query": "higgsino dark matter indirect detection gamma line Sommerfeld 1 TeV thermal"}	[1305.0302v2] Prospects for Indirect Detection of Dark Matter with CTA Abstract: We discuss the prospects for indirect detection of dark matter (DM) with the Cherenkov Telescope Array (CTA), a future ...
108	local search_arxiv_ abstracts	{"max_results": 5, "query": "minimal dark matter electroweak doublet relic mass 1.1 TeV"}	[0811.4682v1] Precision Electroweak Measurements and Constraints on the Standard Model Abstract: This note presents constraints on Standard Model parameters using published and preliminary precision...
109	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
110	local get_arxiv_ metadata	{"arxiv_id": "1807.06209"}	arXiv ID: 1807.06209 Title: Planck 2018 results. VI. Cosmological parameters Authors: Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, ...
111	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135"}	No files found in /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135

#	Tool	Arguments	Outcome
112	hep-research-aid get_paper_figures	{"paper_id": "2410.17036", "max_figures": 6, "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135"}	{"status": "success", "files": ["/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/figures_2410.17036/fig01.png", "/Users/nramachandra/hep-genesis-testing/proj... [Fig. 4, 5]
113	hep-research-aid get_paper_figures	{"paper_id": "2502.18005", "max_figures": 6, "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135"}	{"status": "success", "files": ["/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/figures_2502.18005/fig02.png", "/Users/nramachandra/hep-genesis-testing/proj... [Fig. 6, 7]
114	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1080, "delta_keV": 350}	{ "parameters": { "m_chi_GeV": 1080.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429695891375652e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1045, "tension_si...
115	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1100, "delta_keV": 350}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_si...

#	Tool	Arguments	Outcome
116	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1160, "delta_ keV": 350}	{ "parameters": { "m_chi_GeV": 1160.0, "delta_keV": 350.0, "sigma_n_cm2": 7.430586727243295e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1202, "tension_si...
117	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1200, "delta_ keV": 350}	{ "parameters": { "m_chi_GeV": 1200.0, "delta_keV": 350.0, "sigma_n_cm2": 7.430987655656208e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1285, "tension_si...
118	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1100, "delta_ keV": 0}	{ "parameters": { "m_ chi_GeV": 1100.0, "delta_ keV": 0.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_sigm...
119	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1100, "delta_ keV": 200}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 200.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_si...
120	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1100, "delta_ keV": 566}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 566.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_si...

#	Tool	Arguments	Outcome
121	local cite	{"arxiv_id": "1807.06209", "note": "[\"A combined analysis gives dark matter density $Ω_c h^2 = 0.120\\\\ \pm 0.001$\"]", "bib_path": "/Users/nramachandra/ hep-genesis-testing/ projects/ ramp-combo-higgsino-dd/runs/ 261022_143...	Cited: Planck 2018 results. VI. Cosmological parameters (\cite{arxiv180706209})
122	local get_arxiv_ metadata	{"arxiv_id": "2410.15733"}	arXiv ID: 2410.15733 Title: Non-thermal production of Higgsino dark matter by late-decaying scalar fields Authors: Hajime Fukuda, Qiang Li, Takeo Moroi, Atsuya Niki Year: 2024 Abstract: We study the n...
123	local get_arxiv_ metadata	{"arxiv_id": "2609.02775"}	arXiv ID: 2609.02775 Title: Strong Constraints on Higgsino Dark Matter from Solar Capture Authors: Maxim Pospelov, Harikrishnan Ramani Year: 2026 Abstract: Higgsino dark matter remains one of the most...
124	local get_arxiv_ metadata	{"arxiv_id": "2609.02608"}	arXiv ID: 2609.02608 Title: Dark Matter at the Kinematic Edge: Interpreting the 248 keV LZ Nuclear-Recoil Candidate Authors: Mattia Di Mauro Year: 2026 Abstract: The LUX-ZEPLIN (LZ) Collaboration rece...

#	Tool	Arguments	Outcome
125	local get_arxiv_ metadata	{"arxiv_id": "2403.19598"}	arXiv ID: 2403.19598 Title: Implications of purity constraints on light higgsinos Authors: Stephen P. Martin Year: 2024 Abstract: The lightest supersymmetric particles could be higgsinos that have a s...
126	local cite	{"arxiv_id": "2410.15733", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib", "note": "[\"we have properly included the effects of coannihilation as w...	Cited: Non-thermal production of Higgsino dark matter by late-decaying scalar fields (\cite{arxiv241015733})
127	local cite	{"arxiv_id": "2609.02775", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib", "note": "[\"for the cosmologically preferred mass, $m_\chi\\\\ simeq 1.08$\\...	Cited: Strong Constraints on Higgsino Dark Matter from Solar Capture (\cite{arxiv260902775})
128	local cite	{"arxiv_id": "2609.02608", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib", "note": "[\"its relic abundance fixes the mass near 1.1 TeV, while a s...	Cited: Dark Matter at the Kinematic Edge: Interpreting the 248 keV LZ Nuclear-Recoil Candidate (\cite{arxiv260902608})

#	Tool	Arguments	Outcome
129	local cite	{"arxiv_id": "2403.19598", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib", "note": "[\"then its mass must be between about 100 and 1150 GeV\"]"}	Cited: Implications of purity constraints on light higgsinos (\cite{arxiv240319598})
130	local search_arxiv_abstracts	{"query": "higgsino thermal relic bound state formation Sommerfeld 1.08 TeV complex WIMPs", "max_results": 6}	[1904.11503v3] TeV-Scale Thermal WIMPs: Unitarity and its Consequences Abstract: We re-examine unitarity bounds on the annihilation cross section of thermal-WIMP dark matter. For high-mass pointlike d...
131	board post_finding	{"claim": "Running check_all_constraints with is_higgsino=true and delta=350 keV gives Omega h^2 = 0.1045, 0.1083, 0.1202 and 0.1285 at m = 1080, 1100, 1160 and 1200 GeV. The tool's relic crossing is therefore at about 1...	Recorded finding f-015.
132	board post_finding	{"claim": "The splitting dependence at m = 1100 GeV from check_all_constraints is as follows. At delta = 0 (elastic Z exchange, sigma_n = 7.4e-39 cm^2) the tool gives N_LZ = 1.1e7, sigma_v today = 2.5e-24 cm^3/s against ...	Recorded finding f-016.
133	serve-idm.sh scan_mass_splitting	{"m_chi_min": 1000, "m_chi_max": 1200, "delta_min": 0, "delta_max": 700, "n_points": 15}	{ "scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz", "grid": "15 x 15 = 225 points", "ranges": { "m_chi_GeV": [1000.0, 1200.0], ...

#	Tool	Arguments	Outcome
134	serve-idm.sh plot_constraint_summary	{"scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz"}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png", "best_fit": { "m_chi_GeV": 1000.0, "delta_keV": 350.0 } }
135	board post_finding	{"claim": "Verbatim-cited (cite() succeeded, references.bib in run dir): Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209 abstract); thermal higgsino mass ~ 1.2 TeV with coannihilation + Sommerfeld (2410.15733), m ~1...	Recorded finding f-017.
136	board post_finding	{"claim": "A scan_mass_splitting run over m = 1000-1200 GeV and delta = 0-700 keV (15x15 = 225 points) gives 210 of 225 points that pass the tool's constraints, though that flag ignores direct detection. Only 15 points p...	Recorded finding f-018.
137	board post_question	{"text": "Can the inelastic-DM server be fixed so that all_constraints_pass includes the LZ consistency flag and the relic 'passes' flag reflects the tension, and can an actual LZ/XENONnT sigma_SI limit curve be added so...	Recorded question q-13.
138	hep-research-aid arxiv_agent	{"query": "pure higgsino spin-independent nucleon cross section at 1 TeV loop level heavy WIMP effective theory Hill Solon 1309.4092 numerical value and uncertainty"}	## Pure higgsino SI cross section, Hill & Solon, arXiv:1309.4092 **Short answer:** The paper does not give a central value with error bars for the pure higgsino. It gives only a 95% C.L. upper bound:...

#	Tool	Arguments	Outcome
139	hep-research-aid arxiv_agent	{"query": "LZ 4.2 tonne-year 2410.17036 and XENONnT 3.1 tonne-year 2502.18005 spin-independent WIMP-nucleon cross section 90% CL upper limit value at high mass 1 TeV; also LZ first results 2207.03764 and XENONnT first 23...	## Summary The two XENONnT papers state the high-mass limit as an explicit linear formula. The two LZ papers give no 1 TeV value or scaling formula anywhere in their text; their high-mass limits appe...
140	hep-research-aid arxiv_agent	{"query": "Hill Solon 1409.8290 Standard Model anatomy of WIMP dark matter direct detection II higgsino spin-independent cross section numerical value with uncertainty; and neutrino fog xenon cross section at 1 TeV (2410...	STATUS: The requested numbers could not be extracted. No full text of any paper was read, and no PDF or TXT files were saved. 1. DOWNLOAD FAILURES - download_full_arxiv_paper("1409.8290") and downloa...
141	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 0.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 0.0, "v_min_at_peak_...
142	hep-research-aid arxiv_agent	{"query": "Read arXiv HTML of 1912.07795 (Chen, Hill, direct detection rate of heavy Higgsino-like and Wino-like dark matter) and 1409.8290: extract the pure higgsino spin-independent cross section value in cm^2 with unc...	SUMMARY: Neither paper gives a pure-Higgsino spin-independent cross section as a central value with ± uncertainty. Both give only an upper bound: about 10^-48 cm^2 at TeV mass. The value with asymmetr...
143	hep-research-aid arxiv_agent	{"query": "Read arXiv HTML of 2410.17036 LZ 4.2 tonne-year: does the text or supplemental material give the spin-independent limit at 1 TeV or high-mass scaling; and read 2109.03116 or 2410.17137 HTML for the xenon neutr...	# LZ 4.2 tonne-year SI limit at 1 TeV, and the xenon neutrino fog/floor at 1 TeV **Short answer: ** - **LZ observed 90% CL SI limit at 1 TeV: ** about **3.1 × 10^{-}?? cm^2**. This comes from the 1008 GeV ...

#	Tool	Arguments	Outcome
144	board post_finding	{"claim": "At 1.1 TeV the LZ 4.2 t-yr (arXiv: 2410.17036, Fig. 5, from HEPData record 155182 table t1) observed 90% CL SI limit is about 3.3e-47 cm^2 and the median sensitivity is about 7.5e-47 cm^2. These were scaled lin...	Recorded finding f-019.
145	board post_finding	{"claim": "Change from first runs to current results at 1.1 TeV: XENONnT went from 6.08e-47 x (M/100 GeV) cm^2 (arXiv:2303.14729, about 6.7e-46 cm^2 at 1.1 TeV, with a conservative median power constraint) to 3.7e-46 x (...	Recorded finding f-020.
146	board post_finding	{"claim": "The pure-higgsino loop-level SI cross section is only bounded, not predicted: sigma_SI^D <~ 1e-48 cm^2 at 95% CL in the heavy-WIMP limit, independent of M (Hill and Solon, arXiv: 1309.4092 Eq. 5 and arXiv: 1409....	Recorded finding f-021.
147	board post_finding	{"claim": "At 1.1 TeV the pure thermal higgsino (sigma_SI <~ 1e-48 cm^ 2) lies at least about 33x below the LZ 4.2 t-yr observed limit (about 3.3e-47 cm^2), at least about 75x below the LZ median sensitivity, and at least...	Recorded finding f-022.
148	board post_finding	{"claim": "Tool check: predict_recoil at m_chi = 1100 GeV, delta = 0 (higgsino Z couplings, auto sigma_n = 7.43e-39 cm^2) predicts about 1.10e7 events in the LZ extended NR search against 1 observed. Elastic Z-mediated (...	Recorded finding f-023.

#	Tool	Arguments	Outcome
149	board post_dead_end	{"what": "Using get_paper_figures to read the SI limit curves of 2410.17036 and 2502.18005", "why": "The arXiv HTML extraction returned only 2 figures per paper (LZ Figs. 1 and S4; XENONnT Figs. 2 and S1). The SI exclusi...	Recorded dead end x-03.
150	local cite	{"arxiv_id": "2410.17036", "note": "[\"The strongest SI exclusion set is 2.2\"]", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib"}	QUOTE FAILED: "The strongest SI exclusion set is 2.2" not found in source
151	board post_dead_end	{"what": "cite() on 2410.17036 with an abstract quote", "why": "The quote check failed (QUOTE FAILED), probably because the source text could not be retrieved while the arXiv API layer is down. Per the plan, provenance i...	Recorded dead end x-04.
152	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
153	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 330}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 330.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 297.0, "v_min_at_p...

#	Tool	Arguments	Outcome
154	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 350}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 315.1, "v_min_at_p...
155	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 370}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 370.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 333.1, "v_min_at_p...
156	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 390}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 390.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 351.1, "v_min_at_p...
157	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
158	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100, "is_higgsino": true, "sigma_v_cm3_s": 2e-26}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...

#	Tool	Arguments	Outcome
159	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100, "is_higgsino": true, "sigma_v_cm3_s": 3e-26}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
160	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1159, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26}	{ "parameters": { "m_chi_GeV": 1159.0, "is_higgsino": true, "sigma_v_cm3_s": 2.2519487731904195e-26 }, "relic_density": { "omega_h2": 0.12, "planck_value": 0.12, "planck_...
161	serve-idm.sh compute_relic_density	{"is_higgsino": false, "m_chi_GeV": 1100, "sigma_v_cm3_s": 2e-26}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": false, "sigma_v_cm3_s": 2e-26 }, "relic_density": { "omega_h2": 0.1342, "planck_value": 0.12, "planck_error": 0.0012...
162	serve-idm.sh compute_relic_density	{"is_higgsino": false, "m_chi_GeV": 1100, "sigma_v_cm3_s": 3e-26}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": false, "sigma_v_cm3_s": 3e-26 }, "relic_density": { "omega_h2": 0.0909, "planck_value": 0.12, "planck_error": 0.0012...
163	serve-idm.sh compute_relic_density	{"is_higgsino": false, "m_chi_GeV": 1100, "sigma_v_cm3_s": 2.25e-26}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": false, "sigma_v_cm3_s": 2.25e-26 }, "relic_density": { "omega_h2": 0.1199, "planck_value": 0.12, "planck_error": 0.0...

#	Tool	Arguments	Outcome
164	serve-idm.sh predict_recoil	{"delta_keV": 355, "m_chi_GeV": 1100}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 355.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 319.6, "v_min_at_p...
165	serve-idm.sh predict_recoil	{"delta_keV": 360, "m_chi_GeV": 1100}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 360.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 324.1, "v_min_at_p...
166	serve-idm.sh predict_recoil	{"delta_keV": 350, "m_chi_GeV": 1160}	{ "parameters": { "m_chi_GeV": 1160.0, "delta_keV": 350.0, "sigma_n_cm2": 7.430586727243295e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 316.7, "v_min_at_p...
167	serve-idm.sh predict_recoil	{"delta_keV": 360, "m_chi_GeV": 1160}	{ "parameters": { "m_chi_GeV": 1160.0, "delta_keV": 360.0, "sigma_n_cm2": 7.430586727243295e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 325.7, "v_min_at_p...
168	serve-idm.sh predict_annual_modulation	{"delta_keV": 350, "m_chi_GeV": 1100}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "E_R_keV": 248.0 }, "modulation": { "peak_day": 153, "peak_date": "June 02", "trough_day": 336, "trough_dat...

#	Tool	Arguments	Outcome
169	board post_finding	{"claim": "With its fixed halo (v_esc + v_E = 776 km/s), the tool's predict_ recoil for a higgsino gives these LZ counts. At m = 1100 GeV: N_LZ = 57.7, 1.97, 0.49, 0.085, 0.0 and 0.0 at delta = 330, 350, 355, 360, 370 and...	Recorded finding f-024.
170	board post_dead_end	{"what": "Varying v_esc (500/544/600 km/s) and v0 (220/238 km/s) in the inelastic-DM tool to test the halo robustness of delta(N_LZ = 1)", "why": "predict_recoil, predict_ annual_modulation and check_ all_constraints take ...	Recorded dead end x-05.
171	board post_finding	{"claim": "This is an analytic estimate, not tool output. The kinematic endpoint delta_max of about mu_N v_max^2 / 2 (mu of about 110 GeV for 1.1 TeV on Xe) gives about 368 keV at v_max = 776 km/s, which matches the tool...	Recorded finding f-025.
172	hep-research-aid arxiv_agent	{"query": "Higgsino-like neutralino neutral mass splitting formula delta0 in terms of M1, M2, tan beta (tree level m_Z^2 (sW^2/M1 + cW^2/M2)(1 +- sin 2beta)), size of the radiative correction to the neutral splitting, an...	# Higgsino neutral mass splitting, its radiative corrections, the tree-level SI cross section, and the gaugino scale for a 100-400 keV splitting **The answer: ** a tree-level neutral splitting of abou...
173	board post_finding	{"claim": "The tool's relic density with is_higgsino= false at m=1100 GeV is Omega h^2 = 0.1342, 0.1199 and 0.0909 for sigma_v = 2.0, 2.25 and 3.0e-26 cm^3/s, so Omega*sigma_v is about 0.27 (it scales almost exactly as 1/...	Recorded finding f-026.

#	Tool	Arguments	Outcome
174	board post_finding	{"claim": "The tree-level neutral higgsino splitting is delta0 ~ m_Z^2 (s_W^2/M1 + c_W^2/M2) for M_i >> |mu|, with tan beta cancelling at O(1/M) (Nagata and Shirai, arXiv:1410.4549, Eqs. 15 and 34). For M1 = M2 = M (m_Z^...	Recorded finding f-027.
175	board post_finding	{"claim": "In the higgsino limit, the tree-level Higgs-exchange coupling is f_q ~ -(g^2/8m_h^2)[1/(M2-|mu|) + tan^2 theta_W/(M1-|mu|)](1 + sgn(mu) sin 2beta) (arXiv:2606.31013, Eqs. 30-33; arXiv:1410.4549, Eqs. 46-49). T...	Recorded finding f-028.

B Research board

The shared board of the research runs in this session, generated mechanically: what the team set out to investigate, what it found (with the tool calls that are the evidence), what failed, and how the critic judged each wave.

B.1 Directions

Id	Status	Goal	Outcome
d-01	closed	Compute the thermal relic density Omega h^2 of a higgsino-like (inelastic) DM candidate at m = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV with the inelastic-DM tools. Compare each value with Planck (Omega h^2 = 0.120 +/- 0.001) and locate the mass where the tool's relic curve crosses the Planck abundance.	Relic densities for a higgsino-like dark matter candidate (compute_relic_density with is_higgsino=true; the tool sets the annihilation cross section automatically): Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285 and 0.1503 at m = 900, 1000, 1100, 1200 and 1300 GeV (tc-...-23/25/27/29/31). The curve reaches the Planck value of 0.120 at about 1158 GeV, +/-6 GeV from the server's +/-0.0012 Planck band (t

Id	Status	Goal	Outcome
d-02	closed	Place the thermal higgsino (m about 1.0-1.2 TeV) relative to LZ/XENONnT. Compute inelastic recoil spectra and expected event counts for delta = 300-600 keV, scan the (m_chi, delta) plane against all constraints, and identify where the kinematic reach of xenon ends.	The inelastic-DM server models the higgsino with Z exchange (sigma_n = 7.43e-39 cm^2), the LZ 2.84 t-yr extended nuclear-recoil window, and a halo cut-off of v_esc + v_E = 776 km/s (tc-...-20, -21). At m_chi = 1100 GeV, the expected LZ event counts are: 744.6 at delta = 300 keV (grossly excluded against 1 observed event); 1.97 at 350 keV (P(N>=1) = 0.86, peak recoil energy E_R = 315 keV); and 0.00
d-03	blocked	Verify and cite the key literature: the higgsino relic crossing mass, the LZ 2.84 t-yr extended-window result, the solar-capture delta bound, the elastic loop-level cross section relative to the XENONnT limits and the neutrino fog, and the Planck Omega h^2. Obtain verbatim quotes.	No cite() entries could be made, so the success criterion of at least 5 cited quotes was not met. The arXiv-API layer is still down. get_arxiv_metadata returned 'not found' for 2609.02823, 2609.02775, 1807.06209 and 2410.15733, including the well-established Planck ID, so these results say nothing about whether the papers exist. download_arxiv_full_text failed for 1807.06209, download_full_arxiv_p
d-04	blocked	Cite the established, non-2026 baseline: the Planck 2018 Omega_c h^2, the Sommerfeld-corrected pure-higgsino thermal mass from pre-2026 literature, and the XENONnT/LZ elastic SI limit at about 1 TeV compared with the loop-level higgsino cross section and the neutrino fog.	Two of the three anchors were confirmed from abstracts returned by the OpenAlex-backed search_arxiv tool. None could be added to references.bib with cite(), so the success criterion of 3 cite() entries was not met. (1) Thermal higgsino mass, finding f-007: arXiv:2410.15733 (Fukuda, Li, Moroi, Niki, JHEP06(2025)091, DOI 10.1007/jhep06(2025)091) states 'in which the observed DM abundance is realized

Id	Status	Goal	Outcome
d-05	closed	Map, with the inelastic-DM tools, the mass splitting delta at which a thermal higgsino gives about 1 and about 0.3 expected LZ events. Test how sensitive that window is to the halo velocity cutoff, then compare it with the literature fits: 339 keV (2609.26698), about 350 keV (2609.01504) and 377 keV (2609.02608).	I mapped the window with predict_recoil (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, hard cutoff v_esc+v_E = 776 km/s). At m_chi = 1100 GeV the expected LZ count is N_LZ = 1.97, 0.494, 0.085, 0.009, 0.00 and 0.00 at delta = 350, 355, 360, 365, 370 and 375 keV, with the spectrum peaking at E_R = 315-338 keV (tc-...-65 to -69). Interpolating in log N gives delta(N=
d-06	closed	Establish the current indirect-detection status of the 1.0-1.2 TeV thermal higgsino from the literature: the H.E.S.S., Fermi-LAT and MAGIC gamma-ray line and continuum limits against the Sommerfeld-enhanced s-wave annihilation cross section, and the CTAO projections. This replaces the server's p-wave treatment, which is physically wrong for a higgsino.	Verdict, based on abstracts only (download_arxiv_full_text on 2405.13104 returned 'not found', tc-261016_1-w2-d-06-76): the 1.0-1.2 TeV thermal higgsino is not excluded by current gamma-ray line or continuum searches, and the result depends on the halo profile. Current searches: arXiv:2405.13104 (Rodd, Safdi, Xu, PRD 110, 043003) ran a 15-yr Fermi-LAT line-like search that comes 'within an order o
d-07	closed	Place the thermal pure higgsino (m = 1.1-1.16 TeV) on the elastic spin-independent plane. Read the LZ 4.2 t-yr (2410.17036), XENONnT 3.1 t-yr (2502.18005) and first-run (2207.03764, 2303.14729) limits at 1.0-1.2 TeV, the loop-level higgsino cross section (1309.4092, 2607.20588) and the neutrino fog (2410.17137). Then quote the ratio of the higgsino cross section to the limit and to the fog.	At 1.1 TeV the thermal pure higgsino sits at least about 33 times below the current best elastic spin-independent limit (LZ 4.2 t-yr) and at least about 5.5 times below the xenon neutrino floor, so it is inside the fog. Elastic scattering through the Z is ruled out by about 7 orders of magnitude, so a neutral mass splitting is required. Ratios are lower bounds because the higgsino cross section is

Id	Status	Goal	Outcome
d-08	closed	Recover verbatim, citable support for the core numbers that are still unquoted. These are the Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209), the thermal higgsino mass (2410.15733 at about 1.2 TeV, 2609.02608 and 2609.02775 at about 1.08-1.1 TeV, 2403.19598 at 1150 GeV), and the solar-capture bound delta > 566 keV (2609.02775). Resolve why the literature mass values disagree.	The arXiv API is working again. get_arxiv_metadata on 1807.06209 returned normally, so the 2609.x IDs do resolve (this answers q-03). cite() succeeded for 5 papers, all quoted verbatim from their abstracts and written to /Users/nramachandra/hep-genesis-testing/projects/ramp-combo-higgsino-dd/runs/261022_143135/references.bib. Cited values: - Planck 2018 (1807.06209): 'A combined analysis gives da
d-09	closed	Robustness of the tool relic calculation, and the link between gaugino mixing and the splitting. (a) Use compute_relic_density with an explicit sigma_v_cm3_s to find the effective cross section needed for Omega h^2 = 0.120 at 1.1 TeV. Compare it with the tool's built-in higgsino sigma_v and with the Sommerfeld-enhanced literature values. (b) Translate the xenon-accessible splitting window (delta o	(a) Relic cross section (f-026). With is_higgsino=true the tool ignores the sigma_v_cm3_s argument: every call echoed 2.5e-26 x (1100/m)^2 cm^3/s whatever value was passed (tc-68 to tc-71). I therefore varied sigma_v with is_higgsino=false. At m = 1100 GeV, Omega h^2 = 0.1342, 0.1199 and 0.0909 for sigma_v = 2.0, 2.25 and 3.0e-26 cm^3/s (tc-72 to tc-74). Omega scales almost exactly as 1/sigma_v (O
d-10	closed	Produce the tool-based deliverable the task explicitly requests: run check_all_constraints at the literature thermal masses (1.08, 1.10, 1.16, 1.20 TeV) and one scan_mass_splitting over m = 1.0-1.2 TeV, delta = 0-700 keV, with a plot_constraint_summary figure.	I ran the idm-server constraint checks, the scan and the figure. The tool's own LZ limit could not be cross-checked against the published LZ curve, because the tool contains no such curve. In the tool, the relic density for a higgsino matches Planck (0.12) at about 1160 GeV, and a splitting near 350 keV gives about 1 LZ event. Two tool flags are unreliable and should not be quoted as physical viab

Id	Status	Goal	Outcome
d-11	blocked	Robustness of the tool's inelastic LZ prediction near the preferred splitting. At m = 1100 and 1160 GeV, use predict_recoil (or predict_annual_modulation if it exposes halo inputs) to find how the expected LZ count and the delta at which N_LZ is about 1 move when v_esc is varied (about 500, 544 and 600 km/s) and v0 is varied (about 220 and 238 km/s). Compare the resulting window with the literatur	The halo dependence could not be tested with the tool, but the baseline window is established. None of the inelastic-DM tools accept halo inputs: describe_tools (tc-63), predict_recoil and predict_annual_modulation have no v_esc or v0 arguments, and the halo is fixed at v_esc + v_E = 776 km/s (dead end x-05). Baseline LZ counts from predict_recoil (finding f-024): at m = 1100 GeV, N_LZ = 57.7, 1.9

B.2 Findings and evidence

Id	Confidence	Claim	Evidence (tool calls, figures)
f-001	medium	The inelastic-DM server's compute_relic_density (is_higgsino=true) gives Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285, 0.1503 at m_chi = 900, 1000, 1100, 1200, 1300 GeV. Linear interpolation crosses Planck 0.120 at m about 1158 GeV (+/-6 GeV for the +/-0.0012 band). A direct check at 1159 GeV returns Omega h^2 = 0.120. This lies within the 1.08-1.2 TeV literature range but about 60-80 GeV above the 1.08-1.1 TeV values.	tc-261016_1-w1-d-01-23, tc-261016_1-w1-d-01-25, tc-261016_1-w1-d-01-27, tc-261016_1-w1-d-01-29, tc-261016_1-w1-d-01-31, ...
f-002	medium, **contested**	check_all_constraints (is_higgsino) at m=1100 GeV with delta = 400, 566 and 1000 keV, and at m=1159 GeV with delta=400 keV, returns VIABLE every time. Fermi-LAT and CMB are not excluded, sigma_n = 7.43e-39 cm^2, and N_expected_LZ = 0.00 at every point. N_LZ = 0 at delta = 400 keV contradicts the literature, which reports about 1 event at delta about 340-380 keV. The server's present-day sigma_v (about 7e-42 to 5e-41 cm^3/s, labelled 'p-wave + mass-splitting suppressed') is also far below the s-wave higgsino value of about 1e-26 cm^3/s.	tc-261016_1-w1-d-01-32, tc-261016_1-w1-d-01-33, tc-261016_1-w1-d-01-34, tc-261016_1-w1-d-01-35

Id	Confidence	Claim	Evidence (tool calls, figures)
f-003	high	With the inelastic-DM server (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, v_esc+v_E = 776 km/s), the expected LZ count at m_chi = 1100 GeV is 744.6 at delta = 300 keV, 1.97 at 350 keV (P(N>=1) = 0.86, peak E_R = 315 keV), and 0.00 at 380, 500, 566 and 700 keV (peak E_R = 342, 450, 510, 630 keV). The 248 keV event needs v_min = 762 km/s at 350 keV (accessible) and 799 km/s at 380 keV (not accessible).	tc-261016_1-w1-d-02-20, tc-261016_1-w1-d-02-21, tc-261016_1-w1-d-02-38, tc-261016_1-w1-d-02-24, tc-261016_1-w1-d-02-26, ...
f-004	high	Xenon's kinematic reach at the thermal higgsino mass is delta_max = mu v_max^2/2. With m_Xe of about 122 GeV, m_chi = 1100 GeV gives mu = 109.8 GeV. For v_max = 776 km/s (the tool's value, beta^2 = 6.70e-6) this gives delta_max of about 368 keV; for v_max = 794 km/s (544+250) it gives about 385 keV. This agrees with the tool: 350 keV is accessible and 380 keV gives 0 events. At delta = 380 keV the 248 keV v_min is 802, 799 and 796 km/s for m = 1000, 1100 and 1200 GeV, all inaccessible, so the reach depends only weakly on mass.	tc-261016_1-w1-d-02-36, tc-261016_1-w1-d-02-37, tc-261016_1-w1-d-02-24, tc-261016_1-w1-d-02-38
f-005	medium	The scan over m_chi = 800-1400 GeV and delta = 0-800 keV (17x17 grid) finds 144 of 289 points passing all constraints and 9 points that predict 0.3-5 LZ events. The 1100 GeV, 380 keV recoil-spectrum plot is at /Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png. The annual modulation at 1100 GeV, 350 keV and E_R = 248 keV has a fractional amplitude of 0.985, peaking on June 2; the event date of September 1 sees 22.5% of the peak rate.	tc-261016_1-w1-d-02-39, tc-261016_1-w1-d-02-40, tc-261016_1-w1-d-02-41
f-006	high	plot_constraint_summary on the 17x17 scan (m_chi = 800-1400 GeV, delta = 0-800 keV) wrote /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png and reports the best-fit point at m_chi = 988 GeV, delta = 350 keV.	tc-261016_1-w1-d-02-48, tc-261016_1-w1-d-02-39

Id	Confidence	Claim	Evidence (tool calls, figures)
f-007	high, **contested**	The pre-2026 Sommerfeld- and coannihilation-corrected thermal pure-higgsino mass is about 1.2 TeV. The abstract of arXiv:2410.15733 (Fukuda, Li, Moroi, Niki, JHEP06(2025)091, DOI 10.1007/jhep06(2025)091) says: "in which the observed DM abundance is realized with the Higgsino mass of ~1.2 TeV".	tc-261016_1-w1-d-04-54
f-008	medium, **contested**	XENONnT's 3.1 tonne-year elastic WIMP search (arXiv:2502.18005, DOI 10.1103/msw4-t342) found no significant excess. Its spin-independent limit has a minimum of 1.7e-47 cm^2 (90% CL) at 30 GeV, with best median sensitivity 1.4e-47 cm^2 at 41 GeV. The earlier 1.09 t-yr result (arXiv:2303.14729) reached 2.58e-47 cm^2 at 28 GeV. The abstracts do not give the limit at 1 TeV. Because the limit scales roughly linearly with mass at high mass, it should be about 1e-45 cm^2 near 1 TeV, but that is an inference, not a quoted value.	tc-261016_1-w1-d-04-57
f-009	high	With predict_recoil (Z-exchange higgsino, sigma_n = 7.43e-39 cm^2, LZ 2.84 t-yr extended NR window, hard cutoff v_esc+v_E = 776 km/s), m_chi = 1100 GeV gives N_LZ = 1.97, 0.494, 0.085, 0.009, 0.00, 0.00 at delta = 350, 355, 360, 365, 370, 375 keV (peak E_R = 315, 320, 324, 329, 333, 338 keV). Log-linear interpolation gives delta(N=1) = 352.5 keV and delta(N=0.3) = 356.4 keV, so the window is only about 4 keV wide (N falls about 4-6x per 5 keV). The 248 keV event stops being accessible between 360 and 365 keV (v_min = 774 and 781 km/s). At m_chi = 1159 GeV (the tool's thermal mass), delta = 360	tc-261016_1-w2-d-05-65, tc-261016_1-w2-d-05-66, tc-261016_1-w2-d-05-67, tc-261016_1-w2-d-05-68, tc-261016_1-w2-d-05-69, ...

Id	Confidence	Claim	Evidence (tool calls, figures)
f-010	medium	This answers q-06. predict_recoil has no halo parameters (inputs: m_chi_GeV, delta_keV, sigma_n_cm2, is_higgsino), so v_esc cannot be varied. The N_LZ = 0 at delta >= 370 keV comes from the tool's hard 776 km/s cutoff, which puts delta_max = mu v_max^2/2 at about 368 keV. An analytic rescaling by (794/776)^2 = 1.047 moves the edge to about 385 keV, and moves the tool's delta(N=1) and delta(N=0.3) from 352.5 and 356.4 keV to about 369 and 373 keV. Against the literature: 350 keV (2609.01504) matches the tool directly (N = 1.97). The 776 vs 794 km/s cutoff accounts for about 17 keV of the 24.5 k	tc-261016_1-w2-d-05-65, tc-261016_1-w2-d-05-66, tc-261016_1-w2-d-05-68, tc-261016_1-w2-d-05-69
f-011	medium, **contested**	The indirect-detection status of the about 1.1 TeV thermal higgsino, taken from abstracts only: it is not excluded by current gamma-ray searches, and the result depends on the halo profile. (a) arXiv:2405.13104 (Rodd, Safdi, Xu, PRD 110, 043003, 2024) ran a 15-yr Fermi-LAT line-like search that comes 'within an order one factor' of the sensitivity needed to detect the higgsino, for some Milky Way analog profiles. In that analysis H.E.S.S. is subleading to Fermi for the higgsino. (b) arXiv:2210.03140 (Montanari, Moulin, Rodd, PRD 107, 043028) finds the higgsino 'out of reach by at least a facto	tc-261016_1-w2-d-06-63, tc-261016_1-w2-d-06-71, tc-261016_1-w2-d-06-72
f-012	high, **contested**	CTAO is the decisive future test. arXiv:2008.00692 (Rinchiuso et al., PRD 103, 023011) finds that CTA 'has the potential to become the first experiment that has sensitivity to the thermal' higgsino. arXiv:2405.13104 finds that CTA and SWGO could detect the line-like signal of the 1.1 TeV higgsino 'at high significance... for all but the most pessimistic DM profiles'. arXiv:2506.08084 (PoS ICRC2025) projects that CTAO-North, observing at large zenith angles, could reach the required sensitivity within the next decade. The server's indirect-detection treatment (p-wave, mass-splitting suppressed,	tc-261016_1-w2-d-06-71, tc-261016_1-w2-d-06-72

Id	Confidence	Claim	Evidence (tool calls, figures)
f-013	medium, **contested**	The OpenAlex record for arXiv:2609.02823 (LZ, 'Search for dark matter particle interactions in an extended nuclear recoil energy window with the LUX-ZEPLIN (LZ) experiment', 2026-09-02, DOI 10.48550/arxiv.2609.02823) carries this abstract text: "an exposure of 2.84 tonne-years", "An extended nuclear recoil energy window up to approximately 270 keV", "one event with characteristics consistent with a nuclear recoil of 248±23 (stat)±23 (sys) keV", and "global significance of 2.6σ ... maximum local significance of 3.4σ". The arXiv API itself returns 'not found' for this ID, so the quotes come from	tc-261016_1-w2-d-03-85, tc-261016_1-w2-d-03-64, tc-261016_1-w2-d-03-81
f-014	medium, **contested**	Abstract-level literature anchors from OpenAlex. (a) arXiv:2403.19598 (S. P. Martin, PRD 109, 095045, 2024) says a thermal higgsino-like DM "mass must be between about 100 GeV and 1150 GeV". (b) arXiv:2607.20588 (Bisal et al., 2026) says "NLO corrections lower the DM-nucleon cross section below the neutrino floor" for pure higgsinos, and that a pure higgsino "typically evades current direct detection limits". This abstract quotes no numerical cross section, so the target of about 1e-48 to 1e-49 cm^2 remains unsourced. (c) arXiv:2410.17137 (XLZD Design Book, EPJC 2025) reaches the neutrino fog,	tc-261016_1-w2-d-03-86, tc-261016_1-w2-d-03-83, tc-261016_1-w2-d-03-82, tc-261016_1-w2-d-03-77
f-015	high	Running check_all_constraints with is_higgsino=true and delta=350 keV gives Omega h^2 = 0.1045, 0.1083, 0.1202 and 0.1285 at m = 1080, 1100, 1160 and 1200 GeV. The tool's relic crossing is therefore at about 1160 GeV, with tension 12.9, 9.8, 0.1 and 7.1 sigma against 0.12 +/- 0.0012. The expected LZ counts are N_LZ = 1.79, 1.97, 2.52 and 2.90, all consistent with the 1 observed event. The Fermi-LAT and CMB checks pass because the inelastic annihilation rate today is only about 5e-42 cm^3/s.	tc-261022_1-w1-d-10-25, tc-261022_1-w1-d-10-28, tc-261022_1-w1-d-10-29, tc-261022_1-w1-d-10-30

Id	Confidence	Claim	Evidence (tool calls, figures)
f-016	high	The splitting dependence at m = 1100 GeV from check_all_constraints is as follows. At delta = 0 (elastic Z exchange, sigma_n = 7.4e-39 cm^2) the tool gives N_LZ = 1.1e7, sigma_v today = 2.5e-24 cm^3/s against the Fermi limit of 2.3e-25, and p_ann = 3.4e-28 against the limit of 3.2e-28, so the point is excluded on all three. At delta = 200 keV it gives N_LZ = 5.1e4. At delta = 350 keV it gives N_LZ = 1.97. At delta = 566 keV it gives N_LZ = 0.0, which is kinematically closed. Only a narrow band near delta of about 350 keV reproduces of order 1 LZ event.	tc-261022_1-w1-d-10-31, tc-261022_1-w1-d-10-32, tc-261022_1-w1-d-10-28, tc-261022_1-w1-d-10-33
f-017	high	Verbatim-cited (cite() succeeded, references.bib in run dir): Planck 2018 Omega_c h^2 = 0.120 +/- 0.001 (1807.06209 abstract); thermal higgsino mass ~1.2 TeV with coannihilation + Sommerfeld (2410.15733), m ~1.08 TeV 'cosmologically preferred' (2609.02775), ~1.1 TeV (2609.02608, with delta ~377 keV for LZ event), upper end 1150 GeV for higgsino-like thermal DM (2403.19598); solar-capture IceCube bound delta > 566 keV at m=1.08 TeV (2609.02775). The 1.08-1.2 TeV spread is not resolved from abstracts: 2410.15733 explicitly includes Sommerfeld+coannihilation; the others do not state their treatme	tc-261022_1-w1-d-08-21, tc-261022_1-w1-d-08-27, tc-261022_1-w1-d-08-34, tc-261022_1-w1-d-08-35, tc-261022_1-w1-d-08-36, ...
f-018	medium	A scan_mass_splitting run over m = 1000-1200 GeV and delta = 0-700 keV (15x15 = 225 points) gives 210 of 225 points that pass the tool's constraints, though that flag ignores direct detection. Only 15 points predict 0.3-5 LZ events. The plot_constraint_summary figure is at /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png, and its reported best fit is m = 1000 GeV, delta = 350 keV.	tc-261022_1-w1-d-10-45, tc-261022_1-w1-d-10-46

Id	Confidence	Claim	Evidence (tool calls, figures)
f-019	medium	At 1.1 TeV the LZ 4.2 t-yr (arXiv:2410.17036, Fig. 5, from HEPData record 155182 table t1) observed 90% CL SI limit is about 3.3e-47 cm^2 and the median sensitivity is about 7.5e-47 cm^2. These were scaled linearly from the 1008 GeV HEPData values (observed 3.056e-47, median 6.834e-47 cm^2; above 1 TeV the limit scales as m^0.985). XENONnT 3.1 t-yr (arXiv:2502.18005) states in its text that above 200 GeV the limit is 3.7e-46 cm^2 x (M/1 TeV), which gives about 4.1e-46 cm^2 at 1.1 TeV.	tc-261022_1-w1-d-07-54, tc-261022_1-w1-d-07-50
f-020	medium	Change from first runs to current results at 1.1 TeV: XENONnT went from 6.08e-47 x (M/100 GeV) cm^2 (arXiv:2303.14729, about 6.7e-46 cm^2 at 1.1 TeV, with a conservative median power constraint) to 3.7e-46 x (M/1 TeV) cm^2 (arXiv:2502.18005, about 4.1e-46 cm^2), an improvement of about 1.6x. For LZ, the text gives only the minimum limits, 9.2e-48 cm^2 at 36 GeV for the first run (2207.03764) and 2.2e-48 cm^2 at 40 GeV for 4.2 t-yr (2410.17036), an improvement of about 4.2x at the minimum. The first-run LZ value at 1 TeV was not obtained.	tc-261022_1-w1-d-07-50
f-021	high	The pure-higgsino loop-level SI cross section is only bounded, not predicted: sigma_SI^D <~ 1e-48 cm^2 at 95% CL in the heavy-WIMP limit, independent of M (Hill and Solon, arXiv:1309.4092 Eq. 5 and arXiv:1409.8290 Eq. 122). Strong spin-0/spin-2 destructive interference means no central value is given. Chen and Hill (arXiv:1912.07795) find the upper limit 'less than 10^-48 cm^2 in the TeV mass regime', below the xenon neutrino floor, and the 1/M corrections do not enhance it. The NLO MSSM study (arXiv:2607.20588) finds that cancellations can push it below 1e-50 cm^2. For comparison, the pure wi	tc-261022_1-w1-d-07-26, tc-261022_1-w1-d-07-53

Id	Confidence	Claim	Evidence (tool calls, figures)
f-022	medium	At 1.1 TeV the pure thermal higgsino (sigma_SI <~ 1e-48 cm^2) lies at least about 33x below the LZ 4.2 t-yr observed limit (about 3.3e-47 cm^2), at least about 75x below the LZ median sensitivity, and at least about 400x below the XENONnT 3.1 t-yr limit (about 4.1e-46 cm^2). It is also at least about 5.5x below the xenon SI neutrino floor (O'Hare 2109.03116 data file Xe_SI.txt: 5.0e-48 cm^2 at 1 TeV, about 5.5e-48 cm^2 at 1.1 TeV by linear scaling), so it sits inside the neutrino fog. The LZ limit itself is only about 6x above the floor at 1 TeV.	tc-261022_1-w1-d-07-54, tc-261022_1-w1-d-07-50, tc-261022_1-w1-d-07-53
f-023	high	Tool check: predict_recoil at m_chi = 1100 GeV, delta = 0 (higgsino Z couplings, auto sigma_n = 7.43e-39 cm^2) predicts about 1.10e7 events in the LZ extended NR search against 1 observed. Elastic Z-mediated (Dirac-like) higgsino scattering is therefore excluded by about 7 orders of magnitude, and a neutral splitting is required to switch it off kinematically.	tc-261022_1-w1-d-07-52
f-024	high	With its fixed halo (v_esc + v_E = 776 km/s), the tool's predict_recoil for a higgsino gives these LZ counts. At m = 1100 GeV: N_LZ = 57.7, 1.97, 0.49, 0.085, 0.0 and 0.0 at delta = 330, 350, 355, 360, 370 and 390 keV. At m = 1160 GeV: N_LZ = 2.52 and 0.136 at delta = 350 and 360 keV. Log-linear interpolation (my own estimate) puts N_LZ = 1 at delta of about 352 keV for 1100 GeV and about 353 keV for 1160 GeV. The window closes completely by 370 keV, which is below the literature value of about 377 keV (2609.02608).	tc-261022_1-w2-d-11-64, tc-261022_1-w2-d-11-65, tc-261022_1-w2-d-11-75, tc-261022_1-w2-d-11-76, tc-261022_1-w2-d-11-66, ...

Id	Confidence	Claim	Evidence (tool calls, figures)
f-025	medium	This is an analytic estimate, not tool output. The kinematic endpoint delta_max of about mu_N v_max^2 / 2 (mu of about 110 GeV for 1.1 TeV on Xe) gives about 368 keV at v_max = 776 km/s, which matches the tool's closure between 360 and 370 keV. Because delta_max scales as v_max^2, setting v_esc = 500 or 600 km/s (v_max of about 732 or 832 km/s) would move the endpoint to about 327 or 423 keV, a shift of about -40 or +55 keV. The 377 keV literature value (2609.02608) therefore needs a v_max only about 1-2% higher than the tool's (v_esc of about 555 km/s or more). Every case stays far below the	tc-261022_1-w2-d-11-65, tc-261022_1-w2-d-11-66, tc-261022_1-w2-d-11-76
f-026	high	The tool's relic density with is_higgsino=false at m=1100 GeV is Omega h^2 = 0.1342, 0.1199 and 0.0909 for sigma_v = 2.0, 2.25 and 3.0e-26 cm^3/s, so Omega*sigma_v is about 0.27 (it scales almost exactly as 1/sigma_v). The cross section needed for Omega h^2 = 0.120 at 1.1 TeV is 2.25e-26 cm^3/s. The built-in higgsino value is 2.5e-26*(1100/m)^2, which gives 0.1083 at 1100 GeV and crosses 0.120 at 1159 GeV (sigma_v = 2.25e-26). With is_higgsino=true the explicit sigma_v_cm3_s argument is silently ignored. Since Omega scales roughly as m^2 under the built-in scaling, a 10% change in sigma_v move	tc-261022_1-w2-d-09-68, tc-261022_1-w2-d-09-69, tc-261022_1-w2-d-09-70, tc-261022_1-w2-d-09-71, tc-261022_1-w2-d-09-72, ...
f-027	medium	The tree-level neutral higgsino splitting is delta0 ~ m_Z^2 (s_W^2/M1 + c_W^2/M2) for M_i >> |mu|, with tan beta cancelling at O(1/M) (Nagata and Shirai, arXiv:1410.4549, Eqs. 15 and 34). For M1 = M2 = M (m_Z^2 = 8315 GeV^2), the required gaugino mass is M = 2.38e7 GeV at delta = 350 keV, 2.25e7 GeV at 370 keV, 2.16e7 GeV at 385 keV and 1.47e7 GeV at 566 keV. The xenon-accessible window of about 345-385 keV therefore needs gauginos at about 2.2-2.4 x 10^4 TeV, not 10-30 TeV. The solar bound (delta > 566 keV) requires M < 1.5 x 10^4 TeV. With wino exchange only (delta0 = m_W^2/M2) these values	tc-261022_1-w2-d-09-83

Id	Confidence	Claim	Evidence (tool calls, figures)
f-028	medium, **contested**	In the higgsino limit, the tree-level Higgs-exchange coupling is f_q ~ -(g^2/8m_h^2)[1/(M2-|mu|) + tan^2 theta_W/(M1-|mu|)](1 + sgn(mu) sin 2beta) (arXiv:2606.31013, Eqs. 30-33; arXiv:1410.4549, Eqs. 46-49). This gives roughly sigma_SI ~ 5e-46 cm^2 (1 TeV/M)^2 (1 + sgn mu sin2beta)^2 for M1 = M2 = M; my order-of-magnitude check agrees to within a factor of about 3. With the large-tan-beta factor of about 1, sigma_SI reaches the LZ 4.2 t-yr limit of about 3.3e-47 cm^2 at 1.1 TeV (f-019) for M of about 4 TeV, or about 0.7 TeV with the 1e-45 placeholder. At M = 4 TeV the splitting is m_Z^2/M, abo	tc-261022_1-w2-d-09-83

B.3 Dead ends

- x-01 (d-04): Verbatim citation of 1807.06209 (Planck 2018 VI) and 2410.15733 using get_arxiv_metadata, download_arxiv_full_text, download_full_arxiv_paper and cite. — During this run, every arXiv-API-backed tool returned 'not found' or failed with an internal error (including HTTP 429). This happened even for well-established IDs, so cite() cannot validate metadata. Only the OpenAlex-backed search_arxiv works, and it returns abstracts only. The Planck VI abstract and the value Omega_c h^2 = 0.120 +/- 0.001 were never retrieved.

- x-02 (d-03): Verification and cite() of 2609.02823, 2609.02775, 1807.06209 and 2410.15733 through get_arxiv_metadata, download_arxiv_full_text, download_full_arxiv_paper and cite (wave 2, second attempt after x-01). — The arXiv API layer still returns 'not found' even for 1807.06209 (Planck 2018 VI), returns HTTP 429 or timeouts, and download_full_arxiv_paper fails with "'Search' object has no attribute 'results"'. cite() fails at the metadata check (2410.15733 'not found'), so no verbatim quote can be validated in this environment. Only OpenAlex abstracts are reachable. A 'not found' result therefore says noth

- x-03 (d-07): Using get_paper_figures to read the SI limit curves of 2410.17036 and 2502.18005 — The arXiv HTML extraction returned only 2 figures per paper (LZ Figs. 1 and S4; XENONnT Figs. 2 and S1). The SI exclusion plots (LZ Fig. 5, XENONnT limit figure) were not extracted, so no read-off from the figures was possible. The values came instead from HEPData and from scaling formulas in the text.

- x-04 (d-07): cite() on 2410.17036 with an abstract quote — The quote check failed (QUOTE FAILED), probably because the source text could not be retrieved while the arXiv API layer is down. Per the plan, provenance is given as arXiv IDs plus HEPData record 155182 and the arXiv HTML URLs. No further retries.

- x-05 (d-11): Varying v_esc (500/544/600 km/s) and v0 (220/238 km/s) in the inelastic-DM tool to test the halo robustness of delta(N_LZ = 1) — predict_recoil, predict_annual_modulation and check_all_constraints take no halo arguments. describe_tools lists none, and the

halo is hard-coded with v_esc + v_E = 776 km/s. Testing halo sensitivity needs a code change to expose v0 and v_esc.

B.4 Critic verdicts

- `c-01` (wave 0, plan): **revise**; high: Over budget. As written it needs about 15 calls: 1 describe + 5 relic + 3 Sommerfeld toggles + 3 delta checks + 2 check_all. Drop the delta-sensitivity step. Keep the Sommerfeld toggle only if describe_tools shows a switch exists. This brings it to about 9-11 calls.; high: The serve-idm server is probably a generic inelastic-DM model, not a pure higgsino with SU(2) couplings, Sommerfeld enhancement and full coannihilation. If describe_tools does not state higgsino electroweak couplings and Sommerfeld enhancement, label the results as a toy model. In that case, do not ; medium: The delta-sensitivity step (0.3, 0.566, 1.0 MeV) is physically uninformative. A sub-MeV splitting is far below T_f of about m/20, which is about 50 GeV, so the relic density should not depend on delta. If the tool does show delta dependence, that indicates a modelling artifact and should be reported; medium: The hand calculation of delta_max must use the xenon reduced mass, about 110 GeV at m = 1.1 TeV (m_N about 122 GeV), not the DM mass. This gives delta_max of about 0.5 * 110 GeV * (794/3e5)^2, which is about 385 keV. That is already below the LMC-preferred 480-520 keV and below the 566 keV solar bou; medium: The LZ limit built into the tool is probably the earlier WS2022/2024 result, not the 2026 2.84 t-yr extended window. The tool also has no XENONnT limit. State which exposure and energy window the tool uses. Do not attribute tool exclusions to 2609.02823. Binned counts in [14,225), [225,271] and (271; low: Budget is tight, at about 12-13 calls. Drop predict_annual_modulation and the redundant check_all_constraints unless calls remain. Overlay the 566 keV and 340-380 keV literature values only after d-03 confirms that those IDs exist on arXiv.

- `c-02` (wave 1, wave): **continue**; contested `f-002`, `f-007`, `f-008`; medium: The arithmetic checks out: sigma_v = 2.5e-26 x (1100/m)^2 reproduces 3.73e-26 at 900 GeV, and interpolating between 1100 and 1200 GeV gives about 1158 GeV. However, the tool is a single-channel Kolb-Turner scaling calibrated to 1.1 TeV. The 1158 GeV crossing is a calibration artefact, not a higgsino; high: The scan counts 144 of 289 points as viable. That count uses the same relic 'passes' flag that returns True at 7-25 sigma tension, so it probably includes points that are far from the Planck abundance. It also relies on the p-wave-suppressed indirect-detection model, which is wrong for a higgsino. T; medium: The best fit at m = 988 GeV, delta = 350 keV fits the LZ event, not the thermal relic. With the tool's own relic curve, 988 GeV gives Omega h^2 of about 0.087, which is under-abundant. The best fit is therefore not a full-abundance thermal higgsino, and this should be stated.; high: The count falls from 744.6 at 300 keV to 1.97 at 350 keV to 0 at 380 keV. That steep drop comes from the hard cutoff v_esc + v_E = 776 km/s. The literature uses about 544 + 250 km/s, roughly 794 km/s (giving delta_max of about 385 keV), and fits delta of about 377 keV (2609.02608) or 339 keV (2609.2; medium: The elastic leg of the question has not been answered with sourced numbers: the 1 TeV LZ/XENONnT spin-independent limit, the loop-level higgsino cross section, and the neutrino fog. The 2026 IDs (q-03) are still unverified. The indirect-detection conclusion relies on a tool that treats the higgsino

- `c-03` (wave 2, wave): **continue**; contested `f-011`, `f-012`, `f-013`, `f-014`; medium: The (794/776)^2 rescaling of delta(N=1) and delta(N=0.3) from 352.5/356.4 keV to about 369/373 keV is a heuristic. It assumes the whole N(delta) curve stretches with v_max^2, but near the kinematic edge the count depends on how the halo tail is populated between v_min and v_max, not only on where th; low: The 350 keV point (N = 1.97, peak E_R

= 315 keV) and the 300 keV point (744.6) come from wave-1 calls under d-02 and are not among this finding's evidence ids. The interpolations check out: ln-linear between 1.97 and 0.494 gives 352.45 keV, and between 0.494 and 0.085 gives 356.4 keV. The finding sh; high: The board brief still says 'I did not compute any relic densities... No inelastic-DM or relic-density tools were available'. This contradicts d-01, which computed Omega h^2 at five masses (crossing at about 1158 GeV), and d-02 and d-05, which computed LZ event counts. The brief must be rewritten bef; high: The solar-capture bound delta > 566 keV (2609.02775) carries the bottom line: it is the reason the sub-MeV LZ interpretation is disfavoured. Yet no abstract excerpt containing it has been posted. 2609.06760, 2609.15321 and 2610.00226 are also unverified. OpenAlex search_arxiv does work, so abstract-; medium: Neither the elastic loop-level pure-higgsino SI cross section (about 1e-48 to 1e-49 cm^2) nor the xenon elastic limit at 1.1 TeV is sourced. The roughly 1e-45 cm^2 figure is an extrapolation. As a result, the elastic leg of 'where does the higgsino sit relative to current limits' is unsupported.

- `c-04` (wave 2, report): **revise**; high: unsupported: | 900 | 3.73e-26 | 0.0731 | 39.1σ | — The board gives no per-point tension values. The only figures it states are 9.8σ at 1100 GeV and an overall range of 7–25σ. A 39.1σ tension falls outside that range, so the 'as reported by the tool' column is not supported by the ledger. The 25.1σ, ; high: unsupported: | 300 | 744.6 | 270 | 701 | — The board gives N_LZ = 744.6 at 300 keV but no peak E_R (270 keV) and no v_min (701 km/s) for that splitting. Board v_min values exist only for 350 keV (762), 360 keV (774), 365 keV (781) and 380 keV (799).; high: unsupported: | 355 | 0.494 | 320 | 768 | — The v_min of 768 km/s at 355 keV is not on the board. Likewise, v_min = 787 (370 keV), 793 (375 keV), 945 (500 keV), 1025 (566 keV) and 1189 km/s (700 keV) in the same column appear nowhere in the findings or direction results.; high: unsupported: x_f ≈ 24.4–24.8 — The board states only x_f of about 24.5. The 24.4–24.8 range is not sourced.; high: unsupported: It also shows how narrow and cutoff-dependent that window is, which explains why the literature fits scatter between 339 and 377 keV. — This is an overclaim. The board says the 339 keV value is a normalization-free sideband optimum and is not comparable to the tool's count. It also says

- `c-05` (wave 0, plan): **revise**; high: The gaugino-scale estimate in the rationale is off by about three orders of magnitude. Using delta0 ~ m_Z^2/M, a splitting of 350-570 keV needs M ~ 1e7 GeV (about 1e4 TeV), not 10-30 TeV. Gauginos at 10-30 TeV give splittings of order 0.3-1 GeV. The formula also depends on tan(beta) and the relative; medium: Step 4 ('Optionally call predict_sterile_nu_grid ... no') is not a step; delete it. Before spending about 10 calls on compute_relic_density, confirm with describe_tools that it accepts a sigma_v_cm3_s argument. Also add the Planck-uncertainty propagation the brief asks for. With Omega proportional t; medium: The direction calls get_paper_figures on 7 papers, plus search, predict_recoil and cite. That is about 11-12 calls with no room for retries. Order the papers by priority: LZ 2024, then the higgsino loop sigma_SI source, then the fog, then XENONnT 2025, then the first runs. The first-run comparison m; medium: The predict_recoil check at delta=0 is only meaningful if the tool includes Z-mediated vector scattering. Confirm this with describe_tools first. The phrase 'forces a splitting of at least about 100 keV' is inconsistent with the 345-385 keV kinematic window used in d-09. For m = 1.1 TeV on xenon (mu; medium: The budget is exceeded. Five papers at download+read+cite each is about 15 calls, before any fallback. Prioritize 1807.06209 and two of the mass papers that use different physics (for example Sommerfeld-only versus Sommerfeld plus bound-state formation), and mark the rest as optional. Quote the Plan; low: The cite-failure fallback logic duplicates d-07. Assign the single cite() API probe to one direction and have the other reuse its result to save calls.

- `c-06` (wave 1, wave): **continue**

- `c-07` (wave 2, wave): **continue**; contested `f-028`; medium: The 1-2% (v_esc about 555 km/s) requirement uses the overall kinematic endpoint, mu v_max^2/2 = 368 keV. That endpoint applies at the optimal recoil energy E_R = mu*delta/m_N, not at 248 keV. The relevant quantity is v_min at the 248 keV event, which is (m_N E_R/mu + delta)/sqrt(2 m_N E_R). This for; medium: The v0 dependence is not quantified. Near the endpoint, the rate depends exponentially on the tail of the velocity distribution, so v0, v_esc truncation and the choice of SHM versus non-Maxwellian halo can shift the N_LZ = 1 splitting by several keV even at fixed v_max. Without an independent rate c; low: Defining the window by N_LZ about 1 is a heuristic. With 1 observed event, a proper statement would use the Poisson likelihood plus the event's energy-reconstruction uncertainty: an approximate band in delta of N_LZ between about 0.05 and 4.7, not a single crossing point. The window width of about 1; medium: The formula keeps 1/(M_i - |mu|), but the estimate uses 1/M. At M = 4 TeV and mu = 1.1 TeV this is a correction of about 40% in amplitude and about 2x in sigma_SI. The tanbeta factor (1 + sgn(mu) sin2beta)^2 ranges from 0 to 4, which moves the M at which LZ is reached between 0 and about 8 TeV. The ; low: cite() was not run for 1410.4549, 2606.31013 or 2609.09830, so the equation references are unverified paraphrases from arxiv_agent. The radiative correction to the neutral splitting ('tens of percent') was not quantified, although it maps directly onto the inferred gaugino scale.; low: The evidence excerpts show the echoed sigma_v but not the input value. The claim that sigma_v_cm3_s is silently ignored with is_higgsino=true therefore rests on the agent's report of what was passed. It is plausible, since tc-71 echoes 2.5e-26*(1100/1159)^2, but it is not directly visible. The tool

- `c-08` (wave 2, report): **revise**; high: unsupported: where freeze-out occurs at T_f ≈ 47 GeV and x_f ≈ 24.5 — No freeze-out temperature or x_f appears in any finding, direction result or critique. The board records only Omega h^2 values and the 1159-1160 GeV crossing.; high: unsupported: with peak on June 2 and trough on December 2. The rate on the event date (September 1) is 0.225 of the peak rate. — d-11 reports only the 98.5% fractional amplitude and the June 2 peak (tc-79). The trough date, the September 1 event date and the 0.225 rate ratio are not on the board.; high: unsupported: The local significance is 3.4σ and the global significance 2.6σ. — The board (d-08, f-017) cites 2609.02608 only for the mass near 1.1 TeV and for delta ≈ 377 keV reproducing the 248 keV event. No local or global significance is recorded.; high: unsupported: Searches also returned work on bound-state formation and unitarity for TeV WIMPs (arXiv:1904.11503, arXiv:1407.7874) — Neither arXiv ID appears anywhere on the board or ledger.; high: unsupported: The paper's main subject is non-thermal production by late-decaying scalars. — The board characterises 2410.15733 only as giving about 1.2 TeV with coannihilation and Sommerfeld. Nothing on the board describes its main subject.; high: unsupported: Chatterjee et al. (arXiv:2609.09830) — The board never names the authors of 2609.09830. It is listed only by title and as 'a 2026 paper'.

C Figures not shown in the main text

Produced during the session; captions carry their provenance.




Figure 1: Figure 1: Data from WS2024, in black with all analysis cuts applied, and in gray outside the FV without veto coincidence cuts. Red crosses and blue circles represent events that are vetoed by the prompt and delayed veto coincidence cuts, respectively. The dashed line shows the active volume, averaged over azimuth, and the two solid lines depict the FV, at the azimuths of its smallest and largest radial extent. The drift field affects the position reconstruction such that the observed radius of the TPC varies as a function of both drift time and azimuthal angle. [arXiv:2410.17036] Reproduced from arXiv:2410.17036 (extracted with `get_paper_figures`).




Figure 2: Figure S4: Distribution from which the salt events are drawn. The parameters are described in the text. [arXiv:2410.17036] Reproduced from arXiv:2410.17036 (extracted with `get_paper_figures`).




Figure 3: Figure 2: Distribution of events in \mathrm{cS1}–\mathrm{cS2} for \mathrm{SR1a} (top) and \mathrm{SR1b} (bottom). All data points are represented as pie charts indicating the fraction of the likelihood from the best-fit model including a 200 \mathrm{GeV}/c^{2} WIMP signal, evaluated at the data point. The scatter size is scaled according to the WIMP likelihood fraction for visualization only. In addition, the one (dark shaded, solid line) and two (light shaded, dotted line) \sigma contours of the ER, AC, surface background, and 200 \mathrm{GeV}/c^{2} WIMP signal are shown. The NR background fo [arXiv:2502.18005] Reproduced from arXiv:2502.18005 (extracted with `get_paper_figures`).




Figure 4: Figure S1: The cS1–cS2 distribution for \mathrm{SR1}, from Figure 2. The model was modified to include the LM+LN and LL 2\upnu\mathrm{ECEC} peaks with unconstrained \mathrm{CY} while excluding the WIMP component. The regions corresponding to the first (solid gray) and second (hatched gray) steps of unblinding are also indicated. [arXiv:2502.18005] Reproduced from arXiv:2502.18005 (extracted with `get_paper_figures`).