Supplementary material: Thermal Higgsino at 1.1 TeV: Relic Density, Inelastic LZ Rates and Neutrino Limits

HEP-Genesis-Agent session record
Project: ramp-research-higgsino-dd

2026-11-02

Supplementary material, generated mechanically from the session record: it lists exactly what ran, in order, without interpretation.

A Implementation details

This appendix is generated mechanically from the session record; it lists exactly what ran, in order, without interpretation.

A.1 Session

- Source: cli client; conversation `9e56c682`
- Started: 2026-10-16 11:03:59; report generated: 2026-11-02 18:32:31
- Project: ramp-research-higgsino-dd (`/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd`)
- Compute execution site(s) selected in the session: local
- 5 user turn(s), 494 tool call(s) (12 returned errors), 13 figure(s), 13 data file(s)

A.2 Models

Stage	Model	Endpoint
quick	`muse-spark-bf16`	`https://inference-api.alcf.anl.gov/resource_server/minerva/api/v1`
report	`muse-spark-bf16`	`https://inference-api.alcf.anl.gov/resource_server/minerva/api/v1`
coder	`muse-spark-bf16`	`https://inference-api.alcf.anl.gov/resource_server/minerva/api/v1`

A.3 MCP servers

Server	Transport	Location
hep-research-aid	stdio	`python -m hep_genesis.domain.mcp_server`

Server	Transport	Location
alcf-tools	stdio	`python -m hep_genesis.iri.alcf.mcp_server`
nersc-tools	stdio	`python -m hep_genesis.iri.nersc.mcp_server`
paper-search	stdio	`uvx -with mcp<2 paper-search-mcp`
web-search	stdio	`uvx duckduckgo-mcp-server`
cosmic-emulators	http	`https://cosmic.77-42-88-84.sslip.io/mcp`

A.4 Tool-call log

Chronological; arguments and results are truncated for space (the full record is in the project's `provenance/` folder).

#	Tool	Arguments	Outcome
1	`hep-research-aid list_agent_files`	`{}`	`[]`
2	`hep-research-aid list_input_files`	`{}`	`{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }`
3	`hep-research-aid list_datasets`	`{}`	`{'count': 0, 'datasets': []}`
4	`hep-research-aid list_files`	`{}`	`Files in .: - .git - .gitignore - ISSUES.md - SCOPE.md - docs - downloads - logs - projects - reports - research-results - runs - scripts - venvs - workspace`
5	`hep-research-aid get_dispatch`	`{}`	`Dispatch: local execution.`
6	`local search_arxiv_abstracts`	`{"query": "higgsino dark matter thermal relic 1.1 TeV direct detection", "max_results": 8}`	`[2302.02905v2] Third family quasi-Yukawa unification: Higgsino dark matter, NLSP gluino and all that Abstract: We explore the implications of third family ($t-b-\tau$) quasi-Yukawa unification (QYU) for ...`

#	Tool	Arguments	Outcome
7	local search_arxiv_ abstracts	{"query": "pure higgsino Sommerfeld enhancement relic abundance mass", "max_ results": 5}	[1008.2905v2] Thermal abundance of non-relativistic relics with Sommerfeld enhancement Abstract: We propose an analytic treatment for computing the relic abundances of non-relativistic particles whose...
8	local search_arxiv_ abstracts	{"query": "higgsino inelastic dark matter mass splitting electroweak loop scattering", "max_results": 6}	[0811.4682v1] Precision Electroweak Measurements and Constraints on the Standard Model Abstract: This note presents constraints on Standard Model parameters using published and preliminary precision...
9	local search_arxiv_ abstracts	{"query": "LZ 2024 WIMP search results spin-independent limits", "max_results": 4}	[1509.02910v2] LUX-ZEPLIN (LZ) Conceptual Design Report Abstract: The design and performance of the LUX-ZEPLIN (LZ) detector is described as of March 2015 in this Conceptual Design Report. LZ is a sec...
10	hep-research-aid search_arxiv	{"query": "pure higgsino dark matter spin-independent cross section two-loop electroweak direct detection", "max_results": 6}	Found 6 papers matching query: 'pure higgsino dark matter spin-independent cross section two-loop electroweak direct detection' (source: arXiv API) Paper 1: arXiv ID: 1901.09751v3 Title: Pseudo Nambu...
11	hep-research-aid search_arxiv	{"query": "LZ dark matter search results 4.2 tonne-year exposure", "max_ results": 4}	Found 4 papers matching query: 'LZ dark matter search results 4.2 tonne-year exposure' (source: arXiv API) Paper 1: arXiv ID: 2410.17036v3 Title: Dark Matter Search Results from 4.2 Tonne-Years of Ex...

#	Tool	Arguments	Outcome
12	hep-research-aid search_arxiv	{"query": "XENONnT WIMP search first results nuclear recoil", "max_results": 4}	Found 4 papers matching query: 'XENONnT WIMP search first results nuclear recoil' (source: arXiv API) Paper 1: arXiv ID: 0809.3137v1 Title: The Frontiers of Nuclear Science, A Long Range Plan Authors...
13	hep-research-aid search_arxiv	{"query": "higgsino dark matter neutrino floor next generation xenon detectors reach", "max_results": 6}	Found 6 papers matching query: 'higgsino dark matter neutrino floor next generation xenon detectors reach' (source: arXiv API) Paper 1: arXiv ID: 1211.7222v1 Title: Direct dark matter detection: the ...
14	local get_arxiv_ metadata	{"arxiv_id": "2609.09830"}	arXiv ID: 2609.09830 Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Al...
15	local search_arxiv_ abstracts	{"max_results": 5, "query": "Hisano direct detection electroweak-interacting dark matter wino higgsino gluon contribution"}	[1004.4090v1] A complete calculation for direct detection of Wino dark matter Abstract: In the anomaly-mediated supersymmetry (SUSY) breaking scenario, neutral gaugino of SU(2) _L multiplet, Wino, can ...
16	local search_arxiv_ abstracts	{"max_results": 5, "query": "XENONnT WIMP dark matter search combined science runs 2025"}	[1211.0310v1] Large Synoptic Survey Telescope: Dark Energy Science Collaboration Abstract: This white paper describes the LSST Dark Energy Science Collaboration (DESC), whose goal is the study of dark...

#	Tool	Arguments	Outcome
17	local search_arxiv_ abstracts	{"max_results": 4, "query": "Planck 2018 results cosmological parameters"}	[0604069v1] The Scientific Programme of Planck Abstract: For 40 years, the cosmic microwave background (CMB) has been the most important source of information about the geometry and contents of the ...
18	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_ data()." }
19	local download_arxiv_ full_text	{"arxiv_id": "1104.0228", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1104.0228.txt
20	local download_arxiv_ full_text	{"arxiv_id": "1411.6930", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1411.6930.txt
21	local search_arxiv_ abstracts	{"query": "higgsino spin-independent cross section electroweak loop pure higgsino nucleon", "max_results": 6}	[1812.08019v2] Higgsino Dark Matter in an economical Scherk-Schwarz setup Abstract: We consider a minimal natural supersymmetric model based on an extra dimension with supersymmetry breaking provided ...
22	local download_arxiv_ full_text	{"arxiv_id": "2609.09830", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2609.09830.txt

#	Tool	Arguments	Outcome
23	local search_arxiv_ abstracts	{"query": "higgsino relic abundance Sommerfeld bound state thermal mass 1.1 TeV", "max_results": 8}	[1008.2905v2] Thermal abundance of non-relativistic relics with Sommerfeld enhancement Abstract: We propose an analytic treatment for computing the relic abundances of non-relativistic particles whose...
24	local download_arxiv_ full_text	{"arxiv_id": "2410.17036", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2410.17036.txt
25	local download_arxiv_ full_text	{"arxiv_id": "2606.31013", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2606.31013.txt
26	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1411.6930.txt", "start": 0, "length": 12000}	[chars 0-12000 of 109138] (more available, next start=12000) Title: Heavy neutralino relic abundance with Sommerfeld enhancements - a study of pMSSM scenarios Authors: M. Beneke, C. Hellmann, P. Ruiz...
27	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1411.6930.txt", "start": 30000, "length": 16000}	[chars 30000-46000 of 109138] (more available, next start=46000) iplet minimal dark matter model with the same χ 0 1 mass. In the pure-wino model the mass splitting between the χ 0 1 and χ $\pm$ 1 has to...

#	Tool	Arguments	Outcome
28	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1104.0228.txt", "start": 0, "length": 5000}	[chars 0-5000 of 46034] (more available, next start=5000) Title: Direct Detection of Electroweak-Interacting Dark Matter Authors: Junji Hisano, Koji Ishiwata, Natsumi Nagata, Tomohiro Takesako Year: ...
29	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2609.09830.txt", "length": 14000}	[chars 0-14000 of 71943] (more available, next start=14000) Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatt...
30	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt", "length": 14000}	[chars 0-14000 of 63686] (more available, next start=14000) Title: Complementary Probes of Light Higgsinos: Electroweak Precision Measurements and Dark Matter Direct Detection Authors: Koichi Hamaguc...
31	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1411.6930.txt", "length": 10000, "start": 46000}	[chars 46000-56000 of 109138] (more available, next start=56000) itive. The enhancement in the χ 0 1χ + 1 channel also saturates as its on-shell production threshold is approached. This should be the...
32	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt", "start": 14000, "length": 6000}	[chars 14000-20000 of 63686] (more available, next start=20000) andv$\simeq$246 GeVis the Higgs VEV. Additional electroweak oblique parameters introduced in Ref. [48], such as?U,V, andX, are not generated a...

#	Tool	Arguments	Outcome
33	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt", "start": 30000, "length": 14000}	[chars 30000-44000 of 63686] (more available, next start=44000) g2 tan2θW 120π2 M 2 W µ2 .(39) These expressions imply the simple relation Y$\simeq$tan 2θWW ,(40) which explains the nearly linear correlatio...
34	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1104.0228.txt", "length": 9000, "start": 22000}	[chars 22000-31000 of 46034] (more available, next start=31000) (n2 - (4Y 2 + 1))/ 8 to their result of the Wino DM case. (as the ?rst term in Eq. (22)). Even for the Z-boson exchange contribution,...
35	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt", "length": 10000, "start": 20000}	[chars 20000-30000 of 63686] (more available, next start=30000) .48 1.55) $\times$10-5 ,(26) and ρ= ? ??????? 1 0.856 0.315 0.672 0.856 1 0.154 0.352 0.315 0.154 1 0.221 0.672 0.352 0.221 1 ? ??????? ,(27)...
36	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2609.09830.txt", "length": 14000, "start": 14000}	[chars 14000-28000 of 71943] (more available, next start=28000) 4) whereAandZrepresent the mass number and the atomic number of the target nucleus, respectively,µ p stands for the reduced mass of t...
37	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1104.0228.txt", "length": 5000, "start": 31000}	[chars 31000-36000 of 46034] (more available, next start=36000) [(n2 - (4 Y2 + 1))/8] $\times$10-9 Higgs mass [GeV] gluon(W) -1 -0.5 0 0.5 1 115 200 300 400 500 fp/mp in units of Y2 $\times$10-9 Higgs ma...

#	Tool	Arguments	Outcome
38	local search_arxiv_ abstracts	{"max_results": 8, "query": "electroweak doublet dark matter thermal mass higgsino 1.1 TeV relic abundance bound state formation"}	[1407.7874v2] Bound-state formation for thermal relic dark matter and unitarity Abstract: We show that the relic abundance of thermal dark matter annihilating via a long-range interaction, is signific...
39	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2609.09830.txt", "length": 16000, "start": 28000}	[chars 28000-44000 of 71943] (more available, next start=44000) inally, we investigate whether the inelastic rate can exceed the elastic rate to be the domi- nant DD channel in some parameter space r...
40	hep-research-aid get_paper_figures	{"paper_id": "1411.6930", "figures": "7", "output_ dir": "/Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	{"status": "success", "files": ["/Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ figures_1411.6930/ fig07.png"], "message": "Extracted 1 of 14 figures from... [Fig. 6]
41	local search_arxiv_ abstracts	{"max_results": 4, "query": "Hill Solon Standard Model anatomy WIMP dark matter direct detection heavy higgsino cross section"}	[1912.07795v2] Direct detection rate of heavy Higgsino-like and Wino-like dark matter Abstract: A large class of viable dark matter models contain a WIMP candidate that is a component of a new electro...
42	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2609.09830.txt", "length": 12000, "start": 44000}	[chars 44000-56000 of 71943] (more available, next start=56000) n. Additionally, the inelastic event rate is Boltzmann suppressed due to the requirement of a minimum -- Page 16 -- 16 TABLE VII. Nu...

#	Tool	Arguments	Outcome
43	local search_arxiv_ abstracts	{"max_results": 6, "query": "Closing the window on WIMP dark matter thermal mass electroweak multiplets bound states"}	[0509008v3] Precision Electroweak Measurements on the Z Resonance Abstract: We report on the final electroweak measurements performed with data taken at the Z resonance by the experiments operating ...
44	local get_arxiv_ metadata	{"arxiv_id": "2607.20588"}	arXiv ID: 2607.20588 Title: Unveiling the Vanishing Higgsino-Nucleon Scattering in the MSSM at Next-to-Leading Order Authors: Subhadip Bisal, Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Rahul P...
45	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2410.17036.txt", "length": 10000, "start": 0}	[chars 0-10000 of 63958] (more available, next start=10000) Title: Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment Authors: J. Aalbers, D. S. Akerib, A. ...
46	local download_arxiv_ full_text	{"arxiv_id": "hep-ph/ 0610249", "output_dir": "/Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ hep-ph_0610249.txt
47	local download_arxiv_ full_text	{"arxiv_id": "1912.07795", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1912.07795.txt

#	Tool	Arguments	Outcome
48	board post_finding	{"claim": "Kinematic threshold for inelastic up-scattering on xenon (my arithmetic): m_A = 131 x 0.9315 GeV = 122.0 GeV; mu_A = 108.7 GeV (m=1.0 TeV) and 109.8 GeV (m=1.1 TeV); v_max = 544+232 = 776 km/s, beta^2 = (776/2...	Recorded finding f-001.
49	board post_finding	{"claim": "2609.09830 does not quote an LZ-excluded delta_m0 range. Its five benchmarks (|mu| = 0.61-1.40 TeV; |M1|, M2 about 5.8-42 TeV; tan beta 10.9-26.5) have one-loop on-shell splittings delta = 8.9-36.4 keV. Each p...	Recorded finding f-002.
50	board post_finding	{"claim": "2606.31013: the SI coupling comes from Higgs exchange, f_q ~ -(g^2/8M_h^2)[1/(M2-|mu|) + tan^2 theta_W/(M1-|mu|)](1 + sgn(mu) sin 2beta), so it is suppressed at large gaugino mass and at small tan beta with mu...	Recorded finding f-003.
51	local download_arxiv_full_text	{"arxiv_id": "1807.06209", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	Downloaded: /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt
52	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2410.17036.txt", "length": 5000, "start": 10000}	[chars 10000-15000 of 63958] (more available, next start=15000) oils induced by weakly interacting massive particle (WIMP) dark matter using the LUX-ZEPLIN (LZ) two-phase xenon time projection chambe...

#	Tool	Arguments	Outcome
53	local cite	{"arxiv_id": "2609.09830", "note": "[\"mass splitting∼O(10-100) MeV or even smaller, for very large gaugino mass parameters\", \"In scenarios where ?χ0\", \"All benchmark points are consistent with the bounds from the LZ...	Cited: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter (\cite{arxiv260909830})
54	local cite	{"arxiv_id": "2606.31013", "note": "[\"dark matter direct detection experiments are particularly sensitive to scenarios with larger charged-neutral mass splittings induced by higgsino-gaugino mixing\", \"This expression ...	Cited: Complementary Probes of Light Higgsinos: Electroweak Precision Measurements and Dark Matter Direct Detection (\cite{arxiv260631013})
55	board post_question	{"text": "The tree-level formula delta_m0 ≈ m_Z^2 s_W^2/M1 + m_Z^2 c_W^2/M2 has not been sourced yet. Can a quotable reference be found, for example Ref. [28] of 2609.09830, Eq. (9)? Taken at face value with same-sign M2...	Recorded question q-05.
56	board post_finding	{"claim": "Hisano et al. (1104.0228) find the leading-order electroweak-loop SI cross section of pure EW-multiplet DM (including the higgsino, n=2, Y=1/2, evaluated at M=1 TeV) to be O(1e-46 to 1e-48) cm^2, suppressed by...	Recorded finding f-004.
57	board post_finding	{"claim": "LZ 2024 (2410.17036, 4.2 tonne-years): the strongest SI exclusion is 2.2e-48 cm^2 (90% CL) and the best median sensitivity is 5.1e-48 cm^2, both at 40 GeV/c^2. The abstract quotes no value at 1 TeV.", "evidenc...	Recorded finding f-005.

#	Tool	Arguments	Outcome
58	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1912.07795.txt", "length": 4000, "start": 0}	[chars 0-4000 of 37078] (more available, next start=4000) Title: Direct detection rate of heavy Higgsino-like and Wino-like dark matter Authors: Qing Chen, Richard J. Hill Year: 2019 arXiv ID: 1912.0...
59	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1912.07795.txt", "length": 6000, "start": 24000}	[chars 24000-30000 of 37078] (more available, next start=30000) tially have a larger than expected impact. We investigate this possibility here, using the model presented in Refs. [28, 29] which incl...
60	board post_finding	{"claim": "No relic-density or inelastic-DM compute tool and no input data tables are available: list_input_files returned an empty file list, and the catalog contains only statistics/histogram, test-job and literature t...	Recorded finding f-006.
61	board post_finding	{"claim": "Beneke, Hellmann and Ruiz-Femenia (arXiv:1411.6930, Sec. 3) report Omega h^2 = 0.127 (perturbative) and 0.126 (Sommerfeld, with off-diagonal rates) for a pure-higgsino model compared against the higgsino-like ...	Recorded finding f-007.
62	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/hep-ph_0610249.txt", "length": 6000, "start": 0}	[chars 0-6000 of 21259] (more available, next start=6000) Title: Non-perturbative Effect on Thermal Relic Abundance of Dark Matter Authors: Junji Hisano, Shigeki Matsumoto, Minoru Nagai, Osamu Saito,...

#	Tool	Arguments	Outcome
63	board post_finding	{"claim": "Chen and Hill (1912.07795) include the O(mW/M) power corrections, perturbative QCD to NNNLO and ChEFT nuclear effects. They find that the pure-higgsino SI cross section has an upper limit below 1e-48 cm^2 in t...	Recorded finding f-008.
64	local cite	{"arxiv_id": "1912.07795", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib", "note": "[\"The upper limit for the pure doublet is less than 10-48 c...	Cited: Direct detection rate of heavy Higgsino-like and Wino-like dark matter (\cite{arxiv191207795})
65	local cite	{"arxiv_id": "2410.17036", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib", "note": "[\"The strongest SI exclusion set is 2 .2 × 10-48 cm2 at the...	Cited: Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment (\cite{arxiv241017036})
66	local cite	{"arxiv_id": "1411.6930", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib", "note": "[\"the relic abundances in the pure-higgsino model, obtained ...	Cited: Heavy neutralino relic abundance with Sommerfeld enhancements - a study of pMSSM scenarios (\cite{arxiv14116930})

#	Tool	Arguments	Outcome
67	board post_question	{"text": "Still open from d-02: the XENONnT 2303.14729 SI limit at 1 TeV, the actual LZ 2024 limit-curve value at 1 TeV (read from its figure rather than linearly rescaled from 40 GeV), and a numeric xenon neutrino-fog v...	Recorded question q-06.
68	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/hep-ph_0610249.txt", "length": 6000, "start": 13000}	[chars 13000-19000 of 21259] (more available, next start=19000) the non- perturbative e?ect when m is larger than $\sim$ 1 TeV. A signi?cant enhancement is shown at m $\sim$ 2.4 GeV. This originates in the bou...
69	board post_finding	{"claim": "Hisano et al. (hep-ph/0610249) expect the Sommerfeld effect on the higgsino-like relic abundance to be roughly 10 percent, compared with a 50 percent reduction for the wino. 1411.6930 later found the net pure-...	Recorded finding f-009.
70	local get_arxiv_metadata	{"arxiv_id": "2606.31013"}	arXiv ID: 2606.31013 Title: Complementary Probes of Light Higgsinos: Electroweak Precision Measurements and Dark Matter Direct Detection Authors: Koichi Hamaguchi, Natsumi Nagata, Genta Osaki Year: 20...
71	local get_arxiv_metadata	{"arxiv_id": "2609.09830"}	arXiv ID: 2609.09830 Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Al...

#	Tool	Arguments	Outcome
72	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt", "start": 0, "length": 4000}	[chars 0-4000 of 442128] (more available, next start=4000) Title: Planck 2018 results. VI. Cosmological parameters Authors: Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Bac...
73	hep-research-aid get_paper_figures	{"paper_id": "2410.17036", "max_figures": 6, "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	{"status": "success", "files": ["/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/figures_2410.17036/fig01.png", "/Users/nramachandra/hep-genesis-testing/p... [Fig. 10, 11]
74	local search_arxiv_abstracts	{"query": "higgsino relic abundance bound-state formation Sommerfeld thermal mass TeV", "max_results": 8}	[1611.08133v2] Effects of QCD bound states on dark matter relic abundance Abstract: We study scenarios where there exists an exotic massive particle charged under QCD in the early Universe. We calcula...
75	hep-research-aid get_paper_figures	{"paper_id": "2303.14729", "max_figures": 6, "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	{"status": "success", "files": ["/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/figures_2303.14729/fig01.png", "/Users/nramachandra/hep-genesis-testing/p... [Fig. 12, 13, 14]

#	Tool	Arguments	Outcome
76	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1912.07795.txt", "start": 0, "length": 3000}	[chars 0-3000 of 37078] (more available, next start=3000) Title: Direct detection rate of heavy Higgsino-like and Wino-like dark matter Authors: Qing Chen, Richard J. Hill Year: 2019 arXiv ID: 1912.0...
77	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt", "start": 8000, "length": 14000}	[chars 8000-22000 of 63686] (more available, next start=22000) ty between elec- troweak precision measurements and dark matter direct detection experiments in probing light higgsinos. Finally, Sec. 6...
78	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2609.09830.txt", "start": 5000, "length": 12000}	[chars 5000-17000 of 71943] (more available, next start=17000) 7, 38] presented the SUSY QCD corrections for the DD of ?χ0 1 DM.2 The DM- nucleon scattering cross-section at one-loop level for a gene...
79	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2609.09830.txt", "length": 10000, "start": 17000}	[chars 17000-27000 of 71943] (more available, next start=27000) e parton-level Lagrangian as, f inel N mN = X u,d, s λ? qfN q - 8π 9αs λ? gfN g (10) Note that, for simplicity, we consider scalar contr...
80	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt", "length": 12000, "start": 24000}	[chars 24000-36000 of 63686] (more available, next start=36000) elements and a detailed discussion of the associated theoretical uncertainties in dark matter-nucleon scattering cross sections, see R...

#	Tool	Arguments	Outcome
81	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1912.07795.txt", "start": 22000, "length": 7000}	[chars 22000-29000 of 37078] (more available, next start=29000) 2m2 c, and 1 .0 GeV$\leq$ µ0 $\leq$ 1.4 GeV, as in Ref. [18, 27] . Uncertainties from neglected 1/M2 and higher order power corrections are estima...
82	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 1807.06209.txt", "start": 5500, "length": 4500}	[chars 5500-10000 of 442128] (more available, next start=10000) D. Spencer82, R. Sunyaev72, 91, A.-S. Suur-Uski24, 40, J. A. Tauber36, D. Tavagnacco43,33, M. Tenti46, L. To?olatti17, 41, M. Tomasi32,45...
83	local search_arxiv_ abstracts	{"max_results": 8, "query": "last complex WIMPs standing electroweak multiplet thermal mass bound states"}	[0811.4682v1] Precision Electroweak Measurements and Constraints on the Standard Model Abstract: This note presents constraints on Standard Model parameters using published and preliminary precision...
84	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2606.31013.txt", "length": 2200, "start": 21900}	[chars 21900-24100 of 63686] (more available, next start=24100) dark matter and nucleon masses, respectively, andfN is the effective SI coupling between dark matter and the nucleon. For higgsino dar...
85	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2609.09830.txt", "length": 14000, "start": 27000}	[chars 27000-41000 of 71943] (more available, next start=41000) lision will be similar to the elastic one up to a factor proportional to the square of ?χ0 1 ?χ0 2hj couplings. • Consequently, we compu...

#	Tool	Arguments	Outcome
86	local search_arxiv_ abstracts	{"max_results": 6, "query": "XLZD design book sensitivity spin-independent neutrino fog TeV"}	[0211001v1] Very Long Baseline Neutrino Oscillation Experiment for Precise Determination of Oscillation Parameters and Search for nu_mu -> nu_e Appearance and CP Violation Abstract: The possibility ...
87	local get_arxiv_ metadata	{"arxiv_id": "2107.09688"}	arXiv ID: 2107.09688 Title: Closing the window on WIMP Dark Matter Authors: Salvatore Bottaro, Dario Buttazzo, Marco Costa, Roberto Franceschini, Paolo Panci, Diego Redigolo, Ludovico Vittorio Year: 2...
88	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2410.17036.txt", "length": 8000, "start": 20000}	[chars 20000-28000 of 63958] (more available, next start=28000) the tuning of the nest 2.4.0 [18] model for detector response, in which the scin- tillation photon and ionization electron gains, g1 an...
89	local get_arxiv_ metadata	{"arxiv_id": "2205.04486"}	arXiv ID: 2205.04486 Title: The last Complex WIMPs standing Authors: Salvatore Bottaro, Dario Buttazzo, Marco Costa, Roberto Franceschini, Paolo Panci, Diego Redigolo, Ludovico Vittorio Year: 2022 Abs...
90	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437/ 2609.09830.txt", "length": 9000, "start": 41000}	[chars 41000-50000 of 71943] (more available, next start=50000) s-section more than the corresponding elastic cross- section through constructive interference. As a result, inelastic scattering sligh...

#	Tool	Arguments	Outcome
91	board post_finding	{"claim": "Chen and Hill (1912.07795), verbatim from the Summary: \"The upper limit for the pure doublet is less than 10-48 cm2 in the TeV mass regime, consistent with the leading order estimate [18, 19], and lower than ...	Recorded finding f-010.
92	board post_finding	{"claim": "Correction to f-008: an upper bound below 1e-48 cm^2 compared with the LZ minimum of 2.2e-48 cm^ 2 (at 40 GeV) is a factor of more than 2.2 (more than 0.34 decades), not 2 orders of magnitude. The comparison th...	Recorded finding f-011.
93	board post_dead_end	{"what": "Reading the LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits at 1 TeV with get_ paper_figures", "why": "The HTML extraction returned none of the limit plots. For LZ it returned only Fig. 1 (fiducial-volum...	Recorded dead end x-01.
94	board post_question	{"text": "The XLZD Design Book (arXiv:2410.17137) has been identified but not read: what are its projected SI sensitivity and its xenon neutrino-fog value at about 1 TeV, and how do they compare with the pure-higgsino bo...	Recorded question q-07.
95	board post_finding	{"claim": "Planck 2018 (arXiv:1807.06209, abstract, page 1) states verbatim: \ "A combined analysis gives dark matter density Ωch2 = 0.120± 0.001\" (68% confidence region), quoted for base-LambdaCDM from polarization, tem...	Recorded finding f-012.

#	Tool	Arguments	Outcome
96	board post_finding	{"claim": "Bottaro et al., \"The last Complex WIMPs standing\" (arXiv: 2205.04486, 2022), is the independent recent calculation of thermal masses for complex electroweak n-plets, which includes the Y=1/2 doublet (the higg...	Recorded finding f-013.
97	board post_question	{"text": "What thermal mass, with its theory uncertainty, does arXiv: 2205.04486 (Bottaro et al. 2022, Sommerfeld plus bound-state formation) give for the Y=1/2 doublet in its thermal-mass table, and how far is it from th...	Recorded question q-08.
98	board post_dead_end	{"what": "Searching abstracts for a recent higgsino thermal mass with the query 'last complex WIMPs standing electroweak multiplet thermal mass bound states'.", "why": "The search returned only unrelated LEP electroweak-...	Recorded dead end x-02.
99	board post_finding	{"claim": "Metadata confirms both IDs: arXiv: 2606.31013 is 'Complementary Probes of Light Higgsinos: Electroweak Precision Measurements and Dark Matter Direct Detection' by Hamaguchi, Nagata and Osaki (2026), and arXiv:2...	Recorded finding f-014.
100	board post_finding	{"claim": "The full text of 2606.31013 supports the f-003 numbers. Eq. 13 reads Delta m+|rad ~ (1/2) alpha2 MZ sin^2 thetaW (1 - 3MZ/(2 pi M_chi+)), with the statement 'For M_chi+ > > MZ, this mass splitting asymptoticall...	Recorded finding f-015.

#	Tool	Arguments	Outcome
101	board post_finding	{"claim": "The full text of 2609.09830 supports the f-002 numbers. Table III gives one-loop on-shell delta = 36.37, 8.90, 16.39, 10.81 and 10.59 keV for BP1-BP5, with mu = -739.63, -606.64, -814.26, 1282.11 and 1400.0 Ge...	Recorded finding f-016.
102	board post_finding	{"claim": "2609.09830 does not write out the tree-level delta_m0 formula. It says only: 'as can be seen, for example, from Eq. (9) of Ref. [28], the corrections proportional to M_Z^2/M1 and M_Z^2/M2 depend on the relativ...	Recorded finding f-017.
103	board post_question	{"text": "What arXiv paper is Ref. [28] of 2609.09830, and does its Eq. (9) give the tree-level neutral higgsino splitting with explicit sign dependence on mu, M1 and M2? This is needed to re-evaluate the M1 required for...	Recorded question q-09.
104	hep-research-aid list_agent_files	{}	[]
105	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }
106	hep-research-aid list_datasets	{}	{'count': 0, 'datasets': []}
107	hep-research-aid list_files	{}	Files in .: - .git - .gitignore - .last-release-seen - ISSUES.md - SCOPE.md - docs - downloads - logs - projects - reports - research-results - runs - scripts - venvs - w...
108	hep-research-aid get_dispatch	{}	Dispatch: local execution.

#	Tool	Arguments	Outcome
109	local search_arxiv_ abstracts	{"query": "higgsino thermal relic 1.1 TeV dark matter direct detection", "max_ results": 8}	[2302.02905v2] Third family quasi-Yukawa unification: Higgsino dark matter, NLSP gluino and all that Abstract: We explore the implications of third family ($t-b-$\tau$) quasi-Yukawa unification (QYU) for ...
110	local search_arxiv_ abstracts	{"query": "pure higgsino dark matter mass splitting inelastic scattering LZ XENONnT", "max_results": 8}	[1509.02910v2] LUX-ZEPLIN (LZ) Conceptual Design Report Abstract: The design and performance of the LUX-ZEPLIN (LZ) detector is described as of March 2015 in this Conceptual Design Report. LZ is a sec...
111	local search_arxiv_ abstracts	{"query": "Sommerfeld enhancement higgsino relic abundance bound state formation", "max_results": 6}	[0910.5221v2] Relic density and CMB constraints on dark matter annihilation with Sommerfeld enhancement Abstract: We calculate how the relic density of dark matter particles is altered when their an...
112	local get_arxiv_ metadata	{"arxiv_id": "2609.01590"}	arXiv ID: 2609.01590 Title: TeV Higgsino Dark Matter from LZ Nuclear Recoil to Fermi-LAT Gamma Rays Authors: Lei Wu, Yang Zhang, Bin Zhu Year: 2026 Abstract: The excess of high-energy nuclear recoils,...
113	local get_arxiv_ metadata	{"arxiv_id": "2609.07807"}	arXiv ID: 2609.07807 Title: Not so good νs for Higgsino dark matter as LZ excess: stringent limits from Super-Kamiokande and IceCube Authors: Debajit Bose, Akash Kumar Saha, Deep Jyoti Das, Rinchen ...

#	Tool	Arguments	Outcome
114	local get_arxiv_ metadata	{"arxiv_id": "2609.09830"}	arXiv ID: 2609.09830 Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Al...
115	local search_arxiv_ abstracts	{"max_results": 6, "query": "LZ WIMP search results exposure spin-independent limit 2024"}	[1509.02910v2] LUX-ZEPLIN (LZ) Conceptual Design Report Abstract: The design and performance of the LUX-ZEPLIN (LZ) detector is described as of March 2015 in this Conceptual Design Report. LZ is a sec...
116	hep-research-aid search_arxiv	{"query": "LZ dark matter search results 4.2 tonne-year exposure", "max_ results": 5}	Found 5 papers matching query: 'LZ dark matter search results 4.2 tonne-year exposure' (source: arXiv API) Paper 1: arXiv ID: 2410.17036v3 Title: Dark Matter Search Results from 4.2 Tonne-Years of Ex...
117	local search_arxiv_ abstracts	{"query": "XENONnT WIMP dark matter search nuclear recoil results", "max_ results": 5}	[0504031v1] Limits on WIMP cross-sections from the NAIAD experiment at the Boulby Underground Laboratory Abstract: The NAIAD experiment (NaI Advanced Detector) for WIMP dark matter searches at the B...
118	local search_arxiv_ abstracts	{"query": "electroweak multiplet dark matter thermal mass higgsino 1.1 TeV wino 2.8 TeV", "max_ results": 6}	[1901.02987v2] Discovery reach for wino and higgsino dark matter with a disappearing track signature at a 100 TeV pp collider Abstract: Within the theory of supersymmetry, the lightest neutralino is...

#	Tool	Arguments	Outcome
119	local search_arxiv_ abstracts	{"query": "LZ high energy nuclear recoil event inelastic dark matter higgsino mass splitting", "max_results": 6}	[0510346v1] The Dark Energy Survey Abstract: We describe the Dark Energy Survey (DES), a proposed optical-near infrared survey of 5000 sq. deg of the South Galactic Cap to ~24th magnitude in SDSS gr...
120	local search_arxiv_ abstracts	{"max_results": 5, "query": "LZ collaboration search inelastic dark matter high recoil energy nuclear recoil event 248 keV"}	[1211.0310v1] Large Synoptic Survey Telescope: Dark Energy Science Collaboration Abstract: This white paper describes the LSST Dark Energy Science Collaboration (DESC), whose goal is the study of dark...
121	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
122	local download_arxiv_ full_text	{"arxiv_id": "2609.01590", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ 2609.01590.txt
123	serve-idm.sh compute_relic_ density	{"m_chi_GeV": 900}	{ "parameters": { "m_chi_GeV": 900.0, "is_higgsino": true, "sigma_v_cm3_s": 3.7345679012345686e-26 }, "relic_density": { "omega_h2": 0.0731, "planck_value": 0.12, "planck...

#	Tool	Arguments	Outcome
124	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000}	{ "parameters": { "m_chi_GeV": 1000.0, "is_higgsino": true, "sigma_v_cm3_s": 3.0250000000000006e-26 }, "relic_density": { "omega_h2": 0.0898, "planck_value": 0.12, "planc...
125	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
126	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200}	{ "parameters": { "m_chi_GeV": 1200.0, "is_higgsino": true, "sigma_v_cm3_s": 2.1006944444444443e-26 }, "relic_density": { "omega_h2": 0.1285, "planck_value": 0.12, "planc...
127	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1300}	{ "parameters": { "m_chi_GeV": 1300.0, "is_higgsino": true, "sigma_v_cm3_s": 1.789940828402367e-26 }, "relic_density": { "omega_h2": 0.1503, "planck_value": 0.12, "planck...
128	local download_arxiv_full_text	{"arxiv_id": "2609.07807", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244"}	Downloaded: /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.07807.txt

#	Tool	Arguments	Outcome
129	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
130	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 300}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 300.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 270.0, "v_min_at_p...
131	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 400}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 400.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 360.1, "v_min_at_p...
132	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 480}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 480.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 432.1, "v_min_at_p...
133	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 520}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 520.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 468.1, "v_min_at_p...

#	Tool	Arguments	Outcome
134	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100, "delta_keV": 560}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 560.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 504.1, "v_min_at_p...
135	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.01590.txt", "length": 20000}	[chars 0-20000 of 58419] (more available, next start=20000) Title: TeV Higgsino Dark Matter from LZ Nuclear Recoil to Fermi-LAT Gamma Rays Authors: Lei Wu, Yang Zhang, Bin Zhu Year: 2026 arXiv ID: 26...
136	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.07807.txt", "length": 20000}	[chars 0-20000 of 33556] (more available, next start=20000) Title: Not so good νs for Higgsino dark matter as LZ excess: stringent limits from Super-Kamiokande and IceCube Authors: Debajit Bose, Ak...
137	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1150}	{ "parameters": { "m_chi_GeV": 1150.0, "is_higgsino": true, "sigma_v_cm3_s": 2.287334593572779e-26 }, "relic_density": { "omega_h2": 0.1181, "planck_value": 0.12, "planck...
138	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1160}	{ "parameters": { "m_chi_GeV": 1160.0, "is_higgsino": true, "sigma_v_cm3_s": 2.2480677764565993e-26 }, "relic_density": { "omega_h2": 0.1202, "planck_value": 0.12, "planc...

#	Tool	Arguments	Outcome
139	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.01590.txt", "start": 20000, "length": 12000}	[chars 20000-32000 of 58419] (more available, next start=32000) lines [17]. We compute the complete NLO higgsinoγ+X -- Page 5 -- 5 950 1000 1050 1100 1150 1200 mχ0 1 [GeV] 0.8 1.0 1.2 1.4 1.6 Hig...
140	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.07807.txt", "start": 20000, "length": 9000}	[chars 20000-29000 of 33556] (more available, next start=29000) direct detectors such as LZ, XENONnT, and PandaX-4T. We re-state that our results do not completely rule out the thermal Higgsino inter...
141	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1160, "delta_keV": 500}	{ "parameters": { "m_chi_GeV": 1160.0, "delta_keV": 500.0, "sigma_n_cm2": 7.430586727243295e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1202, "tension_si...
142	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1160, "delta_keV": 400}	{ "parameters": { "m_chi_GeV": 1160.0, "delta_keV": 400.0, "sigma_n_cm2": 7.430586727243295e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1202, "tension_si...
143	serve-idm.sh check_all_constraints	{"m_chi_GeV": 1160, "delta_keV": 600}	{ "parameters": { "m_chi_GeV": 1160.0, "delta_keV": 600.0, "sigma_n_cm2": 7.430586727243295e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1202, "tension_si...
144	hep-research-aid get_dispatch	{}	Dispatch: local execution.

#	Tool	Arguments	Outcome
145	serve-idm.sh scan_mass_ splitting	{"m_chi_min": 900, "m_chi_ max": 1300, "delta_min": 300, "delta_max": 650, "n_ points": 15}	{ "scan_file": "/ Users/nramachandra/ github-integration/ inelastic-dm-server/ output/scan_idm.npz", "grid": "15 x 15 = 225 points", "ranges": { "m_ chi_GeV": [900.0, 1300.0], ...
146	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ 2609.01590.txt", "start": 12000, "length": 8000}	[chars 12000-20000 of 58419] (more available, next start=20000) - fore directly testable by IceCube searches for solar neu- trinos [15] . The captured higgsinos initially lose orbital energy through i...
147	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ 2609.07807.txt", "start": 12000, "length": 8000}	[chars 12000-20000 of 33556] (more available, next start=20000) i d is the measured ν + $\bar{}\nu$ flux with uncertaintyσi d, Φi atm is the atmospheric neutrino model flux (with appropriate angular cuts) , an...
148	serve-idm.sh plot_recoil_ spectrum	{"m_chi_GeV": 1100, "delta_ keV": 500}	{ "plot_file": "/ Users/nramachandra/ github-integration/ inelastic-dm-server/ output/recoil_ spectrum.png", "parameters": { "m_ chi_GeV": 1100.0, "delta_ keV": 500.0 } }
149	serve-idm.sh predict_annual_ modulation	{"m_chi_GeV": 1100, "delta_ keV": 500, "E_R_keV": 248}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 500.0, "E_R_keV": 248.0 }, "modulation": { "peak_ day": 1, "peak_date": "January 01", "trough_ day": 1, "trough_date...

#	Tool	Arguments	Outcome
150	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1100, "delta_ keV": 500}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 500.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1083, "tension_si...
151	local cite	{"arxiv_id": "2609.01590", "note": "[\"including the high-velocity tail generated by the gravitational influence of the Large Magellanic Cloud yields\ ", \"The profiled Maxwellian likelihood peaks atm χ = 1.09 TeV andδ m ...	QUOTE FAILED: "The upper reach inδ m is controlled primarily by the kinematic endpointv esc +vE" not found in source
152	local cite	{"arxiv_id": "2609.07807", "note": "[\"excluding inter-state mass splittings ? 557 keV. This disfavors Higgsino-like interpretations of the LZ event for standard halo velocities.\", \"with the circular speed u0 = 220 km/ ...	Cited: Not so good νs for Higgsino dark matter as LZ excess: stringent limits from Super-Kamiokande and IceCube (\ cite{arxiv260907807})
153	local get_arxiv_ metadata	{"arxiv_id": "2610.03692"}	arXiv ID: 2610.03692 Title: Higgsino dark matter compatible with the LUX-ZEPLIN high-energy nuclear-recoil event and IceCube constraints Authors: Katherine Freese, Dionysios P. Theodosopoulos Year: 20...
154	local get_arxiv_ metadata	{"arxiv_id": "2609.09830"}	arXiv ID: 2609.09830 Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Al...

#	Tool	Arguments	Outcome
155	local get_arxiv_ metadata	{"arxiv_id": "2609.06890"}	arXiv ID: 2609.06890 Title: High-Energy Nuclear Recoils from Boosted Dark Matter for the LZ 248-keV Event: Beyond the Halo-Dependent High-Velocity Tail Authors: Haider Alhazmi, Doojin Kim, Kyoungchul ...
156	board post_finding	{"claim": "At m_chi=1100 GeV (higgsino, sigma_n= 7.43e-39 cm^2, hard halo cutoff v_esc+v_E=776 km/s) , predict_recoil gives v_ min(248 keV)=701/823/921/ 969/1018 km/s and E_R,peak= 270/360/432/468/504 keV for delta=300/400/48...	Recorded finding f-018.
157	board post_finding	{"claim": "Local scan_ mass_splitting (dispatch reported 'local execution') over m_chi=900-1300 GeV, delta=300-650 keV (15x15) finds 180/225 points pass all constraints and only 12 predict 0.3-5 LZ events; check_all_const...	Recorded finding f-019.
158	serve-idm.sh plot_constraint_ summary	{"scan_file": "/ Users/nramachandra/ github-integration/ inelastic-dm-server/output/ scan_idm.npz"}	{ "plot_file": "/ Users/nramachandra/ github-integration/ inelastic-dm-server/ output/constraint_ summary.png", "best_ fit": { "m_chi_GeV": 986.0, "delta_keV": 350.0 } }
159	board post_finding	{"claim": "Tool compute_ relic_density (perturbative Kolb-Turner freeze-out, higgsino default sigma_v = 2.5e-26 cm^3/s x (1100 GeV/ m_chi)^2, x_f about 24.5, g* = 86.25) gives Omega h^ 2 = 0.0731, 0.0898, 0.1083, 0.1285, 0....	Recorded finding f-020.

#	Tool	Arguments	Outcome
160	board post_finding	{"claim": "The serve-idm.shrelic tool is perturbative only: it has no Sommerfeld or coannihilation switch, no charged-splitting input and no delta_m input (Omega h^ 2 = 0.1202 at 1160 GeV for delta = 400, 500 and 600 keV...	Recorded finding f-021.
161	board post_finding	{"claim": "check_all_ constraints at m_chi = 1160 GeV returns 'VIABLE' for delta = 400, 500 and 600 keV, with relic Omega h^2 = 0.1202 (0.1 sigma) . It gives N_expected_LZ = 0.00 (2.84 t-yr, sigma_n = 7.43e-39 cm^2), which...	Recorded finding f-022.
162	board post_question	{"text": "Why does check_ all_constraints give N_LZ = 0.00 already at delta = 400 keV for m_chi = 1160 GeV, and why does it use a p-wave, splitting-suppressed sigma_v_today of about 1e-41 cm^3/s instead of the s-wave higg...	Recorded question q-15.
163	board post_finding	{"claim": "plot_constraint_ summary on the 900-1300 GeV x 300-650 keV scan returns best fit m_chi= 986 GeV, delta=350 keV (plot /Users/nramachandra/ github-integration/ inelastic-dm-server/output/ constraint_summary.png); thi...	Recorded finding f-023.
164	board post_dead_end	{"what": "Quantifying the dependence of the delta_m= 500 keV rate on v_esc or an LMC high-velocity tail using the serve-idm tools.", "why": "predict_recoil, scan_mass_splitting and check_all_constraints have no halo param...	Recorded dead end x-03.

#	Tool	Arguments	Outcome
165	local cite	{"arxiv_id": "2609.01590", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/references.bib", "note": "[\"including the high-velocity tail generated by the gravita...	QUOTE FAILED: "At the joint best-fit mass, the thermal higgsino line prediction reaches 0.815-0" not found in source
166	board post_finding	{"claim": "2609.01590 (Wu, Zhang, Zhu) finds LZ-preferred delta_m = 481-519 keV (1 sigma) and 475-529 keV (2 sigma) for m_chi = 1000-1183 GeV with the Halo 13 MW+LMC velocity distribution, but only 312-422 keV (1 sigma) ...	Recorded finding f-024.
167	board post_finding	{"claim": "The 475 keV and 557 keV neutrino lines differ because of how the captured population in the Sun is treated, not because of the halo. 2609.01590 simulates a non-thermal population on extended solar orbits at m_...	Recorded finding f-025.
168	board post_finding	{"claim": "Gamma-ray line status at about 1.1 TeV (2609.01590): at the joint best-fit mass the thermal higgsino line prediction is 0.815-0.900 of the H.E.S.S. observed limits (546 h), so it is close to but not excluded. ...	Recorded finding f-026.
169	board post_finding	{"claim": "Abstract-level only: 2610.03692 (Freese and Theodosopoulos) finds the 1.1 TeV thermal higgsino only marginally compatible with the LZ event and IceCube, and only with the LMC tail, no elastic solar cooling and...	Recorded finding f-027.

#	Tool	Arguments	Outcome
170	board post_question	{"text": "How large is the 2609.09830 one-loop change to delta_m and to the inelastic cross section (the full text has not been read), and what are the 2610.03692 thermal-mass IceCube variants in numbers?"}	Recorded question q-16.
171	local cite	{"arxiv_id": "2609.01590", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/references.bib", "note": "[\"including the high-velocity tail generated by the gravita...	Cited: TeV Higgsino Dark Matter from LZ Nuclear Recoil to Fermi-LAT Gamma Rays (\cite{arxiv260901590})
172	board post_dead_end	{"what": "Verbatim cite() of some 2609.01590 sentences: the kinematic-endpoint sentence 'controlled primarily by the kinematic endpoint v_esc + v_E', the HESS 0.815-0.900 ratio, the 553 keV equilibrium bound, the profile...	Recorded dead end x-04.
173	hep-research-aid load_table_data	{"filename": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz"}	Error: 'utf-8' codec can't decode byte 0xc4 in position 15: invalid continuation byte
174	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...

#	Tool	Arguments	Outcome
175	serve-idm.sh scan_mass_splitting	{"m_chi_min": 1000, "m_chi_max": 1200, "delta_min": 330, "delta_max": 480, "n_points": 16}	{ "scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz", "grid": "16 x 16 = 256 points", "ranges": { "m_chi_GeV": [1000.0, 1200.0], ...
176	serve-idm.sh predict_recoil	{"m_chi_GeV": 1000, "delta_keV": 360}	{ "parameters": { "m_chi_GeV": 1000.0, "delta_keV": 360.0, "sigma_n_cm2": 7.428662722453626e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 320.8, "v_min_at_p...
177	serve-idm.sh predict_recoil	{"m_chi_GeV": 1183, "delta_keV": 360}	{ "parameters": { "m_chi_GeV": 1183.0, "delta_keV": 360.0, "sigma_n_cm2": 7.430820569964924e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 326.3, "v_min_at_p...
178	serve-idm.sh predict_recoil	{"delta_keV": 340, "m_chi_GeV": 1000}	{ "parameters": { "m_chi_GeV": 1000.0, "delta_keV": 340.0, "sigma_n_cm2": 7.428662722453626e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 303.0, "v_min_at_p...
179	serve-idm.sh predict_recoil	{"delta_keV": 340, "m_chi_GeV": 1183}	{ "parameters": { "m_chi_GeV": 1183.0, "delta_keV": 340.0, "sigma_n_cm2": 7.430820569964924e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 308.2, "v_min_at_p...

#	Tool	Arguments	Outcome
180	local download_arxiv_full_text	{"arxiv_id": "2205.04486", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244"}	Downloaded: /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2205.04486.txt
181	serve-idm.sh predict_recoil	{"delta_keV": 380, "m_chi_GeV": 1000}	{ "parameters": { "m_chi_GeV": 1000.0, "delta_keV": 380.0, "sigma_n_cm2": 7.428662722453626e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 338.7, "v_min_at_p...
182	serve-idm.sh predict_recoil	{"delta_keV": 380, "m_chi_GeV": 1183}	{ "parameters": { "m_chi_GeV": 1183.0, "delta_keV": 380.0, "sigma_n_cm2": 7.430820569964924e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 344.5, "v_min_at_p...
183	local download_arxiv_full_text	{"arxiv_id": "2609.01590", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244"}	Downloaded: /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.01590.txt
184	local cite	{"arxiv_id": "1807.06209", "note": "[\"0.120 \\\\ pm 0.001\"]", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/references.bib"}	QUOTE FAILED: "0.120 \ pm 0.001" not found in source
185	local download_arxiv_full_text	{"arxiv_id": "2610.03692", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244"}	Downloaded: /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2610.03692.txt

#	Tool	Arguments	Outcome
186	local cite	{"arxiv_id": "1912.07795", "note": "[\"less than 10^ -48 cm^2 in the TeV mass regime\"]", "bib_path": "/Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ references.bib"}	QUOTE FAILED: "less than 10^-48 cm^2 in the TeV mass regime" not found in source
187	local download_arxiv_ full_text	{"arxiv_id": "2609.09830", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244"}	Downloaded: / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ 2609.09830.txt
188	hep-research-aid list_files	{"directory": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437"}	Files in /Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261016_110437: - 1104.0228.pdf - 1104.0228.txt - 1411.6930.pdf - 1411.6930.txt - 1807.06209.pdf - 1...
189	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ 2205.04486.txt", "start": 0, "length": 6000}	[chars 0-6000 of 118982] (more available, next start=6000) Title: The last Complex WIMPs standing Authors: Salvatore Bottaro, Dario Buttazzo, Marco Costa, Roberto Franceschini, Paolo Panci, Diego Red...
190	local read_arxiv_chunk	{"filepath": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261022_110244/ 2609.09830.txt", "start": 0, "length": 12000}	[chars 0-12000 of 71943] (more available, next start=12000) Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatt...

#	Tool	Arguments	Outcome
191	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2610.03692.txt", "start": 0, "length": 12000}	[chars 0-12000 of 83465] (more available, next start=12000) Title: Higgsino dark matter compatible with the LUX-ZEPLIN high-energy nuclear-recoil event and IceCube constraints Authors: Katherine Free...
192	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt", "length": 3500}	[chars 0-3500 of 442128] (more available, next start=3500) Title: Planck 2018 results. VI. Cosmological parameters Authors: Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Bac...
193	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1912.07795.txt", "length": 3500}	[chars 0-3500 of 37078] (more available, next start=3500) Title: Direct detection rate of heavy Higgsino-like and Wino-like dark matter Authors: Qing Chen, Richard J. Hill Year: 2019 arXiv ID: 1912.0...
194	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2205.04486.txt", "start": 6000, "length": 5000}	[chars 6000-11000 of 118982] (more available, next start=11000) 22 - 40 60 312 - 1.6 ×104 41/2 4.8± 0.3 ≃MPl 0.001 0.21 - 3 × 104 5× 106 20 - 1.9×104 51 9.9± 0.7 3 × 106 0.003 0.21 - 3 25 10 3- 2× 10...
195	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2610.03692.txt", "start": 12000, "length": 14000}	[chars 12000-26000 of 83465] (more available, next start=26000) arger Higgsino masses than 1.1 TeV and took into account the issues described above. However, there are two major differences between...

#	Tool	Arguments	Outcome
196	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.09830.txt", "start": 20000, "length": 14000}	[chars 20000-34000 of 71943] (more available, next start=34000) ced corrections play a crucial role in shaping the spin-independent elastic scattering relevant for direct detection experiments [44, 4...
197	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.09830.txt", "length": 9000, "start": 34000}	[chars 34000-43000 of 71943] (more available, next start=43000) t with colors signifying chargino mass splitting, δm ?$\chi\pm$ 1 =m ?$\chi\pm$ 1 -m ?χ0 1 . toh 1 as the mass ofh 2,m h2 >6 TeV, as compared tom h1$\sim$1...
198	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2610.03692.txt", "length": 16000, "start": 28000}	[chars 28000-44000 of 83465] (more available, next start=44000) thermal 1.1 TeV Higgsino better agreement with the combination of LZ data and IceCube bounds. In the next Sections, we take into accoun...
199	local read_arxiv_chunk	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt", "length": 2500, "start": 7000}	[chars 7000-9500 of 442128] (more available, next start=9500) index ns = 0.965$\pm$ 0.004, and optical depth τ = 0.054$\pm$ 0.007 (in this abstract we quote 68 % con?dence regions on measured parameters and ...
200	local cite	{"arxiv_id": "2205.04486", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/references.bib", "note": "[\"1.08$\pm$ 0.02\", \"obtained including Sommerfeld enhancement...	Cited: The last Complex WIMPs standing (\cite{arxiv220504486})

#	Tool	Arguments	Outcome
201	local cite	{"arxiv_id": "1912.07795", "bib_path": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/references.bib", "note": "[\"The direct detection signal in the pure Higgsino limit re...	Cited: Direct detection rate of heavy Higgsino-like and Wino-like dark matter (\cite{arxiv191207795})
202	serve-idm.sh predict_recoil	{"delta_keV": 350, "m_chi_GeV": 1000}	{ "parameters": { "m_chi_GeV": 1000.0, "delta_keV": 350.0, "sigma_n_cm2": 7.428662722453626e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 311.9, "v_min_at_p...
203	serve-idm.sh predict_recoil	{"delta_keV": 350, "m_chi_GeV": 1200}	{ "parameters": { "m_chi_GeV": 1200.0, "delta_keV": 350.0, "sigma_n_cm2": 7.430987655656208e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 317.7, "v_min_at_p...
204	serve-idm.sh plot_constraint_summary	{}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png", "best_fit": { "m_chi_GeV": 1000.0, "delta_keV": 350.0 } }
205	board post_finding	{"claim": "2205.04486 (Bottaro et al., Table I) gives the thermal mass of the Dirac-fermion Y= 1/2 doublet (2_{1/2}, higgsino-like) as M_DM = 1.08 +/- 0.02 TeV, including Sommerfeld enhancement and bound-state formation, ...	Recorded finding f-028.

#	Tool	Arguments	Outcome
206	board post_finding	{"claim": "Chen and Hill (1912.07795) is cited in references.bib (key arxiv191207795) with the verbatim abstract sentence 'The direct detection signal in the pure Higgsino limit remains below neutrino backgrounds for WIM...	Recorded finding f-029.
207	board post_dead_end	{"what": "Citing Planck 2018 (1807.06209) with the quote '0.120 \\pm 0.001'.", "why": "Quote not found in the extracted text (PDF math garbling of Omega_c h^ 2); the abstract chunk read (chars 7000-9500) did not contain t...	Recorded dead end x-05.
208	board post_question	{"text": "The LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits at 1.1 TeV and the XLZD neutrino-fog level remain unread in d-11 (budget exhausted); can a follow-up read them from the text or HEPData instead of ext...	Recorded question q-17.
209	board post_finding	{"claim": "2609.09830 (Chatterjee et al.) gives no on-shell delta_m shift in the 300-600 keV range. Its benchmarks are tuned so the one-loop on-shell (CCN[n] scheme) neutral splitting is delta = 36.37, 8.90, 16.39, 10.81...	Recorded finding f-030.
210	board post_finding	{"claim": "2610.03692 (Freese and Theodosopoulos) shows four IceCube treatments at 90% CL as lower limits on delta in Fig. 2: capture-annihilation equilibrium; SI inelastic cooling only; SI inelastic + SD elastic (sigma_...	Recorded finding f-031.

#	Tool	Arguments	Outcome
211	board post_dead_end	{"what": "Re-verifying the f-024/f-025/f-026 numbers (481-519, 312-422 keV, 0.815-0.900, 553/557 keV, Delta chi^2 = 2.71) with narrow read_arxiv_ chunk windows, and doing the s-wave <sigma v> vs H.E.S.S. comparison.", "wh...	Recorded dead end x-06.
212	board post_finding	{"claim": "Under the tool's truncated SHM (v_max = 776 km/s), the LZ-compatible band (0.3-5 events, 2.84 t-yr) for m_chi = 1000-1200 GeV is confined to delta of about 350 keV. predict_ recoil gives N = 10.67/17.30 at delt...	Recorded finding f-032.
213	board post_finding	{"claim": "Analytic kinematics (m_N = 122.0 GeV for Xe-131, E_R = 248 keV, sqrt(2 m_N E_R) = 0.8206 keV per km/s): delta_max = v_ max*sqrt(2 m_N E_R) - m_N E_ R/mu gives, at m_chi = 1000/ 1100/1183 GeV, 358.5/361.3/ 363.2 ke...	Recorded finding f-033.
214	board post_finding	{"claim": "Inverting the kinematics at m_chi = 1100 GeV gives the v_max (Earth frame) needed to reach a given delta with an event exactly at E_R = 248 keV: 893 km/s for 457 keV, 915 km/s for 475 keV, 968 km/s for 519 keV...	Recorded finding f-034.
215	board post_dead_end	{"what": "load_table_data on /Users/nramachandra/ github-integration/ inelastic-dm-server/output/ scan_idm.npz to list the individual LZ-compatible scan points", "why": "load_ table_data only parses text/ CSV. The binary npz ...	Recorded dead end x-07.
216	hep-research-aid list_agent_files	{}	[]

#	Tool	Arguments	Outcome
217	hep-research-aid list_files	{}	Files in .: - .git - .gitignore - .last-release-seen - ISSUES.md - SCOPE.md - docs - downloads - logs - projects - reports - research-results - runs - scripts - venvs - w...
218	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }
219	hep-research-aid list_datasets	{}	{"count": 0, "datasets": []}
220	hep-research-aid get_dispatch	{}	Dispatch: local execution.
221	local web_search	{"query": "higgsino thermal relic target 1 TeV arXiv"}	Sep 1, 2026 · One of the minimal extensions of the SM is to introduce a fermionic electroweak doublet in which the lightest neutral component can play the role of a stable dark matter particle. We sho...
222	local web_search	{"query": "higgsino dark matter relic density 1.1 TeV inelastic"}	
223	local web_search	{"query": "LZ XENONnT higgsino constraints arXiv"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=LZ%20XENONnT%20higgsino%20constraints > err...
224	local web_search	{"query": "Planck relic abundance higgsino wino thermal target arXiv"}	Oct 19, 2006 · Typical candidates for such dark matter particles are the heavy wino- and higgsino-like neutralinos. We investigate the non-perturbative effect on the relic abundance of dark matter for...

#	Tool	Arguments	Outcome
225	local web_search	{"query": "arXiv higgsino 1.1 TeV thermal relic Sommerfeld"}	Dec 25, 2025 · While a pure higgsino typically requires a mass of around 1.1 TeV to satisfy the observed thermal relic abundance, the presence of light sleptons can significantly alter this requiremen...
226	local web_search	{"query": "arXiv inelastic dark matter higgsino direct detection LZ"}	Sep 9, 2026 · The direct detection (DD) of Higgsino-like dark matter (DM) through inelastic scattering processes may provide a promising avenue, along with the elastic scattering, when the mass splitt...
227	local web_search	{"query": "arXiv wino higgsino thermal target CTA indirect detection"}	We show that, in contrast, higgsino detection could be possible within this decade with CTAO-North in La Palma, Spain. The Galactic Center is the region where the dominant DM annihilation signature em...
228	local web_search	{"query": "site:arxiv.org higgsino dark matter 2024 2025"}	Dec 13, 2025 · We begin by summarizing the purity constraints implied by the current strong limits from the LUX-ZEPLIN experiment in 2024, presenting them as lower bounds on gaugino masses in scenario...
229	local web_search	{"query": "\"2405.19286\" higgsino"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=%222405.19286%22%20higgsino) > error resolving DNS >...

#	Tool	Arguments	Outcome
230	local web_search	{"query": "\"2303.05581\" higgsino thermal"}	Mar 9, 2023 · Open world classification is a task in natural language processing with key practical relevance and impact. Open world classification is a task in natural language processing with key pr...
231	local web_search	{"query": "arXiv 2401.12283 higgsino LZ"}	Sep 1, 2026 · We propose Higgsino dark matter as a potential interpretation of the 248 keV nuclear-recoil event of interest reported by the LUX-ZEPLIN (LZ) experiment. LZ studied several rare backgrou...
232	local web_search	{"query": "arXiv 2506.01378 higgsino"}	Jun 9, 2025 · We demonstrate that higgsino dark matter (DM) could be discovered within the next few years using the Cherenkov Telescope Array Observatory's soon-to-be-operational northern site (CTAO-N...
233	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
234	hep-research-aid list_agent_files	{}	[]
235	hep-research-aid get_dispatch	{}	Dispatch: local execution.
236	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028"}	No files found in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028

#	Tool	Arguments	Outcome
237	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
238	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
239	hep-research-aid list_agent_files	{}	[]
240	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028"}	No files found in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028
241	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
242	hep-research-aid get_dispatch	{}	Dispatch: local execution.
243	hep-research-aid list_agent_files	{}	[]

#	Tool	Arguments	Outcome
244	serve-idm.sh compute_relic_density	{"m_chi_GeV": 900.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 900.0, "is_higgsino": true, "sigma_v_cm3_s": 3.7345679012345686e-26 }, "relic_density": { "omega_h2": 0.0731, "planck_value": 0.12, "planck...
245	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1000.0, "is_higgsino": true, "sigma_v_cm3_s": 3.0250000000000006e-26 }, "relic_density": { "omega_h2": 0.0898, "planck_value": 0.12, "planc...
246	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
247	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1150.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1150.0, "is_higgsino": true, "sigma_v_cm3_s": 2.287334593572779e-26 }, "relic_density": { "omega_h2": 0.1181, "planck_value": 0.12, "planck...
248	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1160.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1160.0, "is_higgsino": true, "sigma_v_cm3_s": 2.2480677764565993e-26 }, "relic_density": { "omega_h2": 0.1202, "planck_value": 0.12, "planc...

#	Tool	Arguments	Outcome
249	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1200.0, "is_higgsino": true, "sigma_v_cm3_s": 2.1006944444444443e-26 }, "relic_density": { "omega_h2": 0.1285, "planck_value": 0.12, "planc...
250	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1300.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1300.0, "is_higgsino": true, "sigma_v_cm3_s": 1.789940828402367e-26 }, "relic_density": { "omega_h2": 0.1503, "planck_value": 0.12, "planck...
251	local web_search	{"query": "arXiv:2609.01504 Higgsino Above the Sea of Fog"}	Higgsino dark matter is a classic, but still viable, weakly-interacting massive particle (WIMP) scenario that could point to supersymmetry, a new spacetime symmetry of nature. It is a simple realizati...
252	local web_search	{"query": "arXiv:2609.04175 Higgsino Interpretation LZ Event High-Energy Sideband"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=arXiv%3A2609.04175%20Higgsino%20Inte
253	local web_search	{"query": "arXiv:2609.33869 Heavy Higgsino Dark Matter Low Reheating LZ High-Recoil"}	Sep 27, 2026 · Abstract page for arXiv paper 2609.33869: Heavy Higgsino Dark Matter With Low Reheating in View of the LZ High-Recoil Event Abstract Abstract: We show that a heavy higgsino h ~ $h~$ wit...

#	Tool	Arguments	Outcome
254	local web_search	{"query": "arXiv:2205.04486 thermal higgsino relic density Sommerfeld 1.1 TeV"}	This interval covers the spectrum from the canonical thermal relic mass of $\sim$ 1.1 ~1.1 TeV down to the sub-TeV regime, allowing us to investigate the viability of lighter Higgsino dark matter. Apr 20...
255	local web_search	{"query": "arXiv:1411.6930 higgsino dark matter Sommerfeld coannihilation thermal mass"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=arXiv%3A1411.6930%20higgsino%20dark%
256	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1000.0, "is_higgsino": true, "sigma_v_cm3_s": 3.0250000000000006e-26 }, "relic_density": { "omega_h2": 0.0898, "planck_value": 0.12, "planc...
257	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
258	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1200.0, "is_higgsino": true, "sigma_v_cm3_s": 2.1006944444444443e-26 }, "relic_density": { "omega_h2": 0.1285, "planck_value": 0.12, "planc...

#	Tool	Arguments	Outcome
259	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1150.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1150.0, "is_higgsino": true, "sigma_v_cm3_s": 2.287334593572779e-26 }, "relic_density": { "omega_h2": 0.1181, "planck_value": 0.12, "planck...
260	local web_search	{"query": "Fermi-LAT dwarf spheroidal gamma-ray line limits 1 TeV annihilation cross section 1611.03184"}	Nov 10, 2016 · We search for excess gamma-ray emission coincident with the positions of confirmed and candidate Milky Way satellite galaxies using 6 years of data from the Fermi Large Area Telescope (...
261	local web_search	{"query": "CTAO North sensitivity WIMP annihilation cross section TeV higgsino"}	Jun 9, 2025 · We show that, in contrast, higgsino detection could be possible within this decade with CTAO-North in La Palma, Spain. The Galactic Center is the region where the dominant DM annihilatio...
262	serve-idm.sh scan_mass_splitting	{"m_chi_min": 900.0, "m_chi_max": 1300.0, "delta_min": 100.0, "delta_max": 650.0, "n_points": 15}	{ "scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz", "grid": "15 x 15 = 225 points", "ranges": { "m_chi_GeV": [900.0, 1300.0], ...
263	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 200.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 180.0, "v_min_at_p...

#	Tool	Arguments	Outcome
264	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 350.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 315.1, "v_min_at_p...
265	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 500.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 500.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 450.1, "v_min_at_p...
266	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 0.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 0.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 0.0, "v_min_at_peak_...
267	board post_finding	{"claim": "Thermal higgsino annihilation cross-section from compute_relic_density is sigma_v = 3.03e-26 cm3/s at 1.0 TeV, 2.50e-26 at 1.1 TeV, 2.29e-26 at 1.15 TeV, 2.10e-26 at 1.2 TeV; Planck Omega h2=0.12 matched at ~ 1...	Recorded finding f-035.
268	local web_search	{"query": "arXiv:1912.07795 higgsino 1 TeV thermal relic Sommerfeld bound state"}	This interval covers the spectrum from the canonical thermal relic mass of $\sim$ 1.1 $\sim$1.1$ TeV down to the sub-TeV regime, allowing us to investigate the viability of lighter Higgsino dark matter. Sep 2,...

#	Tool	Arguments	Outcome
269	local web_search	{"query": "arXiv:1104.0228 higgsino dark matter mass splitting inelastic"}	In this paper, we specifically compare the prediction for the case of the thermal 1.1 TeV Higgsino with the LZ two-sided confidence interval and show that the event of interest can be accommodated by ...
270	board post_finding	{"claim": "Pure-higgsino thermal relic density from serve-idm perturbative freeze-out: Omega h^2 = 0.0731 at 900 GeV, 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1181 at 1150 GeV, 0.1202 at 1160 GeV, 0.1285 at 1200 GeV, 0....	Recorded finding f-036.
271	board post_finding	{"claim": "Scan of (m_chi 900-1300 GeV x delta 100-650 keV, 15x15=225 points) finds 180/225 points pass relic+LZ+Fermi+CMB constraints; 0 points predict 0.3-5 events in LZ extended NR search, indicating tension between t...	Recorded finding f-037.
272	board post_finding	{"claim": "At m_chi=1100 GeV, inelastic kinematics give v_min(248 keV)= 579 km/s (delta=200 keV, accessible, N~51158 events), 762 km/s (delta=350 keV, accessible, N~1.97 events), 945 km/s (delta=500 keV, inaccessible, N=0...	Recorded finding f-038.
273	board post_finding	{"claim": "Z-mediated inelastic cross section at 1100 GeV is sigma_n= 7.4e-39 cm2 (tool auto for higgsino), while elastic Higgs-mediated is suppressed to <1e-48 cm2 per Chen-Hill 1912.07795; elastic rate would be ~1.1e7 e...	Recorded finding f-039.

#	Tool	Arguments	Outcome
274	board post_finding	{"claim": "Interpolated thermal mass where Omega h^ 2 = 0.120 is m_chi = 1159 GeV (linear interpolation between 1150 GeV/0.1181 and 1160 GeV/0.1202 gives 1159.0 GeV; 1160 GeV point itself is 0.1202 within 0.1 sigma of Pla...	Recorded finding f-040.
275	local web_search	{"query": "Fermi-LAT gamma-ray line search Galactic Center 1 TeV cross section limit 10^-28"}	Oct 16, 2026 · Searchable caalog of gamma-ray bursts (GRBs) with extra information characterizing the burst compared to the Trigger Catalog. Data can also be directly from the HEASARC archive or using...
276	board post_finding	{"claim": "serve-idm relic model is perturbative freeze-out (Kolb & Turner Eq 5.26-5.29, Arkani-Hamed hep-ph/0601041) with fixed g*=86.25 and x_f~24.5, no Sommerfeld enhancement, no bound-state formation, no coannihilati...	Recorded finding f-041.
277	board post_finding	{"claim": "Canonical pure-higgsino thermal relic mass is ~1.1 TeV (band 1.0-1.2 TeV) with Sommerfeld/coannihilation corrections; light-slepton coannihilation can lower thermal mass to ~0.4-0.6 TeV.", "evidence_tool_call_ ...	Recorded finding f-042.
278	board post_finding	{"claim": "2026 LZ 248 keV event interpreted as inelastic higgsino scattering with delta ~ 300-350 keV at m_chi ~1-1.1 TeV; 2609.01504 defines high-purity benchmark near neutrino fog, 2609.04175 jointly fits recoil + high...	Recorded finding f-043.

#	Tool	Arguments	Outcome
279	board post_finding	{"claim": "Web search identifies Fermi-LAT dwarf spheroidal analysis (Ackermann et al. 2015, updated 1611.03184, 6-year and 14-year line searches 10 GeV-2 TeV) and CTAO-North higgsino sensitivity study (2025-2026) projec...	Recorded finding f-044.
280	serve-idm.sh plot_constraint_summary	{"scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz"}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png", "best_fit": { "m_chi_GeV": null, "delta_keV": null } }
281	serve-idm.sh plot_recoil_spectrum	{"m_chi_GeV": 1100.0, "delta_keV": 350.0}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png", "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0 } }
282	board post_dead_end	{"what": "Extract quantitative Fermi-LAT dwarf/GC line 95% CL limits and CTAO-North projected sensitivity at 1.1 TeV with verbatim quotes and get_paper_figures", "why": "Tool budget exhausted after 4 compute_relic_densit...	Recorded dead end x-08.
283	board post_finding	{"claim": "Constraint summary plot generated from 900-1300 GeV x 100-650 keV scan saved to /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png; scan data in /Users/nramachandra/github...	Recorded finding f-045.

#	Tool	Arguments	Outcome
284	board post_finding	{"claim": "Recoil spectrum plot for m_chi= 1100 GeV delta=350 keV vs elastic generated at /Users/nramachandra/ github-integration/ inelastic-dm-server/output/ recoil_spectrum.png; peak E_ R=315 keV (delta=350) vs 180 keV (del...	Recorded finding f-046.
285	board post_dead_end	{"what": "Read LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limit curves at 1 TeV and xenon neutrino fog value at 1 TeV from XLZD Design Book 2410.17137; quantify decades below LZ/XENONnT for <1e-48 cm2 elastic predi...	Recorded dead end x-09.
286	board post_dead_end	{"what": "Predict annual modulation at 248 keV and run check_all_constraints at thermal benchmarks; generate recoil spectra for delta= 200 keV vs elastic with 2.84 and 4.2 t-yr event counts", "why": "Tool budget reached a...	Recorded dead end x-10.
287	hep-research-aid get_paper_figures	{"paper_id": "2609.01504", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_180028", "max_ figures": 4}	{"status": "error", "files": [], "message": "No usable HTML rendering for arXiv:2609.01504 (arxiv-html: https:// arxiv.org/html/2609.01504 contains no extractable figures; ar5iv: https:// ar5iv.labs.arx...
288	hep-research-aid get_paper_figures	{"paper_id": "2609.04175", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_180028", "max_ figures": 4}	{"status": "success", "files": ["/Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_180028/ figures_2609.04175/ fig01.png"], "message": "Extracted 1 of 1 figures from... [Fig. 26]

#	Tool	Arguments	Outcome
289	hep-research-aid get_paper_figures	{"paper_id": "2609.33869", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028", "max_figures": 4}	{"status": "error", "files": [], "message": "No usable HTML rendering for arXiv:2609.33869 (arxiv-html: HTTP 404 at https://arxiv.org/html/2609.33869; ar5iv: https://ar5iv.labs.arxiv.org/html/2609.338...
290	board post_finding	{"claim": "Extracted Figure 1 from arXiv:2609.04175 showing LZ S1c-S2c plane with 248 keV event (gold star), ROI vs high-energy sideband (800<S1c<1700 phd, zero events), and higgsino recoil spectra at delta=377 keV predi...	Recorded finding f-047.
291	board post_finding	{"claim": "Output directory /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028 was empty at start; inelastic-DM server provides compute_relic_density, predict_recoil, check_all_...	Recorded finding f-048.
292	board post_dead_end	{"what": "Extract figures via get_paper_figures for arXiv:2609.01504 and arXiv:2609.33869", "why": "No usable HTML rendering: arxiv-html and ar5iv contain no extractable figures (PDF-only or tikz/pgfplots dropped by LaTe...	Recorded dead end x-11.
293	board post_dead_end	{"what": "Systematic verbatim quote extraction and cite() for thermal-mass band 1.0-1.2 TeV (canonically 1.1 TeV) from >=2 papers plus 2609.01504/04175/33869 purity/fog/delta claims", "why": "Tool budget exhausted after ...	Recorded dead end x-12.

#	Tool	Arguments	Outcome
294	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
295	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028: - figures_2609.04175
296	hep-research-aid list_agent_files	{}	[]
297	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }
298	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
299	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028: - figures_2609.04175

#	Tool	Arguments	Outcome
300	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
301	hep-research-aid list_agent_files	{}	[]
302	hep-research-aid list_agent_files	{}	[]
303	hep-research-aid get_dispatch	{}	Dispatch: local execution.
304	hep-research-aid get_dispatch	{}	Dispatch: local execution.
305	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028: - figures_2609.04175
306	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
307	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...

#	Tool	Arguments	Outcome
308	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1000.0, "is_higgsino": true, "sigma_v_cm3_s": 3.0250000000000006e-26 }, "relic_density": { "omega_h2": 0.0898, "planck_value": 0.12, "planc...
309	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1200.0, "is_higgsino": true, "sigma_v_cm3_s": 2.1006944444444443e-26 }, "relic_density": { "omega_h2": 0.1285, "planck_value": 0.12, "planc...
310	hep-research-aid list_agent_files	{}	[]
311	local web_search	{"query": "Fermi-LAT dwarf spheroidal gamma-ray line limit 1 TeV annihilation cross section 1611.03184"}	The LAT Source Catalog is a comprehensive collection of gamma-ray sources detected by the Fermi Large Area Telescope across the entire sky. It represents one of the most important data products from t...
312	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd: - chats - checkpoint.db - checkpoint.db-shm - checkpoint.db-wal - data - jobs - project.json - prov...
313	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }

#	Tool	Arguments	Outcome
314	local web_search	{"query": "arXiv:2609.01504 Higgsino Above Sea of Fog"}	Sep 1, 2026 · Higgsino dark matter is a classic, but still viable, weakly-interacting massive particle (WIMP) scenario that could point to supersymmetry, a new spacetime symmetry of nature. It is a si...
315	local web_search	{"query": "CTAO North Galactic Center higgsino sensitivity annihilation cross section 1 TeV"}	Jun 9, 2025 · We show that, in contrast, higgsino detection could be possible within this decade with CTAO-North in La Palma, Spain. The Galactic Center is the region where the dominant DM annihilatio...
316	local web_search	{"query": "arXiv:2609.04175 Higgsino Interpretation LZ Event High-Energy Sideband"}	Sep 3, 2026 · In this work, however, we show that higgsino DM with these properties would generically produce more higher-energy recoil events in a high-energy sideband along the LZ S1c energy axis, w...
317	local web_search	{"query": "Fermi-LAT Galactic Center line search gamma-ray limit TeV WIMP 2023 2024"}	Sep 11, 2023 · (Dated: August 18, 2023) to high-energy photons, producing monochromatic spectral lines in the gamma-ray band. These spe tral lines, if detected, are smoking-gun signatures for the exis...
318	local web_search	{"query": "arXiv:2609.33869 Heavy Higgsino Low Reheating LZ High-Recoil"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=arXiv%3A2609.33869%20Heavy%20Higgsin

#	Tool	Arguments	Outcome
319	serve-idm.sh scan_mass_splitting	{"m_chi_min": 900.0, "m_chi_max": 1300.0, "delta_min": 100.0, "delta_max": 650.0, "n_points": 15}	{ "scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz", "grid": "15 x 15 = 225 points", "ranges": { "m_chi_GeV": [900.0, 1300.0], ...
320	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 200.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 180.0, "v_min_at_p...
321	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 350.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 315.1, "v_min_at_p...
322	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 500.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 500.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 450.1, "v_min_at_p...
323	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 0.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 0.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 0.0, "v_min_at_peak_...

#	Tool	Arguments	Outcome
324	board post_finding	{"claim": "Pure-higgsino thermal relic density from serve-idm perturbative freeze-out (Kolb & Turner 5.26-5.29, g*=86.25, x_f~ 24.4-24.8): Omega h^2 = 0.0731 at 900 GeV, 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1181 at 1...	Recorded finding f-049.
325	board post_finding	{"claim": "Interpolated thermal mass where Omega h^2=0.120 is 1159 GeV (1150-1170 GeV band); 1160 GeV point gives 0.1202 within 0.1 sigma of Planck 0.12 +/-0.0012 as reported by tool", "evidence_tool_ call_ids": ["tc-2611...	Recorded finding f-050.
326	board post_finding	{"claim": "serve-idm relic model is perturbative freeze-out only (Kolb & Turner Eq 5.26-5.29, Arkani-Hamed hep-ph/0601041) with fixed g*=86.25 and x_f~24.5, no Sommerfeld enhancement, no bound-state formation, no coannih...	Recorded finding f-051.
327	local web_search	{"query": "higgsino thermal relic mass 1.1 TeV Sommerfeld coannihilation 2205.04486 1411.6930"}	This interval covers the spectrum from the canonical thermal relic mass of \(\sim\) 1.1 TeV down to the sub-TeV regime, allowing us to investigate the viability of lighter Higgsino dark matter. May 25...
328	hep-research-aid list_files	{"directory": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_180028/figures_ 2609.04175"}	Files in /Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_180028/ figures_2609.04175: - fig01.png

#	Tool	Arguments	Outcome
329	board post_finding	{"claim": "Thermal higgsino annihilation cross-section from perturbative freeze-out (compute_relic_density, is_higgsino=true) is sigma_v = 3.03e-26 cm3/s at 1.0 TeV (Omega h2=0.0898), 2.50e-26 at 1.1 TeV (0.1083), 2.10e-...	Recorded finding f-052.
330	board post_finding	{"claim": "Web search identifies Fermi-LAT dwarf spheroidal (Ackermann 2015, 1611.03184, 14-yr line search 10 GeV-2 TeV) and CTAO-North Galactic Center higgsino sensitivity (2025-2026) as relevant references for thermal ...	Recorded finding f-053.
331	board post_finding	{"claim": "Canonical pure-higgsino thermal relic mass is ~1.1 TeV within band 1.0-1.2 TeV including Sommerfeld enhancement and coannihilation corrections; perturbative freeze-out without Sommerfeld gives ~ 1.15-1.16 TeV (...	Recorded finding f-054.
332	board post_finding	{"claim": "(m_chi, delta) scan 900-1300 GeV x 100-650 keV (15x15=225 pts) yields 180/225 points passing relic+LZ+Fermi+CMB; 0 points predict 0.3-5 LZ events in extended NR search, scan saved to scan_idm.npz", "evidence_t...	Recorded finding f-055.
333	board post_finding	{"claim": "At m_chi=1100 GeV, inelastic kinematics: delta=200 keV v_min(248keV)= 579 km/s accessible N=51158, delta=350 keV v_min=762 km/s accessible N=1.97 peak 315 keV, delta=500 keV v_min=945 km/s inaccessible N=0 peak...	Recorded finding f-056.

#	Tool	Arguments	Outcome
334	board post_finding	{"claim": "Z-mediated inelastic sigma_n=7.43e-39 cm2 at 1100 GeV vs elastic Higgs-mediated suppressed to <1e-48 cm2 (>9 dex suppression); elastic rate would be 1.1e7 events if unsuppressed", "evidence_ tool_call_ids": ["t...	Recorded finding f-057.
335	hep-research-aid list_files	{"directory": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ data"}	Files in /Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ data: - downloads
336	hep-research-aid list_files	{"directory": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ research"}	Files in /Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ research: - default
337	hep-research-aid list_files	{"directory": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ data/downloads"}	No files found in / Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ data/downloads
338	hep-research-aid list_files	{"directory": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ research/default"}	Files in /Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ research/default: - board.json - runs - tool-calls.jsonl
339	local web_search	{"query": "CTAO North higgsino thermal dark matter Galactic Center sensitivity arxiv 2506 2503"}	Jun 9, 2025 · We show that, in contrast, higgsino detection could be possible within this decade with CTAO-North in La Palma, Spain. The Galactic Center is the region where the dominant DM annihilatio...

#	Tool	Arguments	Outcome
340	serve-idm.sh plot_constraint_summary	{"scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz"}	{ "plot_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png", "best_fit": { "m_chi_GeV": null, "delta_keV": null } }
341	local web_search	{"query": "Fermi-LAT dwarf spheroidal 95% CL upper limit annihilation cross section 1 TeV bbbar W+W- value"}	Oct 1, 2015 · Dwarf spheroidal galaxies (dSphs) are promising targets for the indirect detection of dark matter through gamma-ray emission due to their proximity, lack of astrophysical backgrounds and...
342	board post_finding	{"claim": "arXiv:2609.01504 Higgsino Above Sea of Fog derives LZ/XENONnT purity bounds on gaugino mixing and shows 1.1 TeV thermal higgsino survives only at high purity and small delta near neutrino fog with benchmark de...	Recorded finding f-058.
343	board post_finding	{"claim": "arXiv:2609.04175 Higgsino Interpretation of LZ Event shows thermal 1.1 TeV higgsino with delta~ 350 keV would generically produce more high-energy recoil events in LZ high-energy sideband (800< S1c<1700 phd) whe...	Recorded finding f-059.
344	board post_finding	{"claim": "Constraint summary plot from 900-1300 GeV x 100-650 keV scan generated at /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png; no best-fit point (m_chi, delta) predicts 0.3...	Recorded finding f-060.

#	Tool	Arguments	Outcome
345	board post_finding	{"claim": "arXiv:2609.33869 Heavy Higgsino with Low Reheating shows non-standard low-reheating cosmology (T_rh <=595 GeV) can accommodate heavy higgsino DM above TeV while explaining LZ 248 keV high-recoil event, relaxin...	Recorded finding f-061.
346	board post_dead_end	{"what": "Extract verbatim Planck 2018 Omega h^2 = 0.120 +/-0.001 quote from 1807.06209.txt via read_ arxiv_chunk and cite() to .bib", "why": "Tool budget exhausted before read_ arxiv_chunk/cite could be executed; 1807.062...	Recorded dead end x-13.
347	board post_dead_end	{"what": "Cross-check thermal mass against literature Sommerfeld+coannihilation+bound-state values 1411.6930 (~1172 GeV at 0.126/0.127) and 2205.04486 doublet mass with bound states", "why": "Tool budget exhausted after ...	Recorded dead end x-14.
348	board post_dead_end	{"what": "Run check_all_ constraints at thermal mass 1159 GeV and at 1000,1100, 1200 GeV for delta=200,350, 500 keV to record relic/LZ/ Fermi/CMB flags", "why": "Tool budget exhausted before check_all_constraints could be ex...	Recorded dead end x-15.
349	board post_dead_end	{"what": "Extract quantitative Fermi-LAT dwarf/GC line 95% CL limits and CTAO-North projected sensitivity at 1.1 TeV with verbatim quotes, cite() entries, and get_paper_ figures limit plots", "why": "Tool budget exhausted...	Recorded dead end x-16.

#	Tool	Arguments	Outcome
350	board post_dead_end	{"what": "Read LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limit curves at 1 TeV and xenon neutrino fog value at 1 TeV from XLZD Design Book 2410.17137; quantify decades below LZ/XENONnT for <1e-48 cm2 elastic predi...	Recorded dead end x-17.
351	hep-research-aid get_paper_figures	{"paper_id": "2205.04486", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028", "max_figures": 4}	{"status": "success", "files": ["/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028/figures_2205.04486/fig06.png"], "message": "Extracted 1 of 3 figures from... [Fig. 27]
352	board post_finding	{"claim": "Extracted Figure 6 from arXiv:2205.04486 The last Complex WIMPs standing showing direct detection and collider signatures in delta m0 vs delta m_QM plane for 2_1/2 (higgsino doublet) with green excluded region...	Recorded finding f-062.
353	board post_dead_end	{"what": "Systematic verbatim quote extraction and cite() for thermal-mass band 1.0-1.2 TeV (canonically 1.1 TeV) from >=2 papers plus 2609.01504/04175/33869 purity/fog/delta claims", "why": "Tool budget exhausted before...	Recorded dead end x-18.
354	board post_dead_end	{"what": "Extract figures via get_paper_figures for arXiv:2609.01504 and arXiv:2609.33869 and additional figures from 1411.6930/1912.07795/1104.0228", "why": "2609.01504 and 2609.33869 have no usable HTML rendering (PDF-...	Recorded dead end x-19.

#	Tool	Arguments	Outcome
355	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
356	hep-research-aid list_agent_files	{}	[]
357	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }
358	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028: - figures_2205.04486 - figures_2609.04175
359	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
360	serve-idm.sh scan_mass_splitting	{"m_chi_min": 900.0, "m_chi_max": 1300.0, "delta_min": 100.0, "delta_max": 650.0, "n_points": 15}	{ "scan_file": "/Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz", "grid": "15 x 15 = 225 points", "ranges": { "m_chi_GeV": [900.0, 1300.0], ...

#	Tool	Arguments	Outcome
361	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 200.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 180.0, "v_min_at_p...
362	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 350.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429930732553727e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 315.1, "v_min_at_p...
363	serve-idm.sh predict_recoil	{"m_chi_GeV": 1100.0, "delta_keV": 500.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
364	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd"}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
365	serve-idm.sh compute_relic_density	{"m_chi_GeV": 900.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
366	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.

#	Tool	Arguments	Outcome
367	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
368	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1150.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
369	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1160.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
370	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
371	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1300.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
372	board post_finding	{"claim": "(m_chi, delta) scan 900-1300 GeV x 100-650 keV (15x15=225 pts) yields 180/225 points passing relic+LZ+Fermi+CMB constraints and 0 points predicting 0.3-5 LZ events; scan saved to /Users/nramachandra/github-int...	Recorded finding f-063.

#	Tool	Arguments	Outcome
373	board post_finding	{"claim": "At m_chi=1100 GeV, inelastic higgsino with delta=200 keV peaks at E_R=180 keV (v_min= 572 km/s) and requires v_ min=579 km/s at 248 keV (accessible, v_esc+v_E=776 km/s), predicting N=51157.5 events in 2.84 t-yr ...	Recorded finding f-064.
374	board post_finding	{"claim": "At m_chi=1100 GeV, inelastic higgsino with delta=350 keV peaks at E_R=315.1 keV (v_min= 757 km/s) and requires v_ min=762 km/s at 248 keV (accessible, v_esc+v_E=776 km/s), predicting N=1.97 events in LZ extended...	Recorded finding f-065.
375	board post_finding	{"claim": "Z-mediated inelastic cross section at 1100 GeV is sigma_n= 7.43e-39 cm2 (tool auto for higgsino) with v_esc+v_E= 776 km/s threshold; delta= 350 keV is near kinematic limit delta_max~364 keV on Xe, explaining shar...	Recorded finding f-066.
376	board post_finding	{"claim": "serve-idm relic model is perturbative freeze-out (Kolb & Turner Eq 5.26-5.29, Arkani-Hamed hep-ph/0601041) with fixed g*=86.25, x_f~ 24.4-24.8, no Sommerfeld/ bound-state/coannihilation delta-dependence; Planck ...	Recorded finding f-067.
377	board post_dead_end	{"what": "Generate recoil spectrum plots (plot_recoil_ spectrum) for delta=200,350, 500 keV vs elastic at 1100 GeV and constraint summary plot (plot_constraint_ summary) from scan_idm.npz", "why": "Tool budget exhausted aft...	Recorded dead end x-20.

#	Tool	Arguments	Outcome
378	board post_finding	{"claim": "Pure-higgsino thermal relic grid from serve-idm perturbative freeze-out: Omega h^2 = 0.0731 at 900 GeV, 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1181 at 1150 GeV, 0.1202 at 1160 GeV, 0.1285 at 1200 GeV, 0.1503...	Recorded finding f-068.
379	board post_dead_end	{"what": "Read LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits at 1 TeV, xenon neutrino fog at 1 TeV from XLZD 2410.17137, and quantify decades below LZ for elastic <1e-48 cm2 prediction (Chen-Hill 1912.07795)", ...	Recorded dead end x-21.
380	board post_finding	{"claim": "Interpolated thermal mass where Omega h^2=0.120 is 1159 GeV (1150-1170 GeV band; 1160 GeV point 0.1202 within 0.1 sigma of Planck 0.12 +/ -0.0012)", "evidence_tool_ call_ids": ["tc-261102_ 1-w1-d-15-33", "tc-2611...	Recorded finding f-069.
381	board post_dead_end	{"what": "Test annual modulation at 248 keV with predict_annual_modulation and run check_all_ constraints benchmarks for mixing-induced visibility (2606.31013 Eq.33)", "why": "Tool budget exhausted; predict_annual_modulat...	Recorded dead end x-22.
382	board post_finding	{"claim": "Thermal higgsino annihilation cross-section perturbative: sigma_v= 3.03e-26 cm3/s at 1.0 TeV, 2.50e-26 at 1.1 TeV, 2.10e-26 at 1.2 TeV; Planck 0.12 matched at ~1.15-1.16 TeV in tool without Sommerfeld/bound-sta...	Recorded finding f-070.

#	Tool	Arguments	Outcome
383	board post_finding	{"claim": "Output directory /Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_180028 contains only figures_2609.04175 and figures_2205.04486 subdirs from prior waves; no new scan/reli...	Recorded finding f-071.
384	board post_dead_end	{"what": "Recompute pure-higgsino relic grid at 900/1000/1100/1150/1160/ 1200/1300 GeV, interpolate thermal mass, extract verbatim Planck 1807.06209 Omega h^2=0.120 +/-0.001 quote via read_arxiv_ chunk+cite, run check_all_...	Recorded dead end x-23.
385	hep-research-aid list_agent_files	{}	[]
386	hep-research-aid list_files	{}	Files in .: - .git - .gitignore - .last-release-seen - ISSUES.md - SCOPE.md - docs - downloads - logs - projects - reports - research-results - runs - scripts - venvs - w...
387	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_ data()." }
388	hep-research-aid list_datasets	{}	{"count": 0, "datasets": []}
389	hep-research-aid get_dispatch	{}	Dispatch: local execution.
390	local web_search	{"query": "Higgsino thermal target direct detection LZ constraints"}	We examine Higgsino dark matter (DM) as a potential explanation of the 248 keV nuclear-recoil event of interest reported by the LUX-ZEPLIN (LZ) experiment, taking into account IceCube constraints on D...

#	Tool	Arguments	Outcome
391	local web_search	{"query": "Higgsino dark matter thermal relic density 1 TeV inelastic direct detection"}	Sep 2, 2026 · In the MSSM, we consider a Higgsino dark matter that saturates the relic density. Apr 4, 2025 · Direct detection experiments imply purity constraints on a dark matter higgsino, limiting ...
392	local web_search	{"query": "Higgsino mass splitting direct detection loop LZ Xenon"}	Sep 2, 2026 · As we will show in this paper, a TeV Higgsino with mass splitting around 350 keV that has inelastic spin independent scattering with the Xenon target provides a possible interpretation o...
393	local web_search	{"query": "arxiv Higgsino thermal target 1.1 TeV relic density"}	This interval covers the spectrum from the canonical thermal relic mass of $\sim$ 1.1 $\sim$1.1$ TeV down to the sub-TeV regime, allowing us to investigate the viability of lighter Higgsino dark matter. We inv...
394	local web_search	{"query": "arxiv Higgsino inelastic dark matter LZ 248 keV"}	Sep 1, 2026 · We propose Higgsino dark matter as a potential interpretation of the 248 keV nuclear-recoil event of interest reported by the LUX-ZEPLIN (LZ) experiment. LZ studied several rare backgrou...
395	local web_search	{"query": "arxiv Higgsino direct detection loop cross section purity"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=arxiv%20Higgsino%20direct%20detectio

#	Tool	Arguments	Outcome
396	local web_search	{"query": "\"248 keV\" Higgsino LZ arXiv 2609"}	Sep 1, 2026 · We propose Higgsino dark matter as a potential interpretation of the 248 keV nuclear-recoil event of interest reported by the LUX-ZEPLIN (LZ) experiment. LZ studied several rare backgrou...
397	local web_search	{"query": "Higgsino thermal target 1.1 TeV relic density arXiv 2504"}	Sep 2, 2026 · Subject to this mapping, the 500 h exposure, the full-density assumption, and the fixed cuspy Einasto halo, the projected Omega sensitivity reaches the thermal Higgsino target selected b...
398	local web_search	{"query": "Pospelov Ramani solar capture Higgsino arXiv"}	Sep 2, 2026 · Higgsino dark matter remains one of the most convincing dark matter candidates. It can exist in a "quasi-Dirac" regime, when the splitting between two Majorana components is on the orde...
399	local web_search	{"query": "Wen Yin PQ symmetric high scale SUSY LZ arXiv"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=Wen%20Yin%20PQ%20symmetric%20high%20s
400	local web_search	{"query": "arXiv:2609.01583 Higgsino LZ 248 keV"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=arXiv%3A2609.01583%20Higgsino%20LZ%20 > er...

#	Tool	Arguments	Outcome
401	local web_search	{"query": "arXiv Higgsino solar capture IceCube 2026 Pospelov"}	Sep 2, 2026 · Confronting it with the non-observation of high-energy neutrinos from the Sun by the IceCube experiment, we obtain a robust limit, δ> 566 \,keV. These limits exclude the interpretation o...
402	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
403	hep-research-aid list_agent_files	{}	[]
404	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857"}	No files found in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857
405	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437: - 1104.0228.pdf - 1104.0228.txt - 1411.6930.pdf - 1411.6930.txt - 1807.06209.pdf - 1...
406	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857"}	No files found in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857

#	Tool	Arguments	Outcome
407	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
408	hep-research-aid list_agent_files	{}	[]
409	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857"}	No files found in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857
410	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437: - 1104.0228.pdf - 1104.0228.txt - 1411.6930.pdf - 1411.6930.txt - 1807.06209.pdf - 1...
411	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib"}	@article{arxiv260909830, title = {Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter}, author = {Arindam Chatterjee and Debottam Das and ...
412	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2410.17036.txt"}	Title: Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment Authors: J. Aalbers, D. S. Akerib, A. K. Al Musalhi, F. Alder, C. S. Amarasinghe, A. Ames, T. J. An...

#	Tool	Arguments	Outcome
413	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1912.07795.txt"}	Title: Direct detection rate of heavy Higgsino-like and Wino-like dark matter Authors: Qing Chen, Richard J. Hill Year: 2019 arXiv ID: 1912.07795 ==========...===
414	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib"}	@article{arxiv260909830, title = {Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter}, author = {Arindam Chatterjee and Debottam Das and ...
415	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1411.6930.txt"}	Title: Heavy neutralino relic abundance with Sommerfeld enhancements - a study of pMSSM scenarios Authors: M. Beneke, C. Hellmann, P. Ruiz-Femenia Year: 2014 arXiv ID: 1411.6930 ==========...==
416	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt"}	Title: Planck 2018 results. VI. Cosmological parameters Authors: Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Ba...
417	local web_search	{"query": "arXiv 2609.01583 Higgsino LZ 248 keV inelastic dark matter"}	We propose Higgsino dark matter as a potential interpretation of the 248 keV nuclear-recoil event of interest reported by the LUX-ZEPLIN (LZ) experiment. LZ studied several rare background processes a...

#	Tool	Arguments	Outcome
418	local web_search	{"query": "arXiv 2609.01590 solar capture IceCube Higgsino delta 566 keV"}	Sep 1, 2026 · Near elastic-scattering blind spots, solar-captured higgsinos fail to reach thermal equilibrium, weakening their annihilation signal, so that ten years of IceCube data exclude δm ? 475 k...
419	local web_search	{"query": "IceCube solar dark matter W+W- limit 2024 2025 capture rate"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=IceCube%20solar%20dark%20matter%20W%...
420	local web_search	{"query": "LMC high speed tail dark matter halo v_esc 850 km/s inelastic scattering"}	Sep 29, 2026 · In particular, the velocity distribution of DM in the solar neighbourhood determines the fraction of particles capable of producing a given recoil. This becomes especially important for...
421	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/tool-results/tc-261102_1-w1-d-21-29.txt"}	Title: Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment Authors: J. Aalbers, D. S. Akerib, A. K. Al Musalhi, F. Alder, C. S. Amarasinghe, A. Ames, T. J. An...
422	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/tool-results/tc-261102_1-w1-d-21-30.txt"}	Title: Direct detection rate of heavy Higgsino-like and Wino-like dark matter Authors: Qing Chen, Richard J. Hill Year: 2019 arXiv ID: 1912.07795 ========== ========== ===...

#	Tool	Arguments	Outcome
423	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/figures_2410.17036"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/figures_2410.17036: - fig01.png - fig04.png
424	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/figures_2303.14729"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/figures_2303.14729: - fig01.png - fig02.png - fig03.png
425	local web_search	{"query": "Bottaro Strumia 2205.04486 thermal Higgsino mass Sommerfeld bound state"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=Bottaro%20Strumia%202205.04486%20ther
426	local web_search	{"query": "Beneke Hellmann Ruiz 1411.6930 Higgsino relic abundance Sommerfeld thermal mass 2.7 TeV"}	Nov 25, 2014 · We present a detailed discussion of Sommerfeld enhancements in neutralino dark matter relic abundance calculations for several popular benchmark scenarios in the general MSSM. 2085, E-4...
427	local web_search	{"query": "arXiv 2609.01590 Higgsino solar capture IceCube"}	Sep 1, 2026 · Near elastic-scattering blind spots, solar-captured higgsinos fail to reach thermal equilibrium, weakening their annihilation signal, so that ten years of IceCube data exclude δm ? 475 k...

#	Tool	Arguments	Outcome
428	local web_search	{"query": "Higgsino thermal mass bound state formation 2022 2023 2024 Complex WIMP Strumia"}	Search failed: DNSError: DNSError('error sending request for url (https://wt.wikipedia.org/w/api.php?action=opensearch&profile=fuzzy&limit=1&search=Higgsino%20thermal%20mass%20bound%20s
429	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1000.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1000.0, "is_higgsino": true, "sigma_v_cm3_s": 3.0250000000000006e-26 }, "relic_density": { "omega_h2": 0.0898, "planck_value": 0.12, "planc...
430	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1100.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1100.0, "is_higgsino": true, "sigma_v_cm3_s": 2.5e-26 } , "relic_density": { "omega_h2": 0.1083, "planck_value": 0.12, "planck_error": 0.001...
431	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1159.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1159.0, "is_higgsino": true, "sigma_v_cm3_s": 2.2519487731904195e-26 }, "relic_density": { "omega_h2": 0.12, "planck_value": 0.12, "planck_...
432	serve-idm.sh compute_relic_density	{"m_chi_GeV": 1200.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1200.0, "is_higgsino": true, "sigma_v_cm3_s": 2.1006944444444443e-26 }, "relic_density": { "omega_h2": 0.1285, "planck_value": 0.12, "planc...

#	Tool	Arguments	Outcome
433	serve-idm.sh compute_relic_density	{"m_chi_GeV": 2800.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 2800.0, "is_higgsino": true, "sigma_v_cm3_s": 3.858418367346939e-27 }, "relic_density": { "omega_h2": 0.6382, "planck_value": 0.12, "planck...
434	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/tool-results/tc-261102_1-w1-d-21-38.txt"}	Title: Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment Authors: J. Aalbers, D. S. Akerib, A. K. Al Musalhi, F. Alder, C. S. Amarasinghe, A. Ames, T. J. An...
435	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/tool-results/tc-261102_1-w1-d-21-39.txt"}	Title: Direct detection rate of heavy Higgsino-like and Wino-like dark matter Authors: Qing Chen, Richard J. Hill Year: 2019 arXiv ID: 1912.07795 ========== ========== ===...
436	local web_search	{"query": "arXiv 2609.01504 Fan Reece Higgsino inelastic LZ"}	Sep 1, 2026 · It is a simple realization of the "inelastic dark matter" mechanism, where its scattering off a nucleus occurs between two nearly degenerate states. Intriguingly, very recently the LZ co...
437	hep-research-aid get_paper_figures	{"paper_id": "2609.01583", "output_dir": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857", "max_figures": 4}	{"status": "error", "files": [], "message": "No usable HTML rendering for arXiv:2609.01583 (arxiv-html: https://arxiv.org/html/2609.01583 contains no extractable figures; ar5iv: https://ar5iv.labs.arx...

#	Tool	Arguments	Outcome
438	board post_finding	{"claim": "serve-idm.sh perturbative compute_relic_density gives thermal Higgsino mass 1159 GeV for Omega h2=0.120, with Omega scaling as (m/1159)^2: 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1285 at 1200 GeV, 0.6382 at ...	Recorded finding f-072.
439	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1411.6930.txt"}	Title: Heavy neutralino relic abundance with Sommerfeld enhancements - a study of pMSSM scenarios Authors: M. Beneke, C. Hellmann, P. Ruiz-Femenia Year: 2014 arXiv ID: 1411.6930 ========== ==========...
440	serve-idm.sh predict_recoil	{"m_chi_GeV": 1080.0, "delta_keV": 350.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1080.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429695891375652e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 314.5, "v_min_at_p...
441	serve-idm.sh predict_recoil	{"m_chi_GeV": 1080.0, "delta_keV": 500.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1080.0, "delta_keV": 500.0, "sigma_n_cm2": 7.429695891375652e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 449.2, "v_min_at_p...
442	serve-idm.sh predict_recoil	{"m_chi_GeV": 1080.0, "delta_keV": 566.0, "is_higgsino": true}	{ "parameters": { "m_chi_GeV": 1080.0, "delta_keV": 566.0, "sigma_n_cm2": 7.429695891375652e-39, "is_higgsino": true }, "kinematics": { "E_R_peak_keV": 508.5, "v_min_at_p...

#	Tool	Arguments	Outcome
443	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1080.0, "delta_keV": 350.0, "is_ higgsino": true}	{ "parameters": { "m_chi_GeV": 1080.0, "delta_keV": 350.0, "sigma_n_cm2": 7.429695891375652e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1045, "tension_si...
444	serve-idm.sh check_all_ constraints	{"m_chi_GeV": 1080.0, "delta_keV": 566.0, "is_ higgsino": true}	{ "parameters": { "m_chi_GeV": 1080.0, "delta_keV": 566.0, "sigma_n_cm2": 7.429695891375652e-39, "is_higgsino": true } , "relic_density": { "omega_h2": 0.1045, "tension_si...
445	hep-research-aid get_paper_figures	{"paper_id": "2410.17036", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_181857", "max_ figures": 4}	{"status": "success", "files": ["/Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_181857/ figures_2410.17036/ fig01.png", "/ Users/nramachandra/ hep-genesis-testing/ p... [Fig. 28, 29]
446	hep-research-aid get_paper_figures	{"paper_id": "2303.14729", "output_dir": "/ Users/nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_181857", "max_ figures": 4}	{"status": "success", "files": ["/Users/ nramachandra/ hep-genesis-testing/ projects/ ramp-research-higgsino-dd/ runs/261102_181857/ figures_2303.14729/ fig01.png", "/ Users/nramachandra/ hep-genesis-testing/ p... [Fig. 30, 31, 32]

#	Tool	Arguments	Outcome
447	board post_finding	{"claim": "Planck 2018 base LambdaCDM gives dark matter density Omega_c h2 = 0.120 ± 0.001 (68% CL) from combined TT,TE, EE+lowE+lensing, as stated in abstract of 1807.06209", "evidence_tool_call_ids": ["tc-261102_1-w1-d-...	Recorded finding f-073.
448	board post_finding	{"claim": "1411.6930 finds pure-higgsino relic abundances with perturbative vs Sommerfeld-enhanced rates almost coincide, leading to almost vanishing net Sommerfeld correction, implying perturbative thermal mass ~1.14-1....	Recorded finding f-074.
449	board post_finding	{"claim": "serve-idm.sh compute_relic_density is perturbative only: sigma_v = 3.025e-26*(1100/m)^2 cm3/s, x_f 24.5-24.7, g_star 86.25, no Sommerfeld, no bound-state formation, no thermal mass corrections, no running coup...	Recorded finding f-075.
450	local web_search	{"query": "Fermi-LAT dwarf spheroidal combined limit annihilation cross section 1 TeV 2023 2024"}	Nov 8, 2023 · Previous studies using the Fermi-Large Area Telescope (LAT) of DM emission from dSphs have provided the most robust and stringent constraints on the DM annihilation cross section and mas...
451	local web_search	{"query": "XLZD DARWIN design book 2410.17137 projected sensitivity 1 TeV neutrino fog"}	Oct 22, 2024 · It is based on the mature liquid xenon time projection chamber technology used in current-generation experiments, LZ and XENONnT. The report discusses the baseline design and opportunit...

#	Tool	Arguments	Outcome
452	board post_dead_end	{"what": "Fetch and verbatim quote 2205.04486 (Bottaro et al Complex WIMPs / thermal Higgsino mass with Sommerfeld+bound-state formation) and recent 2022-2025 Beneke NLO Sommerfeld papers for quantitative shift vs pertur...	Recorded dead end x-24.
453	board post_dead_end	{"what": "Add verbatim Planck 2018 Omega_c h2 quote to references.bib via cite() tool", "why": "cite() requires tool call with validated verbatim quote; budget reached before cite could be executed. Quote identified in 1...	Recorded dead end x-25.
454	board post_finding	{"claim": "predict_recoil at m_chi=1080 GeV, delta=350 keV gives v_min(248 keV)=763 km/s <776 km/s cutoff, N_ expected=1.79 events in 2.84 t-yr (Poisson P>=1=0.83) consistent with LZ 1 event; at delta=500 keV v_min=946 km...	Recorded finding f-076.
455	board post_finding	{"claim": "check_all_ constraints at 1.08 TeV, delta=350 keV returns VIABLE (Omega h2=0.1045, N_LZ=1.79, Fermi sigma_v=6.8e-42 < 2.24e-25, CMB p_ann=9.5e-46 <3.2e-28); at delta=566 keV also returns VIABLE on relic/indirect...	Recorded finding f-077.
456	board post_finding	{"claim": "Standard halo delta_max = mu_A v_max^2/2 ~365 keV at v_max=776 km/s vs ~438 keV at LMC-boosted 850 km/s for m_chi=1080 GeV on xenon (mu~109 GeV) , so even 850 km/s tail cannot make delta=500-566 keV kinematical...	Recorded finding f-078.

#	Tool	Arguments	Outcome
457	board post_finding	{"claim": "Literature tension confirmed via web search: arXiv:2609.01583 proposes ~1 TeV Higgsino inelastic DM for LZ 248 keV event (2.84 t-yr, 2.6 sigma), while solar-capture counterpart (2609.01590 / IceCube 10-yr) der...	Recorded finding f-079.
458	board post_dead_end	{"what": "Extract verbatim quotes and figures from arXiv:2609.01583 and solar-capture paper (2609.01590) via get_paper_figures/read_arxiv_chunk/cite, and run full 5-delta table plus annual modulation and scan_mass_splitt...	Recorded dead end x-26.
459	board post_finding	{"claim": "Chen & Hill 1912.07795 pure-Higgsino (pure doublet) SI cross section upper limit is <1e-48 cm2 in TeV mass regime, below xenon neutrino floor, with amplitude-level cancellation", "evidence_tool_call_ids": ["tc...	Recorded finding f-080.
460	board post_finding	{"claim": "LZ 2410.17036 4.2 tonne-year SI limit: strongest exclusion 2.2e-48 cm2 at 40 GeV/c2 (90% CL) , best median sensitivity 5.1e-48 cm2 at 40 GeV/c2; limit at 1 TeV not quoted in text excerpt and requires figure rea...	Recorded finding f-081.
461	board post_finding	{"claim": "LZ 2410.17036 and XENONnT 2303.14729 figure extraction completed: LZ returned 2 figures (Fig1 detector data, Fig S4 salt distribution), XENONnT returned 3 figures (Fig1 calibration, Fig A2/A3 spatial distribut...	Recorded finding f-082.

#	Tool	Arguments	Outcome
462	board post_finding	{"claim": "Prior f-008 linear extrapolation ~6e-47 cm2 at 1.1 TeV is invalid; correct comparison requires direct read of LZ/XENONnT limit at 1 TeV from figure (expected ~2-3e-47 cm2 scale) vs Chen-Hill bound <1e-48 cm2, ...	Recorded finding f-083.
463	board post_dead_end	{"what": "Direct read of LZ 2410.17036 and XENONnT 2303.14729 SI limits at 1.0-1.2 TeV from text/table/ figure caption", "why": "Extracted text contains only 40 GeV point (2.2e-48 cm2); get_paper_figures HTML rendering re...	Recorded dead end x-27.
464	board post_dead_end	{"what": "cite() addition of Chen & Hill 1912.07795 verbatim quotes to references.bib and XLZD 2410.17137 / Fermi-LAT dwarf verbatim quotes", "why": "Tool budget exhausted before cite() and read_ arxiv_chunk could be exec...	Recorded dead end x-28.
465	board post_dead_end	{"what": "Numeric xenon neutrino fog/floor at 1 TeV and XLZD projection at 1 TeV", "why": "No numeric fog value in extracted LZ or Chen & Hill text excerpts; Chen & Hill only states \ "below neutrino backgrounds/ floor [26...	Recorded dead end x-29.
466	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
467	hep-research-aid list_agent_files	{}	[]

#	Tool	Arguments	Outcome
468	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857: - figures_2303.14729 - figures_2410.17036 - tool-results
469	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437: - 1104.0228.pdf - 1104.0228.txt - 1411.6930.pdf - 1411.6930.txt - 1807.06209.pdf - 1...
470	hep-research-aid list_input_files	{}	{ "status": "success", "files": [], "usage": "Pass any of these filenames to load_table_data()." }
471	engine describe_tools	{}	{ "server": "inelastic-dm-server", "description": "Bridges particle physics (WIMP mass, mass splitting, cross section) with cosmological observables (relic density, recoil spectra, indirect detect...
472	hep-research-aid get_dispatch	{}	Dispatch: local execution.
473	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857: - figures_2303.14729 - figures_2410.17036 - tool-results
474	hep-research-aid list_agent_files	{}	[]

#	Tool	Arguments	Outcome
475	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437: - 1104.0228.pdf - 1104.0228.txt - 1411.6930.pdf - 1411.6930.txt - 1807.06209.pdf - 1...
476	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/tool-results"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/tool-results: - tc-261102_1-w1-d-19-32.txt - tc-261102_1-w1-d-19-33.txt - tc-261102_1-w1-d-...
477	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd"}	Files in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd: - chats - checkpoint.db - checkpoint.db-shm - checkpoint.db-wal - data - jobs - project.json - prov...
478	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2609.09830.txt"}	Title: Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter Authors: Arindam Chatterjee, Debottam Das, Syed Adil Pasha, Alexander Pukhov, Rahul...

#	Tool	Arguments	Outcome
479	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt"}	Title: Complementary Probes of Light Higgsinos: Electroweak Precision Measurements and Dark Matter Direct Detection Authors: Koichi Hamaguchi, Natsumi Nagata, Genta Osaki Year: 2026 arXiv ID: 2606.310...
480	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib"}	@article{arxiv260909830, title = {Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter}, author = {Arindam Chatterjee and Debottam Das and ...
481	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib"}	@article{arxiv260909830, title = {Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter}, author = {Arindam Chatterjee and Debottam Das and ...
482	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1411.6930.txt"}	Title: Heavy neutralino relic abundance with Sommerfeld enhancements - a study of pMSSM scenarios Authors: M. Beneke, C. Hellmann, P. Ruiz-Femenia Year: 2014 arXiv ID: 1411.6930 ========== ==========...
483	hep-research-aid read_text_file	{"filepath": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt"}	Title: Planck 2018 results. VI. Cosmological parameters Authors: Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Ba...

#	Tool	Arguments	Outcome
484	hep-research-aid list_files	{"directory": "/Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd"}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
485	board post_finding	{"claim": "Perturbative serve-idm.sh thermal Higgsino mass is 1159 GeV for Omega h2=0.120 (Planck 2018), with Omega scaling (m/1159)^2: 0.0898 at 1000 GeV (25.1 sigma low), 0.1083 at 1100 GeV, 0.1285 at 1200 GeV, using s...	Recorded finding f-084.
486	board post_finding	{"claim": "1411.6930 Sec 4-5 verbatim: pure-higgsino relic abundances with perturbative vs Sommerfeld-enhanced rates almost coincide, leading to almost vanishing net Sommerfeld correction, so NLO Sommerfeld without bound...	Recorded finding f-085.
487	board post_finding	{"claim": "Planck 2018 base LambdaCDM gives Omega_c h2 = 0.120 ± 0.001 (68% CL) from TT,TE,EE+lowE+lensing, abstract of 1807.06209, defining thermal target Omega h2=0.12 ±0.0012 used for 1159 GeV anchor", "evidence_tool_...	Recorded finding f-086.
488	serve-idm.sh predict_recoil	{"m_chi_GeV": 1080.0, "delta_keV": 400.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.
489	serve-idm.sh predict_recoil	{"m_chi_GeV": 1080.0, "delta_keV": 600.0, "is_higgsino": true}	**error**: Tool error: the run's tool budget is exhausted -- this call was not executed. Post your findings now (board posts are free) and reply with the final JSON.

#	Tool	Arguments	Outcome
490	board post_dead_end	{"what": "Extract verbatim thermal mass table from 2205.04486 Sec 3-4 (Sommerfeld only vs Sommerfeld+bound-state with uncertainty) via read_arxiv_chunk", "why": "Tool budget exhausted before any read_arxiv_chunk/web_sear...	Recorded dead end x-30.
491	board post_dead_end	{"what": "Run compute_relic_density at 1159 GeV and 1000 GeV in wave 2 to anchor perturbative baseline and add cite() for Planck 2018 Omega_c h2=0.120±0.001 and 2205.04486 thermal-mass table to references.bib", "why": "T...	Recorded dead end x-31.
492	board post_finding	{"claim": "predict_recoil at m_chi=1080 GeV gives N_expected=1.79 (v_min=763 km/s) at delta=350 keV, N=0.0 (v_min=946 km/s) at delta=500 keV, N=0.0 (v_min=1026 km/s) at delta=566 keV for 248 keV recoil in 2.84 t-yr xenon...	Recorded finding f-087.
493	board post_finding	{"claim": "Analytic delta_max = mu*v_max^2/2 gives ~365 keV at v_max=776 km/s vs ~438 keV at LMC-boosted 850 km/s for m_chi=1080 GeV on xenon (mu~109 GeV), so even 850 km/s tail cannot make delta=500-566 keV kinematicall...	Recorded finding f-088.
494	board post_finding	{"claim": "check_all_constraints at m_chi=1.08 TeV returns VIABLE for both delta=350 keV (Omega h2=0.1045, N_LZ=1.79) and delta=566 keV (N_LZ=0.0) on relic+Fermi+CMB, because tool omits solar-capture/IceCube constraint; ...	Recorded finding f-089.

A.5 Files produced

- 1104.0228.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1104.0228.txt
- 1411.6930.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1411.6930.txt
- 2609.09830.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2609.09830.txt
- 2410.17036.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2410.17036.txt
- 2606.31013.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/2606.31013.txt
- hep-ph_0610249.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/hep-ph_0610249.txt
- 1912.07795.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1912.07795.txt
- 1807.06209.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt
- 2609.01590.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.01590.txt
- 2609.07807.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.07807.txt
- 2205.04486.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2205.04486.txt
- 2610.03692.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2610.03692.txt
- 2609.09830.txt (text, download_arxiv_full_text): /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/2609.09830.txt

B Research board

The shared board of the research runs in this session, generated mechanically: what the team set out to investigate, what it found (with the tool calls that are the evidence), what failed, and how the critic judged each wave.

B.1 Directions

Id	Status	Goal	Outcome
d-01	closed	Establish the pure-higgsino thermal relic density as a function of mass (about 0.9, 1.0, 1.1, 1.2 and 1.3 TeV), with and without Sommerfeld enhancement, and find the mass at which Omega h^2 equals the Planck 2018 value of 0.120 +/- 0.001.	Not finished: only one published pure-higgsino relic-density point was found, so the success criteria (published values at three or more masses, and a thermal mass from two independent papers) are not met. (1) No relic-density or inelastic-DM compute tool exists here, and list_input_files returned no data files (tc-261016_1-w1-d-01-19, finding f-006). Every value below comes from the literature.
d-02	closed	Place the pure-higgsino elastic spin-independent cross section at about 1.0-1.2 TeV against the LZ 2024 (2410.17036) and XENONnT (2303.14729) limits at the same mass, and against the xenon neutrino fog.	Answer: a thermal pure higgsino at about 1.1 TeV lies about 2 or more orders of magnitude below the LZ 2024 limit and below the xenon neutrino floor. Three of the planned inputs were not read: the XENONnT limit, the LZ limit read from its figure at 1 TeV, and a numeric neutrino-fog value. All are recorded as open question q-06. (1) Prediction. Hisano et al. 1104.0228 work at leading order in elec
d-03	closed	Assess the routes by which a higgsino becomes visible to direct detection: the mass splittings and the inelastic Z-exchange threshold, LZ inelastic constraints (2609.09830), mixing-induced Higgs couplings in the (mu, M1/M2, tan beta) plane (2606.31013), and the (Omega h^2/0.12) rescaling for sub-thermal higgsinos.	Two of the three success criteria are met; the third is only partly met. (1) The inelastic threshold, worked by hand (f-001): m_Xe = 131 x 0.9315 = 122.0 GeV. The reduced mass mu_A is 108.7 GeV at m = 1.0 TeV and 109.8 GeV at 1.1 TeV. With v_max = 544 + 232 = 776 km/s, (v_max/c)^2 = 6.70e-6, so delta_max = mu_A v_max^2/2 is about 364 keV at 1.0 TeV and 368 keV at 1.1 TeV. The minimum-velocity form

Id	Status	Goal	Outcome
d-04	blocked	Pin down the pure-higgsino thermal mass from at least two independent published calculations, including recent work with bound-state formation, and quote the Planck 2018 Omega_c h^2 verbatim. Confirm which mass the 1411.6930 pure-higgsino values (0.126/0.127) refer to.	One of the three success criteria is met. (1) Met: the Planck 2018 value is quoted verbatim from the abstract on page 1 of arXiv:1807.06209: 'A combined analysis gives dark matter density Ωch2 = 0.120± 0.001' (68% CL). It is at roughly characters 6500-7000 of /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/1807.06209.txt (finding f-012, tc-261016_1-w2-
d-05	blocked	Replace the linear 40 GeV extrapolation with direct reads of the LZ 2024 and XENONnT SI limits at about 1 TeV, a numeric xenon neutrino-fog value at about 1 TeV, and one next-generation projection (XLZD/DARWIN). Correct the f-008 arithmetic and quote Chen and Hill verbatim.	Two things were established: the Chen and Hill quotes and the corrected f-008 arithmetic. The 1 TeV limits, the neutrino-fog value and the XLZD projection were not obtained. (1) Chen and Hill (arXiv:1912.07795), verbatim from the Summary: 'The upper limit for the pure doublet is less than 10^-48 cm2 in the TeV mass regime, consistent with the leading order estimate [18, 19], and lower than the est
d-06	closed	Verify 2606.31013 and 2609.09830 with metadata, and trace the contested f-002 and f-003 numbers to quoted excerpts: Eq. 13 (356 MeV), the f_q formula, Eq. 38, the Table VII NLO rows, and the tree-level delta_m0 formula (Eq. 9 / Ref. [28]).	Both papers are real, and every contested f-002/f-003 number except the tree-level delta_m0 formula is now traced to a verbatim excerpt. That formula is still unsourced, so the M1 needed for delta < 364 keV was not rederived. No cite() calls were made: the quotes are recorded on the board as f-015 and f-016, not in references.bib. (1) IDs confirmed by metadata (f-014, closes q-02). 2606.31013 is
d-07	open	Identify Ref. [28] of arXiv:2609.09830 and quote its Eq. (9) for the neutral splitting	
d-08	open	Add a sourced statement on indirect-detection limits for TeV higgsinos	

Id	Status	Goal	Outcome
d-09	closed	Use the serve-idm.sh tools to compute the pure-higgsino thermal relic density Omega h^2 at m_chi = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV, find the mass where Omega h^2 = 0.120, and run the combined constraint check at the thermal point.	Done. compute_relic_density (serve-idm.sh) gives Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285 and 0.1503 at m_chi = 900, 1000, 1100, 1200 and 1300 GeV. It gives 0.1181 at 1150 GeV and 0.1202 at 1160 GeV, so the thermal mass (Omega h^2 = 0.120, Planck 0.12 +/- 0.0012) is about 1159 GeV, bracketed between 1150 and 1160 GeV (f-020). This is about 1 percent below the 1172 GeV inferred from 1411.6930 and
d-10	closed	Compute the inelastic Z-mediated recoil spectrum, the LZ event expectation and the (m_chi, delta_m) constraint map for the thermal higgsino. Reproduce or test the 2609.01590 preferred region (delta_m = 481-519 keV, m_chi = 1000-1183 GeV) against the 475 keV (IceCube) and 557 keV (Super-K + IceCube) neutrino lines.	All runs were local; get_dispatch reported 'local execution'. The tool uses a standard halo with a hard cutoff at v_esc+v_E = 776 km/s, and none of its calls accept a v_esc or LMC-tail option. At m_chi = 1100 GeV (sigma_n = 7.43e-39 cm^2), predict_recoil gives these values for delta = 300/400/480/520/560 keV: peak E_R = 270/360/432/468/504 keV, v_min at the peak = 701/809/886/922/957 km/s, and v_m
d-11	blocked	Close the remaining literature gaps: the 2205.04486 doublet thermal mass, the LZ 2024 and XENONnT SI limits near 1 TeV, and verbatim citations for Planck 2018 and Chen and Hill.	Two of the five gaps are closed; the other three are still open. (1) The doublet thermal mass from 2205.04486 (Bottaro et al., Table I) is quoted verbatim. For the Dirac-fermion Y=1/2 doublet it is M_DM = 1.08 +/- 0.02 TeV, computed with Sommerfeld enhancement and bound-state formation, normalised to Omega_DM h^2 = 0.11933 +/- 0.00091. The paper also says the doublet's direct-detection cross secti

Id	Status	Goal	Outcome
d-12	closed	Extract verbatim, citable inputs from the 2026 LZ-event papers: the halo models, rates and preferred regions, the neutrino exclusion lines, the radiative corrections and the rescue scenarios.	I read the full texts of 2609.01590 and 2609.07807 and only the abstracts of 2610.03692, 2609.09830 and 2609.06890. The validated citations are in /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261022_110244/references.bib (keys arxiv260901590 and arxiv260907807). Not every number has a validated quote: the cite tool rejected several 2609.01590 sentences (the v_esc
d-13	closed	Run an analytic and tool-based kinematic check of delta_max(m_chi, v_max) at E_R = 248 keV. Identify the scan points compatible with the LZ event, and reconcile the tool's 776 km/s cutoff with the upper delta edges of 2609.01590 (profiled Maxwellian 422/457 keV; Halo 13 519/529 keV).	Main result: under the tool's halo, which cuts off at 776 km/s, no point that reproduces the LZ event has delta of 475 keV or more; the compatible band sits at about 350 keV. Everything ran locally. Points in the 0.3-5 event band were identified by bracketing with predict_recoil, because the scan file could not be read (x-07). Event counts from predict_recoil (LZ, 2.84 t-yr): - delta 340 keV: 10.
d-14	blocked	Recover quoted evidence for the contested literature numbers (f-024, f-025, f-026, f-027), and replace the tool's unphysical p-wave indirect-detection pass with an s-wave higgsino comparison at 1.1 TeV.	Two of the four goals are settled from full texts. The other two, plus the indirect-detection comparison, were not reached. (1) 2609.09830 does not give an on-shell delta_m shift for the 475-557 keV window (f-030). Its benchmark points are tuned to small one-loop on-shell splittings: delta = 36.37, 8.90, 16.39, 10.81 and 10.59 keV for BP1-BP5, at |mu| = 0.61-1.40 TeV, all with Omega h^2 < 0.12. Co

Id	Status	Goal	Outcome
d-15	blocked	Compute pure-higgsino thermal relic density Omega h^2 at m_chi = 0.9, 1.0, 1.1, 1.15, 1.2, 1.3 TeV with serve-idm tools, interpolate thermal mass where Omega h^2 = 0.120 +/- 0.001 (Planck 2018), and quantify offset from literature Sommerfeld+coannihilation+bound-state values.	d-15 pure-higgsino thermal relic grid established in prior waves and re-documented this wave via describe_tools (tc-261102_1-w3-d-15-140): serve-idm is perturbative Kolb & Turner 5.26-5.29 with fixed g*=86.25, x_f~24.4-24.8, sigma_v~1/m_chi^2, no Sommerfeld/bound-state/coannihilation delta-dependence; Planck relic reported as 0.12 +/-0.0012. Grid (f-036/f-049/f-068): Omega h^2=0.0731 (900 GeV, 3.7
d-16	closed	Survey recent arXiv literature on higgsino thermal relic target 1.0-1.2 TeV: canonical 1.1 TeV with Sommerfeld/coannihilation, light-slepton coannihilation lowering to 0.4-0.6 TeV, low-reheating raising mass, and 2026 LZ 248 keV inelastic interpretation (2609.01504, 2609.04175, 2609.33869).	Survey of higgsino thermal relic target 1.0-1.2 TeV (canonically 1.1 TeV) partially completed. Established via web_search + describe_tools: canonical pure-higgsino thermal mass ~1.1 TeV band 1.0-1.2 TeV with Sommerfeld/coannihilation (2205.04486 context) vs perturbative freeze-out ~1.15-1.16 TeV (Planck Omega h2=0.12±0.0012, Kolb & Turner). 2609.01504 Higgsino Above Sea of Fog: inelastic DM nearly
d-17	blocked	Place thermal higgsino (1.0-1.2 TeV) relative to LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits and xenon neutrino fog, distinguishing elastic (suppressed to <1e-48 cm2) vs Z-mediated inelastic (delta threshold ~364 keV on Xe) and mapping (m_chi, delta) allowed region.	d-17 partial: (1) scan_mass_splitting 900-1300 GeV x 100-650 keV 15x15=225 pts saved to /Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz with 180/225 passing relic+LZ+Fermi+CMB and 0 pts predicting 0.3-5 LZ events (f-063, tc-145). (2) predict_recoil at 1100 GeV: delta=200 keV peak 180 keV v_min 572 km/s, v_min(248keV)=579 km/s accessible N=51157.5 (f-064, tc-146); del

Id	Status	Goal	Outcome
d-18	closed	Quantify thermal higgsino annihilation cross-section <sigma v> vs Fermi-LAT dwarf/GC line limits and CTAO-North projected sensitivity	Thermal higgsino <sigma v> extracted from compute_relic_density (is_higgsino=true, perturbative Kolb & Turner, g*=86.25): sigma_v=3.03e-26 cm3/s at 1.0 TeV (Omega h2=0.0898, x_f=24.7, T_f=40.53 GeV), 2.50e-26 at 1.1 TeV (0.1083, 24.6, 44.76 GeV), 2.10e-26 at 1.2 TeV (0.1285, 24.5, 48.98 GeV); Planck Omega h2=0.12 matched at ~1.15-1.16 TeV in this tool (no Sommerfeld/bound-state/coannihilation). Fi
d-19	closed	Deepen weakest claim 1: pure-Higgsino thermal relic mass and its uncertainty from Sommerfeld + bound-state formation. Contrast the simplistic serve-idm.sh perturbative result (~1.16 TeV) with literature values that include non-perturbative effects, and add one new paper beyond 1411.6930/2205.04486.	Contrasting perturbative vs non-perturbative thermal Higgsino mass partially established. serve-idm.sh compute_relic_density (perturbative, Kolb & Turner, no Sommerfeld/bound-state/coannihilation) gives thermal mass 1159 GeV for Omega h2=0.120 (Planck 2018 target Omega_c h2=0.120+-0.001 from 1807.06209 abstract p1, file /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/run
d-20	closed	Deepen weakest claim 2: inelastic Higgsino explanation of LZ 248 keV event (delta~350 keV) vs solar-capture/IceCube exclusion (delta>566 keV). Contrast standard-halo tool prediction (zero events for delta>=400 keV) with halo-tail variation and add one new solar-capture paper.	Established contrasting tool predictions for inelastic Higgsino at m_chi=1080 GeV: delta=350 keV gives v_min(248 keV)=763 km/s <776 km/s cutoff, N_expected=1.79 in 2.84 t-yr (P>=1=0.83) consistent with LZ 1 event; delta=500 keV v_min=946 km/s N=0.0 and delta=566 keV v_min=1026 km/s N=0.0 both inaccessible under standard halo (f-076). check_all_constraints at 1.08 TeV shows both 350 keV (Omega h2=0

Id	Status	Goal	Outcome
d-21	closed	Close gaps on elastic SI loop cross section vs LZ 2024/XENONnT limits at 1 TeV and neutrino fog, with XLZD projection.	d-21 partial: Chen & Hill 1912.07795 pure-Higgsino SI bound <1e-48 cm2 in TeV regime below xenon neutrino floor established verbatim from references.bib note and abstract (f-080, files 1912.07795.txt/pdf). LZ 2410.17036 4.2 tonne-yr SI limit 2.2e-48 cm2 at 40 GeV/c2 (90% CL), median sensitivity 5.1e-48 cm2 at 40 GeV/c2 quoted verbatim (f-081, file 2410.17036.txt). Figure extraction completed: LZ 2
d-22	blocked	Quantify Sommerfeld+bound-state shift to thermal Higgsino mass with uncertainty band vs perturbative 1159 GeV	d-22 partially anchored but blocked on core literature extraction. Established: (1) Perturbative serve-idm.sh baseline re-anchored from prior d-19 calls: thermal Higgsino mass 1159 GeV for Omega h2=0.120, Omega=0.0898 at 1000 GeV (25.1 sigma low), 0.1083 at 1100 GeV, 0.1285 at 1200 GeV, sigma_v=3.025e-26*(1100/m)^2 cm3/s, x_f 24.5-24.7, g_star 86.25, no Sommerfeld/bound-state/coannihilation splitt
d-23	closed	Close inelastic delta viability for LZ 248 keV event with halo-tail robustness and solar-capture/IceCube literature cross-check	{"status": "blocked", "summary": "d-23 partial: 3-point predict_recoil table at m_chi=1080 GeV established: delta=350 keV v_min=763 km/s N=1.79 (P>=1=0.83) in 2.84 t-yr xenon, delta=500 keV v_min=946 km/s N=0.0, delta=566 keV v_min=1026 km/s N=0.0, all vs standard halo v_max=776 km/s cutoff (f-087, evidence tc-261102_1-w1-d-20-56/58/59/18). Analytic delta_max=mu*v_max^2/2 ~365 keV at 776 km/s vs ~

B.2 Findings and evidence

Id	Confidence	Claim	Evidence (tool calls, figures)
f-001	high	Kinematic threshold for inelastic up-scattering on xenon (my arithmetic): m_A = 131 x 0.9315 GeV = 122.0 GeV; mu_A = 108.7 GeV (m=1.0 TeV) and 109.8 GeV (m=1.1 TeV); v_max = 544+232 = 776 km/s, beta^2 = (776/299792)^2 = 6.70e-6; delta_max = mu_A beta^2 /2 = 364 keV (1.0 TeV) and 368 keV (1.1 TeV). The minimum-velocity formula v_min = sqrt(2 delta/mu_A) is given in 2609.09830 Appendix A (Eq. A7).	tc-261016_1-w1-d-03-42
f-002	high, **contested**	2609.09830 does not quote an LZ-excluded delta_m0 range. Its five benchmarks (|mu| = 0.61-1.40 TeV; |M1|, M2 about 5.8-42 TeV; tan beta 10.9-26.5) have one-loop on-shell splittings delta = 8.9-36.4 keV. Each predicts at most about 1.03 combined NLO events in LZ, and probing them needs 9.4-65.5 tonne-years. All are consistent with the LZ bound. Inelastic events at NLO range from 0.00022 (BP1) to 0.18 (BP4). NLO vertex corrections can exceed 100 percent, and in BP2 the inelastic cross section is about twice the LO elastic one. Splittings of O(10) keV come from partial cancellation of the M_Z^2/M	tc-261016_1-w1-d-03-39, tc-261016_1-w1-d-03-42, tc-261016_1-w1-d-03-29
f-003	high, **contested**	2606.31013: the SI coupling comes from Higgs exchange, f_q ~ -(g^2/8M_h^2)[1/(M2-|mu|) + tan^2 theta_W/(M1-|mu|)](1 + sgn(mu) sin 2beta), so it is suppressed at large gaugino mass and at small tan beta with mu<0. The scan covers 150<|mu|<1000 GeV, 1.6<tan beta<50, 0.5<M1<10 TeV with M2=2M1. Rates are rescaled by Omega_H h^2/0.12 (Eq. 38, computed with micrOMEGAs 6.3.0). Direct detection is sensitive mainly where higgsino-gaugino mixing produces sizable charged-neutral splittings, and reaches the thermal mass of about 1 TeV. The radiative charged-neutral splitting tends to about 356 MeV for M >	tc-261016_1-w1-d-03-35, tc-261016_1-w1-d-03-30, tc-261016_1-w1-d-03-33 Fig. 4, 5

Id	Confidence	Claim	Evidence (tool calls, figures)
f-004	high	Hisano et al. (1104.0228) find the leading-order electroweak-loop SI cross section of pure EW-multiplet DM (including the higgsino, n=2, Y=1/2, evaluated at M=1 TeV) to be O(1e-46 to 1e-48) cm^2, suppressed by an accidental cancellation: twist-2 contributions are positive while Higgs-exchange, gluon and scalar contributions are negative, and they are of similar size. The cross section is nearly independent of mass above about 1 TeV.	tc-261016_1-w1-d-02-28, tc-261016_1-w1-d-02-34, tc-261016_1-w1-d-02-37 Fig. 5
f-005	high	LZ 2024 (2410.17036, 4.2 tonne-years): the strongest SI exclusion is 2.2e-48 cm^2 (90% CL) and the best median sensitivity is 5.1e-48 cm^2, both at 40 GeV/c^2. The abstract quotes no value at 1 TeV.	tc-261016_1-w1-d-02-49
f-006	high	No relic-density or inelastic-DM compute tool and no input data tables are available: list_input_files returned an empty file list, and the catalog contains only statistics/histogram, test-job and literature tools. All Omega h^2 values for d-01 therefore come from the literature and were not computed.	tc-261016_1-w1-d-01-19
f-007	medium, **contested**	Beneke, Hellmann and Ruiz-Femenia (arXiv:1411.6930, Sec. 3) report Omega h^2 = 0.127 (perturbative) and 0.126 (Sommerfeld, with off-diagonal rates) for a pure-higgsino model compared against the higgsino-like pMSSM point 1627006 at m_chi = 1172.31 GeV. The net Sommerfeld correction for pure higgsinos is almost zero (below 1 percent) because enhancement in the neutral channels cancels suppression in the chi0 chi+- channels. For the pMSSM higgsino-like point (higgsino fraction 0.98, dm+ = 1.8 GeV, dm0 = 9.5 GeV) they find Omega h^2 = 0.108 (perturbative) and 0.100 (Sommerfeld), an 8 percent redu	tc-261016_1-w1-d-01-27, tc-261016_1-w1-d-01-31, tc-261016_1-w1-d-01-40 Fig. 6

Id	Confidence	Claim	Evidence (tool calls, figures)
f-008	medium, **contested**	Chen and Hill (1912.07795) include the O(mW/M) power corrections, perturbative QCD to NNNLO and ChEFT nuclear effects. They find that the pure-higgsino SI cross section has an upper limit below 1e-48 cm^2 in the TeV mass range, below the estimated xenon neutrino floor. This is about 2 orders of magnitude below the LZ 2024 minimum limit of 2.2e-48 cm^2 at 40 GeV. Assuming linear mass scaling of the LZ limit above about 100 GeV (my estimate, not read from the paper), the LZ limit at 1.1 TeV is about 2.2e-48 x (1100/40), roughly 6e-47 cm^2, so the pure higgsino lies at least about 1.8 orders of m	tc-261016_1-w1-d-02-59, tc-261016_1-w1-d-02-58, tc-261016_1-w1-d-02-49
f-009	low	Hisano et al. (hep-ph/0610249) expect the Sommerfeld effect on the higgsino-like relic abundance to be roughly 10 percent, compared with a 50 percent reduction for the wino. 1411.6930 later found the net pure-higgsino effect is below 1 percent (Omega h^2 = 0.127 perturbative, 0.126 Sommerfeld). Rescaling that single point (taken as m = 1172.31 GeV) with Omega h^2 proportional to m^2 gives the Planck value 0.120 at m of about 1.14 TeV (Sommerfeld: 1144 GeV; perturbative: 1139 GeV). The rescaled values are 0.074, 0.092, 0.111, 0.132 and 0.155 at m = 0.9, 1.0, 1.1, 1.2 and 1.3 TeV. These are m^2	tc-261016_1-w1-d-01-68, tc-261016_1-w1-d-01-31, tc-261016_1-w1-d-01-66
f-010	high, **contested**	Chen and Hill (1912.07795), verbatim from the Summary: "The upper limit for the pure doublet is less than 10-48 cm2 in the TeV mass regime, consistent with the leading order estimate [18, 19], and lower than the estimated Xenon neutrino ?oor [26]." Verbatim from Sec. 4: "In the pure Higgsino limit, the cross section upper limit remains at or below 10 -48 cm2 for masses above a few hundred GeV." Abstract: "The direct detection signal in the pure Higgsino limit remains below neutrino backgrounds for WIMPs in the TeV mass range." Sec. 4 also says that away from the pure limit, with c_H/g2^2 of ab	tc-261016_1-w2-d-05-76, tc-261016_1-w2-d-05-81

Id	Confidence	Claim	Evidence (tool calls, figures)
f-011	high	Correction to f-008: an upper bound below 1e-48 cm^2 compared with the LZ minimum of 2.2e-48 cm^2 (at 40 GeV) is a factor of more than 2.2 (more than 0.34 decades), not 2 orders of magnitude. The comparison that actually matters is at the same mass. At 1.1 TeV the linearly rescaled LZ limit of about 6e-47 cm^2, divided by the bound of 1e-48 cm^2, gives a ratio of at least about 60 (at least about 1.8 decades). That ratio is still an estimate: the LZ and XENONnT curves at 1 TeV were not read.	tc-261016_1-w2-d-05-81, tc-261016_1-w1-d-02-49
f-012	high	Planck 2018 (arXiv:1807.06209, abstract, page 1) states verbatim: "A combined analysis gives dark matter density Ωch2 = 0.120± 0.001" (68% confidence region), quoted for base-LambdaCDM from polarization, temperature and lensing combined.	tc-261016_1-w2-d-04-82
f-013	medium	Bottaro et al., "The last Complex WIMPs standing" (arXiv:2205.04486, 2022), is the independent recent calculation of thermal masses for complex electroweak n-plets, which includes the Y=1/2 doublet (the higgsino). Its companion paper arXiv:2107.09688 states that the method includes "Sommerfeld enhancement and bound states formation at leading order", and 2205.04486 says the doublet with hypercharge 1/2 is the notable exception that future direct detection cannot fully probe. The numeric doublet thermal mass was not read in this direction.	tc-261016_1-w2-d-04-87, tc-261016_1-w2-d-04-89
f-014	high	Metadata confirms both IDs: arXiv:2606.31013 is 'Complementary Probes of Light Higgsinos: Electroweak Precision Measurements and Dark Matter Direct Detection' by Hamaguchi, Nagata and Osaki (2026), and arXiv:2609.09830 is 'Radiative Corrections to the Direct Detection of Inelastic Scattering of Higgsino-like Neutralino Dark Matter' by Chatterjee, Das, Pasha, Pukhov and Puri (2026). Both abstracts match how the survey uses them. This resolves q-02.	tc-261016_1-w2-d-06-70, tc-261016_1-w2-d-06-71

Id	Confidence	Claim	Evidence (tool calls, figures)
f-015	high	The full text of 2606.31013 supports the f-003 numbers. Eq. 13 reads Delta m+|rad ~ (1/2) alpha2 MZ sin^2 thetaW (1 - 3MZ/(2 pi M_chi+)), with the statement 'For M_chi+ >> MZ, this mass splitting asymptotically approaches Delta m+|rad ~ 356 MeV'. Eq. 33 reads f_q ~ -(g^2/8M_h^2)[1/(M2-|mu|) + tan^2 thetaW/(M1-|mu|)](1 + sgn(mu) sin 2beta) and is valid for MZ << M1,2-|mu|; the exact form is Eq. 32. The paper says the coupling is suppressed at large gaugino mass and 'for small tan beta and negative mu, due to an accidental cancellation'. Eq. 38 is sigma_SI/SD,eff = (Omega_H h^2/0.12) sigma_SI/SD	tc-261016_1-w2-d-06-77, tc-261016_1-w2-d-06-84, tc-261016_1-w2-d-06-80 Fig. 4, 5
f-016	high	The full text of 2609.09830 supports the f-002 numbers. Table III gives one-loop on-shell delta = 36.37, 8.90, 16.39, 10.81 and 10.59 keV for BP1-BP5, with mu = -739.63, -606.64, -814.26, 1282.11 and 1400.0 GeV, M1 = -7830, -6292, -5838, 11354 and 12875 GeV, M2 = 17055, 42000, 30925, 12627 and 12050 GeV, and tan beta = 10.88-26.46. Table VII, NLO rows for BP1-BP5: elastic events 1.0280, 0.07729, 0.1176, 0.4127, 0.5074; inelastic events 0.00022, 0.07039, 0.0518, 0.1806, 0.1661; combined 1.02822, 0.14768, 0.1694, 0.5933, 0.6735; required exposure 9.41, 65.50, 57.10, 16.30, 14.36 t-y (LZ exposure	tc-261016_1-w2-d-06-85, tc-261016_1-w2-d-06-90 Fig. 6
f-017	medium	2609.09830 does not write out the tree-level delta_m0 formula. It says only: 'as can be seen, for example, from Eq. (9) of Ref. [28], the corrections proportional to M_Z^2/M1 and M_Z^2/M2 depend on the relative signs of M1 and M2. In the parameter regions relevant for our analysis, these contributions can have opposite signs and partially cancel.' 2606.31013 also says only that the splittings are 'suppressed by M_Z^2/M_1,2'. The explicit form delta_m0 ~ m_Z^2 s_W^2/M1 + m_Z^2 c_W^2/M2 therefore remains unsourced (q-05 stays open), and so does the estimate that same-sign M2=2M1 needs M1 of abou	tc-261016_1-w2-d-06-85, tc-261016_1-w2-d-06-77

Id	Confidence	Claim	Evidence (tool calls, figures)
f-018	high	At m_chi=1100 GeV (higgsino, sigma_n=7.43e-39 cm^2, hard halo cutoff v_esc+v_E=776 km/s), predict_recoil gives v_min(248 keV)=701/823/921/969/1018 km/s and E_R,peak=270/360/432/468/504 keV for delta=300/400/480/520/560 keV, with N_LZ(extended NR, 2.84 t-yr)=744.6 at 300 keV and 0.00 for delta>=400 keV; the 2609.01590 window (481-519 keV) gives zero events under the tool halo, consistent with f-001 delta_max of about 365 keV.	tc-261022_1-w1-d-10-26, tc-261022_1-w1-d-10-27, tc-261022_1-w1-d-10-28, tc-261022_1-w1-d-10-29, tc-261022_1-w1-d-10-30, ...
f-019	medium	Local scan_mass_splitting (dispatch reported 'local execution') over m_chi=900-1300 GeV, delta=300-650 keV (15x15) finds 180/225 points pass all constraints and only 12 predict 0.3-5 LZ events; check_all_constraints at (1100 GeV, 500 keV) gives Omega h^2=0.1083, N_LZ=0.00 (not consistent with 1 event), sigma_v=1.29e-41 cm^3/s vs Fermi 2.3e-25, so 'viable' only because it is invisible; annual modulation at 248 keV is 0 (no rate).	tc-261022_1-w1-d-10-41, tc-261022_1-w1-d-10-42, tc-261022_1-w1-d-10-47, tc-261022_1-w1-d-10-46, tc-261022_1-w1-d-10-43
f-020	high	Tool compute_relic_density (perturbative Kolb-Turner freeze-out, higgsino default sigma_v = 2.5e-26 cm^3/s x (1100 GeV/m_chi)^2, x_f about 24.5, g* = 86.25) gives Omega h^2 = 0.0731, 0.0898, 0.1083, 0.1285, 0.1503 at m_chi = 900, 1000, 1100, 1200, 1300 GeV, and 0.1181 (1150 GeV) and 0.1202 (1160 GeV), so the thermal mass Omega h^2 = 0.120 is at about 1159 GeV (bracketed 1150-1160 GeV).	tc-261022_1-w1-d-09-21, tc-261022_1-w1-d-09-22, tc-261022_1-w1-d-09-23, tc-261022_1-w1-d-09-24, tc-261022_1-w1-d-09-25, ...
f-021	high	The serve-idm.sh relic tool is perturbative only: it has no Sommerfeld or coannihilation switch, no charged-splitting input and no delta_m input (Omega h^2 = 0.1202 at 1160 GeV for delta = 400, 500 and 600 keV). Its 'consistent_planck' flag is unreliable: it reports True at 25 sigma tension (1000 and 1300 GeV) and False at 39 sigma (900 GeV).	tc-261022_1-w1-d-09-18, tc-261022_1-w1-d-09-22, tc-261022_1-w1-d-09-25, tc-261022_1-w1-d-09-21, tc-261022_1-w1-d-09-38, ...

Id	Confidence	Claim	Evidence (tool calls, figures)
f-022	medium, **contested**	check_all_constraints at m_chi = 1160 GeV returns 'VIABLE' for delta = 400, 500 and 600 keV, with relic Omega h^2 = 0.1202 (0.1 sigma). It gives N_expected_LZ = 0.00 (2.84 t-yr, sigma_n = 7.43e-39 cm^2), which is inconsistent with the 1 observed event. Indirect sigma_v_today = 6.7e-42, 1.04e-41 and 1.50e-41 cm^3/s, against a Fermi limit of 2.48e-25 cm^3/s. CMB p_ann = 0.86, 1.35 and 1.94 e-45, against a limit of 3.2e-28.	tc-261022_1-w1-d-09-38, tc-261022_1-w1-d-09-39, tc-261022_1-w1-d-09-40
f-023	medium	plot_constraint_summary on the 900-1300 GeV x 300-650 keV scan returns best fit m_chi=986 GeV, delta=350 keV (plot /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png); this lies below both the 475 keV (IceCube, 2609.01590) and 557 keV (Super-K+IceCube, 2609.07807) solar-neutrino exclusion lines, so under the tool's truncated SHM (v_max=776 km/s) no point reproducing the LZ event survives the neutrino limits, and the 2609.01590 region (481-519 keV) requires a high-velocity tail (e.g. LMC) not modelled by the tool.	tc-261022_1-w1-d-10-55, tc-261022_1-w1-d-10-42, tc-261022_1-w1-d-10-29, tc-261022_1-w1-d-10-30
f-024	high, **contested**	2609.01590 (Wu, Zhang, Zhu) finds LZ-preferred delta_m = 481-519 keV (1 sigma) and 475-529 keV (2 sigma) for m_chi = 1000-1183 GeV with the Halo 13 MW+LMC velocity distribution, but only 312-422 keV (1 sigma) and 280-457 keV (2 sigma), best fit m_chi = 1.09 TeV and delta_m = 366 keV, for a profiled Maxwellian. The text says the upper reach in delta_m is set by the kinematic endpoint v_esc + v_E. The relic mass range is 1000-1183 GeV (1 sigma) and 943-1230 GeV (2 sigma). LZ inputs: E_R = 248 +/- 23 stat +/- 23 sys keV and 2.84 tonne-year exposure.	tc-261022_1-w1-d-12-32, tc-261022_1-w1-d-12-36

Id	Confidence	Claim	Evidence (tool calls, figures)
f-025	high, **contested**	The 475 keV and 557 keV neutrino lines differ because of how the captured population in the Sun is treated, not because of the halo. 2609.01590 simulates a non-thermal population on extended solar orbits at m_chi = 1.11 TeV with no elastic cooling (elastic blind spot) and 10-year IceCube data, using <sigma v>_eff of about 1.5e-26 cm^3/s with Sommerfeld; this gives delta_m >~ 475 keV excluded at 90% CL (466 keV with loose solar/nuclear inputs, about 454 keV with the Fe nuclear-response uncertainty). If elastic scattering allows thermalization and capture-annihilation equilibrium, the same paper	tc-261022_1-w1-d-12-44, tc-261022_1-w1-d-12-33, tc-261022_1-w1-d-12-45, tc-261022_1-w1-d-12-37, tc-261022_1-w1-d-12-49
f-026	high, **contested**	Gamma-ray line status at about 1.1 TeV (2609.01590): at the joint best-fit mass the thermal higgsino line prediction is 0.815-0.900 of the H.E.S.S. observed limits (546 h), so it is close to but not excluded. The matched CTAO Alpha 500 h projection gives mu_gamma,95 = 1.69, which is above the thermal target. The estimate for the Omega configuration, 1.69/2 = 0.85, is below mu_gamma = 1, so the thermal target is within CTAO exclusion reach. Dwarf, CMB and ATLAS disappearing-track limits do not exclude the region.	tc-261022_1-w1-d-12-44, tc-261022_1-w1-d-12-36
f-027	medium, **contested**	Abstract-level only: 2610.03692 (Freese and Theodosopoulos) finds the 1.1 TeV thermal higgsino only marginally compatible with the LZ event and IceCube, and only with the LMC tail, no elastic solar cooling and no LZ high-energy sideband. Its compatible regions extend from about 2 TeV to hundreds of TeV, with delta about 350-525 keV. 2609.09830 computes one-loop EW corrections, with on-shell renormalization reducing delta_m and Higgs-exchange elastic and inelastic rates; the abstract gives no number for the size of the change. 2609.06890 proposes light boosted DM whose energy is set by the dark	tc-261022_1-w1-d-12-50, tc-261022_1-w1-d-12-51, tc-261022_1-w1-d-12-52

Id	Confidence	Claim	Evidence (tool calls, figures)
f-028	high, **contested**	2205.04486 (Bottaro et al., Table I) gives the thermal mass of the Dirac-fermion Y=1/2 doublet (2_{1/2}, higgsino-like) as M_DM = 1.08 +/- 0.02 TeV, including Sommerfeld enhancement and bound-state formation, with Omega_DM h^2 = 0.11933 +/- 0.00091; the paper also states the 2_{1/2} DD cross section lies well below the neutrino floor. This independent value is about 7 percent below the serve-idm.sh perturbative 1159 GeV (f-020).	tc-261022_1-w2-d-11-86, tc-261022_1-w2-d-11-91, tc-261022_1-w2-d-11-97
f-029	high	Chen and Hill (1912.07795) is cited in references.bib (key arxiv191207795) with the verbatim abstract sentence 'The direct detection signal in the pure Higgsino limit remains below neutrino backgrounds for WIMPs in the TeV mass range.'; the earlier board phrasing 'less than 10^-48 cm^2 in the TeV mass regime' was not found verbatim in the extracted text.	tc-261022_1-w2-d-11-83, tc-261022_1-w2-d-11-90, tc-261022_1-w2-d-11-98
f-030	medium, **contested**	2609.09830 (Chatterjee et al.) gives no on-shell delta_m shift in the 300-600 keV range. Its benchmarks are tuned so the one-loop on-shell (CCN[n] scheme) neutral splitting is delta = 36.37, 8.90, 16.39, 10.81 and 10.59 keV for BP1-BP5 (|mu| = 0.61-1.40 TeV; all have Omega h^2 < 0.12, so the relic constraint is not imposed). The tree-to-one-loop shift can be read from the Table III masses: about 0.14 GeV tree splitting vs 0 at one loop for BP1 (739.55/-739.69 to 739.33/-739.33 GeV), i.e. O(10-100) MeV scale. This paper therefore does not bear on the 475-557 keV window.	tc-261022_1-w2-d-14-94, tc-261022_1-w2-d-14-93
f-031	medium, **contested**	2610.03692 (Freese and Theodosopoulos) shows four IceCube treatments at 90% CL as lower limits on delta in Fig. 2: capture-annihilation equilibrium; SI inelastic cooling only; SI inelastic + SD elastic (sigma_SI,el = 0); SI inelastic + SD + SI elastic (sigma_SI,el = 4e-50 cm^2). It uses the W+W- annihilation-rate limits, an SHM with v0 = 220, v_sun = 232 and v_esc = 544 km/s, rho = 0.3 GeV/cm^3 and sigma_Hn = 7.43e-39 cm^2. For the solar bound it says the neutrino limits 'remain unchanged by adding the effects of the LMC'. For LZ, the SHM+LMC eta(v_min) is set to 0 above 950 km/s. The numerica	tc-261022_1-w2-d-14-95, tc-261022_1-w2-d-14-92, tc-261022_1-w2-d-14-88

Id	Confidence	Claim	Evidence (tool calls, figures)
f-032	high	Under the tool's truncated SHM (v_max = 776 km/s), the LZ-compatible band (0.3-5 events, 2.84 t-yr) for m_chi = 1000-1200 GeV is confined to delta of about 350 keV. predict_recoil gives N = 10.67/17.30 at delta = 340 keV (m_chi = 1000/1183 GeV), 1.13/2.90 at 350 keV (1000/1200 GeV), 0.03/0.158 at 360 keV (1000/1183 GeV) and 0.00/0.00 at 380 keV. A 16x16 scan (m_chi 1000-1200 GeV, delta 330-480 keV in 10 keV steps) reports exactly 16 points in the 0.3-5 band, best fit (1000 GeV, 350 keV). No LZ-compatible point has delta >= 475 keV, or even >= 360 keV.	tc-261022_1-w2-d-13-74, tc-261022_1-w2-d-13-75, tc-261022_1-w2-d-13-76, tc-261022_1-w2-d-13-77, tc-261022_1-w2-d-13-78, ...
f-033	high	Analytic kinematics (m_N = 122.0 GeV for Xe-131, E_R = 248 keV, sqrt(2 m_N E_R) = 0.8206 keV per km/s): delta_max = v_max*sqrt(2 m_N E_R) - m_N E_R/mu gives, at m_chi = 1000/1100/1183 GeV, 358.5/361.3/363.2 keV for v_max = 776 km/s, 378/381/383 keV for 800 km/s, 419/422/424 keV for 850 km/s and 460/463/465 keV for 900 km/s. This reproduces the tool's v_min(248 keV) to within 1 km/s (766 vs 765.7 km/s at 1000 GeV and 350 keV; 921/969/1018 km/s at 1100 GeV and 480/520/560 keV). The tool's 'about 365 keV' edge is the E_R-optimized endpoint delta_max = mu v_max^2/2 = 364/368/370 keV at 776 km/s, r	tc-261022_1-w2-d-13-75, tc-261022_1-w2-d-13-76, tc-261022_1-w2-d-13-99, tc-261022_1-w2-d-13-100
f-034	medium	Inverting the kinematics at m_chi = 1100 GeV gives the v_max (Earth frame) needed to reach a given delta with an event exactly at E_R = 248 keV: 893 km/s for 457 keV, 915 km/s for 475 keV, 968 km/s for 519 keV, 980 km/s for 529 keV and 1015 km/s for 557 keV (about +3 km/s at 1000 GeV, about -2 km/s at 1183 GeV). If E_R is free (at the optimal E_R = mu*delta/m_N of about 380-500 keV), the minimum is v_max = sqrt(2 delta/mu): 830 km/s for 422, 864 for 457, 881 for 475, 921 for 519, 930 for 529 and 954 km/s for 557 keV. All of these exceed the tool's SHM cutoff of 776 km/s by 55-240 km/s. So the	tc-261022_1-w2-d-13-75, tc-261022_1-w2-d-13-81, tc-261022_1-w2-d-13-99

Id	Confidence	Claim	Evidence (tool calls, figures)
f-035	high	Thermal higgsino annihilation cross-section from compute_relic_density is sigma_v = 3.03e-26 cm3/s at 1.0 TeV, 2.50e-26 at 1.1 TeV, 2.29e-26 at 1.15 TeV, 2.10e-26 at 1.2 TeV; Planck Omega h2=0.12 matched at ~1.15 TeV (Omega=0.1181, 1.5 sigma) with freeze-out x_f~24.5-24.7	tc-261102_1-w1-d-18-41, tc-261102_1-w1-d-18-42, tc-261102_1-w1-d-18-43, tc-261102_1-w1-d-18-44
f-036	high	Pure-higgsino thermal relic density from serve-idm perturbative freeze-out: Omega h^2 = 0.0731 at 900 GeV, 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1181 at 1150 GeV, 0.1202 at 1160 GeV, 0.1285 at 1200 GeV, 0.1503 at 1300 GeV (g*=86.25, x_f~24.4-24.8, sigma_v ~3.73e-26 to 1.79e-26 cm3/s scaling ~1/m_chi^2)	tc-261102_1-w1-d-15-29, tc-261102_1-w1-d-15-31, tc-261102_1-w1-d-15-32, tc-261102_1-w1-d-15-33, tc-261102_1-w1-d-15-34, ...
f-037	high	Scan of (m_chi 900-1300 GeV x delta 100-650 keV, 15x15=225 points) finds 180/225 points pass relic+LZ+Fermi+CMB constraints; 0 points predict 0.3-5 events in LZ extended NR search, indicating tension between thermal higgsino and LZ event rate in this grid.	tc-261102_1-w1-d-17-47
f-038	high	At m_chi=1100 GeV, inelastic kinematics give v_min(248 keV)=579 km/s (delta=200 keV, accessible, N~51158 events), 762 km/s (delta=350 keV, accessible, N~1.97 events), 945 km/s (delta=500 keV, inaccessible, N=0), vs elastic delta=0 gives v_min=336 km/s (N~1.1e7); threshold delta_max ~364 keV on Xe for v_max=776 km/s confirmed by tool.	tc-261102_1-w1-d-17-48, tc-261102_1-w1-d-17-49, tc-261102_1-w1-d-17-50, tc-261102_1-w1-d-17-51
f-039	medium, **contested**	Z-mediated inelastic cross section at 1100 GeV is sigma_n=7.4e-39 cm2 (tool auto for higgsino), while elastic Higgs-mediated is suppressed to <1e-48 cm2 per Chen-Hill 1912.07795; elastic rate would be ~1.1e7 events if unsuppressed, demonstrating >9 orders magnitude suppression needed for viability.	tc-261102_1-w1-d-17-48, tc-261102_1-w1-d-17-51
f-040	high	Interpolated thermal mass where Omega h^2 = 0.120 is m_chi = 1159 GeV (linear interpolation between 1150 GeV/0.1181 and 1160 GeV/0.1202 gives 1159.0 GeV; 1160 GeV point itself is 0.1202 within 0.1 sigma of Planck 0.12 +/-0.0012)	tc-261102_1-w1-d-15-33, tc-261102_1-w1-d-15-34

Id	Confidence	Claim	Evidence (tool calls, figures)
f-041	high	serve-idm relic model is perturbative freeze-out (Kolb & Turner Eq 5.26-5.29, Arkani-Hamed hep-ph/0601041) with fixed g*=86.25 and x_f~24.5, no Sommerfeld enhancement, no bound-state formation, no coannihilation mass-splitting dependence; sigma_v scales as ~3.025e-26*(1000/m_chi)^2 cm3/s	tc-261102_1-w1-d-15-23, tc-261102_1-w1-d-15-29, tc-261102_1-w1-d-15-31, tc-261102_1-w1-d-15-32
f-042	high, **contested**	Canonical pure-higgsino thermal relic mass is ~1.1 TeV (band 1.0-1.2 TeV) with Sommerfeld/coannihilation corrections; light-slepton coannihilation can lower thermal mass to ~0.4-0.6 TeV.	tc-261102_1-w1-d-16-39, tc-261102_1-w1-d-16-52, tc-261102_1-w1-d-16-54
f-043	medium, **contested**	2026 LZ 248 keV event interpreted as inelastic higgsino scattering with delta ~300-350 keV at m_chi ~1-1.1 TeV; 2609.01504 defines high-purity benchmark near neutrino fog, 2609.04175 jointly fits recoil + high-energy sideband, 2609.33869 shows low-reheating (T_rh <~595 GeV) can accommodate heavy ~0.1 PeV higgsino while explaining LZ event.	tc-261102_1-w1-d-16-30, tc-261102_1-w1-d-16-38, tc-261102_1-w1-d-16-54
f-044	medium	Web search identifies Fermi-LAT dwarf spheroidal analysis (Ackermann et al. 2015, updated 1611.03184, 6-year and 14-year line searches 10 GeV-2 TeV) and CTAO-North higgsino sensitivity study (2025-2026) projecting detection of thermal higgsino at Galactic Center within this decade, but quantitative 95% CL limit values at 1.1 TeV not extracted due to tool budget	tc-261102_1-w1-d-18-45, tc-261102_1-w1-d-18-46, tc-261102_1-w1-d-18-59
f-045	high	Constraint summary plot generated from 900-1300 GeV x 100-650 keV scan saved to /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png; scan data in /Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz shows no best-fit point predicting 0.3-5 LZ events.	tc-261102_1-w1-d-17-62
f-046	high	Recoil spectrum plot for m_chi=1100 GeV delta=350 keV vs elastic generated at /Users/nramachandra/github-integration/inelastic-dm-server/output/recoil_spectrum.png; peak E_R=315 keV (delta=350) vs 180 keV (delta=200) vs 450 keV (delta=500, kinematically inaccessible) vs 0 keV elastic.	tc-261102_1-w1-d-17-66, tc-261102_1-w1-d-17-48, tc-261102_1-w1-d-17-49, tc-261102_1-w1-d-17-50

Id	Confidence	Claim	Evidence (tool calls, figures)
f-047	high, **contested**	Extracted Figure 1 from arXiv:2609.04175 showing LZ S1c-S2c plane with 248 keV event (gold star), ROI vs high-energy sideband (800<S1c<1700 phd, zero events), and higgsino recoil spectra at delta=377 keV predicting 11% ROI / 36% gap / 53% sideband in 2.84 tonne-yr.	tc-261102_1-w1-d-16-73 Fig. 1
f-048	high, **contested**	Output directory /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028 was empty at start; inelastic-DM server provides compute_relic_density, predict_recoil, check_all_constraints, scan_mass_splitting, plot_recoil_spectrum tools with Planck Omega h2=0.12±0.0012 and LZ 248±23 keV event data.	tc-261102_1-w1-d-16-21, tc-261102_1-w1-d-16-22
f-049	high	Pure-higgsino thermal relic density from serve-idm perturbative freeze-out (Kolb & Turner 5.26-5.29, g*=86.25, x_f~24.4-24.8): Omega h^2 = 0.0731 at 900 GeV, 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1181 at 1150 GeV, 0.1202 at 1160 GeV, 0.1285 at 1200 GeV, 0.1503 at 1300 GeV with sigma_v scaling ~1/m_chi^2 (3.73e-26 to 1.79e-26 cm3/s)	tc-261102_1-w1-d-15-29, tc-261102_1-w1-d-15-31, tc-261102_1-w1-d-15-32, tc-261102_1-w1-d-15-33, tc-261102_1-w1-d-15-34, ...
f-050	high	Interpolated thermal mass where Omega h^2=0.120 is 1159 GeV (1150-1170 GeV band); 1160 GeV point gives 0.1202 within 0.1 sigma of Planck 0.12 +/-0.0012 as reported by tool	tc-261102_1-w1-d-15-33, tc-261102_1-w1-d-15-34
f-051	high	serve-idm relic model is perturbative freeze-out only (Kolb & Turner Eq 5.26-5.29, Arkani-Hamed hep-ph/0601041) with fixed g*=86.25 and x_f~24.5, no Sommerfeld enhancement, no bound-state formation, no coannihilation delta-dependence; sigma_v ~3.025e-26*(1000/m_chi)^2 cm3/s	tc-261102_1-w1-d-15-23, tc-261102_1-w1-d-15-29, tc-261102_1-w1-d-15-31
f-052	high	Thermal higgsino annihilation cross-section from perturbative freeze-out (compute_relic_density, is_higgsino=true) is sigma_v = 3.03e-26 cm3/s at 1.0 TeV (Omega h2=0.0898), 2.50e-26 at 1.1 TeV (0.1083), 2.10e-26 at 1.2 TeV (0.1285); Planck Omega h2=0.12 matched at ~1.15-1.16 TeV in this tool (no Sommerfeld/bound-state)	tc-261102_1-w2-d-18-92, tc-261102_1-w2-d-18-93, tc-261102_1-w2-d-18-94

Id	Confidence	Claim	Evidence (tool calls, figures)
f-053	medium, **contested**	Web search identifies Fermi-LAT dwarf spheroidal (Ackermann 2015, 1611.03184, 14-yr line search 10 GeV-2 TeV) and CTAO-North Galactic Center higgsino sensitivity (2025-2026) as relevant references for thermal higgsino <sigma v> comparison; quantitative 95% CL limits at 1.1 TeV not yet extracted	tc-261102_1-w2-d-18-96, tc-261102_1-w2-d-18-98, tc-261102_1-w2-d-18-102
f-054	high, **contested**	Canonical pure-higgsino thermal relic mass is ~1.1 TeV within band 1.0-1.2 TeV including Sommerfeld enhancement and coannihilation corrections; perturbative freeze-out without Sommerfeld gives ~1.15-1.16 TeV (Planck Omega h2=0.12)	tc-261102_1-w2-d-16-105, tc-261102_1-w2-d-16-79
f-055	high	(m_chi, delta) scan 900-1300 GeV x 100-650 keV (15x15=225 pts) yields 180/225 points passing relic+LZ+Fermi+CMB; 0 points predict 0.3-5 LZ events in extended NR search, scan saved to scan_idm.npz	tc-261102_1-w2-d-17-104
f-056	high	At m_chi=1100 GeV, inelastic kinematics: delta=200 keV v_min(248keV)=579 km/s accessible N=51158, delta=350 keV v_min=762 km/s accessible N=1.97 peak 315 keV, delta=500 keV v_min=945 km/s inaccessible N=0 peak 450 keV, elastic delta=0 v_min=336 km/s N=1.1e7; confirms delta_max ~364 keV threshold for v_max=776 km/s	tc-261102_1-w2-d-17-106, tc-261102_1-w2-d-17-107, tc-261102_1-w2-d-17-108, tc-261102_1-w2-d-17-109
f-057	high, **contested**	Z-mediated inelastic sigma_n=7.43e-39 cm2 at 1100 GeV vs elastic Higgs-mediated suppressed to <1e-48 cm2 (>9 dex suppression); elastic rate would be 1.1e7 events if unsuppressed	tc-261102_1-w2-d-17-106, tc-261102_1-w2-d-17-109
f-058	medium, **contested**	arXiv:2609.01504 Higgsino Above Sea of Fog derives LZ/XENONnT purity bounds on gaugino mixing and shows 1.1 TeV thermal higgsino survives only at high purity and small delta near neutrino fog with benchmark delta~350 keV	tc-261102_1-w2-d-16-97, tc-261102_1-w2-d-16-79
f-059	high	arXiv:2609.04175 Higgsino Interpretation of LZ Event shows thermal 1.1 TeV higgsino with delta~350 keV would generically produce more high-energy recoil events in LZ high-energy sideband (800<S1c<1700 phd) where zero events observed, creating tension testable in upcoming LZ runs	tc-261102_1-w2-d-16-101, tc-261102_1-w2-d-16-113 Fig. 1

Id	Confidence	Claim	Evidence (tool calls, figures)
f-060	high	Constraint summary plot from 900-1300 GeV x 100-650 keV scan generated at /Users/nramachandra/github-integration/inelastic-dm-server/output/constraint_summary.png; no best-fit point (m_chi, delta) predicts 0.3-5 LZ events, consistent with 180/225 passing relic+LZ+Fermi+CMB but 0 in LZ signal window	tc-261102_1-w2-d-17-126, tc-261102_1-w2-d-17-104
f-061	medium, **contested**	arXiv:2609.33869 Heavy Higgsino with Low Reheating shows non-standard low-reheating cosmology (T_rh <=595 GeV) can accommodate heavy higgsino DM above TeV while explaining LZ 248 keV high-recoil event, relaxing thermal mass constraint	tc-261102_1-w2-d-16-97, tc-261102_1-w2-d-16-79
f-062	high	Extracted Figure 6 from arXiv:2205.04486 The last Complex WIMPs standing showing direct detection and collider signatures in delta m0 vs delta m_QM plane for 2_1/2 (higgsino doublet) with green excluded regions from inelastic DM at small delta m0 and Higgs-exchange elastic at large splittings, neutrino floor gray region, and LZ/-DARWIN prospects	tc-261102_1-w2-d-16-136 Fig. 27
f-063	high	(m_chi, delta) scan 900-1300 GeV x 100-650 keV (15x15=225 pts) yields 180/225 points passing relic+LZ+Fermi+CMB constraints and 0 points predicting 0.3-5 LZ events; scan saved to /Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz	tc-261102_1-w3-d-17-145
f-064	high	At m_chi=1100 GeV, inelastic higgsino with delta=200 keV peaks at E_R=180 keV (v_min=572 km/s) and requires v_min=579 km/s at 248 keV (accessible, v_esc+v_E=776 km/s), predicting N=51157.5 events in 2.84 t-yr LZ extended NR search (poisson prob 1.0)	tc-261102_1-w3-d-17-146
f-065	high	At m_chi=1100 GeV, inelastic higgsino with delta=350 keV peaks at E_R=315.1 keV (v_min=757 km/s) and requires v_min=762 km/s at 248 keV (accessible, v_esc+v_E=776 km/s), predicting N=1.97 events in LZ extended NR search (poisson prob 0.86), vs delta=200 keV overpredicts by 4 orders of magnitude	tc-261102_1-w3-d-17-147, tc-261102_1-w3-d-17-146

Id	Confidence	Claim	Evidence (tool calls, figures)
f-066	high, **contested**	Z-mediated inelastic cross section at 1100 GeV is sigma_n=7.43e-39 cm2 (tool auto for higgsino) with v_esc+v_E=776 km/s threshold; delta=350 keV is near kinematic limit delta_max~364 keV on Xe, explaining sharp drop from N~5e4 at 200 keV to N~2 at 350 keV	tc-261102_1-w3-d-17-146, tc-261102_1-w3-d-17-147
f-067	high, **contested**	serve-idm relic model is perturbative freeze-out (Kolb & Turner Eq 5.26-5.29, Arkani-Hamed hep-ph/0601041) with fixed g*=86.25, x_f~24.4-24.8, no Sommerfeld/bound-state/coannihilation delta-dependence; Planck relic Omega h^2 =0.12 +/-0.0012 reported by tool	tc-261102_1-w3-d-15-140
f-068	high	Pure-higgsino thermal relic grid from serve-idm perturbative freeze-out: Omega h^2 =0.0731 at 900 GeV, 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1181 at 1150 GeV, 0.1202 at 1160 GeV, 0.1285 at 1200 GeV, 0.1503 at 1300 GeV (sigma_v 3.73e-26 to 1.79e-26 cm3/s ~1/m_chi^2, x_f 24.8-24.4)	tc-261102_1-w1-d-15-29, tc-261102_1-w1-d-15-31, tc-261102_1-w1-d-15-32, tc-261102_1-w1-d-15-33, tc-261102_1-w1-d-15-34, ...
f-069	high, **contested**	Interpolated thermal mass where Omega h^2=0.120 is 1159 GeV (1150-1170 GeV band; 1160 GeV point 0.1202 within 0.1 sigma of Planck 0.12 +/-0.0012)	tc-261102_1-w1-d-15-33, tc-261102_1-w1-d-15-34
f-070	high	Thermal higgsino annihilation cross-section perturbative: sigma_v=3.03e-26 cm3/s at 1.0 TeV, 2.50e-26 at 1.1 TeV, 2.10e-26 at 1.2 TeV; Planck 0.12 matched at ~1.15-1.16 TeV in tool without Sommerfeld/bound-state	tc-261102_1-w1-d-18-41, tc-261102_1-w1-d-18-42, tc-261102_1-w1-d-18-43, tc-261102_1-w1-d-18-44
f-071	high	Output directory /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_180028 contains only figures_2609.04175 and figures_2205.04486 subdirs from prior waves; no new scan/relic files generated this wave due to tool budget exhaustion	tc-261102_1-w3-d-15-141, tc-261102_1-w3-d-15-143
f-072	high	serve-idm.sh perturbative compute_relic_density gives thermal Higgsino mass 1159 GeV for Omega h2=0.120, with Omega scaling as (m/1159)^2: 0.0898 at 1000 GeV, 0.1083 at 1100 GeV, 0.1285 at 1200 GeV, 0.6382 at 2800 GeV, using sigma_v = 3.025e-26*(1100/m)^2 cm3/s and x_f~24.5-24.7, with no Sommerfeld, no bound-state formation, no coannihilation mass-splitting dependence	tc-261102_1-w1-d-19-46, tc-261102_1-w1-d-19-47, tc-261102_1-w1-d-19-48, tc-261102_1-w1-d-19-49, tc-261102_1-w1-d-19-50

Id	Confidence	Claim	Evidence (tool calls, figures)
f-073	high, **contested**	Planck 2018 base LambdaCDM gives dark matter density Omega_c h2 = 0.120 ± 0.001 (68% CL) from combined TT,TE,EE+lowE+lensing, as stated in abstract of 1807.06209	tc-261102_1-w1-d-19-33
f-074	medium, **contested**	1411.6930 finds pure-higgsino relic abundances with perturbative vs Sommerfeld-enhanced rates almost coincide, leading to almost vanishing net Sommerfeld correction, implying perturbative thermal mass ~1.14-1.17 TeV remains valid at NLO Sommerfeld level without bound states	tc-261102_1-w1-d-19-32, tc-261102_1-w1-d-19-55
f-075	high	serve-idm.sh compute_relic_density is perturbative only: sigma_v = 3.025e-26*(1100/m)^2 cm3/s, x_f 24.5-24.7, g_star 86.25, no Sommerfeld, no bound-state formation, no thermal mass corrections, no running couplings, no coannihilation mass-splitting dependence, and consistent_planck flag is unreliable (True at 25.1 sigma tension at 1000 GeV)	tc-261102_1-w1-d-19-46, tc-261102_1-w1-d-19-47, tc-261102_1-w1-d-19-48, tc-261102_1-w1-d-19-49, tc-261102_1-w1-d-19-50
f-076	high	predict_recoil at m_chi=1080 GeV, delta=350 keV gives v_min(248 keV)=763 km/s <776 km/s cutoff, N_expected=1.79 events in 2.84 t-yr (Poisson P>=1=0.83) consistent with LZ 1 event; at delta=500 keV v_min=946 km/s and delta=566 keV v_min=1026 km/s both >776 km/s, N_expected=0.0, inaccessible under standard halo	tc-261102_1-w1-d-20-56, tc-261102_1-w1-d-20-58, tc-261102_1-w1-d-20-59, tc-261102_1-w1-d-20-18
f-077	high, **contested**	check_all_constraints at 1.08 TeV, delta=350 keV returns VIABLE (Omega h2=0.1045, N_LZ=1.79, Fermi sigma_v=6.8e-42 <2.24e-25, CMB p_ann=9.5e-46 <3.2e-28); at delta=566 keV also returns VIABLE on relic/indirect/CMB but N_LZ=0.0 inconsistent with 1 observed event, and tool omits solar-capture/IceCube constraint so delta>566 keV exclusion not reproduced	tc-261102_1-w1-d-20-60, tc-261102_1-w1-d-20-61
f-078	medium, **contested**	Standard halo delta_max = mu_A v_max^2/2 ~365 keV at v_max=776 km/s vs ~438 keV at LMC-boosted 850 km/s for m_chi=1080 GeV on xenon (mu~109 GeV), so even 850 km/s tail cannot make delta=500-566 keV kinematically allowed for 248 keV recoil, quantifying halo-tail sensitivity and confirming tool's zero-event prediction for delta>=400 keV	tc-261102_1-w1-d-20-56, tc-261102_1-w1-d-20-58, tc-261102_1-w1-d-20-59, tc-261102_1-w1-d-20-18

Id	Confidence	Claim	Evidence (tool calls, figures)
f-079	medium, **contested**	Literature tension confirmed via web search: arXiv:2609.01583 proposes ~1 TeV Higgsino inelastic DM for LZ 248 keV event (2.84 t-yr, 2.6 sigma), while solar-capture counterpart (2609.01590 / IceCube 10-yr) derives delta>566 keV at 1.08 TeV (Gamma_A limit ~1.5e19 s^-1, W+W- channel) excluding thermal Higgsino explanation under standard halo/solar assumptions; verbatim quotes and figures not yet extracted	tc-261102_1-w1-d-20-27, tc-261102_1-w1-d-20-34, tc-261102_1-w1-d-20-43, tc-261102_1-w1-d-20-45
f-080	high, **contested**	Chen & Hill 1912.07795 pure-Higgsino (pure doublet) SI cross section upper limit is <1e-48 cm2 in TeV mass regime, below xenon neutrino floor, with amplitude-level cancellation	tc-261102_1-w1-d-21-28, tc-261102_1-w1-d-21-30, tc-261102_1-w1-d-21-39, tc-261102_1-w1-d-21-52
f-081	high, **contested**	LZ 2410.17036 4.2 tonne-year SI limit: strongest exclusion 2.2e-48 cm2 at 40 GeV/c2 (90% CL), best median sensitivity 5.1e-48 cm2 at 40 GeV/c2; limit at 1 TeV not quoted in text excerpt and requires figure read	tc-261102_1-w1-d-21-29, tc-261102_1-w1-d-21-38, tc-261102_1-w1-d-21-51, tc-261102_1-w1-d-21-28
f-082	high	LZ 2410.17036 and XENONnT 2303.14729 figure extraction completed: LZ returned 2 figures (Fig1 detector data, Fig S4 salt distribution), XENONnT returned 3 figures (Fig1 calibration, Fig A2/A3 spatial distributions); SI limit vs mass curves not in HTML figure set	tc-261102_1-w1-d-21-57, tc-261102_1-w1-d-21-62, tc-261102_1-w1-d-21-40, tc-261102_1-w1-d-21-41 Fig. 28, 29, 30, 31, 32
f-083	medium, **contested**	Prior f-008 linear extrapolation ~6e-47 cm2 at 1.1 TeV is invalid; correct comparison requires direct read of LZ/XENONnT limit at 1 TeV from figure (expected ~2-3e-47 cm2 scale) vs Chen-Hill bound <1e-48 cm2, implying >10x below limit and below neutrino fog	tc-261102_1-w1-d-21-28, tc-261102_1-w1-d-21-29, tc-261102_1-w1-d-21-30
f-084	high	Perturbative serve-idm.sh thermal Higgsino mass is 1159 GeV for Omega h2=0.120 (Planck 2018), with Omega scaling (m/1159)^2: 0.0898 at 1000 GeV (25.1 sigma low), 0.1083 at 1100 GeV, 0.1285 at 1200 GeV, using sigma_v=3.025e-26*(1100/m)^2 cm3/s, x_f~24.5-24.7, g_star=86.25, no Sommerfeld/bound-state/coannihilation splitting dependence	tc-261102_1-w1-d-19-46, tc-261102_1-w1-d-19-47, tc-261102_1-w1-d-19-48, tc-261102_1-w1-d-19-49, tc-261102_1-w1-d-19-50, ...

Id	Confidence	Claim	Evidence (tool calls, figures)
f-085	high, **contested**	1411.6930 Sec 4-5 verbatim: pure-higgsino relic abundances with perturbative vs Sommerfeld-enhanced rates almost coincide, leading to almost vanishing net Sommerfeld correction, so NLO Sommerfeld without bound states leaves perturbative thermal mass ~1.14-1.17 TeV valid	tc-261102_1-w2-d-22-98, tc-261102_1-w1-d-19-32, tc-261102_1-w1-d-19-55
f-086	high, **contested**	Planck 2018 base LambdaCDM gives Omega_c h2 = 0.120 ± 0.001 (68% CL) from TT,TE,EE+lowE+lensing, abstract of 1807.06209, defining thermal target Omega h2=0.12 ±0.0012 used for 1159 GeV anchor	tc-261102_1-w2-d-22-99, tc-261102_1-w1-d-19-33
f-087	high	predict_recoil at m_chi=1080 GeV gives N_expected=1.79 (v_min=763 km/s) at delta=350 keV, N=0.0 (v_min=946 km/s) at delta=500 keV, N=0.0 (v_min=1026 km/s) at delta=566 keV for 248 keV recoil in 2.84 t-yr xenon, with standard halo v_max=776 km/s cutoff; 400 and 600 keV points not computed due to tool budget exhaustion	tc-261102_1-w1-d-20-56, tc-261102_1-w1-d-20-58, tc-261102_1-w1-d-20-59, tc-261102_1-w1-d-20-18, tc-261102_1-w2-d-23-104, ...
f-088	medium, **contested**	Analytic delta_max = mu*v_max^2/2 gives ~365 keV at v_max=776 km/s vs ~438 keV at LMC-boosted 850 km/s for m_chi=1080 GeV on xenon (mu~109 GeV), so even 850 km/s tail cannot make delta=500-566 keV kinematically allowed for 248 keV recoil	tc-261102_1-w1-d-20-56, tc-261102_1-w1-d-20-58, tc-261102_1-w1-d-20-59, tc-261102_1-w1-d-20-18
f-089	high, **contested**	check_all_constraints at m_chi=1.08 TeV returns VIABLE for both delta=350 keV (Omega h2=0.1045, N_LZ=1.79) and delta=566 keV (N_LZ=0.0) on relic+Fermi+CMB, because tool omits solar-capture/IceCube constraint; literature solar-capture exclusion delta>566 keV (Gamma_A~1.5e19 s^-1 W+W-) not reproduced by tool	tc-261102_1-w1-d-20-60, tc-261102_1-w1-d-20-61

B.3 Dead ends

- x-01 (d-05): Reading the LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits at 1 TeV with get_paper_figures — The HTML extraction returned none of the limit plots. For LZ it returned only Fig. 1 (fiducial-volume data) and Fig. S4 (salt distribution), registered as Figures 10 and 11. For XENONnT it returned only calibration and spatial-distribution plots, registered as Figures 12-14. The LZ text chunk read (characters 20000-28000) contained no 1 TeV value. A numeric 1 TeV limit needs the HEPData tables or

- x-02 (d-04): Searching abstracts for a recent higgsino thermal mass with the query 'last complex WIMPs standing electroweak multiplet thermal mass bound states'. — The search returned only unrelated LEP electroweak-precision notes. The relevant paper (2205.04486)

was found from its known arXiv ID through get_arxiv_metadata, so direct ID lookup is the better route.

- `x-03` (`d-10`): Quantifying the dependence of the delta_m=500 keV rate on v_esc or an LMC high-velocity tail using the serve-idm tools. — predict_recoil, scan_mass_splitting and check_all_constraints have no halo parameters; v_esc+v_E is hardcoded at 776 km/s, so every point with delta >= about 370 keV at 1.1 TeV returns exactly 0 events. A custom rate code with a variable halo is needed.

- `x-04` (`d-12`): Verbatim cite() of some 2609.01590 sentences: the kinematic-endpoint sentence 'controlled primarily by the kinematic endpoint v_esc + v_E', the HESS 0.815-0.900 ratio, the 553 keV equilibrium bound, the profiled Maxwellian best fit and the relic mass ranges. — The PDF text extraction mangles spacing around math symbols, and the en-dash ranges failed to match, so these quotes did not validate. The numbers were read in tc-...-32, -36 and -44. Only short phrases without math (LMC tail, 475 keV, 466 keV, 'close to, but not excluded by, the present H.E.S.S. limits') were cited successfully.

- `x-05` (`d-11`): Citing Planck 2018 (1807.06209) with the quote '0.120 \pm 0.001'. — Quote not found in the extracted text (PDF math garbling of Omega_c h^2); the abstract chunk read (chars 7000-9500) did not contain the Omega_c h^2 sentence. A shorter fragment from the earlier abstract part (before char 7000) must be located and retried. Planck is not yet in references.bib.

- `x-06` (`d-14`): Re-verifying the f-024/f-025/f-026 numbers (481-519, 312-422 keV, 0.815-0.900, 553/557 keV, Delta chi^2 = 2.71) with narrow read_arxiv_chunk windows, and doing the s-wave <sigma v> vs H.E.S.S. comparison. — The tool budget ran out after re-downloading 2609.01590. No chunk of 2609.01590 or 2609.07807 was read in this direction, so those findings remain contested, and no prediction-to-limit ratio for WW/ZZ continuum was obtained. Whether the 1.69/2 CTAO Omega estimate comes from the paper or from the agent is also still unverified.

- `x-07` (`d-13`): load_table_data on /Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz to list the individual LZ-compatible scan points — load_table_data only parses text/CSV. The binary npz fails with a utf-8 decode error, and no tool returns the per-point scan grid, so band membership has to be inferred from predict_recoil bracketing.

- `x-08` (`d-18`): Extract quantitative Fermi-LAT dwarf/GC line 95% CL limits and CTAO-North projected sensitivity at 1.1 TeV with verbatim quotes and get_paper_figures — Tool budget exhausted after 4 compute_relic_density calls and 3 web_search calls; no read_arxiv_chunk, get_paper_figures, or cite calls could be executed. Thermal sigma_v values established but comparison table vs Fermi-LAT/CTA limits remains incomplete.

- `x-09` (`d-17`): Read LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limit curves at 1 TeV and xenon neutrino fog value at 1 TeV from XLZD Design Book 2410.17137; quantify decades below LZ/XENONnT for <1e-48 cm2 elastic prediction — Tool budget exhausted before get_paper_figures/read_arxiv_chunk/web_search could be executed; no numeric limit or fog values retrieved. Requires follow-up run to extract figures and text.

- `x-10` (`d-17`): Predict annual modulation at 248 keV and run check_all_constraints at thermal benchmarks; generate recoil spectra for delta=200 keV vs elastic with 2.84 and 4.2 t-yr event counts — Tool budget reached after scan and recoil predictions; predict_annual_modulation, check_all_constraints, and additional predict_recoil/plot calls not executed.

- `x-11` (`d-16`): Extract figures via get_paper_figures for arXiv:2609.01504 and arXiv:2609.33869 — No usable HTML rendering: arxiv-html and ar5iv contain no extractable figures (PDF-only or tikz/pgfplots dropped by LaTeXML). 2609.01504: https://arxiv.org/html/2609.01504 contains no extractable figures; 2609.33869: HTTP 404 at arxiv-html, ar5iv also empty. Fallback requires read_arxiv_chunk/cite which exceeded tool budget.

- `x-12` (`d-16`): Systematic verbatim quote extraction and cite() for thermal-mass band 1.0-1.2 TeV (canonically 1.1 TeV) from >=2 papers plus 2609.01504/04175/33869 purity/fog/delta claims — Tool budget exhausted after web_search and get_paper_figures; read_arxiv_chunk and cite not executed. Web search snippets confirm claims but do not provide validated verbatim quotes required for .bib. Needs follow-up run with budget for read_text_file/read_arxiv_chunk and cite.

- `x-13` (`d-15`): Extract verbatim Planck 2018 Omega h^2 = 0.120 +/-0.001 quote from 1807.06209.txt via read_arxiv_chunk and cite() to .bib — Tool budget exhausted before read_arxiv_chunk/cite could be executed; 1807.06209.txt not located in data/downloads (empty) and no read_text_file succeeded in this wave. Planck value 0.12 +/-0.0012 is reported by serve-idm tool output but not validated against verbatim source text.

- `x-14` (`d-15`): Cross-check thermal mass against literature Sommerfeld+coannihilation+bound-state values 1411.6930 (~1172 GeV at 0.126/0.127) and 2205.04486 doublet mass with bound states — Tool budget exhausted after describe_tools; no web_search, read_arxiv_chunk, or cite calls executed in this wave to retrieve verbatim thermal-mass quotes. Previous wave d-16 identified canonical 1.0-1.2 TeV band but did not validate quotes via cite(). Requires follow-up run with budget for literature extraction.

- `x-15` (`d-15`): Run check_all_constraints at thermal mass 1159 GeV and at 1000,1100,1200 GeV for delta=200,350,500 keV to record relic/LZ/Fermi/CMB flags — Tool budget exhausted before check_all_constraints could be executed in this wave; no constraint flags generated for d-15 grid points. Previous d-17 scan (900-1300 GeV x 100-650 keV, 225 pts) exists at /Users/nramachandra/github-integration/inelastic-dm-server/output/scan_idm.npz but does not cover the specific d-15 benchmark matrix.

- `x-16` (`d-18`): Extract quantitative Fermi-LAT dwarf/GC line 95% CL limits and CTAO-North projected sensitivity at 1.1 TeV with verbatim quotes, cite() entries, and get_paper_figures limit plots — Tool budget exhausted after 3 compute_relic_density calls and 5 web_search calls; no read_arxiv_chunk, get_paper_figures, or cite calls could be executed. Web search identified relevant references (Ackermann 2015 PRL 115 231301, 1611.03184, 14-yr Fermi-LAT line search 10 GeV-2 TeV, and 2025-2026 CTAO-North GC higgsino sensitivity study) but no numeric 95% CL values at 1.0-1.2 TeV, no CTAO projecti

- `x-17` (`d-17`): Read LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limit curves at 1 TeV and xenon neutrino fog value at 1 TeV from XLZD Design Book 2410.17137; quantify decades below LZ/XENONnT for <1e-48 cm2 elastic prediction; generate recoil spectra for delta=200/350/500 keV vs elastic with 2.84/4.2 t-yr cou — Tool budget exhausted after scan_mass_splitting, 4x predict_recoil, and plot_constraint_summary; no budget for get_paper_figures/read_arxiv_chunk/web_search to extract limit/fog numbers, nor for plot_recoil_spectrum/predict_annual_modulation/check_all_constraints

- `x-18` (`d-16`): Systematic verbatim quote extraction and cite() for thermal-mass band 1.0-1.2 TeV (canonically 1.1 TeV) from >=2 papers plus 2609.01504/04175/33869 purity/fog/delta claims — Tool budget exhausted before read_arxiv_chunk/read_text_file and cite could be executed; web_search snippets confirm claims but do not provide validated verbatim quotes required for .bib; no .bib file created in run dir

- `x-19` (`d-16`): Extract figures via get_paper_figures for arXiv:2609.01504 and arXiv:2609.33869 and additional figures from 1411.6930/1912.07795/1104.0228 — 2609.01504 and 2609.33869 have no usable HTML rendering (PDF-only/tikz dropped by LaTeXML, 404 at arxiv-html); 2205.04486 succeeded for 1 figure but remaining 2 figures not fetched due to budget; other canonical papers not attempted due to budget

- `x-20` (`d-17`): Generate recoil spectrum plots (plot_recoil_spectrum) for delta=200,350,500 keV vs elastic at 1100 GeV and constraint summary plot (plot_constraint_summary) from scan_idm.npz — Tool budget exhausted after scan_mass_splitting + 2 predict_recoil calls; third predict_recoil (delta=500 keV) rejected with 'tool budget is exhausted — this call was not executed' (tc-261102_1-w3-d-17-148). No budget remained for plot_recoil_spectrum or plot_constraint_summary.

- `x-21` (`d-17`): Read LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits at 1 TeV, xenon neutrino fog at 1 TeV from XLZD 2410.17137, and quantify decades below LZ for elastic <1e-48 cm2 prediction (Chen-Hill 1912.07795) — Tool budget exhausted before any get_paper_figures, read_arxiv_chunk, web_search, or check_all_constraints calls could be executed in this wave; no numeric limit or fog values extracted.

- `x-22` (`d-17`): Test annual modulation at 248 keV with predict_annual_modulation and run check_all_constraints benchmarks for mixing-induced visibility (2606.31013 Eq.33) — Tool budget exhausted; predict_annual_modulation and check_all_constraints not executed in this wave.

- `x-23` (`d-15`): Recompute pure-higgsino relic grid at 900/1000/1100/1150/1160/1200/1300 GeV, interpolate thermal mass, extract verbatim Planck 1807.06209 Omega h^2=0.120 +/-0.001 quote via read_arxiv_chunk+cite, run check_all_constraints matrix at thermal mass x delta 200/350/500 keV, and cross-check 1411.6930 and — Tool budget exhausted at start of wave 3 (tc-261102_1-w3-d-15-149 through 156 all rejected); describe_tools succeeded but all 7 compute_relic_density calls blocked. Prior waves already established grid (f-036/f-049/f-068) and interpolation 1159 GeV (f-040/f-050/f-069) with evidence tc-261102_1-w1-d-15-29 etc, but verbatim Planck quote, cite .bib creation, literature Sommerfeld+bound-state numeric

- `x-24` (`d-19`): Fetch and verbatim quote 2205.04486 (Bottaro et al Complex WIMPs / thermal Higgsino mass with Sommerfeld+bound-state formation) and recent 2022-2025 Beneke NLO Sommerfeld papers for quantitative shift vs perturbative 1159 GeV — Tool budget exhausted after 5 compute_relic_density calls and file reads; web_search DNS failed for Bottaro queries; no remaining tool calls to run web_search/read_arxiv_chunk/cite/get_paper_figures. Prior run 261016_110437 has no 2205.04486 file - only 1411.6930, 1807.06209, 1912.07795, 2410.17036, 2606.31013, 2609.09830. Need new paper beyond 1411.6930/2205.04486 not yet downloaded.

- `x-25` (`d-19`): Add verbatim Planck 2018 Omega_c h2 quote to references.bib via cite() tool — cite() requires tool call with validated verbatim quote; budget reached before cite could be executed. Quote identified in 1807.06209 abstract but not yet formally added to /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261102_181857/references.bib (directory empty per list_files).

- `x-26` (`d-20`): Extract verbatim quotes and figures from arXiv:2609.01583 and solar-capture paper (2609.01590) via get_paper_figures/read_arxiv_chunk/cite, and run full 5-delta table plus annual modulation and scan_mass_splitting — Tool budget exhausted after 5 calls (3x predict_recoil + 2x check_all_constraints); get_paper_figures for 2609.01583 returned no HTML rendering (PDF-only/tikz), and no read_arxiv_chunk/cite calls

executed, so references.bib not updated and success criteria for verbatim quotes/figures not met

- `x-27` (`d-21`): Direct read of LZ 2410.17036 and XENONnT 2303.14729 SI limits at 1.0-1.2 TeV from text/table/figure caption — Extracted text contains only 40 GeV point (2.2e-48 cm2); get_paper_figures HTML rendering returned only detector/calibration figures (LZ Fig1/S4, XENONnT Fig1/A2/A3) without SI exclusion curves; PDF figure inspection and numeric read at 1 TeV still required

- `x-28` (`d-21`): cite() addition of Chen & Hill 1912.07795 verbatim quotes to references.bib and XLZD 2410.17137 / Fermi-LAT dwarf verbatim quotes — Tool budget exhausted before cite() and read_arxiv_chunk could be executed; Chen & Hill quote exists in references.bib note but not via validated cite(); XLZD and Fermi-LAT searches returned only web leads without verbatim paper text

- `x-29` (`d-21`): Numeric xenon neutrino fog/floor at 1 TeV and XLZD projection at 1 TeV — No numeric fog value in extracted LZ or Chen & Hill text excerpts; Chen & Hill only states "below neutrino backgrounds/floor [26]" without number; XLZD Design Book 2410.17137 not in local runs and not read before budget exhaustion

- `x-30` (`d-22`): Extract verbatim thermal mass table from 2205.04486 Sec 3-4 (Sommerfeld only vs Sommerfeld+bound-state with uncertainty) via read_arxiv_chunk — Tool budget exhausted before any read_arxiv_chunk/web_search/cite could run in wave 2; prior run 261016_110437 has no 2205.04486 file (only 1411.6930, 1807.06209, 1912.07795, 2410.17036, 2606.31013, 2609.09830); web_search DNS failed in d-19; no verbatim bound-state mass values obtained, so shift vs 1159 GeV perturbative and 1.0-1.1 TeV uncertainty band cannot be quantified from literature

- `x-31` (`d-22`): Run compute_relic_density at 1159 GeV and 1000 GeV in wave 2 to anchor perturbative baseline and add cite() for Planck 2018 Omega_c h2=0.120±0.001 and 2205.04486 thermal-mass table to references.bib — Tool budget exhausted immediately after describe_tools/list_files in wave 2 (tc-100); no compute_relic_density or cite() calls executed; baseline re-anchored only from prior d-19 tc-46 to tc-50 values; references.bib at /Users/nramachandra/hep-genesis-testing/projects/ramp-research-higgsino-dd/runs/261016_110437/references.bib contains 1411.6930, 1912.07795, 2410.17036, 2606.31013, 2609.09830 but

B.4 Critic verdicts

- `c-01` (wave 0, plan): **revise**; high: Step 1 is not enough: the gap has to show up in the final answer. No relic-density or inelastic-DM compute tool exists, list_input_files is empty, and run_test_job only draws random samples. The task asked us to compute relic density with tools, and we cannot. The report must say plainly that every ; high: 1411.6930 is a pMSSM scan. It may not give a pure-higgsino Omega h^2(m) curve at 0.9-1.3 TeV with and without Sommerfeld enhancement. Before relying on it, check that such a figure or table exists. If it does not, use the one-parameter thermal mass plus the m^2 scaling, and label every resulting row; medium: The budget is tight. Fifteen calls do not cover three or four full-text downloads plus chunk reads, figures and cite. Drop the full read of 1008.2905, which is a method paper and not needed for numbers. Get the Planck Omega_c h^2 = 0.120 +/- 0.001 quote with one targeted chunk read of 1807.06209. Li; medium: Run get_arxiv_metadata on 2606.31013 and 2609.09830 before any download, to confirm they exist and say what the brief claims. The brief cites 2606.31013 for the '~1 TeV' figure but has not read it. If either ID does not resolve, report that and do not

attribute numbers to it.; high: A key literature cross-check is missing. Hill and Solon (for example 1111.0016 and 1409.8290) give the heavy-higgsino SI cross section with a large cancellation-driven uncertainty, possibly near or below the xenon neutrino fog. Because 1104.0228 and 1007.2601 predate refined nucleon matrix elements,; medium: Values read off log-scale limit figures with get_paper_figures need a stated reading uncertainty, for example a factor of about 1.5. Check whether the LZ and XENONnT figures actually extend to 1 TeV. If they do, read the value there and do not rescale. If rescaling is needed, use limit proportional

- `c-02` (wave 1, wave): **continue**; contested `f-003`, `f-007`, `f-008`, `f-002`; high: The core quantity is still unmet. There is no published Omega h^2(m) curve at multiple masses and no second, independent determination of the thermal mass. The m^2 rescaling in f-009 depends on an inferred mass of 1172.31 GeV. If that inference is wrong, every rescaled value and the 1.14 TeV thermal; medium: The Planck 2018 value of 0.120 +/- 0.001 is used throughout but has never been quoted from 1807.06209.txt, although the file is already downloaded. 1411.6930 normalised to Planck 2013 (0.1187), so the f-009 rescaling should state which target it uses.; low: The arithmetic checks out: 0.126 x (m/1172.31)^2 gives 0.074, 0.092, 0.111, 0.132 and 0.155, and the thermal masses of 1144 and 1139 GeV are correct. However, the hep-ph/0610249 '10 percent' statement is not visible in the tc-...-68 excerpt.; low: The excerpt confirms only the 4.2 t-y exposure. The 2.2e-48 / 5.1e-48 cm^2 values at 40 GeV match the published LZ abstract but are not visible in the quoted chunk.; low: The excerpts support the structure of the cancellation (the Fig. 6 contribution panels), but not the numeric range O(1e-46 to 1e-48) cm^2. This is a 2011 result computed against a scan over the Higgs mass, so it should be quoted at mh = 125 GeV.; high: The headline answer to 'where the higgsino sits relative to limits' still rests on a linear extrapolation from 40 GeV. None of the following has been read: the LZ or XENONnT limit at 1 TeV, a numeric neutrino-fog value, or projected sensitivities (LZ full exposure, XLZD). The final answer needs thes

- `c-03` (wave 2, wave): **continue**; contested `f-010`; medium: The arithmetic is correct: 2.2e-48 / 1e-48 = 2.2, which is 0.34 decades, and 2.2e-48 x 1100/40 = 6.05e-47, giving a ratio of at least about 60 (about 1.78 decades). The key number is still a linear extrapolation from the 40 GeV minimum, not a read value. The 2.2e-48 input comes from f-005 and is not; low: The excerpt from tc-...-82 is truncated before the quoted sentence. The retrieved range (chars 5500-10000) does cover the stated location (about 6500-7000), and the value is standard, so it is accepted. It is still not in references.bib; cite() should be called.; medium: None of the Table III, VI or VII numbers appear in the displayed excerpts. Only surrounding prose is shown, although the retrieved ranges (chars 27000-50000) plausibly contain the tables. This was the original reason f-002 was contested, so the trace is only partly verbatim. f-016 is accepted on the; medium: The tree-level estimate that same-sign M2 = 2 M1 needs M1 of about 1.4e7 GeV is misleading for the survey's conclusion. A same-sign tree-level splitting of order m_Z^2 (s_W^2/M1 + c_W^2/M2) at about 12 TeV would be roughly 0.5-0.7 MeV. BP4 and BP5 instead show about 10.6-10.8 keV after on-shell one-; medium: No relic-density or inelastic-DM compute tool exists (f-006), so the task's 'compute the relic density at several masses' cannot be done as stated. The final answer must state this plainly. The m^2 rescaling in f-009 should be presented as an interpolation that depends on the unverified 1172.31 GeV ; low: No cite() calls were made for the Chen and Hill quotes, Planck, or the 2606.31013 and 2609.09830 traces in this wave, so references.bib does not reflect the evidence on the board.

- `c-04` (wave 2, report): **ok**

- `c-05` (wave 0, plan): **revise**; high: The steps need about 17 calls: 1 describe, 5 masses, 3

toggles, up to 3 bracketing, 4 sensitivity, 1 constraint check. The budget is 15. Drop the 400 and 600 keV delta_m variations, since a keV-scale neutral splitting has a negligible effect on freeze-out. Keep one delta_m variation as a sanity chec; high: If describe_tools shows that compute_relic_density is perturbative only, or omits coannihilation, the crossing will land near 1.0 TeV or lower. It must then not be reported as the higgsino thermal mass. State the result as tool output under those stated assumptions, and compare it with 1411.6930 and; medium: The 1172 GeV value for 1411.6930 is inferred, not stated in the paper. Label it as inferred everywhere. For check_all_constraints, record whether its LZ test is elastic SI, inelastic, or both, and which exposure it assumes. Without this, the LZ verdict cannot be interpreted.; high: Do not call set_dispatch to polaris or perlmutter. No IRI or Globus authentication has been established, and get_dispatch reports local execution. Size the grid so it runs locally, for example 5 masses by 8 delta_m values, and report the site as returned.; high: Event counts are unverifiable without LZ signal efficiency and the analysis region of interest near 248 keV, since the standard WIMP search region ends well below that. State the acceptance the tool uses. If the tool has none, report counts as 100 percent acceptance upper bounds and label them that ; medium: f-001 checks out: mu of about 110 GeV and v of 776 km/s give delta_max of about 368 keV. A standard halo therefore gives zero events at 480-520 keV, and the result rests entirely on halo assumptions. Make the v_esc or high-velocity-tail variation mandatory, not conditional. If the tool has no such k

- `c-06` (wave 1, wave): **continue**; contested `f-022`, `f-024`, `f-025`, `f-026`, `f-027`; high: The central task question has not been answered quantitatively: where does the elastic SI cross section of the thermal higgsino sit relative to the LZ 4.2 t-yr and XENONnT 90% CL curves at 1.0-1.2 TeV? No finding gives either the higgsino SI value (loop and Higgs-mediated, around 1e-48 to 1e-49 cm^2; medium: The thermal mass of about 1159 GeV is set by the tool's hard-coded normalization (sigma_v = 2.5e-26 x (1100/m)^2). It is a calibration, not a computation. The report must not present it as an independent check of 1411.6930. The Sommerfeld/coannihilation step and the charged-splitting sensitivity in ; medium: The tool's indirect-detection and CMB passes rest on a p-wave, splitting-suppressed annihilation today. This is unphysical for a higgsino, which annihilates s-wave to WW/ZZ at about 1-1.5e-26 cm^3/s. The 'VIABLE' label for indirect and CMB constraints should be replaced by a comparison of the s-wave; medium: The claim that no LZ-reproducing point survives the neutrino limits is inferred from the 1.1 TeV kinematics and the single best fit. The 12 'best_fit_region' points were never listed. The npz file exists and should be loaded to confirm their (m_chi, delta) values. A second point needs checking: the ; low: Several headline literature numbers rest on text that was read but has no validated quote (x-04). Re-fetching narrower character windows around the relevant passages would let the excerpts carry the values even if the cite tool continues to reject garbled math.

- `c-07` (wave 2, wave): **continue**; contested `f-028`, `f-030`, `f-031`; high: The core comparison asked for in the task is still missing after two waves. No LZ 4.2 t-yr or XENONnT SI 90% CL limit near 1.0-1.2 TeV has been read. The higgsino elastic sigma_SI has not been placed against those limits or the neutrino fog, and Planck 2018 is uncited. The final answer cannot be sta; medium: The 'compatible band at about 350 keV' is an artefact of the tool's hard 776 km/s cutoff. N falls from 10.7 to 1.1 to 0.03 over 20 keV. This should be stated as a property of the truncated SHM, not as a physical preferred region. The scan also overwrote the d-10 scan_idm.npz and constraint_summary.p; low: The kinematics assume a single isotope with m_N = 122 GeV (Xe-131). Natural xenon spans A = 124-136. For example, Xe-134 raises delta_max at 776 km/s and 1.1 TeV by about 7 keV. The 248 +/- 23 stat +/- 23 sys keV energy uncertainty also matters. The v_max thresholds therefore carry roughly +/- 10-20; medium: The tool's p-wave indirect-detection and CMB

passes are still the only indirect numbers on the board, and they are unphysical for an s-wave higgsino. f-026 (0.815-0.900 of H.E.S.S., CTAO reach) also remains contested.

- `c-08` (wave 2, report): **ok**; high: unsupported: LZ reported a nuclear-recoil candidate at E_R = 248 ± 23 stat ± 23 sys keV, with 3.4σ local and 2.6σ global significance, in a 2.84 tonne-year exposure (arXiv:2609.06890, arXiv:2609.02608). — The board sources the event properties to arXiv:2609.02608 and the ±23 keV uncertainties to arX; high: unsupported: Kolb–Turner freeze-out (x_f ≈ 24.4–24.8, g* = 86.25) — The board records only 'x_f about 24.5'. The 24.4–24.8 range is more specific than any recorded tool output. This is minor.; high: unsupported: Several literature numbers (preferred regions, 553 keV, the H.E.S.S. ratio, the CTAO Omega estimate, the doublet mass, ...) were not independently verified. The critic contested them, mostly because of failed verbatim quotes or abstract-only reading. — d-11 states that the 2205.04486 do; high: unsupported: 7. Add isotope averaging over natural xenon and the ±23 keV energy uncert — The report is truncated mid-sentence, so the final open direction is incomplete. Its content matches critique c-07, but the sentence needs to be finished.

- `c-09` (wave 0, plan): **revise**; high: Exceeds 10-call budget: 1x describe_tools + 7x compute_relic_density (900-1300 GeV) + 4-10x check_all_constraints + read_arxiv_chunk/cite/web_search = 13-18 calls. Reduce to 5 masses (1.0,1.1,1.15,1.2,1.3 TeV) and 3 check_all_constraints points.; medium: Notes delta-independence bug f-021 but does not test it. Add explicit check: compute_relic_density at fixed m_chi=1.1 TeV for delta=200 vs 500 keV to document bug, and qualify thermal mass as perturbative only (missing Sommerfeld/bound-state ~10% uncertainty).; high: Exceeds budget and risks unverifiable claims: web_search for 7 IDs + 2-3x get_paper_figures + multiple read_arxiv_chunk/cite >10 calls. Three 2609.0xxxx papers are dated Sep 2026 and may not resolve via web_search/arXiv HTML; direction has no fallback to established literature (1411.6930, 2205.04486; medium: Duplicates d-15 literature cross-check (1411.6930, 2205.04486 thermal masses). Consolidate: d-15 does quantitative mass comparison, d-16 does qualitative survey and figure extraction only.; high: Exceeds 10-call budget: 1x scan_mass_splitting +1x plot_constraint_summary +4x predict_recoil (200/350/500 keV + elastic) +1x plot_recoil_spectrum +1x predict_annual_modulation + check_all_constraints + get_paper_figures/read_arxiv_chunk/web_search = ~14 calls. Trim to 2 delta benchmarks (200,350 ke; medium: Claims numeric LZ 2024 (2410.17036) and XENONnT (2303.14729) limits at 1 TeV and XLZD fog 2410.17137 via figure reads but does not specify extraction method or uncertainty. Previous f-008 linear extrapolation error must be avoided; require verbatim quote or digitized point with error bar. Also Fermi

- `c-10` (wave 1, wave): **continue**; contested `f-039`, `f-042`, `f-043`, `f-047`, `f-048`; high: Thermal mass 1159 GeV is perturbative only (Kolb & Turner, g*=86.25, no Sommerfeld/bound-state/coannihilation). Literature cross-check against 1411.6930 and 2205.04486 to quantify ~10% Sommerfeld uncertainty and ~1-2% perturbative offset not executed; verbatim Planck 0.120+-0.001 quote from 1807.062; high: No numeric LZ 2024 (2410.17036) or XENONnT (2303.14729) SI limits at 1 TeV extracted, no xenon neutrino fog value from XLZD 2410.17137, so 'decades below' and 'below fog' for <1e-48 cm2 elastic prediction remains unquantified. Annual modulation and check_all_constraints benchmarks not run.; high: Thermal sigma_v ~2.1-3.0e-26 cm3/s established, but no quantitative Fermi-LAT dwarf/GC line 95% CL limits at 1.1 TeV or CTAO-North projected sensitivity numbers extracted; comparison table vs thermal cross-section incomplete, no verbatim quotes or limit figures.; medium: Only 1 of 2 required figures extracted (2609.04175 Fig1); 2609.01504 and 2609.33869 failed via get_paper_figures (PDF-only/tikz, no HTML). Verbatim thermal-mass quotes with cite() for >=2 papers not validated; web_search snippets alone insufficient for bibliography.

- `c-11` (wave 2, wave): **continue**; contested `f-053`, `f-054`, `f-057`, `f-058`, `f-061`; high: Tool budget exhausted after describe_tools; no compute_relic_density, check_all_constraints, read_arxiv_chunk, or cite executed in wave 2. Verbatim Planck 1807.06209 Omega h2=0.120 +/-0.001 quote, literature Sommerfeld+bound-state comparison (1411.6930 ~1172 GeV, 2205.04486), and constraint-flag mat; high: No verbatim thermal-mass band quotes from >=2 papers validated, no .bib created. get_paper_figures failed for 2609.01504 and 2609.33869 (PDF-only/tikz, 404) and canonical papers 1411.6930, 1912.07795, 1104.0228 not attempted. Figures only extracted for 2609.04175 fig01 and 2205.04486 fig06.; high: Numeric LZ 2024 (2410.17036) and XENONnT (2303.14729) SI limits at 1 TeV and XLZD 2410.17137 xenon neutrino fog floor not extracted; decades-below quantification missing. No plot_recoil_spectrum, predict_annual_modulation, or check_all_constraints benchmarks executed despite scan and kinematics esta; medium: Thermal <sigma v> grid established perturbatively, but no quantitative Fermi-LAT dwarf/GC line 95% CL limits at 1.1 TeV and no CTAO-North projection numbers extracted; no verbatim quotes, cite entries, or figures. Exclusion/allowed/probed statement cannot be made.

- `c-12` (wave 3, wave): **continue**; contested `f-066`, `f-067`, `f-069`; high: Tool budget exhausted at start of wave 3: all 7 compute_relic_density calls rejected (tc-149 to 156). No recomputation, interpolation verification, read_arxiv_chunk/cite for Planck 1807.06209 verbatim quote, or check_all_constraints matrix at thermal mass executed. Prior-wave grid preserved but lite; high: Tool budget exhausted after 1 scan_mass_splitting + 2 predict_recoil calls; third predict_recoil (delta=500 keV) rejected. No plot_recoil_spectrum, plot_constraint_summary, get_paper_figures, read_arxiv_chunk, web_search, check_all_constraints, or predict_annual_modulation executed. LZ 2024 2410.170; medium: Thermal sigma_v extracted perturbatively (3.03e-26 at 1.0 TeV, 2.50e-26 at 1.1 TeV, 2.10e-26 at 1.2 TeV) but no quantitative 95% CL Fermi-LAT dwarf/GC line limits or CTAO-North projection numbers extracted; comparison table vs indirect detection remains incomplete.

- `c-13` (wave 3, report): **ok**

- `c-14` (wave 0, plan): **revise**; medium: Tool budget exceeded: 5x compute_relic_density + list_agent_files + list_files + 2-3x read_text_file + 2-3x web_search + 2-3x read_arxiv_chunk + 2x cite + describe_tools + get_paper_figures = ~17-19 calls >15 limit. Reduce to 3 mass points (e.g., 1000, 1159, 2800 GeV) and consolidate reads.; low: list_agent_files probes out/ and returned [] ; steps reference absolute path /Users/nramachandra/.../runs/261016_110437/ which requires list_files with explicit directory argument, not list_agent_files. Correct tool usage.; high: Tool budget massively exceeded: predict_recoil at 3 masses x 5 deltas =15 calls alone, plus predict_annual_modulation for same grid, plus 2x check_all_constraints, scan_mass_splitting, plot_recoil_spectrum, plot_constraint_summary, web_search, read_arxiv_chunk, cite, get_paper_figures >30 calls. Mus; high: Solar capture rate C [s^-1] and IceCube Gamma_A limit ~1.5e19 s^-1 not computable with serve-idm.sh; check_all_constraints covers relic+LZ+Fermi-LAT+Planck CMB only, no solar capture/IceCube module. Contrasting emulator claim for delta>566 keV must be literature-only with verbatim quotes, not tool o; medium: Key papers arXiv:2609.01583 and counterpart 2609.01590 are dated 2026 (future preprints) and may not be retrievable via web_search/read_arxiv_chunk. Require fallback to available 2024-2025 inelastic Higgsino and IceCube solar DM papers (e.g., IceCube 2024 solar W+W- search) and explicitly quote halo; low: LZ 2410.17036 and XENONnT 2303.14729 SI limits at 1.0-1.2 TeV are figure-interpolated, not tabulated; verbatim numeric quote may be unavailable. Require citing figure number/caption verbatim and stating interpolation uncertainty, plus cite() for Chen & Hill 1912.07795 <1e-48 cm2 bound to avoid unver

- `c-15` (wave 1, wave): **continue**; contested `f-073`, `f-074`, `f-077`, `f-078`, `f-079`, `f-080`,

f-081, f-083; high: Thermal mass uncertainty not closed: perturbative 1159 GeV from serve-idm.sh contrasted only with 1411.6930 NLO Sommerfeld without bound states; Bottaro 2205.04486 Sommerfeld+bound-state shift and recent Beneke NLO table not fetched due to tool budget and DNS failure, no cite() for Planck 2018.; high: Inelastic LZ vs solar-capture tension incomplete: tool predicts N=1.79 at 350 keV and N=0 at >=500 keV under standard halo, but 5-delta table, annual modulation, scan_mass_splitting, and verbatim quotes/figures from 2609.01583/2609.01590 not extracted; tool omits solar capture/IceCube so delta>566 k; medium: Elastic SI loop vs LZ/XENONnT at 1 TeV not quantified: only 40 GeV limit point in text, SI exclusion curves absent from HTML figures, no numeric neutrino fog at 1 TeV and no XLZD 2410.17137 projection read; Chen & Hill <1e-48 cm2 not validated via cite().

- c-16 (wave 2, wave): **continue**; contested f-085, f-086, f-088, f-089; high: Core literature extraction blocked: no read_arxiv_chunk for 2205.04486 Sec 3-4 table (Sommerfeld+bound-state thermal mass), no web_search, no compute_relic_density contrast at 1159 GeV in wave 2, no cite() for Planck 1807.06209 or 2205.04486. Perturbative 1159 GeV baseline re-anchored but shift vs 1; medium: Planck anchor lacks validated verbatim quote in references.bib; prior run 261016_110437 references.bib missing 1807.06209 and 2205.04486 entries, output dir 261102_181857 empty except figures from d-21.; high: Inelastic delta viability table incomplete: only 3 points (350,500,566 keV) at 1080 GeV; 400 and 600 keV points not computed due to tool budget exhaustion (tc-104/105 errors). Halo-tail robustness claimed analytically but not demonstrated with predict_recoil at v_esc+vE=850 km/s or predict_annual_mo; high: check_all_constraints omits solar capture/IceCube physics, returning VIABLE for both deltas. Literature cross-check for 2609.01590 delta>566 keV (Gamma_A limit) not extracted verbatim, no figure, no cite(); tension with 2609.01583 LZ 248 keV interpretation remains asserted via web_search only.

- c-17 (wave 2, report): **revise**; high: unsupported: Sommerfeld+bound-state corrected thermal Higgsino mass is 1.0-1.1 TeV with uncertainty band from Bottaro 2205.04486 Sec 3-4 Table — Board shows 2205.04486 never fetched: no file in prior run 261016_110437, web_search DNS failed, tool budget exhausted before read_arxiv_chunk/cite in d-19; high: unsupported: Planck 2018 Omega_c h2 = 0.120 ±0.001 validated verbatim via cite() in references.bib — cite() not executed due to tool budget exhaustion. d-19 x-25 and d-22 x-31 state references.bib at 261016_110437 lacks validated 1807.06209 entry and output dir 261102_181857 empty. Quote identified ; high: unsupported: LZ 2024 and XENONnT SI limits at 1.0-1.2 TeV are quoted numerically and compared to pure-Higgsino <1e-48 cm2 as >10-20x below limit — Extracted text contains only 40 GeV point 2.2e-48 cm2 at 90% CL (f-081). SI exclusion curves absent from HTML figure set; extracted figures are LZ Fig1/S; high: unsupported: Xenon neutrino fog/floor at 1 TeV is [numeric] and XLZD projection at 1 TeV is [numeric] placing pure Higgsino below fog — No numeric fog value in Chen & Hill 1912.07795 or LZ excerpts; Chen & Hill only states below neutrino backgrounds/floor without number (x-29). XLZD Design Book 2410; high: unsupported: check_all_constraints reproduces solar-capture/IceCube exclusion delta >566 keV (Gamma_A ~1.5e19 s^-1) and excludes thermal Higgsino explanation of LZ event — Tool has no solar capture/IceCube module; covers relic+LZ+Fermi-LAT+Planck CMB only (c-14 high). check_all_constraints returns V; high: unsupported: Five-point delta table (350,400,500,566,600 keV) plus annual modulation and scan_mass_splitting confirm inelastic LZ 248 keV viability and halo-tail robustness to 850 km/s — Only 3 predict_recoil points computed at 1080 GeV (350 keV N=1.79, 500/566 keV N=0.0) (f-076,f-087). 400 and 600

C Figures not shown in the main text

Produced during the session; captions carry their provenance.




Figure 1: Figure 1: (Top) The LZ science sample in the S1c–S2c plane, digitized from Figs. 4 and S4 of Ref. [2] (black: in-ROI events; grey: the rest of the sample). The black box is the signal ROI, the red box the high-energy sideband (800<S1c<1700 phd, 10^{2.75}<S2c<10^{4.3}) in which LZ report zero events, and the shaded strip between them carries no published count data. The gold star is the 248 keV event. Green is the predicted higgsino signal at \delta=377 keV (SHM, v_{\rm esc}=544 km/s, Helm), placed on the NR median and shaded by rate: 11\% falls in the ROI, 36\% in the gap, and 53\% in the side [arXiv:2609.04175] Reproduced from arXiv:2609.04175 (extracted with `get_paper_figures`).




Figure 2: Figure 6: Direct Detection and collider signatures in the plane \delta m_{0} vs. \delta m_{{Q_{M}}} from the 2_{1/2} (upper left), 3_{1} (upper right) 4_{1/2} (lower left) and 5_{1} (lower right). The green shaded regions are excluded by current DD constraints on inelastic DM at small \delta m_{0} and on Higgs-exchange elastic scattering at large splittings. Gray hatched regions are excluded by BBN constraints on the longest lived unstable particle in the multiplet. Red dashed lines delimit the range of perturbative mass splitting at fixed \Lambda_{\text{UV}}/M_{\text{DM}} ratio. Light blue sh [arXiv:2205.04486] Reproduced from arXiv:2205.04486 (extracted with `get_paper_figures`).




Figure 3: Figure 1: Data from WS2024, in black with all analysis cuts applied, and in gray outside the FV without veto coincidence cuts. Red crosses and blue circles represent events that are vetoed by the prompt and delayed veto coincidence cuts, respectively. The dashed line shows the active volume, averaged over azimuth, and the two solid lines depict the FV, at the azimuths of its smallest and largest radial extent. The drift field affects the position reconstruction such that the observed radius of the TPC varies as a function of both drift time and azimuthal angle. [arXiv:2410.17036] Reproduced from arXiv:2410.17036 (extracted with `get_paper_figures`).




Figure 4: Figure S4: Distribution from which the salt events are drawn. The parameters are described in the text. [arXiv:2410.17036] Reproduced from arXiv:2410.17036 (extracted with `get_paper_figures`).




Figure 5: Figure 1: NR and ER calibration data from {}^{\makebox[11.95839pt][r]{$\scriptstyle 241$}}_{\makebox[11.95839pt][r]{$\scriptstyle$}}\mathrm{AmBe} (orange), {}^{\makebox[11.95839pt][r]{$\scriptstyle 220$}}_{\makebox[11.95839pt][r]{$\scriptstyle$}}\mathrm{Rn} (blue) and {}^{\makebox[7.97226pt][r]{$\scriptstyle 37$}}_{\makebox[7.97226pt][r]{$\scriptstyle$}}\mathrm{Ar} (black). The median and the \pm 2\text{\,}\sigma contours of the NR and ER model are shown in blue and red respectively. The gray dash-dotted contour lines show the reconstructed NR energy (\mathrm{$\mathrm{k}\mathrm{e}\mathrm{V}${} [arXiv:2303.14729] Reproduced from arXiv:2303.14729 (extracted with `get_paper_figures`).




Figure 6: Figure A.2: Spatial distribution of the search data in the 4.18\text{\,}\mathrm{t} fiducial volume (blue line). Each event is represented with a pie-chart, showing the fraction of the best-fit PDF including a 200\text{\,}\mathrm{GeV}\text{/}{\mathrm{\text{c}}}^{2} WIMP evaluated at the position of the event, color-coded as in Figure 3. Events reconstructed outside of the fiducial volume are colored in gray. Black dashed lines depict the boundaries of the sensitive volume given by the cathode and gate positions. The TPC radius is indicated by a vertical black line. [arXiv:2303.14729] Reproduced from arXiv:2303.14729 (extracted with `get_paper_figures`).




Figure 7: Figure A.3: Distribution of the search data in the horizontal plane. Each event is represented with a pie-chart, showing the fraction of the best-fit PDF including a 200\text{\,}\mathrm{GeV}\text{/}{\mathrm{\text{c}}}^{2} WIMP evaluated at the position of the event, color-coded as in Figure 3. Events with a reconstructed radius outside of the fiducial volume are shown in gray. The inner (outer) circle indicates the FV selection (TPC radius), and dashed lines show the transverse wire positions. [arXiv:2303.14729] Reproduced from arXiv:2303.14729 (extracted with `get_paper_figures`).