

Research Group Presentation

Overview of Bayesian analysis

Motivating example: Likelihood

- ▶ Suppose X_1, \dots, X_n is an i.i.d. sample with $X_i \sim \text{Exp}(\lambda)$. We would like to estimate λ .

Motivating example: Likelihood

- ▶ Suppose X_1, \dots, X_n is an i.i.d. sample with $X_i \sim \text{Exp}(\lambda)$. We would like to estimate λ .
- ▶ Likelihood approach:

$$\hat{\lambda}_{\text{MLE}} = \frac{n}{\sum_i x_i}$$

$$I(\lambda) = \lambda^{-2}$$

Motivating example: Likelihood

- ▶ Suppose X_1, \dots, X_n is an i.i.d. sample with $X_i \sim \text{Exp}(\lambda)$. We would like to estimate λ .
- ▶ Likelihood approach:

$$\hat{\lambda}_{\text{MLE}} = \frac{n}{\sum_i x_i}$$

$$I(\lambda) = \lambda^{-2}$$

- ▶ If $z_{.025} = \text{qnorm}(.975)$, then approximately and for large n , the interval

$$\left(\hat{\lambda}_{\text{MLE}} - \frac{z_{.025} \lambda_{\text{MLE}}}{\sqrt{n}}, \hat{\lambda}_{\text{MLE}} + \frac{z_{.025} \lambda_{\text{MLE}}}{\sqrt{n}} \right)$$

has 95% coverage, meaning if we repeat the experiment 100 times, the interval will contain the true value of λ in roughly 95 of them.

Motivating example: Bayesian

- ▶ Think of λ as a quantity with its own distribution

Motivating example: Bayesian

- ▶ Think of λ as a quantity with its own distribution
- ▶ Prior distribution: $p(\lambda) = \text{Gamma}(\lambda, \alpha, \beta)$

Motivating example: Bayesian

- ▶ Think of λ as a quantity with its own distribution
- ▶ Prior distribution: $p(\lambda) = \text{Gamma}(\lambda, \alpha, \beta)$
- ▶ Bayes' theorem:

$$\begin{aligned}\pi(\lambda, \mathbf{x}) &\propto L(\lambda)p(\lambda) \\ &= \text{Gamma}(\lambda, \alpha + n, \beta + n\bar{x})\end{aligned}$$

Motivating example: Bayesian

- ▶ Think of λ as a quantity with its own distribution
- ▶ Prior distribution: $p(\lambda) = \text{Gamma}(\lambda, \alpha, \beta)$
- ▶ Bayes' theorem:

$$\begin{aligned}\pi(\lambda, \mathbf{x}) &\propto L(\lambda)p(\lambda) \\ &= \text{Gamma}(\lambda, \alpha + n, \beta + n\bar{x})\end{aligned}$$

- ▶ Based on this we can construct an interval I such that we can literally say “ λ has a 95% chance of being in I ”

Quantile regression

General Framework

► OLS regression: $E[Y|\mathbf{X}] = \mathbf{X}^T \boldsymbol{\beta}$

General Framework

- ▶ OLS regression: $E[Y|\mathbf{X}] = \mathbf{X}^T \boldsymbol{\beta}$
- ▶ Quantile regression: $q_p(Y|\mathbf{X}) = \mathbf{X}^T \boldsymbol{\beta}(p)$

General Framework

- ▶ OLS regression: $E[Y|\mathbf{X}] = \mathbf{X}^T \boldsymbol{\beta}$
- ▶ Quantile regression: $q_p(Y|\mathbf{X}) = \mathbf{X}^T \boldsymbol{\beta}(p)$
- ▶ Frequentist approach:

$$\hat{\boldsymbol{\beta}}(p) = \operatorname{argmin}_{\boldsymbol{\beta}} \sum_i \rho_p(y_i - \mathbf{x}_i^T \boldsymbol{\beta})$$

where $\rho_p(u) = u(p - \mathbf{1}_{\{u < 0\}})$

Bayesian quantile regression with AL

- It can be shown that the previous minimization problem is equivalent to maximizing the likelihood associated to the distribution

$$f_p^{\text{AL}}(u) = p(1 - p) \exp(-\rho_p(u))$$

Bayesian quantile regression with AL

- ▶ It can be shown that the previous minimization problem is equivalent to maximizing the likelihood associated to the distribution

$$f_p^{\text{AL}}(u) = p(1 - p) \exp(-\rho_p(u))$$

- ▶ More generally, we can add a location and a scale parameter:

$$f_p^{\text{AL}}(u|\mu, \sigma) = p(1 - p) \exp(-\rho_p((u - \mu)/\sigma))$$

this distribution is called *asymmetric Laplace* [Yu and Moyeed, 2001]

Bayesian quantile regression with AL

- ▶ It can be shown that the previous minimization problem is equivalent to maximizing the likelihood associated to the distribution

$$f_p^{\text{AL}}(u) = p(1 - p) \exp(-\rho_p(u))$$

- ▶ More generally, we can add a location and a scale parameter:

$$f_p^{\text{AL}}(u|\mu, \sigma) = p(1 - p) \exp(-\rho_p((u - \mu)/\sigma))$$

this distribution is called *asymmetric Laplace* [Yu and Moyeed, 2001]

- ▶ Prior: $p(\beta) \propto 1$ (improper)

Bayesian quantile regression with AL

- ▶ It can be shown that the previous minimization problem is equivalent to maximizing the likelihood associated to the distribution

$$f_p^{\text{AL}}(u) = p(1 - p) \exp(-\rho_p(u))$$

- ▶ More generally, we can add a location and a scale parameter:

$$f_p^{\text{AL}}(u|\mu, \sigma) = p(1 - p) \exp(-\rho_p((u - \mu)/\sigma))$$

this distribution is called *asymmetric Laplace* [Yu and Moyeed, 2001]

- ▶ Prior: $p(\beta) \propto 1$ (improper)
- ▶ Theorem: This leads to a proper posterior distribution

Flexible quantile regression with GAL

- ▶ Problems: AL is very rigid

Flexible quantile regression with GAL

- ▶ Problems: AL is very rigid
 - ▶ Skewness is determined by p , in particular it is symmetric at $p = .5$

Flexible quantile regression with GAL

- ▶ Problems: AL is very rigid
 - ▶ Skewness is determined by p , in particular it is symmetric at $p = .5$
 - ▶ Mode is always at 0

Flexible quantile regression with GAL

- ▶ Problems: AL is very rigid
 - ▶ Skewness is determined by p , in particular it is symmetric at $p = .5$
 - ▶ Mode is always at 0
- ▶ Solution [Yan and Kottas, 2017]:

Flexible quantile regression with GAL

- ▶ Problems: AL is very rigid
 - ▶ Skewness is determined by p , in particular it is symmetric at $p = .5$
 - ▶ Mode is always at 0
- ▶ Solution [Yan and Kottas, 2017]:
 - ▶ We can write

$$f_p^{\text{AL}}(y|\mu, \sigma) = \int_{\mathbb{R}} N(y|\mu + \sigma A(p)z, \sigma^2 B(p)z) \text{Exp}(z|1) dz$$

Flexible quantile regression with GAL

- ▶ Problems: AL is very rigid
 - ▶ Skewness is determined by p , in particular it is symmetric at $p = .5$
 - ▶ Mode is always at 0
- ▶ Solution [Yan and Kottas, 2017]:
 - ▶ We can write

$$f_p^{\text{AL}}(y|\mu, \sigma) = \int_{\mathbb{R}} N(y|\mu + \sigma A(p)z, \sigma^2 B(p)z) \text{Exp}(z|1) dz$$

- ▶ So, let us add a parameter

$$\begin{aligned} f_p^{\text{GAL}}(y|\mu, \sigma) \\ = \int_{\mathbb{R}} N(y|\mu + \sigma \alpha s + \sigma A(p)z, \sigma^2 B(p)z) \text{Exp}(z|1) N^+(s|0, 1) dz ds \end{aligned}$$

Parameter Estimation

Overview of Stan

- ▶ Likelyhood above is very complicated, so sampling from the posterior distribution of parameters becomes an issue

Overview of Stan

- ▶ Likelihood above is very complicated, so sampling from the posterior distribution of parameters becomes an issue
- ▶ Usually, we deal with this using Markov Chain Monte Carlo (MCMC) techniques

Overview of Stan

- ▶ Likelyhood above is very complicated, so sampling from the posterior distribution of parameters becomes an issue
- ▶ Usually, we deal with this using Markov Chain Monte Carlo (MCMC) techniques
- ▶ Coding MCMCs by hand is cumbersome. It can also lead to convergence issues

Overview of Stan

- ▶ Likelyhood above is very complicated, so sampling from the posterior distribution of parameters becomes an issue
- ▶ Usually, we deal with this using Markov Chain Monte Carlo (MCMC) techniques
- ▶ Coding MCMCs by hand is cumbersome. It can also lead to convergence issues
- ▶ Software packages like BUGS and JAGS can automatically code MCMCs based on model structure

Overview of Stan

- ▶ Likelyhood above is very complicated, so sampling from the posterior distribution of parameters becomes an issue
- ▶ Usually, we deal with this using Markov Chain Monte Carlo (MCMC) techniques
- ▶ Coding MCMCs by hand is cumbersome. It can also lead to convergence issues
- ▶ Software packages like BUGS and JAGS can automatically code MCMCs based on model structure
- ▶ Stan [Stan development team, 2012] improves on those packages by implementing HMC and NUTS

Overview of BRMS

- ▶ Stan is low level: it's a programming language on it's own, built on top of C++

Overview of BRMS

- ▶ Stan is low level: it's a programming language on it's own, built on top of C++
- ▶ BRMS [Buckner, 2017] is a high level interface for Stan on R

Overview of BRMS

- ▶ Stan is low level: it's a programming language on it's own, built on top of C++
- ▶ BRMS [Buckner, 2017] is a high level interface for Stan on R
 - ▶ It uses the same formula syntax as `lme4`

Overview of BRMS

- ▶ Stan is low level: it's a programming language on it's own, built on top of C++
- ▶ BRMS [Buckner, 2017] is a high level interface for Stan on R
 - ▶ It uses the same formula syntax as `lme4`
 - ▶ Under the hood, it automatically generates and compiles Stan code, which we can retrieve