




Machine Learning

CS60050

Introduction




Learning From Data

Yaser S. Abu-Mostafa
California Institute of Technology

Lecture 2: **Is Learning Feasible?**



Sponsored by Caltech's Provost Office, E&AS Division, and IST • Thursday, April 5, 2012



Feasibility of learning - Outline

- Probability to the rescue
- Connection to learning
- Connection to *real* learning
- A dilemma and a solution


A related experiment

- Consider a 'bin' with red and green marbles.

$$\mathbb{P}[\text{picking a red marble}] = \mu$$

$$\mathbb{P}[\text{picking a green marble}] = 1 - \mu$$

- The value of μ is unknown to us.
- We pick N marbles independently.
- The fraction of red marbles in sample = ν



Does ν say anything about μ ?


No!

Sample can be mostly green while bin is mostly red.

Yes!

Sample frequency ν is likely close to bin frequency μ .

possible versus probable



What does ν say about μ ?

In a big sample (large N), ν is probably close to μ (within ϵ).

Formally,


$$\mathbb{P} [|\nu - \mu| > \epsilon] \leq 2e^{-2\epsilon^2 N}$$

This is called **Hoeffding's Inequality**.

In other words, the statement " $\mu = \nu$ " is P.A.C.

$$\mathbb{P} [|\nu - \mu| > \epsilon] \leq 2e^{-2\epsilon^2 N}$$

- Valid for all N and ϵ
- Bound does not depend on μ
- Tradeoff: N , ϵ , and the bound.
- $\nu \approx \mu \implies \mu \approx \nu$ ☺



Connection to learning


Bin: The unknown is a number μ

Learning: The unknown is a function $f : \mathcal{X} \rightarrow \mathcal{Y}$

Each marble \bullet is a point $\mathbf{x} \in \mathcal{X}$



● : Hypothesis got it **right** $h(\mathbf{x}) = f(\mathbf{x})$

● : Hypothesis got it **wrong** $h(\mathbf{x}) \neq f(\mathbf{x})$



Back to the learning diagram

The bin analogy:



Are we done?


Not so fast! h is fixed.

For this h , ν generalizes to μ .

‘**verification**’ of h , not **learning**


No guarantee ν will be small.

We need to **choose** from multiple h 's.



Multiple bins

Generalizing the bin model to more than one hypothesis:



Notation for learning


Both μ and ν depend on which hypothesis h

ν is 'in sample' denoted by $E_{\text{in}}(h)$


μ is 'out of sample' denoted by $E_{\text{out}}(h)$

The Hoeffding inequality becomes:

$$\mathbb{P} [|E_{\text{in}}(h) - E_{\text{out}}(h)| > \epsilon] \leq 2e^{-2\epsilon^2 N}$$








Notation with multiple bins



Are we done already? 😊

Not so fast!! Hoeffding doesn't apply to multiple bins.

What?



Coin analogy


Question: If you toss a fair coin 10 times, what is the probability that you will get 10 heads?

Answer: $\approx 0.1\%$

Question: If you toss 1000 fair coins 10 times each, what is the probability that some coin will get 10 heads?

Answer: $\approx 63\%$

From coins to learning



A simple solution

$$\begin{aligned}\mathbb{P}[|E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon] &\leq \mathbb{P}[|E_{\text{in}}(h_1) - E_{\text{out}}(h_1)| > \epsilon \\ &\quad \textbf{or } |E_{\text{in}}(h_2) - E_{\text{out}}(h_2)| > \epsilon \\ &\quad \dots \\ &\quad \textbf{or } |E_{\text{in}}(h_M) - E_{\text{out}}(h_M)| > \epsilon] \\ &\leq \sum_{m=1}^M \mathbb{P}[|E_{\text{in}}(h_m) - E_{\text{out}}(h_m)| > \epsilon]\end{aligned}$$

The final verdict

$$\begin{aligned}\mathbb{P}[|E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon] &\leq \sum_{m=1}^M \mathbb{P}[|E_{\text{in}}(h_m) - E_{\text{out}}(h_m)| > \epsilon] \\ &\leq \sum_{m=1}^M 2e^{-2\epsilon^2 N}\end{aligned}$$

$$\mathbb{P}[|E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon] \leq 2Me^{-2\epsilon^2 N}$$

Thank You!

