

# 2015 Undiagnosed Estimates for WA State

*Jeanette Birnbaum*

*09 November, 2016*

## Contents

<b>1</b>	<b>Summary of Results</b>	<b>2</b>
1.1	2015 . . . . .	2
1.2	2014 . . . . .	2
<b>2</b>	<b>Diagnoses</b>	<b>3</b>
2.1	Analytic sample . . . . .	3
2.2	Diagnoses over time by MSM/Area subgroups . . . . .	3
<b>3</b>	<b>Subgroup Sizes and Testing Histories</b>	<b>3</b>
<b>4</b>	<b>Time from Infection to Diagnosis (TID)</b>	<b>4</b>
<b>5</b>	<b>Incidence and undiagnosed counts</b>	<b>5</b>
<b>6</b>	<b>Undiagnosed fractions</b>	<b>6</b>

# 1 Summary of Results

The undiagnosed fraction declined slightly overall, and in every group. However, note that the number of both undiagnosed and total cases rose slightly in all groups.

## 1.1 2015

Table 1: In both tables, the MSM and KC results are separate marginal results, i.e. they are two different ways of dividing the statewide results.

Group	PLWHA	Undiagnosed Cases	True Prevalence	Undiagnosed Fraction (%)
WA State	11985	1360.0	13345.0	10.2
MSM	9071	648.2	9719.2	6.7
non-MSM	2914	711.7	3625.7	19.6
Inside KC	6661	656.5	7317.5	9.0
Outside KC	5324	703.4	6027.4	11.7

## 1.2 2014


Group	PLWHA	Undiagnosed Cases	True Prevalence	Undiagnosed Fraction (%)
WA State	11637	1369.0	13006.0	10.5
MSM	8803	650.5	9453.5	6.9
non-MSM	2834	718.9	3552.9	20.2
Inside KC	6567	678.5	7245.5	9.4
Outside KC	5070	691.0	5761.0	12.0

## 2 Diagnoses

### 2.1 Analytic sample

Analytic data set has 5596 cases for 2005-2015. From the original file provided by Jason, we excluded 14776 cases based on year (excluding diagnoses prior to 2005) restrictions, and an additional 79 cases who were 16 or younger and had no observed date of LNT.

### 2.2 Diagnoses over time by MSM/Area subgroups



## 3 Subgroup Sizes and Testing Histories


Table 3: Column Percent shows the composition of the total sample. The Percent Yes, Percent No and Percent Missing columns indicate the row percents of the three possible testing history statuses within each subgroup

King County	Mode	N	Column Percent	Percent Yes	Percent No	Percent Missing
All	All	5596	100	46	12	42
Inside KC	MSM	2285	41	69	8	23
Inside KC	non-MSM	869	16	29	24	47
Outside KC	MSM	1415	25	38	11	50
Outside KC	non-MSM	1027	18	21	13	66

As we would expect, the presence of observed LNTs is higher for MSM than non-MSM and inside KC vs outside KC. MSM outside of KC have much lower observed LNTs (only 25%) than MSM inside KC (41%). This means that the outside-KC estimates, regardless of mode subgroup, are based on low levels of observed LNTs. This makes them particularly dependent on our ‘missing at random’ assumption.

## 4 Time from Infection to Diagnosis (TID)

The plots below show the TID curves for the four MSM/Area subgroups. BC=Base Case and UB=Upper Bound. For each subgroup, the upper plot panel is the probability of diagnosis curve (the pdf) and the lower plot panel is the survivor curve (1-cdf). The lower plot panel’s y-axis thus indicates the fraction remaining undiagnosed at a given number of years since infection (x-axis).



## 5 Incidence and undiagnosed counts

Figure 1 displays time series plots of KC cases, comparing 'Inside KC' and 'Outside KC' scenarios. The figure is organized into two columns, each containing two vertically stacked plots. The top row of plots shows 'Diagnoses and Incidence' (Counts vs. Time), and the bottom row shows 'Undiagnosed Cases' (Counts vs. Time). The x-axis for all plots is 'Time' (Year), ranging from 2005 to 2015. The y-axis for the top row is 'Counts' (0 to 100 for Inside KC, 0 to 80 for Outside KC). The y-axis for the bottom row is 'Counts' (0 to 1500 for both). The legend indicates three data series: '# Diagnosed' (black crosses), 'Base Case' (blue line), and 'Upper Bound' (orange line).

**Inside KC (Left Column):**

- Diagnoses and Incidence (Top Plot):** Shows a general downward trend in diagnoses and incidence from approximately 75 counts in 2005 to around 55 counts in 2015. The base case (blue line) and upper bound (orange line) are closely aligned, with the upper bound slightly higher.
- Undiagnosed Cases (Bottom Plot):** Shows a steady decline in undiagnosed cases from approximately 1550 in 2005 to around 1250 in 2015. The base case (blue line) and upper bound (orange line) are closely aligned, with the upper bound slightly higher.

**Outside KC (Right Column):**

- Diagnoses and Incidence (Top Plot):** Shows a general downward trend in diagnoses and incidence from approximately 60 counts in 2005 to around 45 counts in 2015. The base case (blue line) and upper bound (orange line) are closely aligned, with the upper bound slightly higher.
- Undiagnosed Cases (Bottom Plot):** Shows a steady decline in undiagnosed cases from approximately 1550 in 2005 to around 1250 in 2015. The base case (blue line) and upper bound (orange line) are closely aligned, with the upper bound slightly higher.

5

outside KC than inside KC.

## 6 Undiagnosed fractions

These plots show the total number of PLWH (x-axis) broken down by diagnosed and undiagnosed (colors) for the Base Case versus Upper Bound estimates (panels). Undiagnosed fractions are indicated by the percent labels. There is clearly a trend of greater diagnosed PLWH over 2010-2015. If this is a function of reporting rather than a real increase, it may be masking true trends in the undiagnosed fraction as well as true differences between inside and outside KC. Again, apologies for different x-scales on the inside versus outside KC plots.

