

**Demonstrating a Standardized IBM Methodology
using Chinook Salmon PSAT Data from the Gulf
of Alaska**

(Proposal)

Marcel Gietzmann-Sanders

2024

Dr. Andrew Seitz

Dr. Curry Cunningham

Michael Courtney, M.S.

College of Fisheries and Ocean Sciences
University of Alaska Fairbanks

1 Purpose

Vulnerability of fish stocks to various human and environmental impacts is tied to the spatio-temporal distribution and movement of those stocks. Therefore understanding these patterns is important to improving management practice today. Central to understanding these patterns is the creation of behavior models that can be used to test movement hypotheses and predict future and current movement behavior. Yet, as of today, there is no common methodology for building and using such models. The purpose of this study is to provide such a common methodology that takes advantage of modern computing solutions such as machine learning and the cloud. It is hoped that by providing such a strategy that (1) the barrier to producing movement models will be greatly reduced and (2) a common language for discussing, evaluating, and using these models will arise.

2 Navigating This Document

This document is organized in order to allow you, the reader, to read as little or as much as you need/want.

It begins with the **objectives** so that you immediately know exactly what this proposal is hoping to achieve and whether this is of interest to you.

Then it dives into how we will go about demonstrating success in those objectives. This will inform you on the specific **outcomes** to be expected.

Finally there is a **timeline** provided to give a sense of when these outcomes can be expected.

If these sections are compelling enough there is no need to read any further.

However if you are interested in the rationales behind these objectives or diving into the related context, the **Rationale** and **Related Context** sections are for you.

Contents

| | |
|---|-----------|
| 1 Purpose | 2 |
| 2 Navigating This Document | 3 |
| 3 Objectives | 5 |
| 4 Demonstrations | 6 |
| 4.1 Chapter 1 | 6 |
| 4.2 Chapter 2 | 7 |
| 5 Timeline | 10 |
| 5.1 Class Work | 10 |
| 5.2 Thesis | 10 |
| 6 Rationale | 11 |
| 6.1 Objective 1 | 11 |
| 6.2 Objective 2 | 15 |
| 6.3 PSAT Data | 18 |
| 6.4 Chinook salmon in the GOA | 19 |
| 7 Related Context | 20 |
| 7.1 Summary | 20 |
| 7.2 Spatial Management | 22 |
| 7.3 Spatial Assessments | 27 |
| 7.4 Modeling Movement | 32 |
| 7.5 Chinook salmon Ecology and Management | 36 |
| 8 Bibliography | 40 |

3 Objectives

Objective 1. *Standardized Decision Models* Movement (and more generally decision) modeling is fundamentally about modeling the selection of outcomes. So, given a set of choices v_k and a set of features n_k describing those choices, we want a standardized method to fit, evaluate, and use models of the form:

$$\psi_k = G(\eta_k)$$

where ψ_k is the "odds for" choice v_k . Such a model should maximize the following objective:

$$\mathcal{L} = \prod_i P'(v_i|\eta_i)$$

where:

$$P'(v_i|\eta_i) = \frac{\psi_i}{\sum_k \psi_k}$$

Objective 2. *Standardized Decision Simulations* Given an individual in a specific situation, the decision models described above produce a set of probabilities for each potential next outcome v_k . However there is nothing preventing us from taking each of those outcomes, considering them each a new situation, and applying the models again. Carried out repeatedly we will end up with a probabilistic diffusion of the original individual across its decision space. Furthermore, if we can do this for one individual we can do it for an entire distribution of individuals. In sum, using this diffusion modeling approach we can simulate the movement through a decision space of an entire population.

Our second objective then is to build such a diffusion model. However, we would like to do it in such a way that life history, mortality, and environmental models can be provided as input as well. This diffusion model will then allow us to explore the spatio-temporal behavior of a stock under both nominal conditions as well as under various exploratory management strategies.

4 Demonstrations

Our demonstrations of the above objectives will be split into two chapters.

4.1 Chapter 1

Chapter 1 will be concerned with demonstrating the tooling for creating odds models for spatio-temporal movement behavior. It will pursue two specific modeling examples:

Model 1. *A depth occupancy model for Chinook salmon in the Gulf of Alaska (GOA).*

Choices: *Discrete depth ranges.*

Features: *Environmental data derived either from environmental simulations or from remote sensing such as Google Earth, cohort data about the fish themselves, and temporal features such as month, season, or time of day.*

Data: *PSAT tagging data from the GOA.*

Purpose: *To study the heterogeneity in depth occupancy as a function of cohort, environment, and time.*

Model 2. *A movement model for Chinook salmon in the Gulf of Alaska (GOA).*

Choices: *Spatial grid cell selection as movement. Also if clearly different modes of movement are found (migration vs foraging for example) we'll predict the mode as well.*

Features: *Environmental data derived either from environmental simulations or from remote sensing such as Google Earth, cohort data about the fish themselves, temporal features such as month, season, or time of day, and internal state information about the cohort.*

Data: *PSAT tagging data from the GOA.*

Purpose: *To study the heterogeneity in spatial movement/preference as a function of cohort, environment, and time.*

In building these models our primary purpose is to demonstrate the tooling exists to prudently build movement models. We can frame this goal in terms of a "standard" machine learning pipeline and work-flow.

1. Formatting Data for Odds Modeling
2. Class Balancing
3. Splitting Data into Train, Validation, Test
4. Training and Hyperparameter Tuning Using Standard Neural Net Architectures
5. Diagnoses of Fit and Comparison of Different Models
6. Identifying Sample Groups that are More or Less Fit
7. Exploring Discovered Patterns

We will want to explicitly demonstrate each of these processes, provide guidance on using the tools for them, and use them to build the models listed above.

The outcome will be a package in Python for building, diagnosing, and running movement models. Note that at this point the purpose is to demonstrate how to build movement models for the purpose of building simulations. It will be in the next chapter that specific research outcomes directly relevant to management will manifest themselves.

4.2 Chapter 2

Chapter 2 will be concerned with taking these movement models for Chinook salmon in the Gulf of Alaska and building a diffusion model. We will then demonstrate its capabilities by exploring a series of management questions. It is worth noting that these questions have been chosen based on their facility in demonstrating the aspects of the simulation as opposed to their priority to current Chinook salmon management in the GOA.

The overview of the components of this diffusion model are outlined in Fig. 1.

The specific questions we would like to answer with this simulation are:




Figure 1: Simulation Outline

Boxes with rounded edges would be inputs specified by the user. The hexagons are functions provided by the diffusion model and the trapezoids are data generated during the main loop. It is important to note that in this design we are planning for the most data rich case. However, depending on the application, mortality or environmental data may in fact have quite limited availability.

1. **Understanding Connectivity:** by looking at the vector flows of fish throughout the space we will be able to identify which areas are highly connected and which are more or less separated by environment, cohort, or other factors of fish behavior.
2. **Understanding Vulnerability:** by especially looking at changes in depth occupancy and preference for habitat we will attempt to quantify the degree of vulnerability to different kinds of activities as a function of space and time. Specific vulnerabilities of interest might include: vulnerability to specific fishing fleets, vulnerability to naval training activities, vulnerability to marine mammals.
3. **Cumulative Mortality:** by being able to follow cohorts through space and time we should be able to show how different fishing pressure

scenarios might affect the cumulative fishing mortality. Note that the data for specific fleets may be unavailable but that our interest here is to demonstrate that given data we can make these kinds of predictions.

4. **Protected Areas/Times:** Of those fishing pressure scenarios, one of especial interest is modeling Marine Protected Areas (MPAs) or other kinds of closures in space and time. Given the motility of Chinook, how must effective protected areas/times be designed?
5. **Uncertainty:** we will show that such a simulation can be run at scale, bootstrapping can be applied, and therefore uncertainty in simulation outcomes quantified.

To give a specific example of what this might look like in practice, if we were interested in understanding how to protect one specific genetic cohort by creating fishing closures we would:

The outcome will be a package in Python for deploying, running, and analyzing diffusion models.

5 Timeline

5.1 Class Work

1. **Fall 2023:** (University of Florida) FAS 6355c Fisheries Management, FAS 6337c Population Dynamics
2. **Spring 2024:** FISH F621 Estimation of Fish Abundance, FISH F427 Ichthyology
3. **Fall 2024:** STAT F641 Bayesian Statistics, FISH F692 Seminar
4. **Spring 2025:** FISH F631 Data Analysis in Community Ecology, FISH 699 Thesis
5. **Fall 2025:** STAT F651 Statistical Theory I, FISH 699 Thesis
6. **Spring 2026:** FISH F626 Behavioral Ecology of Fishes, STAT F605 Spatial Statistics

5.2 Thesis

1. **Spring 2024:** Depth and Movement Models v0, Thesis Exploration
2. **Fall 2024:** Graduate Study Plan, Self Evaluation, Thesis Proposal, Depth and Movement Models v1
3. **Spring 2025:** Thesis Proposal Defense, Simulation v1
4. **Fall 2025:** Self Evaluation, Write Chapters, Publish Package(s)
5. **Spring 2026:** Thesis Defense

6 Rationale

6.1 Objective 1

6.1.1 Value of Machine Learning in Model Building

One of the functions of science is to build models of the world around us, specifically to look for functions of the form:

$$v = F(\theta)$$

where θ is some vector of information, v is a particular outcome given that information, and F is the model in question. This we might term a *deterministic model* because for any specific θ there is one single outcome v .

However, life is rarely so kind to us. From quantum physics to fisheries science, the world is replete with examples where the same information does not always lead to the same conclusion. Sometimes that is because we do not possess all of the relevant information but sometimes things are just truly random. As such, the model above will in some sense be lying to us in its deterministic certainty - while we will always predict v , v is not always what we will receive.

Instead suppose that for a specific θ there exist a set of possible outcomes $V(\theta) = \{v\}$. We can define a *stochastic model* as:

$$P(v|\theta) = F(v, \theta)$$

where now the model is predicting the probability of v given θ instead of just predicting v itself.

Now what is especially interesting about models of this form is the fact that in some sense for any combination of V and θ they always exist. Whereas to get a deterministic model we need to know precisely what kind of information (θ) we need to make a deterministic prediction, we can always make a stochastic one, even in the presence of no information. The model can always give us a correct answer, it's just a question of how useful that answer is. If you base it off of better data it will become more useful.

Unfortunately, unless we ourselves have built the part of the world we're studying, F is not known to us. Instead our models are really just proposals \hat{F} on what F could be. This presents us with an immediate problem, how do we evaluate any specific \hat{F} if we don't know F itself? Well suppose we've captured a whole series of pairs of θ_i and v_i where i allows us to index the pair in question. Given we know \hat{F} we can directly ask what the likelihood of the data we have collected is, given \hat{F} :

$$\mathcal{L} = \prod_i \hat{F}(v_i, \theta_i)$$

or equivalently (and more easily computed)

$$\ln \mathcal{L} = \sum_i \ln \hat{F}(v_i, \theta_i)$$

\hat{F} 's with higher $\ln \mathcal{L}$ (log likelihood) are better fits to the given data and as we have more and more comprehensive data those data better and better represent F (see Appendix 1).

Therefore finding the "true" F can be summarized with two steps:

1. Collect large amounts of comprehensive data.
2. Find the \hat{F} that maximizes the likelihood of that data.

All of this, however, has been conditional on the form of θ having been chosen. Obviously, the information we choose to collect and build our model with has an impact on how good the model is from a purely predictive perspective. So how can we compare the predictive power of different θ ? Well as we find θ that are more predictive the corresponding F s will have greater and greater likelihoods \mathcal{L} , given the data. But this presents us with a chicken and the egg style problem - to know the value of θ we must know F , but to know F we must have fit \hat{F} which requires a reasonable amount of data to have been collected. I.e., we cannot know in advance if the data we are collecting is going to be useful or not. Instead anyone doing modeling finds themselves in the cycle (Fig. 2).

Each step in this cycle is highly non-trivial. However special interest should be paid to step 3 (finding an \hat{F} that best approximates F).

The standard method here has been to propose a family of possible F which are defined by a set of parameters μ . Then using some choice of likelihood maximization method the set of μ are searched for that maximize the likelihood of the data \mathcal{L} . While this method has borne considerable fruit it still requires a great deal of effort on the side of the scientist to envision various hypotheses on what families of functions would make sense, coding those up, fitting them, and then evaluating them post-fit. Furthermore if the true form of F is not contained in those set of hypotheses tested then F is never found. Lots of effort without any kind of guarantee that F will be discovered.




Figure 2: Modeling Cycle

The four major steps in building good models. Note that while one must always start at (1) this process is indeed a cycle.

The field of Machine Learning (ML) (specifically probabilistic machine learning) on the other hand has been specifically occupied with finding ways to maximize objectives without having to assume much, if anything, about the functional form of the prediction function. Therefore it provides an opportunity to lessen the toil and uncertainty in step 3 and allow scientists to focus more of the efforts on the other steps (and thereby accelerate the whole cycle).

6.1.2 Issues with Probabilistic Machine Learning when Applied to Behavior

Probabilistic machine classification (Oliver Durr, 2020) of the kind we were pointing to in the last section has two pitfalls when it comes to modeling behavior.

The first comes from the fact that such models predict on the same set of outcomes each time, regardless of θ . For example, if we were trying to predict which drink a person will buy at their local cafe we would not only have to include everything currently on the menu but also everything that

could be on the menu across all time points we are interested in predicting. If suddenly a new menu item appears that we have not trained on, our probabilistic classifier will be at a loss.

The second issue comes from the "curse of dimensionality." Basically this is an issue where as the dimensionality of θ increases, the amount of data required to find real relationships in that data increases exponentially. This is a particular problem for behavioral modeling because we in all likelihood need to include data on each of the possible decisions before us. So if I can choose between five different drinks I need to provide features on each of those five drinks. This leads to very high dimensional spaces very quickly.

We can get around both of these problems by introducing what we will term an "odds model." Specifically if we imagine θ now as a variable length vector composed of vectors η_k for each of our possible outcomes v_k then the odds model is given by the function:

$$\psi_k = G(\eta_k)$$

where we now say that:

$$P'(v_k|\theta) = \frac{\psi_k}{\sum_j \psi_j}$$

We can see why we chose the name "odds model" as the ψ_k are simply the "odds for" outcome v_k .

Now note it is not necessarily true that:

$$P'(v_k|\eta_k) = P(v_k|\theta)$$

because it is possible that information in the other η_j informs the probability of v_k . However it should be possible to include such information in the η_k if necessary.

By making this sacrifice, however, we have achieved two things:

1. So long as we can provide a η_k for a newly seen outcome we can predict on it. Therefore we need not have a fixed set of possible outcomes (or a fixed number of such outcomes per θ).
2. Compared to the corresponding probabilistic machine learning classifier, if our total number of outcomes were $|V|$ we've now reduced the dimensionality of our feature space by $|V|$.

And this means that unless we have to pass large amounts of information between the η_k the amount of data we need to collect to fit our model has decreased tremendously. And in cases where data is hard to come by (like animal behavior), any reduction in data requirements is of the utmost importance.

So finally, to model behavior we'd like to fit an odds model of the form:

$$\psi_k = G(\eta_k)$$

that maximizes:

$$\mathcal{L} = \prod_i P'(v_i|\eta_i)$$

where

$$P'(v_i|\eta_i) = \frac{\psi_i}{\sum_k \psi_k}$$

6.2 Objective 2

From studies that started with Beverton and Holt's work on recruitment in fisheries, we know, with a reasonable degree of certainty, that as a general rule fish stocks operate according to an asymptotic relationship between spawning biomass and recruits (Carl Walters, 2005) (Fig. 3). What we see is a clear pattern in recruitment verses spawners for low spawners and then eventually we reach a point where this curve asymptotes and large changes to the number of spawners creates little to no change (especially when considering the actual variability in this data) in the number of recruits (there is some variation on this theme depending on the precise stock recruitment model used). Given the real variability in nature, this pattern looks more like a scatter shot of points but plenty of studies over several species have shown this overall pattern in the mean borne out (Carl Walters, 2005).

The implication for fisheries management is pretty clear - sustainable fishing is in large part driven by just maintaining the spawning stock biomass (SSB) above that knee in the curve (Carl Walters, 2005) (King, 2007). Indeed this is well noted as the basis of why fisheries *can* exist in the first place. If taking spawners always led to significant decreases in recruits, we'd quickly fish a stock out of existence. It is only through the compensation we get




Figure 3: Recruitment

on the asymptote that sustainable fisheries are possible. However, two very simple observations of fish biology will show how actually managing SSB can be made difficult.

The first, and simplest, is age structure. Obviously we're talking here about maintaining a specific level of spawning stock biomass. But today's juveniles are tomorrow's SSB and as fish grow they typically become more fecund. This means that in order to maintain our SSB over time we have to be careful about how we target each age group. This gets even more complicated for fish like Pacific salmon (*Oncorhynchus*) or many species of groupers (*Epinephelinae*) where life history is intimately entwined with spawning. The salmon only breed once (NOAA, 2024a) and some species of groupers are protogynous hermaphrodites (NOAA, 2024b) meaning that if you target a specific age group you are effectively targeting a specific sex. Clearly understanding and managing our age class selectivity matters when it comes to maintaining a healthy SSB for years to come.

The other observation is that many species of fishes actually form multiple sub stocks within what we may consider a single fishery (or sometimes many fisheries distributed across different countries) (Kai Lorenzen, 2010) (PFMC, 2024) (Prince, 2010). Pacific salmon are once again a good example in that they home to specific streams and only breed with other salmon in those streams and many other species of fish form fish spawning aggregations as well (Brad Erisman, 2017). Research also indicates that even fish like Atlantic herring likely form distinct genetic stocks due to differences in homing during breeding and school behavior (Michael Armstrong, 2001).

What does this mean? It means that if we trust our Beverton-Holt style model, instead of having a single SSB to manage we really have several SSBs - one for each sub stock. Obviously, if we set one global piece of management like an Allowable Biological Catch (ABC) that makes sense for a single SSB but then have biased fishing mortality where one sub stock is getting fished out far more than the others we may drop below the SSB for that one stock without actually exceeding our expected ABC for the whole fishery. And this can certainly happen as different sub stocks will often exhibit quite different spatial behaviors in movement (S. Tucker, 2011) (Punt, 2019) (Cadrin, 2020).

All of this to say that understanding heterogeneity in selectivity by things like age class and sub stock is of the utmost importance for managing a fishery and truly ensuring an appropriate amount of SSB remains year over year. It is also worth noting that this knowledge around changes in selectivity in space and time is not only useful for considerations around SSB but also for other management issues such as bycatch avoidance and growth overfishing.

What kind of tooling do we need in order to address these issues and test out ideas for fisheries management? Clearly we need some kind of reliable simulation.

Suppose we built models of spatio-temporal behavior (specifically movement) as expressed in Objective 1. These models we know can be fed information that captures both environmental variables but also factors we'd like to categorize by in order to understand heterogeneity in selectivity. Such variables could include features like the age and genetic cohorts that we just described. Then we could take some baseline assumptions about abundance in space at a given time (say providing recruits at spawning grounds or taking a spatio-temporal assessment as a starting point) and use our behavioral models to simulate time-steps forward. As we do, we could account for natural mortality as well to complete the "environment" portion of our

simulation.

It is important to note that while this may seem like a Lagrangian approach to simulation it actually falls more into the Eulerian camp as, in order to reduce our computational costs, we'd simply have cohorts as a function of space. Remember that our model is only taking the categories that separate one group from the next thereby allowing us to model cohorts instead of single individuals. Furthermore instead of modeling the specific singular behaviors our behavior model as outlined in Objective 1 allows us to understand the "many world" probabilistic outcomes of behavior. We are not following a single fish but instead are watching as that cohort diffuses across its decision space. Therefore we'll term this simulation a diffusion model.

Into our diffusion model we can now incorporate whatever management policy we like as a spatio-temporal specification of fishing mortality - which can now be modulated by our understanding of, say, gear selectivity as a function of space and time. So if we want to implement a marine protected area we can take a baseline fishing pattern and exclude specific areas in space. If we want to see what would happen if we redistribute the fleet we can as well. If we just want to understand the selectivity biases already present we can simulate fleet patterns as they occur today.

The point is that with such a simulation we are able to test different hypotheses of management on data driven understanding of fish behavior.

6.3 PSAT Data

Pop-up Satellite Archival Tags (PSATs) are a great way of collecting behavior data (Computers, 2024). They are defined by three key properties:

1. **Archival:** Transmitting data is only possible if animals come to the surface. Therefore these tags archive collected data for later transmission.
2. **Pop Up:** These tags are programmed to release from a fish at a specific time or under specific conditions to facilitate retrieval.
3. **Satellite:** On pop-up the tags then broadcast data (usually summarized) to Argos satellites which is then relayed to a database for retrieval. Such tags usually also broadcast some information that can be used to deduce position as well which allows for direct retrieval and access to the full archived dataset.

The normal lifecycle then is to catch an animal, attach the tag, allow the animal to go about its life all while collecting data, have the tag pop-up, transfer data to satellite or retrieve the tag, and then clean and analyze the data.

For our purposes these tags critically capture:

1. Temperature
2. Depth
3. Light Levels

The latter two are then used along with tag release location to predict animal movement (latitude and longitude) using a proprietary statistical model (Computers, 2024).

We will look at movements on a day by day basis and depths at the transmitted resolution (2-5 minutes). For those tags retrieved, a much higher resolution is available (1-3 seconds) however most data is only guaranteed from transmission and not from retrieval (Computers, 2024).

Clearly then this kind of data suits our modeling and simulation purposes well as we get behavior outcomes directly.

6.4 Chinook salmon in the GOA

At this point in time the main interest here is that Chinook salmon present an interesting model species for testing out the efficacy of these models and simulations. Chinook salmon are highly motile and undergo clear migration behaviors (S. Tucker, 2011) (Joseph Langan, 2024). Therefore they should provide a good challenge to attempts at behavioral modeling and simulation. It is the author's hope that patterns useful to Chinook salmon management issues may arise from this research but it is not a necessity as the intention here is to develop tools that justify additional data collection for species for which spatial management is of interest.

7 Related Context

7.1 Summary

The sum outcome of the work proposed is the creation of a simulation that can be used to explore, propose, and (in simulation) test management options. The advantages of having such a simulation are many (as is described in the following situations) and therefore it comes as no surprise that the notion of using a simulation to drive understanding of management options is in no way novel. Indeed as is demonstrated in the following sections many different kinds of simulation (or analogs) already exist. Assessment paradigms are a form of simulation, movement models based on more traditional tagging methods can be used in simulation, and individual based models of several sorts have lent themselves to simulations used to assess changes to environment or management as well. Therefore the question is - what advantages come from the proposed direction?

1. **Isolating the Study of Movement (§7.3).** In machine learning there is a well known phenomenon where features that are even mildly correlated with one another can compensate for one another in model fitting. A simple example would be something like height representing the "feature impact" of sex in a model of human health. Therefore it is always dubious to treat the relationship between a feature and a target as an actual indication of the true relationship. Within the spatial assessment paradigms in use today many different parameters of life history are fit together - movement, spatial heterogeneity in recruitment or mortality, fishing pressure, and so on. Because these are all fit as one model it is possible if not likely that features are "compensating" for one another and obscuring the actual relationships between the underlying processes and the outcomes. The simulation proposed herein fits movement on its own which should lead to a greater confidence in its representation of actual fish movement patterns. It also makes the debugging and improvement of those models much more clear and straightforward.
2. **Generalized Strategy (§7.4).** Individual based models (IBMs) also solve the problem mentioned above as they directly model movement as well. However IBMs are notoriously specific to the problem at hand. Furthermore most IBMs represent a specific hypothesis of behavior that has been coded up and as was mentioned in Section 6.1 building models this way takes an enormous amount of time. The strategy

proposed herein provides a generalized tool set for going from data, to a model, and then from there to a spatial simulation. And this means that researchers can contribute to building up a common tool set instead of having to start from ground zero each round.

3. **Repeated Usefulness.** There are obviously more straightforward ways to answer specific spatial questions than by using the framework proposed here. For example if trying to understand the value of a Marine Protected Area (MPA) one could simply study general abundances changes over the course of a year or apply more traditional tagging methodologies. The purpose of this framework would be to provide a centralized datastore, model, and simulation that could be used across many different problems.
4. **Interpretation and Communication (§7.3, 7.2).** Current standard spatial assessments depend on a movement matrix X that defines how cohorts move between grids in the assessment at each time point. Such a matrix is ultimately an abstract mathematical concept that while useful in assessment makes the communication and interpretation of the underlying movement model difficult. Questions such as - what is driving the movement pattern seen - require actual parametrization of that movement matrix X . The strategy we are proposing provides this and therefore makes communication and feedback from stakeholders of the fishery far easier. Given engagement is such an important part of management this is paramount.

Obviously each of these individual pros are not unique to the strategy proposed herein. IBMs, spatial assessment models, and more ad-hoc approaches all share some number of these advantages. However the strategy proposed here presents an opportunity to gain all of these advantages at once which should make it a useful addition to the tools fisheries science already possesses.

7.2 Spatial Management

Fisheries management is as diverse a subject as the species and contexts to which it is applied (Carl Walters, 2005) (King, 2007). It simultaneously has to deal with creatures from Bluefin Tuna (*Thunnus thynnus*) who make large oceanic migrations around the globe to various clams whose entire life cycles can be summarized within a few square meters (Prince, 2010). There are species that school and species that consume those schools. Some live entirely at sea, some in inland rivers and lakes, and some, like salmon and eels spend time in both. There are species whose lives last the span of a year to others that can live longer than any us human beings.

And to capture this vast array of diversity are similarly diverse strategies and gears. From hooks to gillnets, from small boats to at sea factories, from recreational fishers to roving commercial fleets, and everything in between the variety in strategies that exist to catch fish are numerous (Carl Walters, 2005) (King, 2007).

And then there are the stakeholders of these fisheries and their multitude of objectives. What kinds of fish sell the best? Is selling fish even the point or is subsistence a big concern? Is this a primary source of livelihood or are fishers also fishing for other things, or occupied with entirely different forms of work during other parts of the year? Where do conservation groups fit in? What about the natural predators and prey lodged nearby in the food web? Is the stock in need of rebuilding? What requirements of management are there? (Carl Walters, 2005) (King, 2007)

Fisheries management is a field in which the phrase "the devil is in the details" is incredibly relevant.

As with any such endeavor that can come in so many forms and captures so many different people, interests, and potential problems and opportunities, the best creative solutions come when everyone is working together to build and understand those solutions (Prince, 2010) (Carl Walters, 2005). And of utmost importance here is the inclusion of those who actually go out and do the fishing. Fisheries management that creates a sense of inclusion and involvement with the fishing community is the best kind of management due to a number of things:

1. Fishers often have incredible amounts of insight into the species they are targeting
2. Fishers have an interest in the long term sustainability of their work and therefore often have ideas on how to achieve this

3. There are plenty of examples of fishers helping to self enforce policies if they agree with said policy
4. Fishers are key when it comes to collecting the kinds of data required to do management
5. Fishers are one of the primary stakeholders impacted and therefore must have a say in what is going to be done

In general, creating cooperation improves the overall solution whereas exclusion simply creates conflict that works against any kind of management. Therefore the first priority on our list when it comes to doing good management (spatial or otherwise) has to be creating engagement.

To this end simulations and models of the kind that we are proposing should be helpful.

7.2.1 Driving Engagement

First, it is clear that engagement is best facilitated by presentations of knowledge or data that are visual and relevant to the ways in which people normally think about their world (Prince, 2010) (Kai Lorenzen, 2010). By having simulations that can simply be shown and laid down on a map that places the spatio-temporal modeling in familiar context, one should be able to make objections or confirmation much easier. Rather than having to attempt to condense down loads of parameters, summary statistics, and the like in a model, one can simply show a map and folks can point at things that line up or are in disagreement with their intuition and experience. In this way mapped out views of the simulation work to be a useful device for consensus building among multiple groups of stakeholders. Not everyone may be familiar with the nuances of a specific kind of model but everyone with an interest can point to a map and understand what's going on.

With the simulation providing an easier to understand jumping off point to create feedback around a model, the models themselves provide a reasonably straightforward method for taking that feedback and working it into the model itself. This is because these models rely only on the information input as opposed to requiring modelers to determine the relationship between the information and the outcomes.

For example, suppose that we have a more standard model that has been hand crafted by a scientist familiar with the species in question. They may

have included specific associations with, say, oceanic currents, temperatures, and primary productivity. But then if someone looking at the simulation points out that they think the species' behavior is also being driven by day/night patterns the scientist then has to go back to the drawing board, figure out how to add this into their overall model, code it up, tune parameters, and then return - all without any real guarantee that whoever provided the feedback is right. And this would be for one piece of feedback. If there is a lot of feedback this only gets more difficult.

Instead, by using machine learning as proposed here, such feedback would amount to playing around with a few day light features, pushing those to the model, retraining using the same automation as for any of the other feature combinations, and then just reporting if that information helped or not. Feedback can be incorporated much more easily, allowing for direct contribution to the model on the part of those who may have no modeling experience but have lots of practical experience that could suggest forms of information that would improve how targeted the model is.

In summary the combination of clear, relevant visuals from the simulation, and ease of feature incorporation model side allow for a nice iterative loop to drive engagement and input from all stakeholders into the underlying spatial understanding of the fishery.

7.2.2 Spatial Management Today

While we've already noted that the creative solutions in fisheries management are as diverse as the people, species, and objectives present, there are some specific kinds of strategies that are used across the world to do spatial management. By and large they are comprised of performing spatial closures, however those closures differ with regards to who they effect and for how long (Elizabeth Selig, 2016) (Alyson Little, 2014).

1. **Marine Protected Areas (MPAs)** - MPAs are usually permanent closures intended to reduce the fishing pressure in specific areas in the hope that those areas can become "sources" of larvae and stock to help build up the surrounding areas or to help scientists understand the "unfished" condition. Sometimes these areas are total no-go zones for all fishing activity, sometimes specific "low pressure" activities are allowed. In general it is clear that the efficacy of MPAs is largely dependent on the spatio-temporal behavior and spatio-temporal aspects

of the species' life cycle as well as the same aspects of behavior and lifestyle of the species' predators and prey. Species that move about dramatically may not be helped by MPAs and species that are "too" sessile may not actually do much to contribute to stocks outside of the MPA itself. Therefore it is clear that a good understanding of movement is extremely important when both thinking about the overall efficacy of an MPA and also where to place them.

2. **Territorial Use Rights for Fisheries (TURFs)** - TURFs delineate a fishery into spatial regions and then give exclusive access rights to specific communities of fishers to each of the defined regions. The intention with exclusivity is to align the incentives of the fishers with that of sustainability and is a way to address "tragedy of the commons" style issues that arise with completely open access. However, as with MPAs, it is clear that understanding the spatio-temporal behavior of the species in question is of the utmost importance here because for species that exhibit considerable movement, small regions would mean that while spatial rights exist those rights don't really have anything to do with the underlying fish.
3. **Temporary Closures** - These kinds of closures are designed to deal with specific issues directly related to the target species (or related bycatch's) ecology. For example seasonal closures in specific areas might exist to protect spawning aggregations or other kinds of aggregations that create a particularly dangerous kind of vulnerability. Or these closures might protect against vulnerability by another species to fishing activities such as preventing hits on whales or bycatch of an endangered species. Here again, clear spatio-temporal understanding is necessary for defining and determining the efficacy of these kinds of measures.
4. **Real Time Management** - This is a particularly interesting kind of management as instead of trying to predict where closures are needed, in this strategy metrics from the fishers themselves help dictate where closures are needed temporarily. For example if bycatch avoidance is the goal then catch logs (that are usually reported with some frequency in these cases) are used to determine areas where bycatch abundance has spiked and then the fishery can call for a temporary closure lasting days to months in order to avoid that aggregation. In general this strategy is comprised of measuring some overall catch statistic and then responding to that statistic in real time as needs be. Here models

of the kind we’re proposing are arguably less relevant as prediction is not *required* for this kind of strategy to work. However they could certainly help in both defining the extent of the closures in both space and time and potentially helping fishers avoid such closures in the first place by helping drive understanding of where things like high bycatch may occur.

Unsurprisingly then, for any kind of spatial management tool in use today, spatio-temporal understanding of the stock is key to both definition and efficacy (Elizabeth Selig, 2016) (Alyson Little, 2014). Therefore it seems likely that bringing high resolution spatio-temporal data to the bear on the issue can only help make such techniques more nuanced, precise, and thereby effective.

7.2.3 ”Weather” Forecasting

Finally it is worth noting that the potential in combining this modeling with fishery dependent information is quite interesting on its own. Any forecasting model is always going to have issues predicting farther and farther into the future but fisheries dependent data could provide a means for making corrections to keep the model in line. This would amount to something like weather forecast where for the next few days the forecast is accurate enough to precisely predict specific kinds of movement helpful to fishers themselves.

For example it could help them predict where to go to avoid high levels of bycatch, help optimize their use of fuel and labor when searching for fish, or, depending on the degree of subtlety in the model, might even be able to help them target catches that are more or less valuable on the market itself.

With such direct incentives to fishers this might provide a jumping off point to bring them into helping collect the kinds and quantities of data that would allow for increased precision in these models. With such ”weather forecasting” any such increases would directly benefit them in addition to providing overall benefits to management.

7.3 Spatial Assessments

7.3.1 Assessments in Short

Fisheries assessments are one way in which unknown parameters important to management are estimated. By and large they are comprised of two models - (1) a state model that represents the underlying stock and contains such things as growth, age composition, recruitment, abundance, and so on and (2) an observation model that represents how data such as catch rates, fishery independent indices, and so on arise from the state model (King, 2007). Depending on how the assessment is being done the parameters in the state model will either be estimated independently or left as parameters in the model to be fit. The observation model on the other hand is encoded as a set of distributions dependent on those parameters (and including some parameters of its own) thus allowing for the likelihood of the parameters given the data to be computed (Tim Sippel, 2014). This, in turn, allows for maximum likelihood estimation (MLE) routines to then fit the likely parameters of the assessment.

So, as an example, you might have your state model be comprised of age cohorts that have a specific growth pattern (fit externally) and then you leave as parameters natural mortality and recruitment. Vulnerability might then be modeled to capture a couple different kinds of gear, provide an expected distribution for catches given your underlying abundance by cohort, and then "simulate" a few years. Matching this back to real catch data would then allow you to estimate your natural mortality and recruitment by year (King, 2007) (NOAA, 2024c).

In short the pattern is (1) design a simulation that has your unknown statistics as parameters, (2) specify how data is drawn from the simulation, (3) fit your parameters to your data using MLE. Then those desired statistics are extracted from the assessment (usually with some notion of uncertainty attached to them) and used to inform management.

7.3.2 Features of Spatial Assessments

As has been noted by several authors, most standard assessments have typically assumed in their state model (or simulation) that the stock of interest is bounded by the area under assessment, is well mixed, and constitutes a single unit (Tim Sippel, 2014) (Punt, 2019) (Daniel Goethel, 2023). However from studies of fish behavior and the fact that fish obviously don't

obey national or state defined boundaries, these assumptions are rarely true and indeed simulation studies have shown that assuming them can lead to significant errors and biases (Kai Lorenzen, 2010).

Therefore in recent years there has been work in creating new assessments that are able to capture spatial heterogeneity and movement (Punt, 2019) (Cadrin, 2020). There are effectively two different kinds of heterogeneity that such models have to capture - that in recruitment and that in post recruitment movement and connectivity (or lack thereof). As such these models are usually structured as having multiple sub regions within the overall assessment region and then model the following:

1. Sub-region specific mortality, growth, recruitment, or other sub-population specific parameters may be estimated. The degree to which these are shared between or constrained by other regions is often related to the data available and how flexible the model is allowed to be.
2. Movement among regions at each time step is also modeled, usually as a matrix X such that the elements $X_{i,j}$ signify movement rates from sub-region i to sub-region j .

Different assessment paradigms will have varying degree of resolution in the sub-regions, will estimate certain parameters freely to not at all, leave X free or constrain it on other kinds of assumptions (indeed some assessments don't model movement at all) (Punt, 2019) (Daniel Goethel, 2023).

Ultimately what is bounding the freedom in parametrizing these assessments is the availability of data to actually fit all of these parameters (Daniel Goethel, 2023) - a subject to which we turn next.

7.3.3 Integrated Assessments

A primary issue with spatial assessments is their hunger for data. This makes sense given the combination of the "curse of dimensionality" and the fact that adding per region parameters and across region movements introduces not just new parameters but many multiples of parameters as compared to a non-spatial assessment model.

Therefore a huge step forward in being able to run these kinds of assessments was the development of "integrated assessments" - assessments that can take advantage of pretty much any kind of data collected from the fishery (Tim Sippel, 2014). The process here falls neatly into our earlier description of how an assessment works - for some new kind of data (say acoustic tag data) one must add a new component to our observation

model that describes the distribution that would generate our acoustic tag data. With this in hand we’ve connected a new data type to our underlying parameters and can therefore use that data to help fit our parameters.

There are of course issues that arise in terms of how to weight one data type against another when doing MLE (Tim Sippel, 2014) (Punt, 2019) but by and large integrated assessments are a boon because they increase the data available to fit assessments and have led to several tag integrated assessments that include electronic tag data in addition to more standard mark and recapture tag data.

7.3.4 Validating Spatial Assessments

With the recency of their development and overall complexity it has been noted that diagnosing goodness of fit and relevance of spatial assessments is an art that is still in development (although several papers exist that give guidance on best practices in choosing and using spatial assessment paradigms (Daniel Goethel, 2023) (Punt, 2019)).

Every assessment paradigm is of course a simplification of reality but what’s important to know is whether those simplifications are obscuring important structure of the stock and sub-stocks, i.e., are the simplifications introducing bias or errors that would lead to contradictory results from sound management based on the advice of the assessment (Cadrin, 2020)?

Given that ultimately we do *not* know the underlying structure (models such as these are built specifically to elucidate such meaningful structure) we find ourselves with a "Catch-22." We need spatial components to understand spatial structure, but need to know the spatial structure in order to understand what simplifications are allowable. Note that it is not enough to simply design the most complicated spatial model possible given the data will likely not exist to fit that model with any degree of confidence.

The way out of this hole, as many authors have noted, is to use simulations (Cadrin, 2020) (Daniel Goethel, 2023). Specifically one can use simulations to do hypothesis testing of the form - if my stock had spatial structure of the form X would my specific assessment capture that at all or would it introduce biases and/or errors as a result? This is a very powerful technique that has been responsible for illustrating just how important spatial assessments are to management in the first place.

Here then is another potential application of the kind of model proposed in this document. As we mentioned earlier these assessments are trying to model at least three things - recruitment, local population dynamics

parameters such as growth or natural mortality, and then movement. By having a movement simulation based on real observations of the species in question we would be able to help build a high quality simulation of at least the last of those three. Rather than having to assume one form of movement or another we'd instead be able to see whether the assessment paradigm in question can capture the relevant aspects of structure in the *real* movement of our stocks and sub-stocks. And this in turn would help in selecting the assessments best suited to the job (as well as illustrate what uncertainties exist if a lack of specific kinds of data requires us to use an over-simplified assessment paradigm).

7.3.5 Comments on Interpretation

Something that any student of statistical or Machine Learning learns early on is that models can be deceiving in interpretation due to correlations in different kinds of data or parameters. For example a common critique of using "feature importance" on random forests is that models like random forests can extract the value of a set of correlated features out of a single feature. So, for example if studying vulnerability to a specific kind of disease, a low feature importance for something like exercise may not mean anything given exercise patterns may be heavily correlated with things like, say, diet. As models get more and more complicated and include more features the likelihood that situations like these arise becomes greater and greater.

For such things as fisheries assessments this is particularly concerning as the specific ranges of parameters derived from the models are used directly to set such things as allowable biological catch (King, 2007). Yet given we are often estimating recruitment, abundance, mortality, and in the case of spatial assessments things like movement and spatial heterogeneity all in the same model the only guarantee we have is that this choice of parameters fits the current data, not that they are parameters that are indicative of reality itself.

In order to get a better guarantee that the parameters of the model are indicative of reality rather than just parameters that cause fit in the overall model one would need to estimate the parameters independently so cross feature "compensation" cannot occur and obscure the interpretation of the results.

It has been noted that one of the most difficult aspects to fit is the movement matrix X described above (Punt, 2019). So here too is another application of the model proposed in this document. Rather than having to depend on fitting X through the assessment MLE routine such a fit could be

taken directly from the behavioral model proposed herein thereby reducing the data requirements for the assessment itself and guaranteeing that X is not being used to improve fit of the model in compensation for other important life history and abundance patterns.

7.3.6 Other Applications of a Movement Model

There are many decisions that have to be made when designing a spatial assessment (Cadrin, 2020) (Daniel Goethel, 2023) and some of these would directly benefit from the kind of view our proposed model would provide:

1. **Choosing sub-regions** - Selecting the scale of sub-regions, the boundaries of those regions, and whether those regions should move over time requires intimate knowledge of fish movement.
2. **Parameters to make regional** - Not all population dynamic parameters need be regional and obviously being able to share information across regions reduces the data requirements for the assessments. By knowing what kinds of covariates actually drive movement, data driven decision about what to keep regional versus not could be made.
3. **Temporal resolution** - What time step to choose or whether only specific seasons need to modeled (and what those season are) would be another place in which data driven decisions would be aided by models of the kind proposed herein.

7.4 Modeling Movement

Modeling movement has traditionally divided into two categories - the Eulerian and Lagrangian approaches. In brief, the former deals with movement as a series of flows throughout a space (that may or may not be discretized) while the latter treats each animal independently (Carl Walters, 2005). We will provide a brief introduction to both approaches.

7.4.1 Eulerian Approach

The typical strategy behind the Eulerian approach is to subdivide the region in question into discrete subregions and then model movements between the boundaries of these regions as the result of advection and diffusion where advection is a directed movement and diffusion is not (John Sibert, 1999) (Patrick Lehodey, 2008).

These advection and diffusion terms are represented by partial differential equations where each may be parametrized by life history or environmental factors depending on the model in question (Patrick Lehodey, 2008). Data is included in the model as boundary conditions, for example where we were fish released and where were they recovered. Finally time too is discretized in order to allow finite difference methods to be used to solve the PDE's in question (John Sibert, 1999).

The result of solution is a description of the advection and diffusion terms parametrized by whatever was included in the original model. With these in hand, simulations can then be run to represent the overall movement modeled.

The advantage of the Eulerian approach is that it is specifically attempting to catch population level movement patterns and therefore does not need as thorough parametrization as the Lagrangian approaches. It is also therefore less data intense in its requirements and generally stock assessments are only interested in population or sub-population level movement anyways (Patrick Lehodey, 2008).

7.4.2 Lagrangian Approach

In contrast to the Eulerian approach the Lagrangian approach attempts to model at the individual level. Therefore you'll see the terms Lagrangian and Individual Based Model (IBM) used interchangeably.

The approaches used to build these kinds of models are extremely variable (Volker Grimm, 2005) (a point we'll get back to momentarily) but follow the general paradigm of:

1. Collect some form of movement data (could be very high resolution or quite low resolution in some cases).
2. Collect environmental data such as temperatures, measures of abundance, measures of productivity, flow/current data, etc.
3. Craft a model that relates the specific queues from the environment into specific movements.
4. Tune any parameters of those models through some kind of fitting paradigm.

All of these steps are presented so vaguely because there really are no general paradigms for how these are done. For example, in the last step - tuning the parameters - strategies range from integration of deterministic models against boundary conditions (Joe Phillips, 2018), to "by-eye" tuning (R. Andrew Goodwin, 2006), to weighted averages across multiple different allowable parameters (Brian Burke, 2016).

Such approaches however have been shown to be able to model very fine scale movement dynamics and have been especially useful in applications in rivers where structures like dams are being designed and the ecological impacts are wanting to be understood (R. Andrew Goodwin, 2006) (Volker Grimm, 2005). These kinds of fine scale dynamics are not possible with Eulerian approaches.

IBM's can also capture much finer detail about *why* specific movements occur because they are effectively modeling the individual itself. Whereas such detail ends up getting washed out by Eulerian approaches the focus on the individual means very specific details of behavior have to be sorted out and understood. A feature which, depending on the situation, can be a pro or a con.

Finally it has been noted that one can derive Eulerian models from Lagrangian ones making the latter perhaps more general (Joe Phillips, 2018).

7.4.3 Struggles in Building IBMs

There are a series of key issues (Volker Grimm, 2005) that have arisen in building IBM's of which we will note a few that relate to our proposal:

1. **High Precision as a Requirement** - IBMs typically predict what an animal *will* do as opposed to the likelihoods of what it *could* do.

This creates the requirement for capturing *all* the meaningful drivers of variability in the behavior of the animal. To see why this is consider the example of a creature that can either choose to stay put or move to another location. If the more likely of the two actions is to stay put a deterministic model *must* predict that the animal will *always* stay put. But this is obviously non-sensical. However, because the model is deterministic in nature the modeler cannot depend on stochasticity to save the day - instead they must find whatever is driving the difference between staying put and moving on. And this can cascade into greater and greater requirements for modeling the "internal state" of an animal.

2. **Obscurity** - given the inherent complexity of IBMs that arises from the point above it gets increasingly difficult to make IBMs clear to new users or potential developers. There can often be many dozens of different parameters, functions, and "internal states" to keep track of without even considering all of the external modeling that has to go in place to provide for all of this. And this makes for not just complicated descriptions but equally if not more complicated code.
3. **Hand Made** - almost all IBMs have been hand-coded in the sense that every functional relationship is specifically crafted and outlined by the modeler. This is obviously a very laborious process but it also means that no two IBMs are the same and that lack of generalization or abstraction makes exchanging lessons learned near difficult to impossible.
4. **No Generalization of Fitting or Diagnostics** - because each IBM is effectively unique the methods for fitting, the data used, and the diagnostics developed are all unique as well. This once again increases the workload of the modeler, makes exchanging lessons learned and best practices extremely difficult, and also makes starting a new model from scratch quite prohibitive.

7.4.4 Comparison to Proposed Odds Model

Our odds modeling approach looks to directly address these issues by:

1. **Modeling Likelihoods** - by modeling likelihoods instead of having a deterministic model we no longer *have* to model each kind of relevant variance, instead the variance can be encoded through stochasticity.

While we would obviously like to know what creates the variance, this allows us to build our model iteratively and to whatever complexity is allowable by the data. It also means we can build models that can be used in contexts where specific kinds of environmental or "internal state" data may be missing.

2. **Information as the Model** - by using machine learning (ML) we are drawing the attention away from the specifics of the model itself to what information was used to train it and what metrics were used to select the best fit. This makes the model far clearer and creates a generalized paradigm for describing any such model.
3. **Machine Learning** - as described in our rationale for creating these models, ML allows us to remove much of the labor in finding the approximate function that relates our inputs and outputs and instead allows us to place our focus on creating quality information to input to the model.
4. **MLE in All Cases** - odds modeling allows us to use the same measure of fitness every single time - likelihood. And because all of the models share the same format, the form of diagnostics and fitting can also be shared.

7.5 Chinook salmon Ecology and Management

Chinook salmon (*Oncorhynchus tshawytscha*) are the largest species of Pacific salmon (Brian Riddell, 2018). Like other Pacific salmon the species is both anadromous and semelparous leading to a very interesting and distinct life history.

Juvenile Chinook salmon are generally born in the rivers where their parents spawned (although some genetic mixing does occur through straying). Sometime in their first or second year of life they transition to the marine environment where they will spend the rest of their life before maturation (Brian Riddell, 2018). This leads to a specific system of enumerating age that includes both years in freshwater and years at sea. For example if a particular fish spends one winter in freshwater and so far 2 at sea it would be a 1.2 age fish. If it spent zero winters in freshwater and 3 at sea it would be a 0.3 age fish. Clearly both of these examples are the same overall age.

The immature fish then spend time (up to five years) growing in the marine environment before maturing and then migrating back to the river where they were born to breed and subsequently die (Brian Riddell, 2018). The timing of such homing behavior is clearly season dependent with species further north having to avoid ice build up during the cooler months (Brian Riddell, 2018). The mechanism behind this natal homing ability is still a question under active research (Chloe Bracis, 2012).

Given this homing behavior, Chinook salmon populations are in fact a series of reasonably separated genetic sub-populations each of which can have its own life history strategy. Indeed using tagging data it has been demonstrated that both the initial and later marine distributions of Chinook salmon is highly dependent on the region of origin for the fish in question (Andrew Shelton, 2019) (S. Tucker, 2011).

Beyond this, large scale analysis of fisheries dependent data has also demonstrated some of the likely migration patterns the species exhibits during its rearing phase at sea (Joseph Langan, 2024).

Studies like these point to Chinook salmon being a highly mobile, spatially heterogeneous species.

Broadly distributed through out the Northern pacific, Chinook salmon are known to spawn in rivers spanning from the Sacramento River in central California to the Kotzebue Sound in Alaska on the eastern side of the pacific and primarily in the Kamchatka Peninsula on the western side (Brian Riddell, 2018).

Marine residency in the North Eastern pacific is primarily coastal from California up to Western Alaska and then spans into much of the Eastern Bering Sea (Brian Riddell, 2018) (Joseph Langan, 2024).

Chinook salmon are well known piscivores with several studies of stomach contents finding primarily fish and squid as their main sources of prey. It has been noted that this diet is similar to that of Coho salmon but that it is unlikely that the two species experience much cross-species competition for these resources (Brian Riddell, 2018).

Chinook salmon themselves are prey for animal groups such as Salmon sharks, whales, and dolphins but it is still unclear how much mortality in later life stages is due to predation as opposed to other sources. Size selective predation on earlier life stages is a well known driver of natural mortality (Brian Riddell, 2018).

Other drivers of natural mortality include a variety of known diseases, starvation during periods of low ocean productivity, and potential competition for food with species like whales (Brian Riddell, 2018).

Chinook salmon have been the subject of hatchery programs since the 1950s with the past decade seeing hatchery releases of 254million individuals on average. The vast majority of these individuals (approx. 200million) have come from hatchery programs in the United States with the remaining significant portion from programs in Canada (Brian Riddell, 2018).

Fish are released at weights of around 6-9g and a large number of these have a micro-wire with an identifiable code etched into them inserted into their snouts. These are the tags driving the Coded Wire Tag program in the Northeastern Pacific that has become a major source of demographic and movement data used in managing and understanding Chinook salmon sub-populations (Andrew Shelton, 2019). The etchings provide a unique label for each cohort from each hatchery.

The initial purpose of these hatchery programs was to supplement losses in juveniles due to targeting of adult fish in river systems, however the programs have not created the kind of recovery that was intended and it is unclear how wild and hatchery populations interact. For example it is well known that there is quite a bit of genetic exchange between wild and hatchery types and concern that hatchery types are less suited to wild environments (Brian Riddell, 2018).

There are valuable tribal, commercial, and recreational fisheries of Chinook salmon throughout the Pacific (PFMC, 2024) yet given the highly

mobile nature of Chinook salmon independent management by sovereign nations is simply not possible and so management is the ultimate responsibility of the Pacific Salmon Commission. Federal management councils (NOAA for the United States) then collaborate with the Pacific Salmon Commission to set harvest quotas and other management objectives. The governments of Canada and the United States then work with states, tribes, and industry to create and manage shared objectives based on these overall goals (NOAA, 2022).

These management goals as well as designations of statuses such as "overfished" have to be stated on the sub-population level given the fact that Chinook salmon form so many different genetic sub-populations (PFMC, 2024) - a fact that further complicates management. Indeed several sub populations are actually listed as endangered and threatened under the Endangered Species Act (NOAA, 2024a).

This combined with other threatened stocks (NOAA, 2024a) means that bycatch avoidance is a serious issue for Chinook salmon management and for other overlapping fisheries such as the Walleye Pollock fishery (James Ianelli, 2015).

To manage this diversity the United States uses a variety of tools including but not limited to - differentiated management zones, minimum size limits, bag limits for recreational fisheries, allowed seasons and quotas, and fishing gear restrictions (PFMC, 2024).

Finally, Chinook salmon are especially prized for their large size which has led to serious questions about size selective mortality from fishing as both recreational and commercial fisheries have incentives for targeting larger fish. While it is not yet known whether fisheries are responsible, there is clear evidence that fish size in Chinook salmon populations has been decreasing steadily over the years which has potential impacts on both the fisheries themselves (in terms of optimizing yields) and for Chinook salmon's primary predators (Neala Kendall, 2011)(Brian Riddell, 2018).

Being an interesting species to model, however, is not enough for our purposes. Given that we want to build machine learning models and model driven simulations we also need a species with available and high quality data. Chinook salmon in the GOA fit this bill nicely. There is quite a lot of electronic tagging data that has been collected for them (CITE), all of which is fisheries independent allowing for closer examination of Chinook salmon behavior, and there is an abundance of other diverse data types including coded wire tag (CWT) data, fishery independent trawl data, and

so on. This availability of data means that we can focus on the modeling aspects of this problem as opposed to data acquisition.

All in all then, given their ecology, data availability, and the corresponding complexities of management, Chinook salmon represent an ideal model species for trying to understand and quantify spatial patterns of movement and behavior.

8 Bibliography

References

- Alyson Little, Coby Needle, R. H. D. H. C. T. M. (2014). Real-time spatial management approaches to reduce bycatch and discards: experiences from europe and the united states. *Fish and Fisheries*.
- Andrew Shelton, William Satterthwaite, E. W. B. F. B. B. (2019). Using hierarchical models to estimate stock-specific and seasonal variation in ocean distribution, survivorship, and aggregate abundance of fall run chinook salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 76:95–108.
- Brad Erisman, William Heyman, S. K. T. E. S. P. O. A.-O. R. N. (2017). Fish spawning aggregations: where well-placed management actions can yield big benefits for fisheries and conservation. *Fish and Fisheries*.
- Brian Burke, James Anderson, J. M. L. T. D. T. N. B.-A. B. (2016). Estimating behavior in a black box: how coastal oceanographic dynamics influence yearling chinook salmon marine growth and migration behaviors. *Environmental Biology of Fishes*, 99:671–686.
- Brian Riddell, Richard Brodeur, A. B. P. M. J. M. J. O.-M. T. L. W. B. W. A. W. (2018). *The Ocean Ecology of Pacific Salmon and Trout*, chapter Ocean Ecology of Chinook Salmon. American Fisheries Society.
- Cadrin, S. (2020). Defining spatial structure for fishery stock assessment. *Fisheries Research*.
- Carl Walters, S. M. (2005). *Fisheries Ecology and Management*. Princeton University Press.
- Chloe Bracis, J. A. (2012). An investigation of the geomagnetic imprinting hypothesis for salmon. *Fisheries Oceanography*, 21(2-3):170–181.
- Computers, W. (2024). Minipat.
- Daniel Goethel, Aaron Berger, S. C. (2023). Spatial awareness: good practices and pragmatic recommendations for developing spatially structured stock assessments. *Fisheries Research*.

- Elizabeth Selig, Kristin Kleisner, O. A. F. A. A. C.-T. R. F. M. H. L. K. P. M. B. R. L. S.-D.-A.-M. S. D. T. E. T. S. T. S. V. (2016). A typology of fisheries management tools: using experience to catalyse greater success. *Fish and Fisheries*.
- James Ianelli, D. S. (2015). Estimating impacts of the pollock fishery bycatch on western alaska chinook salmon. *ICES Journal of Marine Science*, 72(4):1159–1172.
- Joe Phillipsa, Alex Guptaa, I. S. E. v. S. M. L. P. L. J. H. S. N. (2018). An individual-based model of skipjack tuna (*katsuwonus pelamis*) movement in the tropical pacific ocean. *Progress in Oceanography*, 164:63–74.
- John Sibert, John Hampton, D. F. P. B. (1999). An advection–diffusion–reaction model for the estimation of fish movement parameters from tagging data, with application to skipjack tuna (*katsuwonus pelamis*). *Canadian Journal of Fisheries and Aquatic Sciences*, 56:925–938.
- Joseph Langan, Curry Cunningham, J. W. S. M. (2024). Opening the black box: New insights into the role of temperature in the marine distributions of pacific salmon. *Fish and Fisheries*.
- Kai Lorenzen, Robert Steneck, R. W. A. P. F. C. K. L. (2010). The spatial dimensions of fisheries: Putting it all in place. *Bulletin of Marine Science*, 86(2):169–177.
- King, M. (2007). *Fisheries Biology, Assessment and Management*. Blackwell Publishing.
- Michael Armstrong, S. C. (2001). Morphometric variation among spawning groups of the gulf of maine–georges bank herring complex. *International Herring Symposium*.
- Neala Kendall, T. Q. (2011). Length and age trends of chinook salmon in the nushagak river, alaska, related to commercial and recreational fishery selection and exploitation. *Transactions of the American Fisheries Society*, 140:611–622.
- NOAA (2022). Ocean salmon fisheries on the west coast.
- NOAA (2024a). Chinook salmon (protected).
- NOAA (2024b). Red grouper.

- NOAA (2024c). Tool: Age structured assessment program (asap).
- Oliver Durr, Beate Sick, E. M. (2020). *Probabilistic Deep Learning*. Manning Publications.
- Patrick Lehodey, Inna Senina, R. M. (2008). A spatial ecosystem and populations dynamics model (seapodym) – modeling of tuna and tuna-like populations. *Progress in Oceanography*, 78:304–318.
- PFMC (2024). *Pacific Coast Salmon Fishery Management Plan*. Pacific Fishery Management Council.
- Prince, J. (2010). Rescaling fisheries assessment and management: A generic approach, access rights, change agents, and toolboxes. *Bulletin of Marine Science*, 86(2):197–219.
- Punt, A. (2019). Spatial stock assessment methods: a viewpoint on current issues and assumptions. *Fisheries Research*.
- R. Andrew Goodwin, John Nestler, J. A. L. W. D. L. (2006). Forecasting 3-d fish movement behavior using a eulerian–lagrangian–agent method (elam). *Ecological Modelling*, 192:197–223.
- S. Tucker, M. Trudel, D. W. J. C. J. M. M. T. C. W. T. D. B. (2011). Life history and seasonal stock-specific ocean migration of juvenile chinook salmon. *Transactions of the American Fisheries Society*, 140:1101–1119.
- Tim Sippel, J. Paige Eveson, B. G. C. L. S. H. M. M. P. K. F. C. V. T. S. T. A. A.-d.-S.-S. N. (2014). Using movement data from electronic tags in fisheries stock assessment: A review of models, technology and experimental design. *Fisheries Research*.
- Volker Grimm, S. R. (2005). *Individual-based Modeling and Ecology*. Princeton University Press.