

Session 3 (Design: choosing the right study): Exercises

Foundations of Clinical Research · Week 3 · Saturday · Neudata · #ClearDataClearImpact

This handout has three parts. No computer or calculator is needed.

Part	Activity	When	How
A	Name that design (8 abstracts + 1 optional)	End of section 3.1 (15 min)	Pairs
B	Choose the design for 5 questions (4 given + your own; 2 optional bonus)	After section 3.2 (20 min)	Your group
C	Proposal builder 3: design, setting and population	End of the session (35 min)	Your group

Key words of the day: exposure (what comes first) · outcome (what we count later) · comparison group · observational (we watch) · experimental (we assign) · randomised (chance assigns) · cross-sectional (snapshot) · case-control (look back) · cohort (follow forward) · cluster / crossover / factorial trial · non-inferiority · pilot · interrupted time series · qualitative.

Part A: Name that design

Instructions

1. Work in pairs. Read each short abstract.
 2. Ask the three questions from the decision tree:
 - **Did the investigator ASSIGN the exposure or treatment?** If yes: **was it assigned at random?**
 - If not assigned: **is there a comparison group?**
 - If there is: **in which direction does time run?** Exposure first, then outcome (cohort); outcome first, then look back (case-control); both at the same moment (cross-sectional).
 3. Write the design and **one word or phrase from the abstract that proves it.**
 4. Put a star next to the abstract you found hardest.
- > **Design choices:** case report / case series · cross-sectional · case-control · cohort (prospective or retrospective) · non-randomised intervention study · randomised controlled trial (RCT) · systematic review / meta-analysis

Abstract 1. Two hundred and forty adults on mechanical ventilation were randomly allocated by a central computer to oral care with chlorhexidine twice daily or to standard oral care. Ventilator-associated pneumonia was recorded over 28 days by an assessor who did not know the group.

Design: _____ Evidence: _____

Abstract 2. On one day in March, two trained nurses examined all 412 inpatients of a hospital for pressure injuries and recorded the type of mattress each patient was lying on. Pressure injuries were

present in 9% of patients on standard mattresses and 4% on pressure-relieving mattresses.

Design: _____ Evidence: _____

Abstract 3. Forty ICU patients who developed candidaemia were compared with 80 ICU patients from the same units who did not, matched on age and month of admission. The investigators then searched the records of both groups for parenteral nutrition in the 14 days before the culture date.

Design: _____ Evidence: _____

Abstract 4. All 600 adults admitted with acute ischaemic stroke who received thrombolysis were classified on arrival by door-to-needle time (under 60 minutes or 60 minutes and over). Each patient was then followed up to record functional independence at 90 days.

Design: _____ Evidence: _____

Abstract 5. Using electronic records from 2019 to 2022, the investigators identified 1,150 ICU patients and divided them into those who did and did not receive a continuous benzodiazepine infusion for 3 days or more. They then compared how many patients in each group had delirium recorded later in the ICU stay.

Design: _____ Evidence: _____

Abstract 6. We describe five patients who developed severe hypernatraemia within a week of a new fluid protocol being introduced on our unit, with their laboratory results and clinical course.

Design: _____ Evidence: _____

Abstract 7. Central-line infection rates were compared for the six months before and the six months after a unit-wide insertion-bundle education programme, which the research team introduced on all beds on the same day.

Design: _____ Evidence: _____

Abstract 8. We searched three databases for randomised trials comparing restrictive with liberal fluid strategies in adults with septic shock, assessed each trial's risk of bias, and pooled 28-day mortality across nine trials.

Design: _____ Evidence: _____

Abstract 9 (optional). Twelve hospital wards were randomly allocated, ward by ward, to a new hand-hygiene feedback programme or to usual practice. Hospital-acquired infections were counted in all patients on those wards for six months.

Design: _____ Evidence: _____

Extra (if you finish early): for each design, write one strength and one weakness.

Part B: Choose the design for 5 questions

Instructions. For each question, name the best **feasible** design (use the decision tree and the "Match the design to the question" slide), give **one reason**, and name the design's **main weakness**. Question 5 is your group's own research question from Part 1.

#	Question	Design	Reason	Main weakness
1	How many inpatients have a pressure injury today, and is it more common on standard mattresses?			
2	Does daily chlorhexidine bathing reduce bloodstream infections? It will be introduced ward by ward, and nurses on one ward cannot bathe only half their patients.			
3	A sepsis bundle was introduced in our hospital in 2023. We have monthly sepsis mortality for 3 years before and 2 years after. Did mortality fall?			
4	Why do nurses not report near-miss medication errors?			
5	Our own question: _____ _____			

Bonus (optional, if you finish early)

#	Question	Design	Reason
6	What are the risk factors for a rare event: Stevens–Johnson syndrome in hospital patients?		
7	A new 5-day antibiotic course is cheaper and easier than the 10-day course. We only need to show it is not worse for cure.		

For your own question (5), also answer: **at the end, will you be allowed to say "caused" or "associated with"?** _____

Part C: Proposal builder 3 (design, setting and population)

Instructions. As a group, complete the first three rows of Part 2 of the proposal, then copy them into the template (Templates/Research_Proposal_Template.docx). The rest of Part 2 (exposure,

Our research question (from Part 1):

Draw your study on a timeline (exposure → outcome; mark when each is measured):



- Page 4