

FIGURES




Figure 1: 59-year-old male with Multiple biliary hamartomas (MBH) in liver. (2022)

1A: CT reveals multiple round, small, low-density nodules.

1B: CE-MRI shows multiple small, low-density, non-enhancing nodules within the liver.

1C: Coronal MRCP image depicts the typical “starry sky” configuration of MBH in the liver, due to the presence of multiple, small, high signal cystic lesions scattered throughout hepatic parenchyma, not communicating with the non-dilated intra- and extrahepatic biliary tree




Figure 2: 59-year-old male with Multiple biliary hamartomas (MBH) in liver. (2022)

2A: Multiple dark purple nodules of varying sizes (ranging from millimeters to centimeters) are scattered on the liver surface, with smooth surfaces. Their dark purple color suggests possible blood stasis or tissue changes, distinguishing them from normal liver tissue.

2B: Pathological images show multiple irregularly dilated biliary duct-like structures lined by uniform cuboidal epithelial cells, embedded in a dense, grayish-white, tough fibrous stroma. These findings are consistent with biliary hamartoma (magnification, $\times 100$).




Figure 3: 59-year-old male with Multiple biliary hamartomas (MBH) in liver (2024). Postoperative two-year follow-up MRI (3A) and MRCP (3B) showed no significant changes compared to the previous studies two years ago.