

# Voltage-Gated Calcium Channels and Idiopathic Generalized Epilepsies

HOUMAN KHOSRAVANI AND GERALD W. ZAMPONI

*Department of Physiology and Biophysics, Hotchkiss Brain Institute, University of Calgary, Calgary, Canada*

---

I. Introduction: Idiopathic Generalized Epilepsies	941
A. Classification of seizures and epilepsies	941
B. Features of idiopathic generalized epilepsy	943
II. Molecular Physiology of Voltage-Gated Calcium Channels and Associated Ancillary Subunits	944
A. Subtypes and physiological roles of voltage-gated calcium channels	944
B. Molecular structure of voltage-gated calcium channels	944
C. Structural basis of calcium channel function	945
D. Ancillary subunits of calcium channels	946
III. Voltage-Gated Calcium Channels and Ancillary Subunits in Idiopathic Generalized Epilepsy	947
A. T-type calcium channels and spike-wave seizures	947
B. P/Q-type channel defects and spike-wave seizure disorders	953
C. Involvement of ancillary subunits of voltage-gated calcium channels in seizure disorders	955
D. Antiepileptic drugs and calcium channel pharmacology	956
IV. Conclusions	956

---

**Khosravani, Housman, and Gerald W. Zamponi.** Voltage-Gated Calcium Channels and Idiopathic Generalized Epilepsies. *Physiol Rev* 86: 941–966, 2006; doi:10.1152/physrev.00002.2006.—The idiopathic generalized epilepsies encompass a class of epileptic seizure types that exhibit a polygenic and heritable etiology. Advances in molecular biology and genetics have implicated defects in certain types of voltage-gated calcium channels and their ancillary subunits as important players in this form of epilepsy. Both T-type and P/Q-type channels appear to mediate important contributions to seizure genesis, modulation of network activity, and genetic seizure susceptibility. Here, we provide a comprehensive overview of the roles of these channels and associated subunits in normal and pathological brain activity within the context of idiopathic generalized epilepsy.

## I. INTRODUCTION: IDIOPATHIC GENERALIZED EPILEPSIES

Epilepsy is a disorder of recurrent spontaneous seizures and is among the most common neurological conditions, accounting for 0.5% of the whole burden of diseases worldwide (164). One in 10 persons with a normal life span can expect to experience at least 1 seizure (83). Epileptic seizures are characterized as abnormal hyperexcitable, hypersynchronous neuronal population activity and manifest themselves in different ways depending on their site of origin and subsequent spread (85). The clinical diversity observed in epileptic seizure disorders is a reflection of the numerous cellular and network routes to seizure genesis. Seizures can originate in different brain structures that are responsible for motor, sensory, cognitive, and autonomic systems. The transition to seizure activity can be considered as a disruption of normal brain function. This transformation to pathological activity can

be manifested by changes that occur at the level of single neurons (e.g., molecular alterations in ion channels, complements thereof, and regulatory proteins), local circuitry (e.g., alterations in cell-to-cell coupling via chemical and electrical synapses as well as ephaptic communication), or at the level of anatomically defined neuronal networks (e.g., structural changes and alteration between excitatory and inhibitory neuronal, interneuronal, and glial elements) (191). This notion is complicated by the fact that intrinsic neuronal properties can feedback to alter network dynamics, and vice versa. Moreover, activity at the network level can trigger alterations in gene expression that can affect the firing properties of individual neurons.

### A. Classification of Seizures and Epilepsies

From a clinical perspective, seizures can be classified into two broad categories (107). Focal (or partial) seizures are localized to one hemisphere and involve only a

small brain region such as an area within the motor cortex that can result in motor seizures. Partial epileptic seizures may also arise from focal brain lesions caused, for example, by traumatic brain injury (116). However, it is now known that certain focal epilepsies can have a genetic origin (287). The second category comprises generalized seizures, which are characterized by virtually simultaneous onset of seizure activity over both brain hemispheres as seen using field potential recordings (Fig. 1). These generalized epileptic syndromes can further be classified as either symptomatic or idiopathic. Symptomatic generalized epilepsy is thought to be caused by identifiable factors such as anoxia and infection, which can result in widespread brain damage. In contrast, idiopathic generalized epilepsies have no clear etiology and may be at least partly caused by defects at the genetic level (20, 202). Data gathered from several cohort studies suggest that the frequency of idiopathic generalized epilepsies in the general epilepsy population is between 15 and 20% (reviewed in Ref. 130). As outlined above, although epileptic disorders span a continuum of conditions, we focus here predominantly on idiopathic generalized epilepsies to highlight recent developments implicating the involvement of voltage-gated calcium channels in this spectrum of epilepsy disorders.

Absence-type epilepsies account for 3–4% of all seizure disorders (63). As classified by the International League Against Epilepsy (ILAE) (3), typical absence seizures are a defining attribute of several of the idiopathic generalized epilepsies (Table 1). Seizure episodes are typified by a sudden impairment in consciousness, behav-

TABLE 1. *Clinical attributes of generalized idiopathic epilepsy*

<i>Childhood absence epilepsy</i>
Absence seizures manifest between ages 4 and 8
More common in females
Respond well to antiepileptic drugs
<i>Juvenile absence epilepsy</i>
Onset around puberty
No sex predominance
Atypical absence seizures
Often have other seizure types (generalized tonic-clonic, myoclonic, and atonic seizures)
Patients may be intellectually impaired
<i>Juvenile myoclonic epilepsy</i>
Seizure onset between ages 8 and 18
Have generalized tonic-clonic and myoclonic seizures
Myoclonic seizures are characterized by sudden jerks of the neck, shoulders, and arms that may be repetitive without losing awareness
Patients may exhibit photosensitivity
Most common form of idiopathic generalized epilepsy
<i>Generalized tonic-clonic seizures alone</i>
Seizures occur upon awakening
Seizures can occur during sleep
Time of seizure occurrence may not be linked to any external factors

Generalized seizures show bilateral and synchronous onset observed in electroencephalographic (EEG) recordings. Seizures in idiopathic generalized epilepsy are usually one of several types: generalized tonic-clonic, myoclonic, absence, or atypical absence. Idiopathic generalized epilepsy is a subtype of generalized epilepsy that has a genetic predisposition and includes the epileptic disorders (3) shown in the table.

ioral arrest, and may be accompanied by facial clonus or other subtle physical manifestations. These seizures are generally short in duration (lasting typically <10 s) and terminate suddenly without any postseizure alterations in




FIG. 1. Spike-wave discharges in human absence epilepsy. Typical 3- to 4-Hz spike-wave discharges observed over the entire brain, a hallmark of generalized epilepsy. Electroencephalographic (EEG) recordings are presented for a selection of electrodes placed over the scalp covering the entire head. Placement of EEG electrodes is depicted on a head model. Voltage tracings are presented in a bipolar manner with potentials, originally recorded in reference to a noninvolved electrode, represented as a subtraction from two neighboring electrodes to produce a single tracing. [Adapted from Yoshinaga et al. (312).]

behavior. The electrographic hallmark of absence seizures are spike-wave discharges (SWDs) that can be recorded using electroencephalography (EEG). These discharges typically arise synchronously over both brain hemispheres with a frequency of ~3–4 Hz and are maximal over frontal midline regions with minimal involvement of posterior brain regions (85, 122, 234). In generalized epilepsy, SWDs are principally mediated by interactions between thalamic and cortical networks (19) and are known to involve the activity of voltage-gated calcium channels. It must be noted that although absence seizures are always associated with SWDs, the converse is not true as SWDs are observed in other forms of epilepsy (85). This is a relevant point when considering subsequent discussion on models of spike-wave epilepsy.

## B. Features of Idiopathic Generalized Epilepsy

Patients affected by idiopathic generalized epilepsy commonly present with their first seizure between the ages of 3 and 16 (107); however, adult onset has also been reported (187). Patients with idiopathic epilepsy syndromes exhibit no underlying structural or brain lesions when assayed by radiological imaging and are otherwise neurologically normal (84, 107). In the case where absence seizures are the predominant epileptic seizure type (84), childhood absence epilepsy and juvenile absence epilepsy encompass the two main idiopathic epilepsy subtypes. Patients with childhood absence epilepsy are typically neurologically normal. On the other hand, juvenile absence epilepsy patients often have intellectual impairment and they have atypical absence seizures, characterized by a longer duration, less abrupt onset, loss of postural tone, and often the co-occurrence of other seizure types (Table 1). Two other idiopathic generalized epilepsy seizure types, juvenile myoclonic epilepsy and idiopathic generalized epilepsy with tonic-clonic seizures alone (4) (Table 1), are recognized by the international classification (3). Specific diagnosis of different idiopathic generalized epilepsy types occurs best when both clinical and EEG data are available. It is interesting to note that generalized tonic-clonic seizures in idiopathic generalized epilepsy usually occur after awakening, particularly from a brief period of sleep when preceded by sleep deprivation. This relation is particularly clear in patients with juvenile myoclonic epilepsy and implicates the thalamo-cortical network, which is known to be involved in both sleep rhythm and SWD generation.

Of the absence-type idiopathic generalized epilepsy disorders, childhood absence epilepsy is perhaps the one that has been most extensively studied at the genetic level. Patients with childhood absence epilepsy can have a family history of epilepsy that is inherited in an autosomal dominant manner, with concordances of up to 85%

reported in monozygotic twins (15). However, genetic studies have also identified individuals with mutant genes who do not exhibit the epileptic phenotype (244). Moreover, the precise identification of gene loci is complicated by the presence of a large number of genetic polymorphisms in the childhood absence epilepsy population (241). Therefore, it is not entirely clear what the relative contributions of external or endogenous factors (such as involvement of multiple genes) are in bringing about the spectrum of epileptic seizure types observed in idiopathic generalized epilepsies. Nonetheless, there is substantial evidence implicating a genetic component in idiopathic generalized epilepsy, and in particular in absence-type seizure disorders (120). Genetic association studies have identified mutations in a number of different types of voltage- and ligand-gated ion channels, including GABA<sub>A</sub> receptors, as well as voltage-dependent sodium, potassium, and chloride channels, resulting in either a gain or loss of function (102, 201, 285) (see Table 2).

In the context of idiopathic epilepsies, the two brain structures that have received the most attention are the neocortex and the thalamus. The normal connectivity of these two brain structures differs at the level of neuronal/interneuronal cell types and at the level of network organization. Interactions between the thalamus and neocortex bring about synchronized oscillations that, under normal conditions, serve physiological roles such as the generation of sleep spindles, which occur in early stages of sleep (262). During spike-wave seizures, the normal function of this circuitry is somehow altered in a manner that results in the generation of SWDs. As such, voltage-gated calcium channels have increasingly emerged as important players in the generation of SWDs in both humans and in experimental models of epilepsy. In this article, we review the current state of knowledge concerning the role of these channels and their auxiliary subunits in idio-

TABLE 2. *Other voltage- and ligand-gated ion channel genes carrying mutations that have been linked to generalized epilepsies*

Gene	Channel/Subunit	Epilepsy Disorder	Reference Nos.
<i>KCNA1</i>	K <sub>v</sub> 1.1	EA1	28, 319
<i>KCNQ2</i>	K <sub>v</sub> 7.2	BFNC, myokymia	59
<i>KCNQ3</i>	K <sub>v</sub> 7.3	BFNC	42, 252
<i>CLCN2</i>	CLC-2	CAE	113
<i>SCN1A</i>	Na <sub>v</sub> 1.1	GEFS+, SMEI	50, 89
<i>SCN2A</i>	Na <sub>v</sub> 1.2	GEFS+, BFNIS	117, 272
<i>SCN1B</i>	β <sub>1</sub>	GEFS+	295
<i>GABRA1</i>	GABA <sub>A</sub> (α <sub>1</sub> )	JME	53
<i>GABRG2</i>	GABA <sub>A</sub> (γ <sub>2</sub> )	GEFS+, FS, CAE	12, 294
<i>GABRD</i>	GABA <sub>A</sub> (δ)	GEFS+	71

EA, episodic ataxia; BFNC, benign familial neonatal convulsions; CAE, childhood absence epilepsy; GEFS+, generalized epilepsy with febrile seizures plus; SMEI, severe myoclonic epilepsy of infancy; BFNIS, benign familial neonatal and infantile seizures; JME, juvenile myoclonic epilepsy; FS, febrile seizures.

pathic generalized epilepsy and related pathophysiologicals. Specifically, we focus predominantly on T-type and P/Q-type channels because these are the major calcium channel subtypes that have been implicated in seizure disorders in both humans and animal models of epilepsy, in particular in the context of idiopathic generalized epilepsies.

## II. MOLECULAR PHYSIOLOGY OF VOLTAGE-GATED CALCIUM CHANNELS AND ASSOCIATED ANCILLARY SUBUNITS

### A. Subtypes and Physiological Roles of Voltage-Gated Calcium Channels

Voltage-gated calcium channels are key mediators of calcium entry into neurons in response to membrane depolarization. Calcium influx via these channels mediates a number of essential neuronal responses, such as the activation of calcium-dependent enzymes, gene expression (72, 93, 273), the release of neurotransmitters from presynaptic sites (247, 258, 302), and the regulation of neuronal excitability (217). The nervous system expresses a number of different calcium channels with unique cellular and subcellular distributions and specific physiological functions. They have been classified into two major categories (41): low voltage-activated (LVA) calcium channels (i.e., T-type channels) and high voltage-activated (HVA) channels, although this classification should not be applied rigidly, as some of the HVA channel subtypes can, under certain circumstances, be activated at relatively negative voltages (308). As outlined in greater detail below, LVA channels are activated by small depolarizations near typical neuronal resting membrane potentials and are key contributors to neuronal excitability. Their functional identification is aided by their sensitivities to blockers such as nickel ions (95, 161), mibefradil (190), and the scorpion venom-derived peptide kurtotoxin (49, 250); however, the above blockers cannot be considered as truly selective for T-type channels. HVA channels require larger membrane depolarizations to open and can be further subdivided, based on pharmacological and biophysical characteristics, into L-, N-, R-, P-, and Q-types. L-type channels are slow to activate and inactivate with barium as the charge carrier and are defined by their sensitivities to dihydropyridine agonists and antagonists (95). They are typically found on cell bodies where they participate, among other functions, in the activation of calcium-dependent enzymes and in calcium-dependent gene transcription events (8, 72, 298). N-type channels produce inactivating currents that are selectively and potentially inhibited by  $\omega$ -conotoxins GVIA and MVIIA, two peptides isolated from fish-hunting marine snails (2, 91, 212, 230). P- and Q-type channels are identified by their

differential sensitivities to the American funnel web spider toxin  $\omega$ -agatoxin IVA (2), and like N-type channels, they are concentrated at presynaptic nerve terminals where they are linked to the release of neurotransmitters (300, 301). In the context of neurotransmitter release, N-type and P/Q-type channels do not appear to be created equally, as N-type channels tend to support inhibitory neurotransmission, whereas the P/Q-type channels have more frequently been linked to the release of excitatory neurotransmitters but can also support inhibitory release (32, 33, 76, 163, 225). R-type channels were originally termed as such because of their resistance to the above blockers (229). These channels are rapidly inactivating and activate at somewhat more hyperpolarized potentials compared with the other HVA calcium channel subtypes (257). They can potentially, albeit not totally selectively, be inhibited by SNX-482, a peptide toxin isolated from tarantula venom (23, 206). R-type channels are distributed in proximal dendrites and presynaptic nerve termini (288, 311, 316). Their precise physiological function remains enigmatic; however, there is evidence that these channels underlie carbachol-dependent plateau potentials in hippocampal CA1 neurons (152) and may mediate neurotransmitter release at select synapses (296).

### B. Molecular Structure of Voltage-Gated Calcium Channels

HVA calcium channels are heteromultimers that are formed through association of  $\alpha_1$ -,  $\beta$ -,  $\alpha_2\delta$ -, and  $\gamma$ -subunits (Fig. 2). In contrast, LVA channels may contain only the  $\alpha_1$ -subunit; however, this remains an area of controversy. While effects of ancillary subunit coexpression on T-type channel function have been reported in certain expression systems (75, 79, 121), antisense knockdown of calcium channel  $\beta$ -subunits in neurons does not appear to affect T-type channel function (155, 169). Moreover, a biochemical association between T-type channel  $\alpha_1$ - and ancillary subunits has not been demonstrated (79).

The principal pore-forming subunit of both LVA and HVA calcium channels is the  $\alpha_1$ -subunit, which contains the key structural moieties that are required for a functional calcium channel, and which is the sole determinant of the calcium channel subtype. To date, nine different types of neuronal calcium channel  $\alpha_1$ -subunits have been identified and shown to fall into three major classes:  $\text{Ca}_v1$ ,  $\text{Ca}_v2$ , and  $\text{Ca}_v3$  (Fig. 2). The  $\text{Ca}_v1$  family encodes different isoforms of L-type channels (149, 193, 197, 282, 304). Among the  $\text{Ca}_v2$  family, alternate splice isoforms of  $\text{Ca}_v2.1$  encode P- and Q-type channels (22),  $\text{Ca}_v2.2$  represents N-type channels (80), and  $\text{Ca}_v2.3$  corresponds to R-type channels (229, 257, 305) (for review, see Ref. 254). Finally, the  $\text{Ca}_v3$  family represents three different types of T-type channels (i.e.,  $\text{Ca}_v3.1$ ,  $\text{Ca}_v3.2$ , and  $\text{Ca}_v3.3$ ) with




### Ancillary subunits

$\beta_1, \beta_2, \beta_3, \beta_4$

$\gamma_1$  through  $\gamma_8$

$\alpha_2\text{-}\delta_1$  through  $\alpha_2\text{-}\delta_4$



### Neuronal $\alpha_1$ subunits

**HVA**  $\text{Ca}_v1.2$  } L-type  
 $\text{Ca}_v1.3$  }  
 $\text{Ca}_v1.4$  }

$\text{Ca}_v2.1$  P/Q-type  
 $\text{Ca}_v2.2$  N-type  
 $\text{Ca}_v2.3$  R-type

**LVA**  $\text{Ca}_v3.1$  } T-type  
 $\text{Ca}_v3.2$  }  
 $\text{Ca}_v3.3$  }

FIG. 2. Subunit assembly and subtypes of voltage-gated calcium channels. Graphic representation of the high voltage-activated calcium channel complex consisting of the main pore forming  $\alpha_1$ -subunit plus ancillary,  $\beta$ -,  $\gamma$ -, and  $\alpha_2\text{-}\delta$ -subunits. Low voltage-activated calcium channels may consist of only the  $\alpha_1$ -subunit (not separately shown). Different neuronal  $\alpha_1$ -subunits correspond to different calcium channel isoforms identified in native neurons.

distinct kinetic properties (43, 56, 145, 160, 194, 198, 199, 216). These different types of calcium channel  $\alpha_1$ -subunits support specific physiological functions, as evident from findings obtained from calcium channel knockout mice (Table 3), and by studying genetic abnormalities in calcium channels in disease states (for review, see Ref. 219).

### C. Structural Basis of Calcium Channel Function

To provide a context for the effects of mutations in calcium channel genes that have been linked to the etiologies of neurological disorders such as epilepsy, we will briefly touch on the structural determinants of calcium

channel function. The  $\alpha_1$ -subunits are comprised of four major transmembrane domains that are structurally homologous to those found in voltage-gated sodium and potassium channels (reviewed in Refs. 39, 40; see Fig. 2). Each domain consists of six transmembrane helices, with intracellularly localized  $\text{NH}_2$  and  $\text{COOH}$  termini. The fourth membrane-spanning  $\alpha$ -helix (S4 segment) contains positively charged arginine and lysine residues every three to four amino acids. Mutagenesis and crystallization studies involving potassium channels have confirmed early suggestions that this region forms the voltage sensor (38), a structural element that translocates within the membrane in response to changing membrane potential

TABLE 3. Phenotypic consequences of calcium channel  $\alpha_1$ - and ancillary subunit knockout studies

Channel Subunit	Phenotype	Reference Nos.
$\text{Ca}_v1.1$	Mice die at birth due to asphyxiation, due to lack of skeletal muscle contraction (including diaphragm).	270
$\text{Ca}_v1.2$	Phenotype is lethal, mice die around day 12 postcoitum, isolated myocytes spontaneously beat until day 14 postcoitum.	248
$\text{Ca}_v1.3$	Mice are deaf due to loss of cochlear synaptic transmission and also exhibit cardiac arrhythmias.	223
$\text{Ca}_v1.4$	Mice are blind due to loss of rod vision.	185
$\text{Ca}_v2.1$	Mice show ataxia and absence seizures and die ~4 wk after birth.	135, 256
$\text{Ca}_v2.2$	Mice are hyposensitive to neuropathic and inflammatory pain, show reduced levels of anxiety, and show altered state of vigilance and reduced ethanol consumption.	16, 112, 128, 142, 207, 239
$\text{Ca}_v2.3$	Mice exhibit altered sensitivity to pain.	240
$\text{Ca}_v3.1$	The phenotype is behaviorally normal, mice are resistant to certain types of seizures including other genetic models of absence seizures and can rescue the epileptic phenotype of $\text{Ca}_v2.1$ mice. They also exhibit altered sleep structure. These mice may experience hyperperception of visceral pain.	143, 144, 256
$\text{Ca}_v3.2$	Mice are unusually small and show developmental deformities (i.e., trachea). Also, relaxation of vascular smooth muscle is impaired giving rise to deficient control of blood pressure.	44
$\beta_1$	Mice die at birth due to lack of skeletal muscle contraction.	106, 270
$\beta_2$	Embryonic lethal.	9
$\beta_3$	Mice appear visibly normal.	58, 205
$\beta_4$	De facto knockout due to premature stop; mice show absence seizures and ataxia.	31
$\alpha_2\text{-}\delta_2$	De facto knockout due to premature stop; mice have absence seizures.	26
$\gamma_1$	Mice appear normal.	96
$\gamma_2$	De facto knockout due to premature stop; mice have absence seizures and other features such as ataxic gait, paroxysmal dyskinesia.	166

(77, 132, 176) and mediates channel opening. The reentrant P-loops between S5 and S6 segments are believed to line the pore of the channel to allow for the passage of permeant ions. The P-loops each contain a critical glutamic acid residue (4 in total) that together form a ring of negative charge and allow for selectivity of divalent cations over monovalent ions (82, 307, 309). However, other amino acid residues in the outer vestibule of the pore are likely to contribute to ion permeation and selectivity (92, 242). The  $\alpha_1$ -subunits also contain the key structural elements that allow the channel to inactivate upon prolonged membrane depolarization. Voltage-dependent inactivation, a key feature that regulates availability of a calcium channel to open, is thought to involve a physical occlusion of the channel pore by a cytoplasmic gating particle. This inactivation gate appears to be formed at least in part by the cytoplasmic domain I-II linker region. However, other regions of the channel, in particular the S6 helices, are also important (reviewed in Ref. 267), perhaps by forming a docking site for the inactivation gate (268). The COOH terminus region of the  $\alpha_1$ -subunit contains the key loci responsible for calcium-dependent inactivation, a process that is observed only when calcium is used as the charge carrier. Originally thought to occur only with L-type channels, more recent evidence indicates that all types of high voltage-activated channels are subject to this type of regulation by calcium ions. Mechanistically, this process involves a dynamic interaction of the calcium binding protein calmodulin with the COOH-terminal region of the  $\alpha_1$ -subunit (158, 170, 218, 320).

The cytoplasmic linker regions of various types of calcium channel  $\alpha_1$ -subunits form key interaction sites for regulatory proteins and may be substrates for phosphorylation events. For example, the I-II linker regions of  $\text{Ca}_v2.1$  and  $\text{Ca}_v2.2$  subunits interact with G protein  $\beta\gamma$ -subunits (for review see Refs. 70, 74, 315); the II-III linker regions of these two channels interact with a number of synaptic proteins (for review, see Refs. 40, 259). The domain II-III linker region of  $\text{Ca}_v3.2$  interacts with  $\text{G}\beta_2$ -subunits (306) and is a substrate for calmodulin-dependent protein kinase phosphorylation (299). These are but a few examples, as the list of calcium channel interacting proteins is too extensive to comprehensively review here. Nonetheless, in the context of epilepsy-associated genetic mutations, it is important to consider that their physiological effects may manifest themselves by interfering with calcium channel regulatory mechanisms, without necessarily causing alterations in the biophysical properties of the channels per se.

#### D. Ancillary Subunits of Calcium Channels

While calcium channel  $\alpha_1$ -subunit is sufficient to form functional channels, the association of HVA channel

$\alpha_1$ -subunits with ancillary  $\beta$ - (73),  $\alpha_2\text{-}\delta$ - (5, 35, 147), and  $\gamma$ -subunits (18) is known to result in altered functional properties and/or increased plasma membrane expression.

The  $\alpha_2\text{-}\delta$ -subunit is translated as a single gene product, and then posttranslationally cleaved into a membrane-spanning  $\delta$ -peptide (27 kDa) and extracellular  $\alpha_2$ -peptide (143 kDa) (60), which are subsequently relinked via disulfide bonds. Expression studies have shown that the  $\alpha_2\text{-}\delta$ -protein alters current kinetics and current densities when coexpressed with HVA  $\alpha_1$ -subunits, although to different degrees with different channels (147, 310). The  $\alpha_2\text{-}\delta$ -subunit is the only known calcium channel ancillary subunit to interact with a clinically active drug compound, in that its association with gabapentin mediates analgesia (251). To date, four genes encoding for different  $\alpha_2\text{-}\delta$ -proteins ( $\alpha_2\text{-}\delta_1$ ,  $\alpha_2\text{-}\delta_2$ ,  $\alpha_2\text{-}\delta_3$ ,  $\alpha_2\text{-}\delta_4$ ) have been identified, with several potential additional splice variants (146).

Four different types of  $\beta$ -subunits ( $\beta_1$  through  $\beta_4$ ), along with various splice variants (reviewed in Refs. 73, 232), have been identified and characterized. Unlike  $\alpha_2\text{-}\delta$  proteins,  $\beta$ -subunits appear to be exclusively cytosolic in nature, with the exception of  $\beta_{2a}$ , which can become palmitoylated and thus membrane anchored (90, 228). The  $\beta$ -subunit core resembles membrane-associated guanylate kinase homologs with conserved interacting SH3 and guanylate kinase (GK) domains (275). Residues in the GK domain participate in the formation of a hydrophobic groove that is involved in the high-affinity binding to a conserved region within the domain I-II linker region of HVA calcium channels, termed alpha interaction domain (AID) (47, 291). Recently published crystal structure data have revealed that binding of the  $\beta$ -subunit to the channel is critically dependent on a functional association of the SH3 and GK regions (47, 213, 291), which is stabilized by the beta interaction domain, a short amino acid stretch that was originally thought to directly interact with the calcium channel AID region (291). The functional consequences of  $\beta$ -subunit coexpression include increased plasma membrane expression of the  $\alpha_1$ -subunit (17, 48, 226) as well as altered channel kinetics (90, 310) (reviewed in Ref. 5). Consistent with an important role in calcium channel function, knockout of individual  $\beta$ -subunit genes results in severe physiological consequences (see Table 3).

Eight different types of  $\gamma$ -subunits have been identified ( $\gamma_1$  through  $\gamma_8$ ) (5). The  $\gamma$ -subunits are comprised of four transmembrane domains with intracellular  $\text{NH}_2$  and COOH termini (Fig. 2), but the mutual sites of interaction with the  $\alpha_1$ -subunit remain unknown. While it has been observed that some of the  $\gamma$ -subunits have the propensity to alter the functional characteristics of HVA calcium channels (238) (reviewed in Ref. 18), the precise action of these subunits on HVA calcium channels remains to be

completely understood, and there is evidence that these subunits can interact with other membrane proteins such as AMPA receptors (45).

Considering the important roles of these subunits in regulating calcium channel function, one may perhaps expect that naturally occurring mutations in calcium channel subunit genes may have the propensity to alter normal physiological responses in animals and humans. As we will outline below, this does indeed occur within the context of idiopathic generalized epilepsies.

### III. VOLTAGE-GATED CALCIUM CHANNELS AND ANCILLARY SUBUNITS IN IDIOPATHIC GENERALIZED EPILEPSY

#### A. T-Type Calcium Channels and Spike-Wave Seizures

##### 1. T-type channel physiology

T-type channels mediate a spectrum of physiological roles including pacemaker activity and various forms of physiological and pathophysiological neuronal oscillations (99, 217). Moreover, it was recently demonstrated that T-type channel activity can result in calcium-induced calcium release in neurons of the paraventricular nucleus of the thalamus and other midline neurons associated with the thalamocortical system (233). [Note that calcium-induced calcium release is known to occur in thalamocortical neurons but it does not involve the action of T-type channels (29).] This novel role for T-type channels in the thalamocortical system putatively lends itself to numerous calcium-dependent signaling pathways that extend the role of these channels beyond their biophysical attributes and into complex processes such as synaptic plasticity (237).

T-type channels display several unique biophysical features compared with HVA channels. They activate and inactivate near the resting membrane potential of neurons (approximately  $-60$  mV), display more rapid inactivation kinetics, and deactivate more slowly to produce pronounced tail currents (30). There is also an increased overlap in the voltage dependences of activation and inactivation, which gives rise to a phenomenon termed the "window current." At membrane voltages where the window current is operational, a small fraction of T-type channels never fully inactivates and can be rapidly recruited for calcium influx upon depolarization. Interactions between the window current and the  $K^+$  leak current in thalamic (reticular and thalamocortical) and possibly cortical neurons can result in membrane bistability (57). This T-type-mediated potential can interact with other active currents (e.g.,  $I_h$ ) to modulate neuronal output between nonoscillatory and oscillatory modes that

have different physiological correlates such as the slow ( $<1$  Hz) sleep rhythm (57, 124). In response to a membrane hyperpolarization, a large fraction of T-type channels is recovered from the inactivated state, thus priming them for opening events. The ensuing calcium entry is thought to mediate a low-threshold calcium potential, a transient membrane depolarization that is sufficiently large to trigger a succession of sodium spikes (62, 157, 188, 245, 314). This feature is commonly referred to as "rebound bursting" and has been shown to occur in various neuronal subtypes including thalamocortical relay neurons, thalamic reticular neurons (54, 69, 126, 174), and a subpopulation of neocortical cells (61).

##### 2. T-type calcium channel distribution

It is important to note that the precise biophysical characteristics of T-type calcium channels in relation to their physiological roles vary with  $Ca_v3$  channel subtype and can be dramatically affected by alternate splicing of a given T-type channel gene (203). Their unique gating kinetics in addition to their specific regional distribution allows for these channels to be used by different neuronal subtypes to generate a variety of network dynamics. The exact distribution and protein expression levels of T-type calcium channels in the brain are not well understood, with most of the current knowledge having been derived from *in situ* hybridization studies in rat brain slices (55, 138, 276). Although T-type channel mRNA has been detected across all brain regions including the cerebellum, we focus predominantly on the major brain structures thought to be involved in spike-wave seizures, i.e., the thalamus and neocortex. All three T-type isoforms are expressed at relatively high levels in the hippocampus (276), with  $Ca_v3.1$  being the dominantly expressed subtype in thalamocortical neurons (276). In contrast,  $Ca_v3.2$  and  $Ca_v3.3$  appear to be expressed more prominently in the thalamic reticular nucleus (276). Cells in this nucleus are primarily GABAergic interneurons that play a key role in synchronizing thalamic outputs onto thalamocortical neurons that then project to the cortex (262). Diffuse mRNA levels have been detected for  $Ca_v3.1$  and  $Ca_v3.3$  in most cortical areas (276). Transcript levels of  $Ca_v3.2$  in the cortex typically occur at lower levels, except in layer V cortical pyramidal neurons (276). It should be noted, however, that it is not clear whether presence of mRNA correlates well with protein expression levels in the plasma membrane, and what particular splice isoform of the channel may be present (105).

In most types of neurons, the specific subcellular distribution of T-type channels has not yet been identified by immunocytochemical means. However, concordant evidence from numerous electrophysiological and functional imaging studies suggests that they are located both on and near the soma (136), and generally at more distal

dendritic sites (129, 137, 139, 182, 188, 200). Moreover, a recent study using electrophysiological and pharmacological investigation of reticular cells in thalamic brain slices indicates the presence of a slowly inactivating T-type current in the dendrites and a fast inactivating current at the soma (133). Pharmacological manipulations have suggested that the fast inactivating current is likely to be carried by  $\text{Ca}_v3.2$  channels, whereas the slow T-type current may be mediated by  $\text{Ca}_v3.3$  channels. These results are supported by data from *in situ* hybridization studies for reticular neurons where mRNA expression of both isoforms has been detected (276). The subcellular distribution (e.g., soma vs. dendrites) is likely subject to further heterogeneity when considering the different T-type channel isoforms, splice variants, and developmental changes in the lifetime of the organism. For example, previous reports in reticular thalamic neurons have reported differences in the rates of inactivation of T-type currents in intact slices (66) (slow) compared with acutely dissociated neurons (126) (fast).

### 3. T-type channels and the thalamocortical network

The thalamocortical circuit is the primary link between peripheral sensory systems and the cerebral cortex. This circuitry is also one of the most studied in the context of neurophysiological rhythm generation and encompasses structures responsible for the regulation of brain states such as arousal and slow-wave sleep, a hallmark thalamocortical oscillation (263). From a simplistic anatomical perspective, this circuit consists of a network of reticular, thalamocortical (also referred to as relay), and neocortical neurons. Cortical neurons directly innervate reticular and thalamocortical neurons, whereas the reticular neurons provide GABAergic projections onto each other and onto thalamocortical neurons, which in turn synapse onto neocortical neurons (Fig. 3). Indeed, although the thalamus receives many sensory inputs, the primary source of excitatory synapses onto it comes from the cortex (86, 87, 172, 173), which exemplifies the influence of neocortical activity on thalamic information processing. Both neocortical and thalamocortical cells have excitatory projections back onto reticular neurons, and the activity of the circuit is further regulated via thalamic (local circuit) and neocortical interneurons (189, 249). The intraconnectivity between reticular neurons is believed to be a form of lateral inhibition (286) and central to the role of this local network in modulating overall thalamic outputs to cortical areas during both physiological and seizure activity (100, 255).

Thalamic neuronal firing can be broadly categorized into tonic-mode and burst-mode firing. The latter involves the action of T-type calcium channels in the thalamocortical circuitry and is highlighted by their contribution to the propagation of thalamically generated spindles to the neocortex. Bursting in reticular neurons is mediated by

low-threshold calcium potentials (126) in response to corticofugal volleys from neocortical inputs (52). The bursts are preceded by prolonged hyperpolarizing potentials (263) that are triggered by the activity of G protein-coupled inward rectifier potassium channels (101) and lead to the activation of T-type currents in dendrites of reticular neurons (126). Modeling studies have demonstrated a sensitive link between the compartmental distribution of T-type conductances and the ability of thalamocortical and, more so, reticular neurons to fire in burst mode (69). Specifically, for a fixed burst threshold in a model of reticular neurons, GABAergic conductances are required to be relatively larger if the T-type current is localized to the soma versus dendrites.

The low-threshold calcium potential-induced bursts of action potentials occur in a subpopulation of reticular neurons and are manifested as tonic firing with burstlike modulation due to membrane bistability (100). The GABAergic activity of reticular neurons (involving both  $\text{GABA}_A$  and  $\text{GABA}_B$ ) results in hyperpolarization and subsequent low-threshold calcium potential-mediated rebound bursting in thalamocortical neurons, which faithfully track the bursts in the reticular network and manifest themselves as sleep spindles (263). There is some evidence from *in vivo* studies that burst-mode firing in thalamic neurons can occur during wakefulness (109, 110); however, this is not a common occurrence, and this interpretation may be complicated by an intermediate state of brain activity such as drowsiness (264).

Absence-type seizures occur preferentially during drowsiness or slow-wave sleep, which highlights the involvement of common synchronizing elements (such as the reticular network) underlying certain sleep and seizure states. However, unlike the state when sleep spindles are generated (101), the presence and interaction with the cortex is critical for the generation of SWDs (266). During cortically generated SWDs the GABAergic reticular neurons faithfully follow the cortical bursting activity and are synchronized with the activity recorded in the cortex (Fig. 4). In contrast, thalamocortical neurons undergo a sustained hyperpolarization in response to phasic inhibitory postsynaptic potentials (IPSPs), which are however incapable of recovering a sufficiently large fraction of T-type channels. Consequently, thalamocortical neurons do not exhibit rebound bursts during the epoch when SWDs are observed in the cortex (265) (Fig. 4). Reticular neurons are thus driven by the activity in the cortex, which leads to the inhibition of thalamocortical neurons, and prevents the feedback to cortical areas. Therefore, in one possible scenario, increased T-type channel (i.e.,  $\text{Ca}_v3.2$  and  $\text{Ca}_v3.3$ ) activity could result in increased burst-mode firing in thalamic reticular neurons. The persistent activity of reticular neurons during SWDs causes increased membrane conductance in thalamocortical neurons in the form of steady inhibition, which may perhaps explain the



## Thalamocortical Circuit




FIG. 3. The simplified thalamocortical circuit involved in the generation of spike-wave discharges. *Left:* neuronal circuitry implicated in the generation of spike-wave seizures. For simplicity, local-circuit interneurons in the thalamus and cortex are not illustrated. Neurons are indicated in the form of color-coded circles that correspond to neuronal phenotypes obtained from staining experiments in cats that were used in modeling studies (51, 66, 67). Thalamic relay neurons (green) receive sensory inputs and project onto cortical pyramidal neurons in layers III/IV and V/VI in the cerebral cortex (blue). Sensory inputs can also project onto thalamic inhibitory interneurons (black, thalamus) that can in turn modulate the firing of relay neurons. Thalamocortical relay neurons can also synapse onto cortical inhibitory interneurons (black, cortex). Corticothalamic projections from layer VI of the cortex can reciprocally synapse onto relay neurons in addition to neurons in the thalamic reticular nucleus (red). The GABAergic reticular neurons are highly interconnected both via chemical and electrical synapses and form a pacemaker subcircuit within the thalamus. Both reticular and relay neurons express T-type calcium channels and can exhibit rebound bursts.

unconsciousness during absence seizures caused by inhibition of synaptic transmission of signals from the outside world (263). It should be noted that there are a large number of modeling studies in which underlying ionic currents and network dynamics are explored in the context of both physiological and epileptiform thalamocortical oscillations. Although review of these works is beyond the scope of this manuscript, a concise review by Destexhe and Sejnowski (68) is available.

The rhythmic activity that is observed in EEG recordings during an epileptic seizure is a reflection of cortical and thalamic network interactions. In a number of rodent models of epilepsy, the frequencies of SWDs are typically faster than the 3–4 Hz observed with the common form of human absence seizures, which may hint at interspecies differences in these network interactions, or perhaps somewhat different pathways for seizure generation. Indeed, the frequency of SWDs in these rodent models is

complicated by the notion that some of the observed thalamocortical oscillations may be part of normal physiological activity (221, 303). Historically, there has been much debate with regard to the site of initiation for SWD-based seizures (reviewed in Refs. 195, 263); specifically, the question pertaining to which structure, cortex versus thalamus, is the key anatomical substrate for initiating SWDs. Initial evidence for the involvement of both structures came from human studies undergoing invasive intracranial EEG monitoring (reviewed in Ref. 21), which is now deemed unethical for generalized forms of epilepsy due to a consensus that large networks in the brain are involved. However, a body of evidence from *in vivo* experiments in rats and cats suggests that the cortex may be the minimal substrate for generating spike-wave seizures (reviewed in Ref. 263). Moreover, recent studies utilizing multisite recordings in two accepted rat genetic models of absence epilepsy, the Genetic Absence Epilepsy Rat from




FIG. 4. Firing patterns of thalamic and cortical elements in animal models of spike-wave seizures. *A*: simultaneous extracellular and intracellular recording of absence seizure activity in an adult genetic absence epilepsy rat from Strasbourg (GAERS). The intracellular recording is from a thalamic reticular neuron, which exhibits bursting activity in synchrony with spike-wave discharges recorded extracellularly. [Adapted from Slaght et al. (253).] *B*: simultaneous electroencephalographic (EEG), extracellular field, and intracellular recordings of cortical and thalamic neurons during an epoch of spontaneous polyspike-wave seizure activity in a cat under ketamine-xylazine anesthesia. Cortical and thalamocortical (in the ventrolateral nucleus, VL) recordings were obtained simultaneously, and reticular neurons were recorded later and superimposed due to the repeatability of the oscillation. Cortical neurons exhibit bursting during each of the paroxysmal depolarizing shifts (191). This activity is closely tracked by reticular neurons that exhibit T-type mediated bursting activity. The GABAergic influence of reticular neurons on thalamocortical cells can be seen by the series of inhibitory postsynaptic potentials (see dots), which result in tonic hyperpolarization of these neurons for the duration of the spike-wave activity. Similar observations have been reported during spike-wave seizures in the GAERS rat (222). Whether thalamocortical neurons exhibit rebound spikes is a function of their membrane potential caused by inputs from reticular and cortical neurons. A large fraction of these cells, however, cannot fire, which is believed to result in disruption of information flow to the cortex resulting in the absence phenotype during seizures. [Adapted from Steriade (263) and Steriade and Contreras (265).]

Strasbourg (GAERS) and the Wistar Albino Glaxo rats, bred in Rijswijk (WAG/Rij), have suggested that neocortical cell firing temporally precedes (on the order of milliseconds) the activity in the thalamus (195, 221). Taken together, these studies imply that complex interactions involving the entire thalamocortical network are necessary in generating rhythmic activity under both physiological and pathophysiological states, with T-type channels playing a key role in modulating the firing properties of neuronal elements.

### 3. Genetic animal models

Animal studies have provided insights as to whether T-type expression is altered via epileptic activity, or

whether increased T-type expression is a requirement for the development of seizure activity. For example, the GAERS rat exhibits 7- to 11-Hz SWDs, usually after 30 days of life. In this rat model of absence epilepsy, an increase in T-type currents in reticular neurons has been reported after the second postnatal week (284). This increase in T-type currents is putatively mediated by elevated  $Ca_v3.2$  expression in reticular neurons, which as the animal develops to exhibit absencelike seizures, is also accompanied by elevated expression of  $Ca_v3.1$  in relay neurons (277). This suggests that an increase in T-type channel expression can precede the development of seizures, perhaps by altering network dynamics via mecha-

nisms that affect potentiation. In support of this notion, a modeling study employing a strictly thalamic network has demonstrated differential effects on network activity in response to an augmented T-type conductance in reticular neurons (279). Although rebound spiking number in individual reticular neurons was relatively unchanged, the increase in T-type conductance resulted in increased network synchrony by introducing a phase-lag between reticular and thalamocortical firing. It should be noted that a similar effect on synchrony was observed for increased calcium-activated potassium conductance in the model. Nonetheless, a degree of caution is warranted when attempting to interpret these results in relation to *in vivo* recordings that take place in the context of larger and more complexly connected networks. Presently, a precise causal connection between increased T-type channel activity and subsequent development of seizures remains to be established. It is conceivable that developmental causes involving either modulation of the channels, their redistribution with other isoforms, and alternate splicing may be contributing factors. Indeed, such developmental effects occur in the heart where  $\text{Ca}_v3.1$  and  $\text{Ca}_v3.2$  undergo differential developmental expression (208). Such alterations resulting in an epileptic phenotype represent gradual neurophysiological changes over time. In converse to these more slowly developing processes of epileptogenesis, it has been shown that even a single episode of status epilepticus in rat hippocampal brain slices can result in a selective increase in T-type channel activity, indicating that epileptic seizures *per se* may have the propensity to rapidly alter T-type currents (271), thus potentially lowering the threshold for subsequent seizure events.

Recent *in vivo* studies involving T-type ( $\text{Ca}_v3.1$ ) knock-out (KO) mice have provided additional insights into the role of T-type channels in the generation of SWDs and absencelike seizure episodes (144). Ablation of the  $\text{Ca}_v3.1$  gene in mice abolishes rebound spiking in dissociated adult thalamocortical neurons but does not alter their ability to fire tonically. Deep thalamic recordings from  $\text{Ca}_v3.1$  KO mice also revealed a lack of synchronized activity. *In vivo* recordings from the cortical surface in these mice demonstrated a resistance to baclofen-induced 3- to 5-Hz SWDs that could be induced in control animals. This resistance could be due to a loss of rebound bursting in thalamocortical neurons, which are known to express high levels of  $\text{Ca}_v3.1$  channels and to receive strong GABAergic inputs. Intriguingly, these mice are not resistant to SWDs induced by bicuculline, and tonic-clonic seizures were inducible in both KO and control animals with 4-aminopyridine injection. A possible explanation of why these mice experience bicuculline-induced seizures may be due to cortical involvement. Indeed, previous work has shown that the isolated cortex is able to generate bicuculline-induced SWDs, suggesting that the

cortex can act as the seizure-generating zone (266). An alternative explanation for lack of protection against seizures in some models versus others may involve the reticular neurons that are known to predominantly express  $\text{Ca}_v3.2$  and  $\text{Ca}_v3.3$  T-type channels rather than  $\text{Ca}_v3.1$ . Therefore, it is conceivable that other T-type calcium channel isoforms (for example,  $\text{Ca}_v3.2$ ) may be involved in the generation of hypersynchronous activity in some models of epilepsy without requiring the action of  $\text{Ca}_v3.1$  channels in thalamocortical neurons. The findings obtained with  $\text{Ca}_v3.1$  KO mice also indicate that full-blown convulsive seizures may recruit different mechanistic pathways. Moreover, consistent with the functional link between T-types and intrathalamic rhythm generation,  $\text{Ca}_v3.1$  KO mice show altered sleep architecture, with markedly diminished thalamic delta (1–4 Hz) waves and sleep spindles (7–14 Hz) (159). Intriguingly, the slow (<1 Hz) rhythms that can be sustained by the cortex remained relatively intact in KO animals.

More recently, it has been demonstrated that  $\text{Ca}_v3.1$  KO rescues the epileptic phenotypes seen in a number of murine models of absence seizures, including  $\text{Ca}_v2.1$  knockout mice, which (as we discuss in detail in sect. *mBI*) display severe absence seizures (256). Thalamocortical neurons isolated from  $\text{Ca}_v2.1$  KO mice were shown to exhibit an ~50% increase in T-type currents (predominantly due to  $\text{Ca}_v3.1$ ). In contrast,  $\text{Ca}_v2.1^{-/-}/\text{Ca}_v3.1^{+/-}$  mice exhibited a 25% reduction in T-type currents, yet they continued to exhibit SWDs with no changes in seizure frequency or duration. Thalamic mRNA transcript levels for  $\text{Ca}_v3.1$  were not different between  $\text{Ca}_v2.1^{+/+}$  and  $\text{Ca}_v2.1^{-/-}$  mice. These findings suggest that, under certain conditions, even reduced levels of T-type channel activity are capable of sustaining SWDs. Overall, these results indicate that complete KO of  $\text{Ca}_v3.1$  channels can play a protective role against SWDs in these genetic models of absence epilepsy, as well as in a subset of pharmacological seizure models (Fig. 5).

#### 4. T-type channel mutations in epileptic patients

Ion channel defects are considered to be one of the primary etiological causes of idiopathic generalized epilepsy (244). T-type channels have always been likely candidates due to their eminent presence in cortical and thalamic structures and their established physiological role in modulating neuronal firing. Moreover, clinically active antiepileptic drugs have been associated with T-type calcium channel inhibition (see below). However, although long suspected, a direct link involving T-type channels and the generalized spike-wave epilepsies in humans was only recently established. A number of missense mutations have now been identified in the  $\text{Ca}_v3.2$  calcium channel gene in patients diagnosed with childhood absence epilepsy and other forms of idiopathic gen-




FIG. 5. Ablation of  $\text{Ca}_v3.1$  T-type calcium channels can rescue several genetic models of absence seizures from the epileptic phenotype. Electroencephalographic recordings (15 s) are shown from the cortex of  $\text{Ca}_v2.1^{-/-}$ ,  $\text{Ca}_v2.1^{\text{tg/tg}}$  (tottering),  $\beta 4^{\text{lh/lh}}$  (lethargic), and  $\gamma 2^{\text{stg/stg}}$  (stargazer) mice with or without the ablation of  $\text{Ca}_v3.1$  channels, during spike-wave discharges corresponding to absence seizures (asterisks). In each genetic model, a complete ablation ( $\text{Ca}_v3.1^{-/-}$ ) is required to rescue the animal from the epileptic phenotype. [Adapted from Song et al. (256).]

eralized epilepsy (Fig. 6). The first genetic study involved 118 patients with childhood absence epilepsy, and point mutations were detected in 14 of these patients, but not in 230 control individuals (46). Intriguingly, none of the affected individuals had a family history of epilepsy, but carriers inherited a mutant copy of the gene in a heterozygous manner from one parent. A large number of polymorphisms were also reported that were identified in both childhood absence epilepsy and control groups. Of the identified 12 mutations, only two were found in more than one affected individual. A second study investigated a heterogeneous population of patients with idiopathic gen-

eralized epilepsy (including childhood absence epilepsy, juvenile myoclonic epilepsy, and febrile convulsions) and identified three additional missense mutations and one nonsense mutation that occurred in 9 of 192 individuals (118). None of these new mutations segregated with a specific epilepsy phenotype, and their presence was not associated with a single subtype of idiopathic generalized epilepsy. It is important to note that every single one of the reported  $\text{Ca}_v3.2$  mutations could also be found in seizure-free individuals.

The consequences of the entire set of missense mutations on  $\text{Ca}_v3.2$  channel function have recently been examined in transient expression systems (140, 141, 215, 292). Several of these mutations were found to result in small changes in the gating characteristics of both rat and human  $\text{Ca}_v3.2$  channels in a manner consistent with a gain of function (i.e., increased T-type channel activity). Specifically, some of the mutations resulted in a hyperpolarizing shift in the voltage dependence of activation, while others resulted in increased channel availability due to decreased steady-state inactivation. However, the majority of the mutations did not significantly affect the biophysical characteristics of the channel (141, 215), which is intriguing considering that most of the mutations are localized within the domain I-II linker of the channel, a region thought to be involved in voltage-dependent inactivation (at least in HVA calcium channels).

It should be noted that alterations in biophysical properties of mutant channels are not expected to be the sole mechanism for the generation of pathophysiology. It is conceivable that processes such as neuronal development, targeting of channels to appropriate subcellular compartments, and/or cumulative effects from multiple mutations can all contribute to the etiology of idiopathic generalized epilepsies. Moreover, as outlined earlier, it is well known that many intracellular proteins interact with voltage-gated calcium channels and perhaps a number of these “biophysically silent” mutations might, given their localization on intracellular linkers, disrupt important intracellular signaling pathways that bring about, or contribute to, the epileptic phenotype (see also NOTE ADDED IN PROOF). Ultimately, it may prove necessary to perform conditional and region-specific knockin animal studies (both in isolation and combinations thereof) to elucidate their exact functional role in seizure genesis. From a clinical point of view, it appears as if mutant channels with large biophysical changes compared with wild-type channels account for only a small fraction of affected individuals. This is supported by the low incidence of mutations that segregate with idiopathic generalized epilepsy phenotypes (34, 278). Nonetheless, genetically identified mutations in patients and investigation of their function have served to further implicate T-type channels as important players in the idiopathic generalized epilepsies.






FIG. 6. Transmembrane topology of the calcium channel  $\alpha_1$ - and associated ancillary subunits with locations of identified mutations linked to epilepsy. The pore-forming  $\alpha_1$ -subunit that is comprised of four homologous transmembrane domains, flanked by cytoplasmic  $\text{NH}_2$  and  $\text{COOH}$  termini and connected by intracellular linker regions. Each of the four domains consists of six transmembrane spanning  $\alpha$ -helices (termed S1 through S6, indicated by cylindrical segments) and a pore-loop structure that lines the permeation pathway for calcium (see reentrant loop between fifth and sixth transmembrane region in each domain). The S4 segments of each domain carry positively charged amino acid residues in every third position and are thought to form the voltage sensor of the channel (see "+" symbols for voltage sensor). Locations of mutations in  $\text{Ca}_v3.2$  that have been linked to idiopathic generalized epilepsy (circles, idiopathic generalized epilepsy) and epilepsy mutations in  $\text{Ca}_v2.1$  channels in animals and humans (triangles) are indicated. Calcium channel  $\beta$ -subunits are cytoplasmic proteins that tightly associate with the  $\alpha_1$ -subunit domain I-II linker region (232). The  $\alpha_2$ - $\delta$ -subunit is translated as a single protein, which is posttranslationally cleaved into  $\alpha_2$ - and  $\delta$ -subunits that are subsequently relinked via disulfide bonds (5, 147). The  $\delta$ -subunit contains a single transmembrane region connected to the extracellular  $\alpha_2$ -portion. The  $\gamma$ -subunits contain four transmembrane spanning helices flanked by cytoplasmic  $\text{NH}_2$  and  $\text{COOH}$  termini. Approximate locations of mutations in ancillary calcium channel subunits linked to absence seizures in mice and humans (triangles) are also indicated. CAE, childhood absence epilepsy.

Collectively, these findings support the notion that idiopathic generalized epilepsies comprise complex forms of epilepsy where numerous susceptibility alleles can appear and disappear in an individual's genome through factors such as meiotic reshuffling and chance events. Thus these alleles may not segregate in a Mendelian manner and in isolation do not appear to cause a sufficiently large perturbation to the functioning of brain networks to results in the disease phenotype. Instead, they could give rise to a genetic susceptibility by which their effects can combine with other alleles, perhaps involving the same or other proteins, to cause the epileptic condition in affected individuals (201). However, it is possible that future genetic linkage studies may identify calcium channel mutations that clearly segregate, in a Mendelian manner, with an epileptic phenotype.

## B. P/Q-Type Channel Defects and Spike-Wave Seizure Disorders

Central to the mechanisms of spindle and SWD generation are oscillatory rhythms, specifically in the tha-

lamic reticular network (99), which along with bursting activity in areas of the cortex affect the entire thalamocortical circuitry and modulate the firing of thalamocortical neurons (263). Although electrical synapses have recently been implicated in synchronizing the activities of reticular neurons (98, 156, 175), chemical synaptic transmission remains a primary mode of synchronizing the neuronal ensembles required for the generation of SWDs in both the thalamus and cortex. Because  $\text{Ca}_v2.1$  (P/Q-type) calcium channels are key mediators of synaptic transmission in most central and peripheral neurons, alterations in  $\text{Ca}_v2.1$  channel function therefore have the propensity to affect both cellular and network behavior.

### 1. $\text{Ca}_v2.1$ channels and murine models of absence epilepsy

The first evidence implicating  $\text{Ca}_v2.1$  channels in the generation of spike-wave seizures came from several mouse strains that showed absence-like seizure activity. The *tottering* (*tg*) mouse was originally identified based on observations of cerebellar ataxia and intermittent paroxysmal dyskinesia (211). Subsequent examinations re-

vealed the presence of EEG abnormalities that corresponded to absencelike seizures with SWDs in the range of 5–7 Hz (94, 211). Electrophysiological recordings have revealed that CA3 hippocampal pyramidal neurons in *tg* mice are prone to increased burst firing caused by paroxysmal depolarizing shifts when exposed to a high extracellular  $K^+$  model of in vitro seizures, consistent with hyperexcitability (114, 115). The *leaner* ( $tg^{la}$ ) mice display cortical spike-and-wave discharges that also resemble absence seizures. These mice are reported to be severely ataxic and often have a short survival postweaning (94, 177). Finally, the *Rocker* mice show spontaneous bilateral SWDs in the 6- to 7-Hz range while awake at a frequency of about one episode per minute, and which are always accompanied by behavioral arrest (321). These three mouse genotypes and their seizure disorders are recessively inherited and characterized by mild to severe ataxia and, in the case of  $tg^{la}$ , significant loss of cerebellar Purkinje and granular neurons (119).

All three mouse lines show defects in the gene encoding  $Ca_v2.1$  channels (11) (Fig. 6). In *tg* mice, a proline residue normally found in the domain II S5-S6 region of  $Ca_v2.1$  is replaced by leucine (78, 94). This amino acid substitution reportedly results in decreased levels of P/Q-type current levels in native mouse neurons, as well as in transient expression studies (293), whereas the gating characteristics of the mutant channel do not appear to be significantly altered. Given that the mutation is located near the pore region of the channel, this decrease in current activity could perhaps be due to reduced single-channel amplitude, although this has not been confirmed experimentally. The reduction of P/Q-type channel activity in *tg* mice is paralleled by a dysfunction in neurotransmitter release in neocortical neurons (7) and a switch to N-type channel-mediated synaptic transmission (227). Moreover, this aberrant P/Q-type channel function may also account for reduced evoked excitatory synaptic potentials evoked in ventrobasal thalamic neurons (33).

The  $Ca_v2.1$  calcium channel gene of  $tg^{la}$  mice contains a nucleotide substitution in the COOH-terminal region that results in its premature truncation (78, 94). It has been reported that the  $tg^{la}$  mutation causes altered P/Q-type channel currents and, like the *tg* mutation, reduces neurotransmitter release at neocortical synapses (7). It is unlikely that this mutation would affect single-channel conductance, suggesting that the reduced whole cell currents may either be due to a reduced probability of channel opening, or perhaps due to less effective targeting of the channel to the plasma membrane (i.e., reduced channel numbers). The *Rocker* mutation results in the replacement of a lysine residue (T1310K) in the domain III S5-S6 region with threonine (321) (Fig. 6). Its consequences on channel function in neurons or in transient expression systems remain to be determined, although the location near the pore site may again suggest possible effects on

ion permeation. The common theme in the genetic absence models of *tg* and  $tg^{la}$  mice appears to be a reduction in P/Q-type channel-mediated calcium entry, and hence synaptic release. This would be consistent with findings showing that complete genetic ablation of P/Q-type channels in mice results in absence seizures with behavioral arrests (256). However, it is important to note that the absence phenotype in these animal models does not occur in isolation, but rather is accompanied by other neurophysiological defects (i.e., ataxia, cerebellar atrophy), and this complicates our ability to link altered P/Q-type channel activity to the occurrence of seizures (see also sect. v).

## 2. $Ca_v2.1$ channelopathies in humans

In humans, a number of mutations in P/Q-type calcium channels have been associated with conditions such as episodic ataxia type 2 (64, 65, 97, 108, 131, 214, 246, 313) and familial hemiplegic migraine (214; reviewed in Ref. 220). In a few instances, patients carrying these mutations appear to exhibit absence seizures in addition to their ataxic phenotype. For example, a truncation mutation in the COOH-terminal region of  $Ca_v2.1$  has been identified in a juvenile patient diagnosed with episodic ataxia type 2 and afflicted with childhood episodes of absence epilepsy and primary generalized seizures (134). This mutation results in a nonfunctional channel (Fig. 6), which in turn promotes a dominant-negative inhibition of wild-type channels. The same group of investigators recently reported on a family in which five patients exhibit absence epilepsy in combination with cerebellar ataxia (127). A novel point mutation in domain I-S2 region of the  $Ca_v2.1$  calcium channel was identified, shown to segregate with the epileptic/ataxic phenotype, and to result in impairment of channel function. The significance of this study is the apparent association between absence seizures and reduced P/Q-type channel function, which is again consistent with findings in *tg* and  $tg^{la}$  mice. However, it is important to point out that the occurrence of absence seizures in episodic ataxia type 2 patients with P/Q-type channel mutations appears to be rare. Similarly, although familial hemiplegic migraine mutations in P/Q-type channels have been linked with the occurrence of seizures in several case studies (14, 81, 148), these seizures were not classified as absence. Hence, the majority of P/Q-type channel mutations identified in humans, unlike in the rodent counterparts, do not seem to give rise to absence epilepsy.

Considering that loss of P/Q-type channel activity can contribute to the development of absence seizures, one may perhaps speculate that the mutations that give rise to the familial hemiplegic migraine might not result in reduced P/Q-type currents. The functional effects of a number of the familial hemiplegic migraine mutations have been examined following transient expression in systems

such as *Xenopus* oocytes (150, 151), HEK293 cells (111, 283), cultured cerebellar granule neurons (283), and cultured hippocampal neurons (36); however, no clear-cut trend has emerged as to whether these mutations produce gains or loss of P/Q-type function. Most recently, the direct functional consequences of the four original familial hemiplegic migraine mutations (214) on excitatory (36) and inhibitory (37) synaptic transmission have been investigated. In the case of excitatory synaptic transmission, it was observed that all four mutant channels resulted in reduced current densities and diminished current influx for supporting synaptic transmission when compared with constituted wild-type channels in Ca<sub>v</sub>2.1 KO cultured hippocampal neurons (36, 37). In the case of inhibitory synaptic transmission, expression of these mutant channels in inhibitory neurons cultured from Ca<sub>v</sub>2.1 KO neurons reduced the contribution of these channels to supporting postsynaptic GABAergic currents compared with wild-type channels (37). The reduction in P/Q-type channel activity contrasts with the gain of channel function that is observed when certain familial hemiplegic migraine mutations were introduced into a mouse background (289). Given these conflicting results, it remains difficult to say whether the lack of absence seizures seen in familial hemiplegic migraine patients is related to the possibility that P/Q-type channel activity is increased, rather than the type of decrease observed in the case of most episodic ataxia type 2 mutations and in mouse genetic models of absence epilepsy (Fig. 6).

### C. Involvement of Ancillary Subunits of Voltage-Gated Calcium Channels in Seizure Disorders

Ancillary calcium channel subunits are important regulators of HVA calcium channel function. Murine and/or human mutations associated with seizure activity have been identified in all three classes of ancillary subunits (Fig. 6). A frame-shift mutation resulting in a functional knockout of the calcium channel  $\beta_4$ -subunit gives rise to the *lethargic* (*lh*) mouse phenotype, which is known to exhibit absence seizures and ataxia (31). In thalamic neurons isolated from these mice, excitatory, but not inhibitory, neurotransmitter release is reduced similar to what has been observed with *tg* mice (33). In humans, both missense and truncation mutations in the  $\beta_4$ -subunit have been found in idiopathic generalized epilepsy patients (88). Considering that the  $\beta_4$ -subunit is the primary subtype associated with P/Q-type channels, it is possible that these mutations may mediate their physiological effects by altering P/Q-type calcium channel function or targeting; however, since other types of calcium channel  $\alpha_1$ -subunits can also associate with  $\beta_4$ , such a link is not unequivocal. There may also be precedent for changes in calcium channel function via altered  $\beta$ -subunit expres-

sion patterns in (focal) temporal lobe epilepsy as has been shown in a single study (171).

To our knowledge, mutations in either  $\gamma$ - or  $\alpha_2$ - $\delta$ -subunits have so far not been linked to epilepsy in humans; however, there are several mouse phenotypes associated with these subunits. Two different mutations in the calcium channel  $\alpha_2$ - $\delta_2$  subunit (CACNA2D2 gene) result in loss of the full-length proteins giving rise to the *ducky* mouse, with the two identified forms being designated as *du* and *du*<sup>2J</sup>. Both mouse strains are characterized by behavioral arrest, bihemispheric spike-wave seizures with a frequency of ~5–7 Hz, and ataxia and paroxysmal dyskinesia (10, 210). Coexpression of the mutant  $\alpha_2$ - $\delta_2$ -gene in COS cells diminishes Ca<sub>v</sub>2.1 currents (26). Moreover, morphological differences have been observed in Purkinje cells from *du/du* mice, which show reduced size and complexity of dendritic processes rendering these neurons to appear as if they were developmentally immature (26). However, it is important to note that Purkinje cells are not lost as is the case for some of the other genetic epilepsy models that show cerebellar defects (e.g., *tg*<sup>la</sup> mice). Duplication of exon 3 of the  $\alpha_2$ - $\delta_2$ -subunit prevents the disulfide linkage between  $\alpha_2$  and  $\delta_2$ , the *entla* mouse (25). These mice show seizure activity in the 2- to 4-Hz range in cortical and hippocampal regions and exhibit reduced Ca<sub>v</sub>2.1 channel activity in the hippocampus. Although these mutations in the CACNA2D2 gene seem to result in reduced P/Q-type channel activity, the precise mechanism by which they result in absence-like seizure activity in the thalamocortical system remains open for investigation.

The spontaneous *Stargazer* (*stg*) and *Wagglers* mouse strains arise from disruptions in the calcium channel  $\gamma_2$ -subunit. These mice exhibit head tossing and absencelike seizures, with SWDs occurring about once per minute (166, 167). In *Wagglers* mice, seizures are dramatically exacerbated by targeted mutations in the calcium channel  $\gamma_4$ -subunit, suggesting that  $\gamma_4$  may compensate against some of the deleterious effects that result from compromised  $\gamma_2$ -function in these mice (168). As with the  $\alpha_2$ - $\delta$ -subunits, the *stg* mutation reportedly results in inhibition of P/Q-type calcium channel activity by a subtle reduction in channel availability (18). However, the  $\gamma_2$ -subunit also interacts with AMPA receptors (45), which are responsible for fast glutamatergic synaptic transmission and are involved in synaptic plasticity (24, 183). This interaction with the  $\gamma_2$  subunit is involved in the targeting of these receptors to synaptic sites (204, 290) and has recently been shown to modulate AMPA receptor function per se (281). Considering that AMPA receptors are known to undergo modification under epileptic processes (1), it remains unclear as to whether the physiological consequences of the *stg* mutation are due to actions on calcium channels, AMPA receptors, or both.



A role of HVA calcium channel interacting proteins is also exemplified in the case of juvenile myoclonic epilepsy patients that carry a mutation in the calcium binding protein EFHC1 that under normal conditions enhances R-type channel activity, an effect that is abolished by the mutations in this protein (274). However, mice deficient of R-type channels do not display seizure activity (162), and hence, it is not clear if the clinical manifestation of the EFHC1 mutation is directly related to the activity of R-type channels.

Taken together, a common theme in the effects of mutations in ancillary subunits appears to be an inhibition of P/Q-type channel activity, although it is by no means proven that their physiological effects are indeed mediated exclusively by these channels.

#### **D. Antiepileptic Drugs and Calcium Channel Pharmacology**

A number of clinically used antiepileptic drugs have been shown to block LVA and/or HVA calcium channels; however, it is not clear to what extent their inhibition is linked to the clinical efficacy of these drugs. A number of antiepileptic drugs are also used in the treatment of other nonepileptic neurological disorders including migraines and depression (236). Although many patients are seizure-controlled on antiepileptic drugs, their effectiveness can depend on the type of seizure disorder, their history of seizures before initiating treatment, and the efficacy achieved after first treatment (153, 154). The therapeutic benefit of antiepileptic drugs involves elevating seizure threshold for neurons and neuronal networks.

A number of antiepileptic drugs block calcium entry via HVA calcium channels. Gabapentin, a drug used in the treatment of generalized tonic-clonic seizures, is known to physically interact with the calcium channel  $\alpha_2\text{-}\delta$ -subunit (186) to block presynaptic calcium channel activity (13), although this mechanism of action does not seem to apply in all regions of the nervous system (27). Lamotrigine, a drug given to patients with absence, tonic-clonic, and partial seizures, has been shown to inhibit N- and P/Q-type channels (260, 297) without affecting T-type channels (318). However, due to the fact that GABA<sub>A</sub> receptors are also a major target for this drug, it is difficult to attribute its anticonvulsive effects unequivocally to calcium channel block. The antiepileptic drug levetiracetam may also block some HVA channels with its main effects exhibited on N-type channels (179). Recently, it was demonstrated that this compound can also effectively bind to the synaptic vesicle protein SV2A in brain-derived membrane and vesicle isolates (181). This interaction was found to be specific to the SV2A isoform (i.e., no binding was observed to its related proteins SV2B or C). Presently, it is not known if this compound could

mediate its effect on N-type channels indirectly via binding to SV2A. One of the many actions of the antiepileptic drug topiramate, commonly used to treat partial and generalized seizures, is to block cholinergic-dependent plateau potentials by blocking R-type channels (152). This might suggest that R-type channel inhibition could potentially protect against seizure activity. The common theme among some of these drugs is an inhibition of calcium channels at presynaptic sites, which is in some ways counterintuitive considering that knockout or decreased expression of P/Q-type channels gives rise to an epileptic phenotype, indicating that acute inhibition of these channels is not functionally equivalent to long-term gene knockdown.

In the case of LVA calcium channels, ethosuximide is an effective and commonly used antiepileptic drug for the treatment of absence seizures in the pediatric population (107). The pharmacological effects of this drug have been a matter of recent debate (125). Despite the known role of T-type channels in rebound bursting (specifically in the thalamus), it has been questioned as to whether the therapeutic effects of ethosuximide are derived from its action on these channels, since no block of LVA calcium currents in rat and cat thalamocortical neurons could be observed (165). On the other hand, in expression systems, ethosuximide reportedly blocks all three cloned human T-type channel isoforms (104). These different observations may perhaps be explained by the apparent state dependence of ethosuximide block that gives rise to an ~10-fold higher affinity inhibition of inactivated channels (note that T-type channels accumulate inactivation during prolonged bursting and spike-wave activity). Alternatively, the conflicting findings could be due to interspecies differences in channel structure in addition to putative cortical action of this drug (231, 266). The antiepileptic drug valproic acid is highly effective in the treatment of absence epilepsy (103). However, whether the clinical action of this drug is due to T-type channel block remains unresolved (235) given that studies have reported only a moderate amount of current block at therapeutic levels (243, 280).

Collectively, these findings indicate that HVA and LVA calcium channels are targeted by antiepileptic drugs; however, it is not clear to what extent their inhibition is linked to the clinical efficacy of these drugs.

#### **IV. CONCLUSIONS**

Epilepsy is one of the most prominent neurological disorders, and its spectrum of etiologies is a reflection of the many routes to seizure onset in brain networks. A growing body of evidence firmly establishes P/Q-type and T-type channels as important contributors to seizure genesis through modulation of neuronal properties that act to



shape network function, whereas other types of calcium channels do not appear to contribute to the development of seizure activity. Collectively, clinical and basic science studies point to new and exciting avenues for research directed toward dissecting the roles of these channels in different seizure disorders and associated neurological problems such as episodic ataxia.

Evidence from rat and mouse models of absence epilepsy, knockout mice, and characterization of functional effects of mutations found in patients all point to the fact that inhibition of P/Q-type channel activity somehow alters neurons and neuronal networks to result in seizure activity. This may also apply for variants of ancillary subunits that can act to reduce P/Q-type channel function. Conversely, however, it has been suggested that some of the clinically active antiepileptic drugs act by inhibiting presynaptic calcium channels. It is important to reiterate that acute inhibition of these channels is not functionally equivalent to long-term gene knockdown. Alternatively, the possibility that the therapeutic action of these drugs may be unrelated to calcium channel inhibition cannot be excluded.

The precise mechanism by which changes in  $\text{Ca}_v2.1$  channel function result in absence-like seizures has remained unclear. Indeed, interpretations of animal and clinical studies in the context of common forms of absence epilepsy are complicated by the fact that some of the murine P/Q-type channel mutations are associated with other neurological defects such as ataxia and cerebellar atrophy and that the absence phenotype in humans carrying P/Q-type channel mutations appears to be accompanied by primary generalized epilepsy. Furthermore, it is important to consider the perspective that the rodent models involving P/Q-type channel mutations may not be sufficiently focal in their brain defects to allow for a causal interpretation of their phenotype in relation to the human disorder. For example, these mice have cerebellar defects and experience mild to severe ataxia, a condition that is not observed in patients with absence epilepsy (19). However, similar to patients, these genetic models of absence epilepsy involve the thalamocortical circuitry, respond to clinically used antiepileptic drugs such as ethosuximide and valproic acid (123, 256), and do experience absencelike behavioral changes (209).

There are several potential avenues by which reduced P/Q-type channel activity could affect network properties that give rise to seizure activity. First, there is evidence that P/Q-type channel defects preferentially affect excitatory synaptic transmission. Such a disruption may well result in decreased input onto inhibitory neurons. Because inhibitory networks are known to play an important role in synchronizing ensembles of neurons, any inappropriate synaptic input onto the inhibitory network may contribute to pathological synchronization that is incompatible with normal network functioning (6, 261).

Second, it has been shown that P/Q-type channel activity is directly linked to calcium-dependent gene transcription via the CREB pathway (273), and therefore, reduced P/Q-type channel activity may compromise appropriate gene regulation and expression. This may be consistent with abnormal neuronal morphology that is observed with human and murine P/Q-type channel mutations (Fig. 6). Finally, it is important to note that T-type calcium channel activity appears to be increased in at least four different mouse models of absence epilepsy (i.e.,  $\text{Ca}_v2.1$  KO, *tg*, *lh*, *stg*) (256, 317); therefore, it is conceivable that the epileptic phenotypes associated with compromised P/Q-type channel function may arise indirectly from enhanced neuronal excitability mediated by T types (Fig. 5).

In contrast to P/Q-type channels, the appearance of enhanced T-type currents elevates the propensity for seizure activity. T-type channels mediate the rebound bursts in cortical and thalamic reticular neurons that are the physiological substrates for SWDs. Therefore, increased T-type channel expression (such as in the GAERS rat) or activity (as seen with gain-of-function mutations found in humans) can contribute to seizure genesis. It should, however, be noted that increased T-type calcium channel activity does not necessarily have to result from increased T-type channel expression. For example, alterations in ionic currents that interact with T-type currents, such as the H current ( $I_h$ ), which is mediated by HCN channels, can result in absence seizures (180, 224). This was recently demonstrated in the case of HCN2 (found primarily in the thalamus)-deficient mice, which exhibit spontaneous 5-Hz SWDs with absencelike episodes (178). Similarly, the WAG/Rij rat, which exhibits spontaneous absencelike seizures of neocortical origin (196), has been shown to have reduced levels of HCN1 protein expressed in the cortex; interestingly, mRNA levels for HCN1 were unaffected, and protein expression levels did not seem to differ from control animals for the three other HCN genes (269).

Given that cortical, thalamocortical relay, and reticular thalamic neurons express different complements of T-type channel isoforms, all three channel subtypes are potential contributors to the epileptic phenotype, but may do so in a regional- or cell type-specific manner. In this regard, it is interesting to note that in the GAERS rat, intrathalamic administration of ethosuximide (a known T-type blocker, see sect. III D) is only effective in reducing seizures by 70% at concentrations well beyond those known to be effective in blocking T-type channels (231). However, intracortical administration of ethosuximide has been shown to be much more effective in halting seizure activity (184), which further supports the notion that T-type-mediated SWD generation is not exclusively a thalamic phenomenon. This notion is further supported by the observation that specific knockout of one given channel subtype (i.e.,  $\text{Ca}_v3.1$  which is expressed predom-

inantly in thalamocortical relay neurons) is not universally effective in protecting against all pharmacologically induced generalized seizures. The susceptibility of the KO mice to bicuculline-induced seizures indicates that rebound bursting in thalamocortical relay neurons may not be required for this form of seizure activity and serves as a reminder of the many complex routes to seizure genesis (reviewed in Refs. 191, 192). On the other hand, reticular neurons express  $\text{Ca}_v3.2$  and  $\text{Ca}_v3.3$ , and it would thus be interesting to explore seizure susceptibility in knockout mice deficient of these channel subtypes. It also remains to be determined whether knockout or downregulation of  $\text{Ca}_v3.1$  could provide a protective role against absence seizures induced by the putative gain of function in  $\text{Ca}_v3.2$  such as those found in humans with idiopathic generalized epilepsy.

Historically, epilepsy has been a disorder through which much has been learned about the central nervous system and brain function in general. Exploration of voltage-gated calcium channels in epilepsy has proven to be both a challenging and rewarding endeavor. Future work promises to unlock many of the mysteries of how intrinsic cellular properties can interact in a complex network to bring about the intricacies of brain function in health and disease.

#### NOTE ADDED IN PROOF

A recent study by Zhong et al. (317a) has demonstrated that the *CACNA1H* ( $\text{Ca}_v3.2$ ) T-type calcium channel gene can undergo extensive alternative gene splicing. The authors reported that several of the reported idiopathic generalized epilepsy mutations in this gene, although not mediating changes in channel function when introduced into any given cDNA, interfere with splicing events. As a consequence, certain splice isoforms that may have unique biophysical properties can no longer be formed in vivo. We propose that this might be a possible mechanism by which these mutations could affect seizure threshold in certain neurons. Thus the findings of Zhong et al. (317a) further support the concept that the reported mutations can manifest their effects in numerous ways beyond direct biophysical changes.

#### ACKNOWLEDGMENTS

With this manuscript we honor the memory and achievements of Dr. Mircea Steriade (1924–2006). Dr. Steriade was a pioneer in the identification of the neuronal and network substrates of thalamic, cortical, and thalamocortical oscillations and their central roles in sleep and the disease state of epilepsy. We particularly thank Dr. Steriade for insightful comments on an earlier version of the manuscript. We also thank Dr. Paolo Federico for review of the clinical section.

Address for reprint requests and other correspondence: G. W. Zamponi, Dept. of Physiology and Biophysics, Univ. of Calgary, 3330 Hospital Dr. NW, Calgary T2N 4N1, Canada (e-mail: zamponi@ucalgary.ca).

#### GRANTS

G. W. Zamponi is a Senior Scholar of the Alberta Heritage Foundation for Medical Research (AHFMR) and a Canada Research Chair. H. Khosravani holds an AHFMR MD/PhD studentship, a Canada Graduate Scholarship, and an MD/PhD studentship from the Canadian Institutes for Health Research.

#### REFERENCES

1. Abegg MH, Savic N, Ehrenguber MU, McKinney RA, and Gahwiler BH. Epileptiform activity in rat hippocampus strengthens excitatory synapses. *J Physiol* 554: 439–448, 2004.
2. Adams ME, Myers RA, Imperial JS, and Olivera BM. Toxotyping rat brain calcium channels with omega-toxins from spider and cone snail venoms. *Biochemistry* 32: 12566–12570, 1993.
3. Against CoCaTotIL. Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsy Epilepsia* 30: 389–399, 1989.
4. Andermann F and Berkovic SF. Idiopathic generalized epilepsy with generalized and other seizures in adolescence. *Epilepsia* 42: 317–320, 2001.
5. Arikath J and Campbell KP. Auxiliary subunits: essential components of the voltage-gated calcium channel complex. *Curr Opin Neurobiol* 13: 298–307, 2003.
6. Avoli M. GABA-mediated synchronous potentials and seizure generation. *Epilepsia* 37: 1035–1042, 1996.
7. Ayata C, Shimizu-Sasamata M, Lo EH, Noebels JL, and Moskowitz MA. Impaired neurotransmitter release and elevated threshold for cortical spreading depression in mice with mutations in the  $\alpha 1A$  subunit of P/Q type calcium channels. *Neuroscience* 95: 639–645, 2000.
8. Bading H, Ginty DD, and Greenberg ME. Regulation of gene expression in hippocampal neurons by distinct calcium signaling pathways. *Science* 260: 181–186, 1993.
9. Ball SL, Powers PA, Shin HS, Morgans CW, Peachey NS, and Gregg RG. Role of the  $\beta(2)$  subunit of voltage-dependent calcium channels in the retinal outer plexiform layer. *Invest Ophthalmol Vis Sci* 43: 1595–1603, 2002.
10. Barclay J, Balaguero N, Mione M, Ackerman SL, Letts VA, Brodbeck J, Canti C, Meir A, Page KM, Kusumi K, Perez-Reyes E, Lander ES, Frankel WN, Gardiner RM, Dolphin AC, and Rees M. Ducky mouse phenotype of epilepsy and ataxia is associated with mutations in the *Cacna2d2* gene and decreased calcium channel current in cerebellar Purkinje cells. *J Neurosci* 21: 6095–6104, 2001.
11. Barclay J and Rees M. Mouse models of spike-wave epilepsy. *Epilepsia* 40 Suppl 3: 17–22, 1999.
12. Baulac S, Huberfeld G, Gourfinkel-An I, Mitropoulou G, Branger A, Prud'homme JF, Baulac M, Brice A, Bruzzone R, and LeGuern E. First genetic evidence of GABA(A) receptor dysfunction in epilepsy: a mutation in the  $\gamma 2$ -subunit gene. *Nat Genet* 28: 46–48, 2001.
13. Bayer K, Ahmadi S, and Zeilhofer HU. Gabapentin may inhibit synaptic transmission in the mouse spinal cord dorsal horn through a preferential block of P/Q-type  $\text{Ca}^{2+}$  channels. *Neuropharmacology* 46: 743–749, 2004.
14. Beauvais K, Cave-Riant F, De Barace C, Tardieu M, Tournier-Lasserre E, and Furby A. New *CACNA1A* gene mutation in a case of familial hemiplegic migraine with status epilepticus. *Eur Neurol* 52: 58–61, 2004.
15. Berkovic SF. *Epilepsy: A Comprehensive Textbook*, edited by Engel J Jr and Pedley TA. Philadelphia, PA: Lippincott-Williams & Wilkins, 1998.
16. Beuckmann CT, Sinton CM, Miyamoto N, Ino M, and Yanagisawa M. N-type calcium channel  $\alpha 1B$  subunit (*Cav2.2*) knockout mice display hyperactivity and vigilance state differences. *J Neurosci* 23: 6793–6797, 2003.
17. Bichet D, Cornet V, Geib S, Carlier E, Volsen S, Hoshi T, Mori Y, and De Waard M. The I-II loop of the  $\text{Ca}^{2+}$  channel  $\alpha 1A$  subunit contains an endoplasmic reticulum retention signal antagonized by the  $\beta$  subunit. *Neuron* 25: 177–190, 2000.

18. **Black JL III.** The voltage-gated calcium channel gamma subunits: a review of the literature. *J Bioenerg Biomembr* 35: 649–660, 2003.
19. **Blumenfeld H.** Cellular and network mechanisms of spike-wave seizures. *Epilepsia* 46 Suppl 9: 21–33, 2005.
20. **Blumenfeld H.** From molecules to networks: cortical/subcortical interactions in the pathophysiology of idiopathic generalized epilepsy. *Epilepsia* 44 Suppl 2: 7–15, 2003.
21. **Blumenfeld H.** The thalamus and seizures. *Arch Neurol* 59: 135–137, 2002.
22. **Bourinet E, Soong TW, Sutton K, Slaymaker S, Mathews E, Monteil A, Zamponi GW, Nargeot J, and Snutch TP.** Splicing of alpha 1A subunit gene generates phenotypic variants of P- and Q-type calcium channels. *Nat Neurosci* 2: 407–415, 1999.
23. **Bourinet E, Stotz SC, Spaetgens RL, Dayanithi G, Lemos J, Nargeot J, and Zamponi GW.** Interaction of SNX482 with domains III and IV inhibits activation gating of alpha(1E) (Ca(V)2.3) calcium channels. *Biophys J* 81: 79–88, 2001.
24. **Bredt DS and Nicoll RA.** AMPA receptor trafficking at excitatory synapses. *Neuron* 40: 361–379, 2003.
25. **Brill J, Klocke R, Paul D, Boison D, Gouder N, Klugbauer N, Hoffmann F, Becker CM, and Becker K.** entla, a novel epileptic and ataxic Cacna2d2 mutant of the mouse. *J Biol Chem* 279: 7322–7330, 2004.
26. **Brodbeck J, Davies A, Courtney JM, Meir A, Balaguero N, Canti C, Moss FJ, Page KM, Pratt WS, Hunt SP, Barclay J, Rees M, and Dolphin AC.** The ducky mutation in Cacna2d2 results in altered Purkinje cell morphology and is associated with the expression of a truncated alpha 2 delta-2 protein with abnormal function. *J Biol Chem* 277: 7684–7693, 2002.
27. **Brown JT and Randall A.** Gabapentin fails to alter P/Q-type Ca<sup>2+</sup> channel-mediated synaptic transmission in the hippocampus in vitro. *Synapse* 55: 262–269, 2005.
28. **Browne DL, Gancher ST, Nutt JG, Brunt ER, Smith EA, Kramer P, and Litt M.** Episodic ataxia/myokymia syndrome is associated with point mutations in the human potassium channel gene, KCNA1. *Nat Genet* 8: 136–140, 1994.
29. **Budde T, Sieg F, Braunewell KH, Gundelfinger ED, and Pape HC.** Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release supports the relay mode of activity in thalamocortical cells. *Neuron* 26: 483–492, 2000.
30. **Burgess DE, Crawford O, Delisle BP, and Satin J.** Mechanism of inactivation gating of human T-type (low-voltage activated) calcium channels. *Biophys J* 82: 1894–1906, 2002.
31. **Burgess DL, Jones JM, Meisler MH, and Noebels JL.** Mutation of the Ca<sup>2+</sup> channel beta subunit gene Cchb4 is associated with ataxia and seizures in the lethargic (lh) mouse. *Cell* 88: 385–392, 1997.
32. **Burke SP, Adams ME, and Taylor CP.** Inhibition of endogenous glutamate release from hippocampal tissue by Ca<sup>2+</sup> channel toxins. *Eur J Pharmacol* 238: 383–386, 1993.
33. **Caddick SJ, Wang C, Fletcher CF, Jenkins NA, Copeland NG, and Hosford DA.** Excitatory but not inhibitory synaptic transmission is reduced in lethargic (Cacnb4(lh)) and tottering (Cacnalatg) mouse thalami. *J Neurophysiol* 81: 2066–2074, 1999.
34. **Callenbach PM, van den Maagdenberg AM, Frants RR, and Brouwer OF.** Clinical and genetic aspects of idiopathic epilepsies in childhood. *Eur J Paediatr Neurol* 9: 91–103, 2005.
35. **Canti C, Nieto-Rostro M, Foucault I, Heblich F, Wratten J, Richards MW, Hendrich J, Douglas L, Page KM, Davies A, and Dolphin AC.** The metal-ion-dependent adhesion site in the Von Willebrand factor-A domain of alpha2delta subunits is key to trafficking voltage-gated Ca<sup>2+</sup> channels. *Proc Natl Acad Sci USA* 102: 11230–11235, 2005.
36. **Cao YQ, Piedras-Renteria ES, Smith GB, Chen G, Harata NC, and Tsien RW.** Presynaptic Ca<sup>2+</sup> channels compete for channel type-preferring slots in altered neurotransmission arising from Ca<sup>2+</sup> channelopathy. *Neuron* 43: 387–400, 2004.
37. **Cao YQ and Tsien RW.** Effects of familial hemiplegic migraine type 1 mutations on neuronal P/Q-type Ca<sup>2+</sup> channel activity and inhibitory synaptic transmission. *Proc Natl Acad Sci USA* 102: 2590–2595, 2005.
38. **Catterall WA.** Structure and function of voltage-gated sodium and calcium channels. *Curr Opin Neurobiol* 1: 5–13, 1991.
39. **Catterall WA.** Structure and modulation of Na<sup>+</sup> and Ca<sup>2+</sup> channels. *Ann NY Acad Sci* 707: 1–19, 1993.
40. **Catterall WA.** Structure and regulation of voltage-gated Ca<sup>2+</sup> channels. *Annu Rev Cell Dev Biol* 16: 521–555, 2000.
41. **Catterall WA, Perez-Reyes E, Snutch TP, and Striessnig J.** International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev* 57: 411–425, 2005.
42. **Charlier C, Singh NA, Ryan SG, Lewis TB, Reus BE, Leach RJ, and Leppert M.** A pore mutation in a novel KQT-like potassium channel gene in an idiopathic epilepsy family. *Nat Genet* 18: 53–55, 1998.
43. **Chemin J, Monteil A, Perez-Reyes E, Bourinet E, Nargeot J, and Lory P.** Specific contribution of human T-type calcium channel isoforms (alpha(1G), alpha(1H) and alpha(1I)) to neuronal excitability. *J Physiol* 540: 3–14, 2002.
44. **Chen CC, Lamping KG, Nuno DW, Barresi R, Prouty SJ, Lavoie JL, Cribbs LL, England SK, Sigmund CD, Weiss RM, Williamson RA, Hill JA, and Campbell KP.** Abnormal coronary function in mice deficient in alpha1H T-type Ca<sup>2+</sup> channels. *Science* 302: 1416–1418, 2003.
45. **Chen L, Chetkovich DM, Petralia RS, Sweeney NT, Kawasaki Y, Wenthold RJ, Bredt DS, and Nicoll RA.** Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature* 408: 936–943, 2000.
46. **Chen Y, Lu J, Pan H, Zhang Y, Wu H, Xu K, Liu X, Jiang Y, Bao X, Yao Z, Ding K, Lo WH, Qiang B, Chan P, Shen Y, and Wu X.** Association between genetic variation of CACNA1H and childhood absence epilepsy. *Ann Neurol* 54: 239–243, 2003.
47. **Chen YH, Li MH, Zhang Y, He LL, Yamada Y, Fitzmaurice A, Shen Y, Zhang H, Tong L, and Yang J.** Structural basis of the alpha1-beta subunit interaction of voltage-gated Ca<sup>2+</sup> channels. *Nature* 429: 675–680, 2004.
48. **Chien AJ, Zhao X, Shirokov RE, Puri TS, Chang CF, Sun D, Rios E, and Hosey MM.** Roles of a membrane-localized beta subunit in the formation and targeting of functional L-type Ca<sup>2+</sup> channels. *J Biol Chem* 270: 30036–30044, 1995.
49. **Chuang RS, Jaffe H, Cribbs L, Perez-Reyes E, and Swartz KJ.** Inhibition of T-type voltage-gated calcium channels by a new scorpion toxin. *Nat Neurosci* 1: 668–674, 1998.
50. **Claes L, Del-Favero J, Ceulemans B, Lagae L, Van Broeckhoven C, and De Jonghe P.** De novo mutations in the sodium-channel gene SCN1A cause severe myoclonic epilepsy of infancy. *Am J Hum Genet* 68: 1327–1332, 2001.
51. **Contreras D, Destexhe A, and Steriade M.** Intracellular and computational characterization of the intracortical inhibitory control of synchronized thalamic inputs in vivo. *J Neurophysiol* 78: 335–350, 1997.
52. **Contreras D and Steriade M.** Spindle oscillation in cats: the role of corticothalamic feedback in a thalamically generated rhythm. *J Physiol* 490: 159–179, 1996.
53. **Cossette P, Liu L, Brisebois K, Dong H, Lortie A, Vanasse M, Saint-Hilaire JM, Carmant L, Verner A, Lu WY, Wang YT, and Rouleau GA.** Mutation of GABRA1 in an autosomal dominant form of juvenile myoclonic epilepsy. *Nat Genet* 31: 184–189, 2002.
54. **Coulter DA, Huguenard JR, and Prince DA.** Calcium currents in rat thalamocortical relay neurones: kinetic properties of the transient, low-threshold current. *J Physiol* 414: 587–604, 1989.
55. **Craig PJ, Beattie RE, Folly EA, Banerjee MD, Reeves MB, Priestley JV, Carney SL, Sher E, Perez-Reyes E, and Volsen SG.** Distribution of the voltage-dependent calcium channel alpha1G subunit mRNA and protein throughout the mature rat brain. *Eur J Neurosci* 11: 2949–2964, 1999.
56. **Cribbs LL, Lee JH, Yang J, Satin J, Zhang Y, Daud A, Barclay J, Williamson MP, Fox M, Rees M, and Perez-Reyes E.** Cloning and characterization of alpha1H from human heart, a member of the T-type Ca<sup>2+</sup> channel gene family. *Circ Res* 83: 103–109, 1998.
57. **Crunelli V, Toth TI, Cope DW, Blethyn K, and Hughes SW.** The “window” T-type calcium current in brain dynamics of different behavioural states. *J Physiol* 562: 121–129, 2005.
58. **Cullinane AB, Coca-Prados M, and Harvey BJ.** Chloride dependent intracellular pH effects of external ATP in cultured human



- non-pigmented ciliary body epithelium. *Curr Eye Res* 23: 443–447, 2001.
59. **Dedek K, Kunath B, Kananura C, Reuner U, Jentsch TJ, and Steinlein OK.** Myokymia and neonatal epilepsy caused by a mutation in the voltage sensor of the KCNQ2 K<sup>+</sup> channel. *Proc Natl Acad Sci USA* 98: 12272–12277, 2001.
  60. **De Jongh KS, Warner C, and Catterall WA.** Subunits of purified calcium channels. Alpha 2 and delta are encoded by the same gene. *J Biol Chem* 265: 14738–14741, 1990.
  61. **De la Pena E and Geijo-Barrientos E.** Laminar localization, morphology, and physiological properties of pyramidal neurons that have the low-threshold calcium current in the guinea-pig medial frontal cortex. *J Neurosci* 16: 5301–5311, 1996.
  62. **De la Pena E and Geijo-Barrientos E.** Participation of low-threshold calcium spikes in excitatory synaptic transmission in guinea pig medial frontal cortex. *Eur J Neurosci* 12: 1679–1686, 2000.
  63. **Delgado-Escueta AV, Treiman DM, and Walsh GO.** The treatable epilepsies. *N Engl J Med* 308: 1508–1514, 1983.
  64. **Denier C, Ducros A, Durr A, Eymard B, Chassande B, and Tournier-Lasserre E.** Missense CACNA1A mutation causing episodic ataxia type 2. *Arch Neurol* 58: 292–295, 2001.
  65. **Denier C, Ducros A, Vahedi K, Joutel A, Thierry P, Ritz A, Castelnovo G, Deonna T, Gerard P, Devoize JL, Gayou A, Perrouty B, Soisson T, Autret A, Warter JM, Vighetto A, Van Bogaert P, Alamowitch S, Rouillet E, and Tournier-Lasserre E.** High prevalence of CACNA1A truncations and broader clinical spectrum in episodic ataxia type 2. *Neurology* 52: 1816–1821, 1999.
  66. **Destexhe A, Contreras D, Steriade M, Sejnowski TJ, and Huguenard JR.** In vivo, in vitro, and computational analysis of dendritic calcium currents in thalamic reticular neurons. *J Neurosci* 16: 169–185, 1996.
  67. **Destexhe A, Neubig M, Ulrich D, and Huguenard J.** Dendritic low-threshold calcium currents in thalamic relay cells. *J Neurosci* 18: 3574–3588, 1998.
  68. **Destexhe A and Sejnowski TJ.** Interactions between membrane conductances underlying thalamocortical slow-wave oscillations. *Physiol Rev* 83: 1401–1453, 2003.
  69. **Destexhe A and Sejnowski TJ.** The initiation of bursts in thalamic neurons and the cortical control of thalamic sensitivity. *Philos Trans R Soc Lond B Biol Sci* 357: 1649–1657, 2002.
  70. **De Waard M, Hering J, Weiss N, and Feltz A.** How do G proteins directly control neuronal Ca<sup>2+</sup> channel function? *Trends Pharmacol Sci* 26: 427–436, 2005.
  71. **Dibbens LM, Feng HJ, Richards MC, Harkin LA, Hodgson BL, Scott D, Jenkins M, Petrou S, Sutherland GR, Scheffer IE, Berkovic SF, MacDonald RL, and Mulley JC.** GABRD encoding a protein for extra- or peri-synaptic GABA<sub>A</sub> receptors is a susceptibility locus for generalized epilepsies. *Hum Mol Genet* 13: 1315–1319, 2004.
  72. **Dolmetsch RE, Pajvani U, Fife K, Spotts JM, and Greenberg ME.** Signaling to the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase pathway. *Science* 294: 333–339, 2001.
  73. **Dolphin AC.** Beta subunits of voltage-gated calcium channels. *J Bioenerg Biomembr* 35: 599–620, 2003.
  74. **Dolphin AC.** G protein modulation of voltage-gated calcium channels. *Pharmacol Rev* 55: 607–627, 2003.
  75. **Dolphin AC, Wyatt CN, Richards J, Beattie RE, Craig P, Lee JH, Cribbs LL, Volsen SG, and Perez-Reyes E.** The effect of alpha2-delta and other accessory subunits on expression and properties of the calcium channel alpha1G. *J Physiol* 519: 35–45, 1999.
  76. **Doroshenko PA, Woppmann A, Miljanich G, and Augustine GJ.** Pharmacologically distinct presynaptic calcium channels in cerebellar excitatory and inhibitory synapses. *Neuropharmacology* 36: 865–872, 1997.
  77. **Doyle DA, Morais Cabral J, Pfuetzner RA, Kuo A, Gulbis JM, Cohen SL, Chait BT, and MacKinnon R.** The structure of the potassium channel: molecular basis of K<sup>+</sup> conduction and selectivity. *Science* 280: 69–77, 1998.
  78. **Doyle J, Ren X, Lennon G, and Stubbs L.** Mutations in the Cacn1a4 calcium channel gene are associated with seizures, cerebellar degeneration, and ataxia in tottering and leaner mutant mice. *Mamm Genome* 8: 113–120, 1997.
  79. **Dubel SJ, Altier C, Chaumont S, Lory P, Bourinet E, and Nargeot J.** Plasma membrane expression of T-type calcium channel alpha(1) subunits is modulated by high voltage-activated auxiliary subunits. *J Biol Chem* 279: 29263–29269, 2004.
  80. **Dubel SJ, Starr TV, Hell J, Ahljian MK, Enyeart JJ, Catterall WA, and Snutch TP.** Molecular cloning of the alpha-1 subunit of an omega-conotoxin-sensitive calcium channel. *Proc Natl Acad Sci USA* 89: 5058–5062, 1992.
  81. **Ducros A, Denier C, Joutel A, Cecillon M, Lescoat C, Vahedi K, Darcel F, Vicaute E, Bousser MG, and Tournier-Lasserre E.** The clinical spectrum of familial hemiplegic migraine associated with mutations in a neuronal calcium channel. *N Engl J Med* 345: 17–24, 2001.
  82. **Ellinor PT, Yang J, Sather WA, Zhang JF, and Tsien RW.** Ca<sup>2+</sup> channel selectivity at a single locus for high-affinity Ca<sup>2+</sup> interactions. *Neuron* 15: 1121–1132, 1995.
  83. **Engel J.** Epilepsy in the world today: medical point of view. *Epilepsia* 43 Suppl 6: 12–13, 2002.
  84. **Engel J Jr.** A proposed diagnostic scheme for people with epileptic seizures and with epilepsy: report of the ILAE Task Force on Classification and Terminology. *Epilepsia* 42: 796–803, 2001.
  85. **Engel J, Pedley TA, and Aicardi J.** *Epilepsy: A Comprehensive Textbook*. Philadelphia, PA: Lippincott-Williams & Wilkins, 1998.
  86. **Erisir A, Van Horn SC, Bickford ME, and Sherman SM.** Immunocytochemistry and distribution of parabrachial terminals in the lateral geniculate nucleus of the cat: a comparison with corticogeniculate terminals. *J Comp Neurol* 377: 535–549, 1997.
  87. **Erisir A, Van Horn SC, and Sherman SM.** Relative numbers of cortical and brainstem inputs to the lateral geniculate nucleus. *Proc Natl Acad Sci USA* 94: 1517–1520, 1997.
  88. **Escayg A, De Waard M, Lee DD, Bichet D, Wolf P, Mayer T, Johnston J, Baloh R, Sander T, and Meisler MH.** Coding and noncoding variation of the human calcium-channel beta4-subunit gene CACNB4 in patients with idiopathic generalized epilepsy and episodic ataxia. *Am J Hum Genet* 66: 1531–1539, 2000.
  89. **Escayg A, MacDonald BT, Meisler MH, Baulac S, Huberfeld G, An-Gourfinkel I, Brice A, LeGuern E, Moulard B, Chaigne D, Buresi C, and Malafosse A.** Mutations of SCN1A, encoding a neuronal sodium channel, in two families with GEFS+2. *Nat Genet* 24: 343–345, 2000.
  90. **Feng ZP, Arnot MI, Doering CJ, and Zamponi GW.** Calcium channel beta subunits differentially regulate the inhibition of N-type channels by individual Gbeta isoforms. *J Biol Chem* 276: 45051–45058, 2001.
  91. **Feng ZP, Doering CJ, Winkfein RJ, Beedle AM, Spafford JD, and Zamponi GW.** Determinants of inhibition of transiently expressed voltage-gated calcium channels by omega-conotoxins GVIA and MVIIA. *J Biol Chem* 278: 20171–20178, 2003.
  92. **Feng ZP, Hamid J, Doering C, Jarvis SE, Bose GM, Bourinet E, Snutch TP, and Zamponi GW.** Amino acid residues outside of the pore region contribute to N-type calcium channel permeation. *J Biol Chem* 276: 5726–5730, 2001.
  93. **Fields RD, Lee PR, and Cohen JE.** Temporal integration of intracellular Ca<sup>2+</sup> signaling networks in regulating gene expression by action potentials. *Cell Calcium* 37: 433–442, 2005.
  94. **Fletcher CF, Lutz CM, O'Sullivan TN, Shaughnessy JD Jr, Hawkes R, Frankel WN, Copeland NG, and Jenkins NA.** Absence epilepsy in tottering mutant mice is associated with calcium channel defects. *Cell* 87: 607–617, 1996.
  95. **Fox AP, Nowicky MC, and Tsien RW.** Kinetic and pharmacological properties distinguishing three types of calcium currents in chick sensory neurones. *J Physiol* 394: 149–172, 1987.
  96. **Freise D, Held B, Wissenbach U, Pfeifer A, Trost C, Himmerkus N, Schweig U, Freichel M, Biel M, Hofmann F, Hoth M, and Flockerzi V.** Absence of the gamma subunit of the skeletal muscle dihydropyridine receptor increases L-type Ca<sup>2+</sup> currents and alters channel inactivation properties. *J Biol Chem* 275: 14476–14481, 2000.
  97. **Friend KL, Crimmins D, Phan TG, Sue CM, Colley A, Fung VS, Morris JG, Sutherland GR, and Richards RI.** Detection of a novel missense mutation and second recurrent mutation in the



- CACNA1A gene in individuals with EA-2 and FHM. *Hum Genet* 105: 261–265, 1999.
98. **Fuentealba P, Crochet S, Timofeev I, Bazhenov M, Sejnowski TJ, and Steriade M.** Experimental evidence and modeling studies support a synchronizing role for electrical coupling in the cat thalamic reticular neurons in vivo. *Eur J Neurosci* 20: 111–119, 2004.
  99. **Fuentealba P and Steriade M.** The reticular nucleus revisited: intrinsic and network properties of a thalamic pacemaker. *Prog Neurobiol* 75: 125–141, 2005.
  100. **Fuentealba P, Timofeev I, Bazhenov M, Sejnowski TJ, and Steriade M.** Membrane bistability in thalamic reticular neurons during spindle oscillations. *J Neurophysiol* 93: 294–304, 2005.
  101. **Fuentealba P, Timofeev I, and Steriade M.** Prolonged hyperpolarizing potentials precede spindle oscillations in the thalamic reticular nucleus. *Proc Natl Acad Sci USA* 101: 9816–9821, 2004.
  102. **Gardiner M.** Genetics of idiopathic generalized epilepsies. *Epilepsia* 46 Suppl 9: 15–20, 2005.
  103. **Goetz CG.** *Textbook of Clinical Neurology*. Philadelphia, PA: Saunders, 2003, p. 1306.
  104. **Gomora JC, Daud AN, Weiergraber M, and Perez-Reyes E.** Block of cloned human T-type calcium channels by succinimide antiepileptic drugs. *Mol Pharmacol* 60: 1121–1132, 2001.
  105. **Greenbaum D, Colangelo C, Williams K, and Gerstein M.** Comparing protein abundance and mRNA expression levels on a genomic scale. *Genome Biol* 4: 117, 2003.
  106. **Gregg RG, Messing A, Strube C, Beurg M, Moss R, Behan M, Sukhareva M, Haynes S, Powell JA, Coronado R, and Powers PA.** Absence of the beta subunit (cchb1) of the skeletal muscle dihydropyridine receptor alters expression of the alpha 1 subunit and eliminates excitation-contraction coupling. *Proc Natl Acad Sci USA* 93: 13961–13966, 1996.
  107. **Guberman A, Bruni J, and Guberman A.** *Essentials of Clinical Epilepsy*. Boston, MA: Butterworth-Heinemann, 1999, p. 207.
  108. **Guida S, Trettel F, Pagnutti S, Mantuano E, Tottene A, Veneziano L, Fellin T, Spadaro M, Stauderman K, Williams M, Volsen S, Ophoff R, Frants R, Jodice C, Frontali M, and Pietrobon D.** Complete loss of P/Q calcium channel activity caused by a CACNA1A missense mutation carried by patients with episodic ataxia type 2. *Am J Hum Genet* 68: 759–764, 2001.
  109. **Guido W, Lu SM, and Sherman SM.** Relative contributions of burst and tonic responses to the receptive field properties of lateral geniculate neurons in the cat. *J Neurophysiol* 68: 2199–2211, 1992.
  110. **Guido W and Weyand T.** Burst responses in thalamic relay cells of the awake behaving cat. *J Neurophysiol* 74: 1782–1786, 1995.
  111. **Hans M, Luvisetto S, Williams ME, Spagnolo M, Urrutia A, Tottene A, Brust PF, Johnson EC, Harpold MM, Stauderman KA, and Pietrobon D.** Functional consequences of mutations in the human alpha1A calcium channel subunit linked to familial hemiplegic migraine. *J Neurosci* 19: 1610–1619, 1999.
  112. **Hatakeyama S, Wakamori M, Ino M, Miyamoto N, Takahashi E, Yoshinaga T, Sawada K, Imoto K, Tanaka I, Yoshizawa T, Nishizawa Y, Mori Y, Niidome T, and Shoji S.** Differential nociceptive responses in mice lacking the alpha(1B) subunit of N-type Ca(2+) channels. *Neuroreport* 12: 2423–2427, 2001.
  113. **Haug K, Warnstedt M, Alekov AK, Sander T, Ramirez A, Poser B, Maljevic S, Hebeisen S, Kubisch C, Rebstock J, Horvath S, Hallmann K, Dullinger JS, Rau B, Haverkamp F, Beyenburg S, Schulz H, Janz D, Giese B, Muller-Newen G, Propping P, Elger CE, Fahlke C, Lerche H, and Heils A.** Mutations in CLCN2 encoding a voltage-gated chloride channel are associated with idiopathic generalized epilepsies. *Nat Genet* 33: 527–532, 2003.
  114. **Helekar SA and Noebels JL.** A burst-dependent hippocampal excitability defect elicited by potassium at the developmental onset of spike-wave seizures in the Tottering mutant. *Brain Res* 65: 205–210, 1992.
  115. **Helekar SA and Noebels JL.** Analysis of voltage-gated and synaptic conductances contributing to network excitability defects in the mutant mouse tottering. *J Neurophysiol* 71: 1–10, 1994.
  116. **Herman ST.** Epilepsy after brain insult: targeting epileptogenesis. *Neurology* 59: S21–S26, 2002.
  117. **Heron SE, Crossland KM, Andermann E, Phillips HA, Hall AJ, Bleasel A, Shevell M, Mercho S, Seni MH, Guiot MC, Mulley JC, Berkovic SF, and Scheffer IE.** Sodium-channel defects in benign familial neonatal-infantile seizures. *Lancet* 360: 851–852, 2002.
  118. **Heron SE, Phillips HA, Mulley JC, Mazarib A, Neufeld MY, Berkovic SF, and Scheffer IE.** Genetic variation of CACNA1H in idiopathic generalized epilepsy. *Ann Neurol* 55: 595–596, 2004.
  119. **Herrup K and Wilczynski SL.** Cerebellar cell degeneration in the leaner mutant mouse. *Neuroscience* 7: 2185–2196, 1982.
  120. **Hirose S, Mitsudome A, Okada M, and Kaneko S.** Genetics of idiopathic epilepsies. *Epilepsia* 46 Suppl 1: 38–43, 2005.
  121. **Hobom M, Dai S, Marais E, Lacinova L, Hofmann F, and Klugbauer N.** Neuronal distribution and functional characterization of the calcium channel alpha2delta-2 subunit. *Eur J Neurosci* 12: 1217–1226, 2000.
  122. **Holmes MD, Brown M, and Tucker DM.** Are “generalized” seizures truly generalized? Evidence of localized mesial frontal and frontopolar discharges in absence. *Epilepsia* 45: 1568–1579, 2004.
  123. **Hosford DA, Clark S, Cao Z, Wilson WA Jr, Lin FH, Morrisett RA, and Huin A.** The role of GABA<sub>B</sub> receptor activation in absence seizures of lethargic (*lh/lh*) mice. *Science* 257: 398–401, 1992.
  124. **Hughes SW, Cope DW, Blethyn KL, and Crunelli V.** Cellular mechanisms of the slow (<1 Hz) oscillation in thalamocortical neurons in vitro. *Neuron* 33: 947–958, 2002.
  125. **Huguenard JR.** Block of T-type Ca(2+) channels is an important action of succinimide antiabsence drugs. *Epilepsy Curr* 2: 49–52, 2002.
  126. **Huguenard JR and Prince DA.** A novel T-type current underlies prolonged Ca(2+)-dependent burst firing in GABAergic neurons of rat thalamic reticular nucleus. *J Neurosci* 12: 3804–3817, 1992.
  127. **Imbrici P, Jaffe SL, Eunson LH, Davies NP, Herd C, Robertson R, Kullmann DM, and Hanna MG.** Dysfunction of the brain calcium channel Cav2.1 in absence epilepsy and episodic ataxia. *Brain* 127: 2682–2692, 2004.
  128. **Ino M, Yoshinaga T, Wakamori M, Miyamoto N, Takahashi E, Sonoda J, Kagaya T, Oki T, Nagasu T, Nishizawa Y, Tanaka I, Imoto K, Aizawa S, Koch S, Schwartz A, Niidome T, Sawada K, and Mori Y.** Functional disorders of the sympathetic nervous system in mice lacking the alpha 1B subunit (Cav 2.2) of N-type calcium channels. *Proc Natl Acad Sci USA* 98: 5323–5328, 2001.
  129. **Isope P and Murphy TH.** Low threshold calcium currents in rat cerebellar Purkinje cell dendritic spines are mediated by T-type calcium channels. *J Physiol* 562: 257–269, 2005.
  130. **Jallon P and Latour P.** Epidemiology of idiopathic generalized epilepsies. *Epilepsia* 46 Suppl 9: 10–14, 2005.
  131. **Jen J, Yue Q, Nelson SF, Yu H, Litt M, Nutt J, and Baloh RW.** A novel nonsense mutation in CACNA1A causes episodic ataxia and hemiplegia. *Neurology* 53: 34–37, 1999.
  132. **Jiang Y, Lee A, Chen J, Ruta V, Cadene M, Chait BT, and MacKinnon R.** X-ray structure of a voltage-dependent K<sup>+</sup> channel. *Nature* 423: 33–41, 2003.
  133. **Joksovic PM, Bayliss DA, and Todorovic SM.** Different kinetic properties of two T-type Ca<sup>2+</sup> currents of reticular thalamic neurons and their modulation by enflurane. *J Physiol* 566: 125–142, 2005.
  134. **Jouveneau A, Eunson LH, Spauschus A, Ramesh V, Zuberi SM, Kullmann DM, and Hanna MG.** Human epilepsy associated with dysfunction of the brain P/Q-type calcium channel. *Lancet* 358: 801–807, 2001.
  135. **Jun K, Piedras-Renteria ES, Smith SM, Wheeler DB, Lee SB, Lee TG, Chin H, Adams ME, Scheller RH, Tsien RW, and Shin HS.** Ablation of P/Q-type Ca(2+) channel currents, altered synaptic transmission, and progressive ataxia in mice lacking the alpha(1A)-subunit. *Proc Natl Acad Sci USA* 96: 15245–15250, 1999.
  136. **Jung HY, Staff NP, and Spruston N.** Action potential bursting in subicular pyramidal neurons is driven by a calcium tail current. *J Neurosci* 21: 3312–3321, 2001.
  137. **Karst H, Joels M, and Wadman WJ.** Low-threshold calcium current in dendrites of the adult rat hippocampus. *Neurosci Lett* 164: 154–158, 1993.
  138. **Kase M, Kakimoto S, Sakuma S, Houtani T, Ohishi H, Ueyama T, and Sugimoto T.** Distribution of neurons expressing alpha 1G subunit mRNA of T-type voltage-dependent calcium channel in adult rat central nervous system. *Neurosci Lett* 268: 77–80, 1999.

139. Kavalali ET, Zhuo M, Bito H, and Tsien RW. Dendritic  $\text{Ca}^{2+}$  channels characterized by recordings from isolated hippocampal dendritic segments. *Neuron* 18: 651–663, 1997.
140. Khosravani H, Altier C, Simms B, Hamming KS, Snutch TP, Mezeyova J, McRory JE, and Zamponi GW. Gating effects of mutations in the Cav3.2 T-type calcium channel associated with childhood absence epilepsy. *J Biol Chem* 279: 9681–9684, 2004.
141. Khosravani H, Bladen C, Parker DB, Snutch TP, McRory JE, and Zamponi GW. Effects of  $\text{Ca(v)3.2}$  channel mutations linked to idiopathic generalized epilepsy. *Ann Neurol* 57: 745–749, 2005.
142. Kim C, Jun K, Lee T, Kim SS, McEnery MW, Chin H, Kim HL, Park JM, Kim DK, Jung SJ, Kim J, and Shin HS. Altered nociceptive response in mice deficient in the  $\alpha(1B)$  subunit of the voltage-dependent calcium channel. *Mol Cell Neurosci* 18: 235–245, 2001.
143. Kim D, Park D, Choi S, Lee S, Sun M, Kim C, and Shin HS. Thalamic control of visceral nociception mediated by T-type  $\text{Ca}^{2+}$  channels. *Science* 302: 117–119, 2003.
144. Kim D, Song I, Keum S, Lee T, Jeong MJ, Kim SS, McEnery MW, and Shin HS. Lack of the burst firing of thalamocortical relay neurons and resistance to absence seizures in mice lacking  $\alpha(1G)$  T-type  $\text{Ca}^{2+}$  channels. *Neuron* 31: 35–45, 2001.
145. Klockner U, Lee JH, Cribbs LL, Daud A, Hescheler J, Pereverzev A, Perez-Reyes E, and Schneider T. Comparison of the  $\text{Ca}^{2+}$  currents induced by expression of three cloned  $\alpha$  subunits,  $\alpha 1G$ ,  $\alpha 1H$  and  $\alpha 1I$ , of low-voltage-activated T-type  $\text{Ca}^{2+}$  channels. *Eur J Neurosci* 11: 4171–4178, 1999.
146. Klugbauer N, Lacinova L, Marais E, Hobom M, and Hofmann F. Molecular diversity of the calcium channel  $\alpha 2\delta$  subunit. *J Neurosci* 19: 684–691, 1999.
147. Klugbauer N, Marais E, and Hofmann F. Calcium channel  $\alpha 2\delta$  subunits: differential expression, function, and drug binding. *J Bioenerg Biomembr* 35: 639–647, 2003.
148. Kors EE, Melberg A, Vanmolkot KR, Kumlien E, Haan J, Raininko R, Flink R, Ginjaar HB, Frants RR, Ferrari MD, and van den Maagdenberg AM. Childhood epilepsy, familial hemiplegic migraine, cerebellar ataxia, and a new CACNA1A mutation. *Neurology* 63: 1136–1137, 2004.
149. Koschak A, Reimer D, Walter D, Hoda JC, Heinzle T, Grabner M, and Striessnig J. Cav 1.4  $\alpha 1$  subunits can form slowly inactivating dihydropyridine-sensitive L-type  $\text{Ca}^{2+}$  channels lacking  $\text{Ca}^{2+}$ -dependent inactivation. *J Neurosci* 23: 6041–6049, 2003.
150. Kraus RL, Sinnegger MJ, Glossmann H, Hering S, and Striessnig J. Familial hemiplegic migraine mutations change  $\alpha 1A$   $\text{Ca}^{2+}$  channel kinetics. *J Biol Chem* 273: 5586–5590, 1998.
151. Kraus RL, Sinnegger MJ, Koschak A, Glossmann H, Stenirri S, Carrera P, and Striessnig J. Three new familial hemiplegic migraine mutants affect P/Q-type  $\text{Ca}^{2+}$  channel kinetics. *J Biol Chem* 275: 9239–9243, 2000.
152. Kuzmiski JB, Barr W, Zamponi GW, and MacVicar BA. Topiramate inhibits the initiation of plateau potentials in CA1 neurons by depressing R-type calcium channels. *Epilepsia* 46: 481–489, 2005.
153. Kwan P and Brodie MJ. Early identification of refractory epilepsy. *N Engl J Med* 342: 314–319, 2000.
154. Kwan P and Brodie MJ. Effectiveness of first antiepileptic drug. *Epilepsia* 42: 1255–1260, 2001.
155. Lambert RC, Maulet Y, Mouton J, Beattie R, Volsen S, De Waard M, and Feltz A. T-type  $\text{Ca}^{2+}$  current properties are not modified by  $\text{Ca}^{2+}$  channel beta subunit depletion in nodose ganglion neurons. *J Neurosci* 17: 6621–6628, 1997.
156. Landisman CE, Long MA, Beierlein M, Deans MR, Paul DL, and Connors BW. Electrical synapses in the thalamic reticular nucleus. *J Neurosci* 22: 1002–1009, 2002.
157. Larkum ME and Zhu JJ. Signaling of layer 1 and whisker-evoked  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  action potentials in distal and terminal dendrites of rat neocortical pyramidal neurons in vitro and in vivo. *J Neurosci* 22: 6991–7005, 2002.
158. Lee A, Wong ST, Gallagher D, Li B, Storm DR, Scheuer T, and Catterall WA.  $\text{Ca}^{2+}$ /calmodulin binds to and modulates P/Q-type calcium channels. *Nature* 399: 155–159, 1999.
159. Lee J, Kim D, and Shin HS. Lack of delta waves and sleep disturbances during non-rapid eye movement sleep in mice lacking  $\alpha 1G$ -subunit of T-type calcium channels. *Proc Natl Acad Sci USA* 101: 18195–18199, 2004.
160. Lee JH, Daud AN, Cribbs LL, Lacerda AE, Pereverzev A, Klockner U, Schneider T, and Perez-Reyes E. Cloning and expression of a novel member of the low voltage-activated T-type calcium channel family. *J Neurosci* 19: 1912–1921, 1999.
161. Lee JH, Gomora JC, Cribbs LL, and Perez-Reyes E. Nickel block of three cloned T-type calcium channels: low concentrations selectively block  $\alpha 1H$ . *Biophys J* 77: 3034–3042, 1999.
162. Lee SC, Choi S, Lee T, Kim HL, Chin H, and Shin HS. Molecular basis of R-type calcium channels in central amygdala neurons of the mouse. *Proc Natl Acad Sci USA* 99: 3276–3281, 2002.
163. Leenders AG, van den Maagdenberg AM, Lopes da Silva FH, Sheng ZH, Molenaar PC, and Ghijsen WE. Neurotransmitter release from tottering mice nerve terminals with reduced expression of mutated P- and Q-type  $\text{Ca}^{2+}$ -channels. *Eur J Neurosci* 15: 13–18, 2002.
164. Leonardi M and Ustun TB. The global burden of epilepsy. *Epilepsia* 43 Suppl 6: 21–25, 2002.
165. Leresche N, Parri HR, Erdemli G, Guyon A, Turner JP, Williams SR, Asprodingi E, and Crunelli V. On the action of the anti-absence drug ethosuximide in the rat and cat thalamus. *J Neurosci* 18: 4842–4853, 1998.
166. Letts VA, Felix R, Biddlecome GH, Arikath J, Mahaffey CL, Valenzuela A, Bartlett FS II, Mori Y, Campbell KP, and Frankel WN. The mouse stargazer gene encodes a neuronal  $\text{Ca}^{2+}$ -channel gamma subunit. *Nat Genet* 19: 340–347, 1998.
167. Letts VA, Kang MG, Mahaffey CL, Beyer B, Tenbrink H, Campbell KP, and Frankel WN. Phenotypic heterogeneity in the stargazin allelic series. *Mamm Genome* 14: 506–513, 2003.
168. Letts VA, Mahaffey CL, Beyer B, and Frankel WN. A targeted mutation in *Cacng4* exacerbates spike-wave seizures in stargazer (*Cacng2*) mice. *Proc Natl Acad Sci USA* 102: 2123–2128, 2005.
169. Leuranguer V, Bourinet E, Lory P, and Nargeot J. Antisense depletion of beta-subunits fails to affect T-type calcium channel properties in a neuroblastoma cell line. *Neuropharmacology* 37: 701–708, 1998.
170. Liang H, DeMaria CD, Erickson MG, Mori MX, Alseikhan BA, and Yue DT. Unified mechanisms of  $\text{Ca}^{2+}$  regulation across the  $\text{Ca}^{2+}$  channel family. *Neuron* 39: 951–960, 2003.
171. Lie AA, Blumcke I, Volsen SG, Wiestler OD, Elger CE, and Beck H. Distribution of voltage-dependent calcium channel beta subunits in the hippocampus of patients with temporal lobe epilepsy. *Neuroscience* 93: 449–456, 1999.
172. Liu XB, Honda CN, and Jones EG. Distribution of four types of synapse on physiologically identified relay neurons in the ventral posterior thalamic nucleus of the cat. *J Comp Neurol* 352: 69–91, 1995.
173. Liu XB and Jones EG. Predominance of corticothalamic synaptic inputs to thalamic reticular nucleus neurons in the rat. *J Comp Neurol* 414: 67–79, 1999.
174. Llinas R and Jahnsen H. Electrophysiology of mammalian thalamic neurones in vitro. *Nature* 297: 406–408, 1982.
175. Long MA, Landisman CE, and Connors BW. Small clusters of electrically coupled neurons generate synchronous rhythms in the thalamic reticular nucleus. *J Neurosci* 24: 341–349, 2004.
176. Long SB, Campbell EB, and Mackinnon R. Voltage sensor of Kv1.2: structural basis of electromechanical coupling. *Science* 309: 903–908, 2005.
177. Lorenzon NM, Lutz CM, Frankel WN, and Beam KG. Altered calcium channel currents in Purkinje cells of the neurological mutant mouse leaner. *J Neurosci* 18: 4482–4489, 1998.
178. Ludwig A, Budde T, Stieber J, Moosmang S, Wahl C, Holthoff K, Langebartels A, Wotjak C, Munsch T, Zong X, Feil S, Feil R, Lancel M, Chien KR, Konnerth A, Pape HC, Biel M, and Hofmann F. Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. *EMBO J* 22: 216–224, 2003.
179. Lukyanetz EA, Shkryl VM, and Kostyuk PG. Selective blockade of N-type calcium channels by levetiracetam. *Epilepsia* 43: 9–18, 2002.
180. Luthi A and McCormick DA. H-current: properties of a neuronal and network pacemaker. *Neuron* 21: 9–12, 1998.



181. Lynch BA, Lambeng N, Nocka K, Kensel-Hammes P, Bajjalieh SM, Matagne A, and Fuks B. The synaptic vesicle protein SV2A is the binding site for the antiepileptic drug levetiracetam. *Proc Natl Acad Sci USA* 101: 9861–9866, 2004.
182. Magee JC and Johnston D. Characterization of single voltage-gated Na<sup>+</sup> and Ca<sup>2+</sup> channels in apical dendrites of rat CA1 pyramidal neurons. *J Physiol* 487: 67–90, 1995.
183. Malenka RC. Synaptic plasticity and AMPA receptor trafficking. *Ann NY Acad Sci* 1003: 1–11, 2003.
184. Manning JP, Richards DA, Leresche N, Crunelli V, and Bowery NG. Cortical-area specific block of genetically determined absence seizures by ethosuximide. *Neuroscience* 123: 5–9, 2004.
185. Mansergh F, Orton NC, Vessey JP, Lalonde MR, Stell WK, Tremblay F, Barnes S, Rancourt DE, and Bech-Hansen NT. Mutation of the calcium channel gene Cacna1f disrupts calcium signaling, synaptic transmission and cellular organization in mouse retina. *Hum Mol Genet* 14: 3035–3046, 2005.
186. Marais E, Klugbauer N, and Hofmann F. Calcium channel alpha(2)delta subunits-structure and Gabapentin binding. *Mol Pharmacol* 59: 1243–1248, 2001.
187. Marini C, King MA, Archer JS, Newton MR, and Berkovic SF. Idiopathic generalized epilepsy of adult onset: clinical syndromes and genetics. *J Neurol Neurosurg Psychiatry* 74: 192–196, 2005.
188. Markram H and Sakmann B. Calcium transients in dendrites of neocortical neurons evoked by single subthreshold excitatory postsynaptic potentials via low-voltage-activated calcium channels. *Proc Natl Acad Sci USA* 91: 5207–5211, 1994.
189. Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, and Wu C. Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* 5: 793–807, 2004.
190. Martin RL, Lee JH, Cribbs LL, Perez-Reyes E, and Hanck DA. Mibefradil block of cloned T-type calcium channels. *J Pharmacol Exp Ther* 295: 302–308, 2000.
191. McCormick DA and Contreras D. On the cellular and network bases of epileptic seizures. *Annu Rev Physiol* 63: 815–846, 2001.
192. McNamara JO. Emerging insights into the genesis of epilepsy. *Nature* 399: A15–A22, 1999.
193. McRory JE, Hamid J, Doering CJ, Garcia E, Parker R, Hamming KS, Chen L, Hildebrand M, Beedle AM, Feldcamp L, Zamponi GW, and Snutch TP. The CACNA1F gene encodes an L-type calcium channel with unique biophysical properties and tissue distribution. *J Neurosci*. In press.
194. McRory JE, Santi CM, Hamming KS, Mezeyova J, Sutton KG, Baillie DL, Stea A, and Snutch TP. Molecular and functional characterization of a family of rat brain T-type calcium channels. *J Biol Chem* 276: 3999–4011, 2001.
195. Meeren H, van Luijckelaar G, Lopes da Silva F, and Coenen A. Evolving concepts on the pathophysiology of absence seizures: the cortical focus theory. *Arch Neurol* 62: 371–376, 2005.
196. Meeren HK, Pijn JP, Van Luijckelaar EL, Coenen AM, and Lopes da Silva FH. Cortical focus drives widespread corticothalamic networks during spontaneous absence seizures in rats. *J Neurosci* 22: 1480–1495, 2002.
197. Mikami A, Imoto K, Tanabe T, Niidome T, Mori Y, Takeshima H, Narumiya S, and Numa S. Primary structure and functional expression of the cardiac dihydropyridine-sensitive calcium channel. *Nature* 340: 230–233, 1989.
198. Monteil A, Chemin J, Bourinet E, Mennessier G, Lory P, and Nargeot J. Molecular and functional properties of the human alpha(1G) subunit that forms T-type calcium channels. *J Biol Chem* 275: 6090–6100, 2000.
199. Monteil A, Chemin J, Leuranguer V, Altier C, Mennessier G, Bourinet E, Lory P, and Nargeot J. Specific properties of T-type calcium channels generated by the human alpha 1I subunit. *J Biol Chem* 275: 16530–16535, 2000.
200. Mouginot D, Bossu JL, and Gahwiler BH. Low-threshold Ca<sup>2+</sup> currents in dendritic recordings from Purkinje cells in rat cerebellar slice cultures. *J Neurosci* 17: 160–170, 1997.
201. Mulley JC, Scheffer IE, Harkin LA, Berkovic SF, and Dibbens LM. Susceptibility genes for complex epilepsy. *Hum Mol Genet* 14 Suppl 2: R243–R249, 2005.
202. Mulley JC, Scheffer IE, Petrou S, and Berkovic SF. Channelopathies as a genetic cause of epilepsy. *Curr Opin Neurol* 16: 171–176, 2003.
203. Murbartian J, Arias JM, Lee JH, Gomora JC, and Perez-Reyes E. Alternative splicing of the rat Ca(v)3.3 T-type calcium channel gene produces variants with distinct functional properties(1). *FEBS Lett* 528: 272–278, 2002.
204. Nakagawa T, Cheng Y, Ramm E, Sheng M, and Walz T. Structure and different conformational states of native AMPA receptor complexes. *Nature* 433: 545–549, 2005.
205. Namkung Y, Smith SM, Lee SB, Skrypnik NV, Kim HL, Chin H, Scheller RH, Tsien RW, and Shin HS. Targeted disruption of the Ca<sup>2+</sup> channel beta3 subunit reduces N- and L-type Ca<sup>2+</sup> channel activity and alters the voltage-dependent activation of P/Q-type Ca<sup>2+</sup> channels in neurons. *Proc Natl Acad Sci USA* 95: 12010–12015, 1998.
206. Newcomb R, Szoke B, Palma A, Wang G, Chen X, Hopkins W, Cong R, Miller J, Urge L, Tarczy-Hornoch K, Loo JA, Dooley DJ, Nadasdi L, Tsien RW, Lemos J, and Miljanich G. Selective peptide antagonist of the class E calcium channel from the venom of the tarantula *Hysterocrates gigas*. *Biochemistry* 37: 15353–15362, 1998.
207. Newton PM, Orr CJ, Wallace MJ, Kim C, Shin HS, and Messing RO. Deletion of N-type calcium channels alters ethanol reward and reduces ethanol consumption in mice. *J Neurosci* 24: 9862–9869, 2004.
208. Niwa N, Yasui K, Opthof T, Takemura H, Shimizu A, Horiba M, Lee JK, Honjo H, Kamiya K, and Kodama I. Cav3.2 subunit underlies the functional T-type Ca<sup>2+</sup> channel in murine hearts during the embryonic period. *Am J Physiol Heart Circ Physiol* 286: H2257–H2263, 2004.
209. Noebels JL. Exploring new gene discoveries in idiopathic generalized epilepsy. *Epilepsia* 44 Suppl 2: 16–21, 2003.
210. Noebels JL, Fariello RG, Jobe PC, Lasley SM, and Marescaux C. Genetic models of generalized epilepsy. In: *Epilepsy: A Comprehensive Textbook*. Philadelphia, PA: Lippincott-Raven, 1997.
211. Noebels JL and Sidman RL. Inherited epilepsy: spike-wave and focal motor seizures in the mutant mouse tottering. *Science* 204: 1334–1336, 1979.
212. Olivera BM, McIntosh JM, Cruz LJ, Luque FA, and Gray WR. Purification and sequence of a presynaptic peptide toxin from *Conus geographus* venom. *Biochemistry* 23: 5087–5090, 1984.
213. Opatowsky Y, Chomsky-Hecht O, Kang MG, Campbell KP, and Hirsch JA. The voltage-dependent calcium channel beta subunit contains two stable interacting domains. *J Biol Chem* 278: 52323–52332, 2003.
214. Ophoff RA, Terwindt GM, Vergouwe MN, van Eijk R, Oefner PJ, Hoffman SM, Lamerdin JE, Mohrenweiser HW, Bulman DE, Ferrari M, Haan J, Lindhout D, van Ommen GJ, Hofker MH, Ferrari MD, and Frants RR. Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the Ca<sup>2+</sup> channel gene CACNL1A4. *Cell* 87: 543–552, 1996.
215. Peloquin J, Khosravani H, Barr W, Bladen C, Evans R, Mezeyova J, Parker D, Snutch TP, McRory JE, and Zamponi GW. Functional analysis of Cav3.2 T-type calcium channel mutations linked to childhood absence epilepsy. *Epilepsia* 47: 655–658, 2006.
216. Perez-Reyes E. Molecular characterization of a novel family of low voltage-activated, T-type, calcium channels. *J Bioenerg Biomembr* 30: 313–318, 1998.
217. Perez-Reyes E. Molecular physiology of low-voltage-activated T-type calcium channels. *Physiol Rev* 83: 117–161, 2003.
218. Peterson BZ, DeMaria CD, Adelman JP, and Yue DT. Calmodulin is the Ca<sup>2+</sup> sensor for Ca<sup>2+</sup>-dependent inactivation of L-type calcium channels. *Neuron* 22: 549–558, 1999.
219. Pietrobon D. Calcium channels and channelopathies of the central nervous system. *Mol Neurobiol* 25: 31–50, 2002.
220. Pietrobon D and Striessnig J. Neurobiology of migraine. *Nat Rev Neurosci* 4: 386–398, 2003.
221. Pinault D. Cellular interactions in the rat somatosensory thalamo-cortical system during normal and epileptic 5–9 Hz oscillations. *J Physiol* 552: 881–905, 2003.

222. Pinault D, Leresche N, Charpier S, Deniau JM, Marescaux C, Vergnes M, and Crunelli V. Intracellular recordings in thalamic neurones during spontaneous spike and wave discharges in rats with absence epilepsy. *J Physiol* 509: 449–456, 1998.
223. Platzer J, Engel J, Schrott-Fischer A, Stephan K, Bova S, Chen H, Zheng H, and Striessnig J. Congenital deafness and sinoatrial node dysfunction in mice lacking class D L-type  $\text{Ca}^{2+}$  channels. *Cell* 102: 89–97, 2000.
224. Poolos NP. The h-channel: a potential channelopathy in epilepsy? *Epilepsy Behav* 7: 51–56, 2005.
225. Potier B, Dutar P, and Lamour Y. Different effects of omega-conotoxin GVIA at excitatory and inhibitory synapses in rat CA1 hippocampal neurons. *Brain Res* 616: 236–241, 1993.
226. Pragnell M, De Waard M, Mori Y, Tanabe T, Snutch TP, and Campbell KP. Calcium channel beta-subunit binds to a conserved motif in the I-II cytoplasmic linker of the alpha 1-subunit. *Nature* 368: 67–70, 1994.
227. Qian J and Noebels JL. Presynaptic  $\text{Ca}^{2+}$  influx at a mouse central synapse with  $\text{Ca}^{2+}$  channel subunit mutations. *J Neurosci* 20: 163–170, 2000.
228. Qin N, Platano D, Olcese R, Costantin JL, Stefani E, and Birnbaumer L. Unique regulatory properties of the type 2a  $\text{Ca}^{2+}$  channel beta subunit caused by palmitoylation. *Proc Natl Acad Sci USA* 95: 4690–4695, 1998.
229. Randall A and Tsien RW. Pharmacological dissection of multiple types of  $\text{Ca}^{2+}$  channel currents in rat cerebellar granule neurons. *J Neurosci* 15: 2995–3012, 1995.
230. Reynolds IJ, Wagner JA, Snyder SH, Thayer SA, Olivera BM, and Miller RJ. Brain voltage-sensitive calcium channel subtypes differentiated by omega-conotoxin fraction GVIA. *Proc Natl Acad Sci USA* 83: 8804–8807, 1986.
231. Richards DA, Manning JP, Barnes D, Rombola L, Bower NG, Caccia S, Leresche N, and Crunelli V. Targeting thalamic nuclei is not sufficient for the full anti-absence action of ethosuximide in a rat model of absence epilepsy. *Epilepsy Res* 54: 97–107, 2003.
232. Richards MW, Butcher AJ, and Dolphin AC.  $\text{Ca}^{2+}$  channel beta-subunits: structural insights aid our understanding. *Trends Pharmacol Sci* 25: 626–632, 2004.
233. Richter TA, Kolaj M, and Renaud LP. Low voltage-activated  $\text{Ca}^{2+}$  channels are coupled to  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release in rat thalamic midline neurons. *J Neurosci* 25: 8267–8271, 2005.
234. Rodin E and Ancheta O. Cerebral electrical fields during petit mal absences. *Electroencephalogr Clin Neurophysiol* 66: 457–466, 1987.
235. Rogawski MA and Loscher W. The neurobiology of antiepileptic drugs. *Nat Rev Neurosci* 5: 553–564, 2004.
236. Rogawski MA and Loscher W. The neurobiology of antiepileptic drugs for the treatment of nonepileptic conditions. *Nat Med* 10: 685–692, 2004.
237. Rose CR and Konnerth A. Stores not just for storage. Intracellular calcium release and synaptic plasticity. *Neuron* 31: 519–522, 2001.
238. Rousset M, Cens T, Restituito S, Barrere C, Black JL III, McEnery MW, and Charnet P. Functional roles of gamma2, gamma3 and gamma4, three new  $\text{Ca}^{2+}$  channel subunits, in P/Q-type  $\text{Ca}^{2+}$  channel expressed in *Xenopus* oocytes. *J Physiol* 532: 583–593, 2001.
239. Saegusa H, Kurihara T, Zong S, Kazuno A, Matsuda Y, Nonaka T, Han W, Toriyama H, and Tanabe T. Suppression of inflammatory and neuropathic pain symptoms in mice lacking the N-type  $\text{Ca}^{2+}$  channel. *EMBO J* 20: 2349–2356, 2001.
240. Saegusa H, Kurihara T, Zong S, Minowa O, Kazuno A, Han W, Matsuda Y, Yamanaka H, Osanai M, Noda T, and Tanabe T. Altered pain responses in mice lacking alpha 1E subunit of the voltage-dependent  $\text{Ca}^{2+}$  channel. *Proc Natl Acad Sci USA* 97: 6132–6137, 2000.
241. Sander T, Schulz H, Saar K, Gennaro E, Riggio MC, Bianchi A, Zara F, Luna D, Bulteau C, Kaminska A, Ville D, Cieuta C, Picard F, Prud'homme JF, Bate L, Sundquist A, Gardiner RM, Janssen GA, de Haan GJ, Kasteleijn-Nolst-Trenite DG, Bader A, Lindhout D, Riess O, Wienker TF, Janz D, and Reis A. Genome search for susceptibility loci of common idiopathic generalised epilepsies. *Hum Mol Genet* 9: 1465–1472, 2000.
242. Sather WA and McCleskey EW. Permeation and selectivity in calcium channels. *Annu Rev Physiol* 65: 133–159, 2003.
243. Sayer RJ, Brown AM, Schwandt PC, and Crill WE. Calcium currents in acutely isolated human neocortical neurons. *J Neurophysiol* 69: 1596–1606, 1993.
244. Scheffer IE and Berkovic SF. The genetics of human epilepsy. *Trends Pharmacol Sci* 24: 428–433, 2003.
245. Schiller J, Schiller Y, Stuart G, and Sakmann B. Calcium action potentials restricted to distal apical dendrites of rat neocortical pyramidal neurons. *J Physiol* 505: 605–616, 1997.
246. Scoggan KA, Chandra T, Nelson R, Hahn AF, and Bulman DE. Identification of two novel mutations in the CACNA1A gene responsible for episodic ataxia type 2. *J Med Genet* 38: 249–253, 2001.
247. Seagar M and Takahashi M. Interactions between presynaptic calcium channels and proteins implicated in synaptic vesicle trafficking and exocytosis. *J Bioenerg Biomembr* 30: 347–356, 1998.
248. Seisenberger C, Specht V, Welling A, Platzer J, Pfeifer A, Kuhbandner S, Striessnig J, Klugbauer N, Feil R, and Hofmann F. Functional embryonic cardiomyocytes after disruption of the L-type alpha1C (Cav1.2) calcium channel gene in the mouse. *J Biol Chem* 275: 39193–39199, 2000.
249. Sherman SM. Interneurons and triadic circuitry of the thalamus. *Trends Neurosci* 27: 670–675, 2004.
250. Sidach SS and Mintz IM. Kurtosis, a gating modifier of neuronal high- and low-threshold Ca channels. *J Neurosci* 22: 2023–2034, 2002.
251. Sills GJ. The mechanisms of action of gabapentin and pregabalin. *Curr Opin Pharmacol* 6: 108–113, 2006.
252. Singh NA, Charlier C, Stauffer D, DuPont BR, Leach RJ, Melis R, Ronen GM, Bjerre I, Quattlebaum T, Murphy JV, McHarg ML, Gagnon D, Rosales TO, Peiffer A, Anderson VE, and Leppert M. A novel potassium channel gene, KCNQ2, is mutated in an inherited epilepsy of newborns. *Nat Genet* 18: 25–29, 1998.
253. Slaght SJ, Leresche N, Deniau JM, Crunelli V, and Charpier S. Activity of thalamic reticular neurons during spontaneous genetically determined spike and wave discharges. *J Neurosci* 22: 2323–2334, 2002.
254. Snutch TP, Peloquin J, Mathews E, and McRory JE. Molecular properties of voltage-gated calcium channels. In: *Voltage-Gated Calcium Channels*. New York: Plenum, 2005.
255. Sohal VS and Huguenard JR. Inhibitory interconnections control burst pattern and emergent network synchrony in reticular thalamus. *J Neurosci* 23: 8978–8988, 2003.
256. Song I, Kim D, Choi S, Sun M, Kim Y, and Shin HS. Role of the alpha1G T-type calcium channel in spontaneous absence seizures in mutant mice. *J Neurosci* 24: 5249–5257, 2004.
257. Soong TW, Stea A, Hodson CD, Dubel SJ, Vincent SR, and Snutch TP. Structure and functional expression of a member of the low voltage-activated calcium channel family. *Science* 260: 1133–1136, 1993.
258. Spafford JD, Van Minnen J, Larsen P, Smit AB, Syed NI, and Zamponi GW. Uncoupling of calcium channel alpha1 and beta subunits in developing neurons. *J Biol Chem* 279: 41157–41167, 2004.
259. Spafford JD and Zamponi GW. Functional interactions between presynaptic calcium channels and the neurotransmitter release machinery. *Curr Opin Neurobiol* 13: 308–314, 2003.
260. Stefani A, Spadoni F, Siniscalchi A, and Bernardi G. Lamotrigine inhibits  $\text{Ca}^{2+}$  currents in cortical neurons: functional implications. *Eur J Pharmacol* 307: 113–116, 1996.
261. Steriade M. Impact of network activities on neuronal properties in corticothalamic systems. *J Neurophysiol* 86: 1–39, 2001.
262. Steriade M. *Neuronal Substrates of Sleep and Epilepsy*. Cambridge, UK: Cambridge Univ. Press, 2003, p. 522.
263. Steriade M. Sleep, epilepsy and thalamic reticular inhibitory neurons. *Trends Neurosci* 28: 317–324, 2005.
264. Steriade M. To burst, or rather, not to burst. *Nat Neurosci* 4: 671, 2001.
265. Steriade M and Contreras D. Relations between cortical and thalamic cellular events during transition from sleep patterns to paroxysmal activity. *J Neurosci* 15: 623–642, 1995.



266. Steriade M and Contreras D. Spike-wave complexes and fast components of cortically generated seizures. I. Role of neocortex and thalamus. *J Neurophysiol* 80: 1439–1455, 1998.
267. Stotz SC, Jarvis SE, and Zamponi GW. Functional roles of cytoplasmic loops and pore lining transmembrane helices in the voltage-dependent inactivation of HVA calcium channels. *J Physiol* 554: 263–273, 2004.
268. Stotz SC and Zamponi GW. Identification of inactivation determinants in the domain IIS6 region of high voltage-activated calcium channels. *J Biol Chem* 276: 33001–33010, 2001.
269. Strauss U, Kole MH, Brauer AU, Pahnke J, Bajorat R, Rolfs A, Nitsch R, and Deisz RA. An impaired neocortical Ih is associated with enhanced excitability and absence epilepsy. *Eur J Neurosci* 19: 3048–3058, 2004.
270. Strube C, Beur M, Powers PA, Gregg RG, and Coronado R. Reduced  $\text{Ca}^{2+}$  current, charge movement, and absence of  $\text{Ca}^{2+}$  transients in skeletal muscle deficient in dihydropyridine receptor beta 1 subunit. *Biophys J* 71: 2531–2543, 1996.
271. Su H, Sochivko D, Becker A, Chen J, Jiang Y, Yaari Y, and Beck H. Upregulation of a T-type  $\text{Ca}^{2+}$  channel causes a long-lasting modification of neuronal firing mode after status epilepticus. *J Neurosci* 22: 3645–3655, 2002.
272. Sugawara T, Tsurubuchi Y, Agarwala KL, Ito M, Fukuma G, Mazaki-Miyazaki E, Nagafuji H, Noda M, Imoto K, Wada K, Mitsudome A, Kaneko S, Montal M, Nagata K, Hirose S, and Yamakawa K. A missense mutation of the  $\text{Na}^+$  channel alpha II subunit gene Na(v)1.2 in a patient with febrile and afebrile seizures causes channel dysfunction. *Proc Natl Acad Sci USA* 98: 6384–6389, 2001.
273. Sutton KG, McRory JE, Guthrie H, Murphy TH, and Snutch TP. P/Q-type calcium channels mediate the activity-dependent feedback of syntaxin-1A. *Nature* 401: 800–804, 1999.
274. Suzuki T, Delgado-Escueta AV, Aguan K, Alonso ME, Shi J, Hara Y, Nishida M, Numata T, Medina MT, Takeuchi T, Morita R, Bai D, Ganesh S, Sugimoto Y, Inazawa J, Bailey JN, Ochoa A, Jara-Prado A, Rasmussen A, Ramos-Peek J, Cordova S, Rubio-Donnadieu F, Inoue Y, Osawa M, Kaneko S, Oguni H, Mori Y, and Yamakawa K. Mutations in EFHC1 cause juvenile myoclonic epilepsy. *Nat Genet* 36: 842–849, 2004.
275. Takahashi SX, Miriyala J, and Colecraft HM. Membrane-associated guanylate kinase-like properties of beta-subunits required for modulation of voltage-dependent  $\text{Ca}^{2+}$  channels. *Proc Natl Acad Sci USA* 101: 7193–7198, 2004.
276. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, and Bayliss DA. Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci* 19: 1895–1911, 1999.
277. Talley EM, Solorzano G, Depaulis A, Perez-Reyes E, and Bayliss DA. Low-voltage-activated calcium channel subunit expression in a genetic model of absence epilepsy in the rat. *Brain Res* 75: 159–165, 2000.
278. Tan NC, Mulley JC, and Berkovic SF. Genetic association studies in epilepsy: “the truth is out there.” *Epilepsia* 45: 1429–1442, 2004.
279. Thomas E and Grisar T. Increased synchrony with increase of a low-threshold calcium conductance in a model thalamic network: a phase-shift mechanism. *Neural Comput* 12: 1553–1571, 2000.
280. Todorovic SM and Lingle CJ. Pharmacological properties of T-type  $\text{Ca}^{2+}$  current in adult rat sensory neurons: effects of anticonvulsant and anesthetic agents. *J Neurophysiol* 79: 240–252, 1998.
281. Tomita S, Adesnik H, Sekiguchi M, Zhang W, Wada K, Howe JR, Nicoll RA, and Brecht DS. Stargazin modulates AMPA receptor gating and trafficking by distinct domains. *Nature* 435: 1052–1058, 2005.
282. Tomlinson WJ, Stea A, Bourinet E, Charnet P, Nargeot J, and Snutch TP. Functional properties of a neuronal class C L-type calcium channel. *Neuropharmacology* 32: 1117–1126, 1993.
283. Tottene A, Fellin T, Pagnutti S, Luvisetto S, Striessnig J, Fletcher C, and Pietrobon D. Familial hemiplegic migraine mutations increase  $\text{Ca}^{2+}$  influx through single human  $\text{CaV}2.1$  channels and decrease maximal  $\text{CaV}2.1$  current density in neurons. *Proc Natl Acad Sci USA* 99: 13284–13289, 2002.
284. Tsakiridou E, Bertollini L, de Curtis M, Avanzini G, and Pape HC. Selective increase in T-type calcium conductance of reticular thalamic neurons in a rat model of absence epilepsy. *J Neurosci* 15: 3110–3117, 1995.
285. Turnbull J, Lohi H, Kearney JA, Rouleau GA, Delgado-Escueta AV, Meisler MH, Cossette P, and Minassian BA. Sacred disease secrets revealed: the genetics of human epilepsy. *Hum Mol Genet* 14: 2491–2500, 2005.
286. Ulrich D and Huguenard JR. GABA(A)-receptor-mediated rebound burst firing and burst shunting in thalamus. *J Neurophysiol* 78: 1748–1751, 1997.
287. Vadlamudi L, Scheffer IE, and Berkovic SF. Genetics of temporal lobe epilepsy. *J Neurol Neurosurg Psychiatry* 74: 1359–1361, 2003.
288. Vajna R, Klockner U, Pereverzev A, Weiergraber M, Chen X, Miljanich G, Klugbauer N, Hescheler J, Perez-Reyes E, and Schneider T. Functional coupling between “R-type”  $\text{Ca}^{2+}$  channels and insulin secretion in the insulinoma cell line INS-1. *Eur J Biochem* 268: 1066–1075, 2001.
289. Van den Maagdenberg AM, Pietrobon D, Pizzorusso T, Kaja S, Broos LA, Cesetti T, van de Ven RC, Tottene A, van der Kaa J, Plomp JJ, Frants RR, and Ferrari MD. A Cacna1a knockin migraine mouse model with increased susceptibility to cortical spreading depression. *Neuron* 41: 701–710, 2004.
290. Vandenberghe W, Nicoll RA, and Brecht DS. Stargazin is an AMPA receptor auxiliary subunit. *Proc Natl Acad Sci USA* 102: 485–490, 2005.
291. Van Petegem F, Clark KA, Chatelain FC, and Minor DL Jr. Structure of a complex between a voltage-gated calcium channel beta-subunit and an alpha-subunit domain. *Nature* 429: 671–675, 2004.
292. Vitko I, Chen Y, Arias JM, Shen Y, Wu XR, and Perez-Reyes E. Functional characterization and neuronal modeling of the effects of childhood absence epilepsy variants of CACNA1H, a T-type calcium channel. *J Neurosci* 25: 4844–4855, 2005.
293. Wakamori M, Yamazaki K, Matsunodaira H, Teramoto T, Tanaka I, Niidome T, Sawada K, Nishizawa Y, Sekiguchi N, Mori E, Mori Y, and Imoto K. Single tottering mutations responsible for the neuropathic phenotype of the P-type calcium channel. *J Biol Chem* 273: 34857–34867, 1998.
294. Wallace RH, Marini C, Petrou S, Harkin LA, Bowser DN, Panchal RG, Williams DA, Sutherland GR, Mulley JC, Scheffer IE, and Berkovic SF. Mutant GABA(A) receptor gamma2-subunit in childhood absence epilepsy and febrile seizures. *Nat Genet* 28: 49–52, 2001.
295. Wallace RH, Wang DW, Singh R, Scheffer IE, George AL Jr, Phillips HA, Saar K, Reis A, Johnson EW, Sutherland GR, Berkovic SF, and Mulley JC. Febrile seizures and generalized epilepsy associated with a mutation in the  $\text{Na}^+$ -channel beta1 subunit gene SCN1B. *Nat Genet* 19: 366–370, 1998.
296. Wang G, Dayanithi G, Newcomb R, and Lemos JR. An R-type  $\text{Ca}^{2+}$  current in neurohypophyseal terminals preferentially regulates oxytocin secretion. *J Neurosci* 19: 9235–9241, 1999.
297. Wang SJ, Huang CC, Hsu KS, Tsai JJ, and Gean PW. Inhibition of N-type calcium currents by lamotrigine in rat amygdalar neurons. *Neuroreport* 7: 3037–3040, 1996.
298. Weick JP, Groth RD, Isaksen AL, and Mermelstein PG. Interactions with PDZ proteins are required for L-type calcium channels to activate cAMP response element-binding protein-dependent gene expression. *J Neurosci* 23: 3446–3456, 2003.
299. Welsby PJ, Wang H, Wolfe JT, Colbran RJ, Johnson ML, and Barrett PQ. A mechanism for the direct regulation of T-type calcium channels by  $\text{Ca}^{2+}$ /calmodulin-dependent kinase II. *J Neurosci* 23: 10116–10121, 2003.
300. Westenbroek RE, Hell JW, Warner C, Dubel SJ, Snutch TP, and Catterall WA. Biochemical properties and subcellular distribution of an N-type calcium channel alpha 1 subunit. *Neuron* 9: 1099–1115, 1992.
301. Westenbroek RE, Sakurai T, Elliott EM, Hell JW, Starr TV, Snutch TP, and Catterall WA. Immunohistochemical identification and subcellular distribution of the alpha 1A subunits of brain calcium channels. *J Neurosci* 15: 6403–6418, 1995.

302. Wheeler DB, Randall A, and Tsien RW. Changes in action potential duration alter reliance of excitatory synaptic transmission on multiple types of  $\text{Ca}^{2+}$  channels in rat hippocampus. *J Neurosci* 16: 2226–2237, 1996.
303. Wiest MC and Nicoletis MA. Behavioral detection of tactile stimuli during 7–12 Hz cortical oscillations in awake rats. *Nat Neurosci* 6: 913–914, 2003.
304. Williams ME, Feldman DH, McCue AF, Brenner R, Velicelebi G, Ellis SB, and Harpold MM. Structure and functional expression of alpha 1, alpha 2, and beta subunits of a novel human neuronal calcium channel subtype. *Neuron* 8: 71–84, 1992.
305. Williams ME, Marubio LM, Deal CR, Hans M, Brust PF, Philipson LH, Miller RJ, Johnson EC, Harpold MM, and Ellis SB. Structure and functional characterization of neuronal alpha 1E calcium channel subtypes. *J Biol Chem* 269: 22347–22357, 1994.
306. Wolfe JT, Wang H, Howard J, Garrison JC, and Barrett PQ. T-type calcium channel regulation by specific G-protein beta-gamma subunits. *Nature* 424: 209–213, 2003.
307. Wu XS, Edwards HD, and Sather WA. Side chain orientation in the selectivity filter of a voltage-gated  $\text{Ca}^{2+}$  channel. *J Biol Chem* 275: 31778–31785, 2000.
308. Xu W, and Lipscombe D. Neuronal  $\text{Ca(V)}_{1.3\alpha(1)}$  L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. *J Neurosci* 21: 5944–5951, 2001.
309. Yang J, Ellinor PT, Sather WA, Zhang JF, and Tsien RW. Molecular determinants of  $\text{Ca}^{2+}$  selectivity and ion permeation in L-type  $\text{Ca}^{2+}$  channels. *Nature* 366: 158–161, 1993.
310. Yasuda T, Chen L, Barr W, McRory JE, Lewis RJ, Adams DJ, and Zamponi GW. Auxiliary subunit regulation of high-voltage activated calcium channels expressed in mammalian cells. *Eur J Neurosci* 20: 1–13, 2004.
311. Yokoyama CT, Westenbroek RE, Hell JW, Soong TW, Snutch TP, and Catterall WA. Biochemical properties and subcellular distribution of the neuronal class E calcium channel alpha 1 subunit. *J Neurosci* 15: 6419–6432, 1995.
312. Yoshinaga H, Ohtsuka Y, Tamai K, Tamura I, Ito M, Ohmori I, and Oka E. EEG in childhood absence epilepsy. *Seizure* 13: 296–302, 2004.
313. Yue Q, Jen JC, Thwe MM, Nelson SF, and Baloh RW. De novo mutation in CACNA1A caused acetazolamide-responsive episodic ataxia. *Am J Med Genet* 77: 298–301, 1998.
314. Yuste R, Gutnick MJ, Saar D, Delaney KR, and Tank DW.  $\text{Ca}^{2+}$  accumulations in dendrites of neocortical pyramidal neurons: an apical band and evidence for two functional compartments. *Neuron* 13: 23–43, 1994.
315. Zamponi GW. Determinants of G protein inhibition of presynaptic calcium channels. *Cell Biochem Biophys* 34: 79–94, 2001.
316. Zhang JF, Randall AD, Ellinor PT, Horne WA, Sather WA, Tanabe T, Schwarz TL, and Tsien RW. Distinctive pharmacology and kinetics of cloned neuronal  $\text{Ca}^{2+}$  channels and their possible counterparts in mammalian CNS neurons. *Neuropharmacology* 32: 1075–1088, 1993.
317. Zhang Y, Mori M, Burgess DL, and Noebels JL. Mutations in high-voltage-activated calcium channel genes stimulate low-voltage-activated currents in mouse thalamic relay neurons. *J Neurosci* 22: 6362–6371, 2002.
- 317a. Zhong X, Liu JR, Kyle JW, Hanck DA, and Agnew WS. A profile of alternative RNA splicing and transcript variation of CACNA1H, a human T-channel gene candidate for idiopathic generalized epilepsies. *Hum Mol Genet* 15: 1497–1512, 2006.
318. Zona C, Niespodziany I, Marchetti C, Klitgaard H, Bernardi G, and Margineanu DG. Levetiracetam does not modulate neuronal voltage-gated  $\text{Na}^{+}$  and T-type  $\text{Ca}^{2+}$  currents. *Seizure* 10: 279–286, 2001.
319. Zuberi SM, Eunson LH, Spauschus A, De Silva R, Tolmie J, Wood NW, McWilliam RC, Stephenson JP, Kullmann DM, and Hanna MG. A novel mutation in the human voltage-gated potassium channel gene (Kv1.1) associates with episodic ataxia type 1 and sometimes with partial epilepsy. *Brain* 122: 817–825, 1999.
320. Zuhlke RD, Pitt GS, Deisseroth K, Tsien RW, and Reuter H. Calmodulin supports both inactivation and facilitation of L-type calcium channels. *Nature* 399: 159–162, 1999.
321. Zwingman TA, Neumann PE, Noebels JL, and Herrup K. Ricker is a new variant of the voltage-dependent calcium channel gene CACNA1A. *J Neurosci* 21: 1169–1178, 2001.