



Statistical Modeling of Structural Connectomes Reveal High Genetic Influence on Connectivity



Jaewon Chung^{1*}, Eric Bridgeford¹, Michael Powell¹, Joshua T. Vogelstein¹

1 - Johns Hopkins University, * - correspondence:  j1c@jhu.edu  [@j1c\(Github\)](https://github.com/j1c)  [@j1chung\(Twitter\)](https://twitter.com/j1chung)

Summary

- Aimed to define heritability for populations of connectomes using statistical modelling.
- Structural connectomes are heritable without controlling for neuroanatomy.
- Neuroanatomy is also highly heritable
- Connectomes remain heritable after controlling for effects of neuroanatomy on connectomes.
- Provide tools for future analysis on populations of connectomes.

Motivation

- Understanding how brain connectivity is influenced by genetics can improve our understanding of brain function and diseases.
- Current methods of analyzing connectomes or hertability exhibit limitations:
 - Selection Graph theoretic features
 - Multivariate normality assumptions

Overview of Analysis

Fig 1: Overview of the framework for measuring heritability of connectomes.

Do changes in **genome** cause changes in **connectomes**?

Causal Analysis of Effect of Genome on Connectomes

- Genome directly affects the structural connectome.
- Neuroanatomym (e.g. brain volume) indirectly affects the connectome.
- Participant history, such as the shared and non-shared environmental influences, and traits are potential confounders.
- The shared and non-shared environment is controlled by comparing between the same sex individuals.



Fig 2: Directed acyclic graph (DAG) illustrating potential relationships between the genome and connectome.

Human Connectome Project 1200

- Structural connectomes are estimated using structural (sMRI) and diffusion magnetic resonance imaging (dMRI).
- Processed with *m2g* pipeline, which uses Constrained Spherical Deconvolution (CSD) model and deterministic tractography.

	Monozygotic	Dizygotic	Non-twin siblings
N	322	212	490
Sex	196 F, 126 M	125 F, 87 M	237 F, 253 M
Age (mean)	29.6 (3.3)	28.9 (3.4)	28.3 (3.9)

Table 1: Participants and their demographics of HCP1200 Dataset.

Three Models of Connectomes

- Exact:** Are the generative models of connectomes the same?
- Global scale:** Are the generative models same after considering global scaling?
- Vertex scale:** Are the generative models same after considering vertex wise scaling?

Fig 3: Examples of the three different models (exact, global scale, and vertex scale) of connectome heritability visualized as adjacency matrices.

Connectome Validation and Heritability of Neuroanatomy

Fig 4: Validating connectomes by comparing monozygotic and dizygotic twins using Kolmogrov-Smirnov test. Heritability of neuroanamy is tested using unconditional test framework. Red squares indicate significant tests; blue indicate non-significant tests.

Tests for Heritability of Connectomes

Fig 5: Testing for unconditional and conditional heritability of connectomes. Red squares indicate significant tests; blue indicate non-significant tests.

Limitations and extensions

- Other staitistical models to consider (e.g. COSIE [3])
- Repeated analysis on functional MRI or in other twin study datasets.

Code



downloads 161k
Stars 268



downloads 107k
Stars 133



References

- [1] Chung et al. "Connectomic Heritability," In preparation (2022)
- [2] Chung et al. "Statistical connectomics," Ann. Rev. Statistics and its Application (2021)
- [3] Arroyo et al. "Inference for multiple heterogeneous networks with a common invariant subspace," JMLR (2021)

Funding

J.C. supported by the BRAIN Initiative (1RF1MH123233). J.T.V. supported by NSF CAREER Award (1942963). Findings and conclusions expressed are those of the authors and not necessarily those of the funders.

Acknowledgements

NeuroData lab for many ideas and feedback. Many at Microsoft Research for w/ graspolagic.