

A database of the healthy human spinal cord morphometry in the PAM50 template space

- Jan Valošek ¹,2,3,4*, Sandrine Bédard ¹, Miloš Keřkovský ⁵, Tomáš
- Rohan ⁶ , and Julien Cohen-Adad ⁶ ^{1,2,6,7}
- Reproducible Preprint 5 1 NeuroPoly Lab, Institute of Biomedical Engineering, Polytechnique Montreal, Montreal, QC, Canada 2
 - $_{6}$ Mila Quebec Al Institute, Montreal, QC, Canada 3 Department of Neurosurgery, Faculty of Medicine
 - and Dentistry, Palacký University Olomouc, Olomouc, Czechia 4 Department of Neurology, Faculty of
 - 8 Medicine and Dentistry, Palacký University Olomouc, Olomouc, Czechia 5 Department of Radiology and
 - 9 Nuclear Medicine, University Hospital Brno and Masaryk University, Brno, Czechia 6 Functional
 - Neuroimaging Unit, CRIUGM, Université de Montréal, Montreal, QC, Canada 7 Centre de recherche du CHU Sainte-Justine, Université de Montréal, Montreal, QC, Canada * These authors contributed equally.

Technical Screening ♂Submitted Repository ♂

Code

DOI: 10.xxxxx/draft

■ Jupyter Book 🗗

- Reproducibility Assets

 Repository □
 - Dataset 🗗
 - Jupyter Book 🗗
 - Container ☑

Moderator: Agah Karakuzu ♂ Screener(s):

@mathieuboudreau

Submitted: 04 September 2023 ²⁰ **Published:** unpublished ²¹

License

Authors of papers retain copyright³ and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

Spinal cord morphometry measures derived from magnetic resonance imaging (MRI) scans serve as valuable prognostic biomarkers for various spinal cord pathologies. Despite their significance, interpreting these biomarkers is challenging due to substantial variability between subjects. The lack of a standardized normalization method to mitigate this variability and the need for a better understanding of morphometric distribution contribute to the current knowledge gap.

In this work, we present a database of healthy normative values for six commonly used measures of spinal cord morphometry built using a new fully-automatic normalization approach. Morphometric measures were computed from a large open-access dataset of healthy adult volunteers (N=203) and brought to the common space of the PAM50 spinal cord template using a newly proposed normalization method based on linear interpolation Figure 1.

The database is interactive, available online (TODO: add a link to interactive preprint) and allows filtering for sex, age, and MRI vendors. The proposed method is open-source and easily accessible through the Spinal Cord Toolbox (SCT) v6.0 and higher.

This new morphometric database allows researchers to normalize morphometrics based on sex and age, thereby minimizing inter-subject variability associated with demographic and biological factors.



Figures




Figure 1: Schematic representation of the normalization approach. (A) T2-weighted images of 203 participants from the spine-generic dataset (multi-subject) were used. The spinal cord was segmented, and vertebral levels were identified automatically using the Spinal Cord Toolbox (SCT). (B) Six morphometric measures were computed for each axial slice from the single-subject segmentation masks. (C) For each level, the number of slices in the subject native space and the corresponding level in the PAM50 template (D) were identified. Then, the morphometric measures were linearly interpolated to the PAM50 space using the number of slices in the PAM50 template and the subject native space for each vertebral level.

Acknowledgements

37

- Thanks to Nick Guenther and Mathieu Guay-Paquet for their assistance with dataset management. We would like to thank Joshua Newton for his contributions in helping us implement the algorithm to SCT. We express our gratitude to Allan R. Martin for his insightful discussions on the clinical aspects of the work. We would also like to thank Nathan Molinier for providing 35 valuable feedback on the manuscript figures. We acknowledge all participants as well as 36 collaborators of the spine-generic study.
- Funded by the Canada Research Chair in Quantitative Magnetic Resonance Imaging [CRC-2020-38 00179], the Canadian Institute of Health Research [PJT-190258], the Canada Foundation for 39 Innovation [32454, 34824], the Fonds de Recherche du Québec - Santé [322736], the Natural Sciences and Engineering Research Council of Canada [RGPIN-2019-07244], the Canada First Research Excellence Fund (IVADO and TransMedTech), the Courtois NeuroMod project, the 42 Quebec Biolmaging Network [5886, 35450], INSPIRED (Spinal Research, UK; Wings for Life, Austria; Craig H. Neilsen Foundation, USA), Mila - Tech Transfer Funding Program. Supported by the Ministry of Health of the Czech Republic, grant nr. NU22-04-00024. All rights reserved. JV has received funding from the European Union's Horizon Europe research and innovation programme under the Marie Sktodowska-Curie grant agreement No 101107932.