

Emergency plan

A card to complete with your team: signs, steps and key contacts.

Warning signs and emergencies

What to watch, how to respond and when to seek urgent help.

When to seek urgent help

When in doubt, call emergency services

This guide is general and does not replace the plan your team gave you or your country's emergency services. If there is a threat to life, call your local emergency number immediately.

- Sudden weakness in the face, arm or leg, trouble speaking or seeing (possible stroke): act fast, every minute counts.
- A seizure lasting more than 5 minutes, or repeated seizures without regaining consciousness.
- A sudden, extremely severe headache, unlike the usual.
- High fever with a stiff neck or confusion.
- Difficulty breathing, choking or loss of consciousness.

First aid during a seizure

1. Stay calm and protect: move away objects they could hit.
2. Place something soft under the head and loosen anything tight around the neck.
3. Do not restrain them and do not put anything in the mouth.
4. Turn them onto their side (recovery position) to help breathing.
5. Time the seizure; if it passes 5 minutes or repeats, call emergency services.
6. Stay with them until they recover and follow the plan given by their doctor.

After the seizure

Drowsiness or confusion afterward is normal. Speak softly, offer reassurance and record what happened for the medical team.

Preparing in advance

- ✓ A written action plan from your team (what to do and who to call).
- ✓ A list of medications and allergies, visible and up to date.
- ✓ Emergency contacts and the neurologist within reach.
- ✓ A medical ID document or bracelet if recommended.