Universal γ-scaling at the edge of metastability: evidence from three independent biological substrates with simulation validation

Yaroslav Vasylenko
neuron7xLab, Poltava region, Ukraine
Independent researcher (no institutional affiliation)
Contact: github.com/neuron7xLab

Abstract

We report empirical evidence for a universal scaling exponent $\gamma \approx 1.0$ observed across three independent biological substrates with additional simulation validation. **Tier 1 — Evidential (real external data):** zebrafish morphogenesis ($\gamma = 1.055$, $n = 47$, CI: [0.89, 1.20], McGuirl 2020), human heart rate variability ($\gamma = 0.885$, CI: [0.83, 1.08], PhysioNet NSR2DB), and human EEG during motor imagery ($\gamma \approx 1.07$, $n = 20$ subjects, CI: [0.88, 1.25], PhysioNet EEGBCI). **Tier 2 — Simulation-validated:** Gray–Scott reaction-diffusion ($\gamma = 0.938$), Kuramoto oscillators at K_c ($\gamma = 0.980$), and BN-Syn spiking criticality ($\gamma \approx 0.49$, honest finite-size deviation from mean-field prediction). Cross-substrate mean from evidential substrates only: $\bar{\gamma}$ with 95% CI containing unity. All Tier 1 substrates pass surrogate testing ($p < 0.05$), and three negative controls (white noise, random walk, supercritical) show γ clearly separated from unity. The BN-Syn finite-size result ($\gamma \approx 0.49$ for $N = 200$, $k = 10$) is consistent with theoretical predictions of finite-size corrections below the upper critical dimension, validating that $\gamma \approx 1.0$ in biological substrates is a genuine property rather than a methodological artifact. We propose that $\gamma \approx 1.0$ constitutes a scaling signature of metastability — the dynamical regime where complex systems maintain coherent computation at the boundary between order and disorder.

1. Introduction

Complex systems across vastly different substrates — from biological tissues to neural networks to financial markets — share a common dynamical feature: they operate most effectively near critical points, at the boundary between ordered and disordered phases [1, 2]. This regime, termed *metastability*, is characterized by long-range correlations, power-law scaling, and the capacity for flexible reconfiguration [3].

Self-organized criticality (SOC) predicts that driven dissipative systems naturally evolve toward critical states exhibiting power-law distributions [1]. In neural systems, evidence for criticality has been found in the form of neuronal avalanches with power-law size distributions [2]. The branching ratio $\sigma \approx 1.0$ — the mean number of downstream activations per event — serves as a diagnostic for criticality in these systems.

We introduce a complementary diagnostic: the *gamma-scaling exponent* γ, defined through the power-law relation between topological complexity C and thermodynamic cost K:

$$K \sim C^{-\gamma} \tag{1}$$

where C is a measure of the system's structural or information complexity and K is the energetic or computational cost per unit of complexity. We present evidence that $\gamma \approx 1.0$ across five independent substrates, suggesting it may represent a universal signature of metastable computation.

The extended mind thesis [4] proposes that cognitive processes extend beyond the brain into the environment. We test this framework empirically by treating the human–AI interaction loop as a measurable physical system and showing that productive cognitive coupling exhibits the same γ-scaling as biological and physical substrates.

2. Theoretical Framework

2.1 Scaling relation

For a system with topological complexity C (measuring information richness, structural diversity, or phase-space dimensionality) and thermodynamic cost K (measuring energy expenditure, computational effort, or dissipation per unit complexity), we define the gamma-scaling relation:

$$K = A \cdot C^{-\gamma} \tag{2}$$

Taking logarithms:

$$\log K = -\gamma \log C + \log A \tag{3}$$

The exponent γ characterizes the system's efficiency-complexity tradeoff:

- $\gamma > 1$: *over-determined* — cost decreases faster than complexity increases (convergent regime)
- $\gamma < 1$: *under-determined* — cost decreases slower than complexity increases (divergent regime)
- $\gamma = 1$: *critical balance* — cost and complexity scale inversely at unit rate (metastable regime)

2.2 Connection to criticality diagnostics

The gamma-scaling exponent relates to established criticality measures:

- **Branching ratio** σ: In spiking networks, $\sigma \approx 1.0$ indicates criticality [2]. Our BN-Syn substrate measures $\gamma = 0.950$ when σ is tuned to the critical regime.
- **Kuramoto order parameter** r: In coupled oscillator systems, coherence $r \in [0, 1]$ measures synchronization. Our market substrate computes γ from Kuramoto coherence trajectories, yielding $\gamma = 1.081$.[1]
- **Spectral radius** ρ: The largest eigenvalue of the system's Jacobian. In the neosynaptex cross-domain integrator, $\rho \approx 1.0$ and $\gamma \approx 1.0$ co-occur in the METASTABLE phase.

2.3 Information-theoretic complexity

For the human–AI cognitive substrate, we define complexity using Shannon information theory:

$$C = H(W) \cdot \log(1 + |V|) \cdot \mathrm{TTR} \tag{4}$$

where $H(W) = -\sum_i p_i \log_2 p_i$ is the Shannon entropy of the word frequency distribution, $|V|$ is the vocabulary size (unique words), and $\mathrm{TTR} = |V|/|W|$ is the type-token ratio. The thermodynamic cost is:

$$K = \frac{|W|}{C} \tag{5}$$

measuring total words (effort) per unit of information complexity.

[1] $\gamma = 1.081$ refers to the market Kuramoto substrate (financial coherence trajectories, illustrative). $\gamma = 0.980$ in Table 1 refers to simulated Kuramoto oscillators at critical coupling K_c. These are distinct substrates.

2.4 Hypotheses

2.4.1 Hypothesis H1: Intelligence as a Dynamical Regime

Statement: Truth is not inevitable — it is the result of independent verification between autonomous witnesses. Within the framework of H1, intelligence is defined as a dynamical regime verified through synchronous phase shifts (γ) across independent channels of a single substrate, with coherent recovery.

Formalization:

$$\forall\, S_i \in \{\text{substrates at metastability}\} : \ \gamma_{S_i} \in [0.85, 1.15] \tag{6}$$

with 95% CI containing 1.0.

Verification criterion: H1 is supported if $\gamma \in [0.85, 1.15]$ with 95% CI containing 1.0 across $N \geq 3$ independent substrates from distinct physical domains, each passing surrogate testing ($p < 0.05$).

Status: SUPPORTED — three independent biological substrates (zebrafish, HRV PhysioNet, EEG PhysioNet), cross-substrate CI from Tier 1 contains unity. All Tier 1 IAAFT p-values < 0.05. BN-Syn finite-size deviation ($\gamma \approx 0.49$) confirms methodology is not trivially producing $\gamma \approx 1.0$.

2.4.2 Hypothesis H2: Computational Efficiency Is a Regime Property

Statement: The regime $\gamma \approx 1$ corresponds to a state that maximizes computational capacity at minimal cost of plasticity maintenance. This is an open claim requiring separate experimental and theoretical verification.

Energy-Regime Conjecture ($\mathcal{C}_E$):

$$\mathcal{C}_E : \quad \gamma \approx 1 \Longleftrightarrow \text{local min of dissipation preserving plasticit} \tag{7}$$

Status of $\mathcal{C}_E$: CONJECTURE — not theorem, not derivation. Qualitative argument only. Formal derivation of the $\beta \to \varepsilon$ bridge is open work.

2.5 Theoretical basis: $\gamma = 1.0$ in mean-field criticality

The result $\gamma = 1.0$ follows from mean-field theory of critical phenomena in multiple universality classes.

Branching process at $\sigma = 1$. In a critical branching process, each event generates on average $\sigma = 1$ successor. The cost of propagating one unit of topological information is exactly one unit of energy [7]. This gives $K = C^{-1}$ directly, yielding $\gamma = 1$.

Self-organized criticality. In the mean-field BTW sandpile [1], avalanche size S and duration T satisfy $\langle S \rangle \sim T^{d_f/d}$. In mean-field ($d \geq d_c$), $d_f = d$, giving $\langle S \rangle \sim T^1$. The cost-complexity ratio $K/C \sim S/T = \text{const}$, yielding $\gamma = 1$.

Directed percolation universality. Neural criticality belongs to the directed percolation universality class [8, 2]. In mean-field DP, the branching ratio $\sigma = 1$ at the critical point, and $\tau = 3/2$ (avalanche size exponent). The scaling relation $\gamma = (\tau_T - 1)/(\tau_S - 1)$ evaluates to exactly 1.0 in mean field.

Finite-size corrections. Below the upper critical dimension d_c, corrections of order $\varepsilon = d_c - d$ appear, pushing γ away from 1.0. Our BN-Syn simulation ($N = 200$ neurons, $k = 10$ sparse connectivity) yields $\gamma \approx 0.49$, consistent with finite-size deviations from the mean-field prediction.

Spectral connection. At SOC, the power spectral density follows $S(f) \sim f^{-\beta}$ with $\beta = 1$ ($1/f$ noise) [1]. The spectral exponent β is related to the Hurst exponent H via $\beta = 2H + 1$ (for fractional Brownian motion), giving $H = 0$ at criticality. In the HRV VLF range and EEG aperiodic component, $\beta \approx 1.0$ during healthy/active states corresponds to $\gamma_{\text{PSD}} \approx 1.0$, consistent with the topo-cost framework.

3. Methods

3.1 Regression

We use Theil–Sen regression [5] — a robust, non-parametric estimator of the slope in the $(\log C, \log K)$ plane. Unlike ordinary least squares, Theil–Sen is resistant to outliers (breakdown point of 29.3%), making it appropriate for noisy empirical data.

Note on fitting method. We fit a deterministic scaling relation $K = A \cdot C^{-\gamma}$ in log-log space, not a power-law probability distribution. For scaling relations between two measured quantities, Theil–Sen regression on $(\log C, \log K)$ pairs is the appropriate estimator. The MLE framework of Clauset, Shalizi & Newman [9] applies to probability distributions $P(x) \sim x^{-\alpha}$; it is not applicable to scaling relations between paired observables.

3.2 Confidence intervals

Bootstrap confidence intervals are computed by resampling with replacement ($B = 500$ iterations) and taking the 2.5th and 97.5th percentiles of the bootstrap distribution of γ.

3.3 Quality gates

We apply three gates before accepting a γ estimate:

1. **Minimum data**: $n \geq 8$ valid pairs
2. **Dynamic range**: $\text{range}(\log C) \geq 0.5$ (sufficient variation)
3. **Fit quality**: $R^2 \geq 0.3$ for individual substrates (relaxed for noisy behavioral data)

3.4 Unified topo-cost framework: spatial and temporal domains

A critical methodological point: the power spectral density $S(f) \sim f^{-\gamma}$ IS a topo-cost relationship. In this temporal formulation:

- **Topo (complexity):** Frequency f — finer temporal structure = higher topological complexity
- **Cost (energy):** $S(f)$ — power spectral density at frequency f, measuring energy invested at that complexity scale

The scaling relation $S(f) = A \cdot f^{-\gamma}$ has exactly the form $K = A \cdot C^{-\gamma}$ from §2.1, where $C = f$ and $K = S(f)$.

3.5 Surrogate testing

To verify that observed γ values are not artifacts of sample structure, we employ IAAFT (Iterative Amplitude Adjusted Fourier Transform) surrogates. For each substrate, we generate $M = 199$ surrogates preserving the amplitude distribution and power spectrum but destroying temporal ordering. We compute a two-tailed p-value:

$$p = \frac{1 + \#\{|\gamma_{\text{null}}| \geq |\gamma_{\text{obs}}|\}}{M + 1} \tag{8}$$

3.6 Negative controls (falsification boundary)

To demonstrate that $\gamma \approx 1.0$ is not a trivial outcome of the methodology, we compute γ for four classes of systems that should NOT exhibit metastable scaling:

1. **White noise**: uniform random topo and cost (no structure)
2. **Random walk**: cumulative random walk topo, independent random cost
3. **Supercritical**: exponential growth with cost $\sim$ topo^2
4. **Subcritical ordered**: cost $\sim$ topo^{-3}

4. Results

4.1 Tier 1: Evidential substrates (real external data)

Table 1 summarizes γ-scaling across all substrates. All three Tier 1 substrates fall within the metastable band ($|\gamma - 1| < 0.15$) with cross-substrate mean $\bar{\gamma} = 1.003 \pm 0.083$.

DFA cross-validation (HRV). DFA exponent $\alpha = 1.107 \pm 0.047$ ($n = 10$ subjects, range $[1.04, 1.18]$), confirming $1/f$ scaling. For stationary processes, $\alpha = (1 + \beta)/2$; $\alpha \approx 1.1$ corresponds to $\beta \approx 1.2$, consistent with our PSD-based $\gamma = 0.885$.

4.2 Tier 2: Simulation-validated substrates

Gray–Scott and Kuramoto yield γ near 1.0, consistent with mean-field predictions (Table 1). BN-Syn ($N = 200$, $k = 10$) yields $\gamma \approx 0.49$, a finite-size deviation confirming the methodology does not trivially produce $\gamma \approx 1.0$.

Alternative model comparison. For each Tier 1 substrate, we compare the power-law scaling model against lognormal (quadratic in log-log) and exponential ($K = Ae^{-\lambda C}$) alternatives using AIC. Zebrafish: power-law preferred ($\Delta\text{AIC}_{\text{ln}} = +1.2$, $\Delta\text{AIC}_{\text{exp}} = +60.9$). HRV: power-law preferred ($\Delta\text{AIC}_{\text{ln}} = +1.5$, $\Delta\text{AIC}_{\text{exp}} = +29.7$). EEG: lognormal preferred ($\Delta\text{AIC} = -76.4$), consistent with the spectral knee in broad-band EEG PSD; the power-law fit remains valid within the aperiodic range but the full 2–35 Hz PSD shows curvature.

4.3 Effect sizes and statistical power

All Tier 1 substrates have Cohen's $d > 11$ (vs null $\gamma = 0$), indicating overwhelming evidence for non-zero scaling. Cross-substrate: $\bar{\gamma} = 1.003$, $t(2) = 0.06$, $p > 0.9$ for $H_0\colon \gamma = 1.0$ (Table 2).

Power caveat. The zebrafish substrate ($n = 47$) has CI width $\sim$0.41 and MDE of $\Delta\gamma = 0.35$ at 80% power. This means values in the range $\gamma \in [0.70, 1.35]$ cannot be reliably distinguished from $\gamma = 1.0$ with this sample size. The convergence of three independent substrates to the metastable band mitigates this limitation, but replication with larger n is needed to narrow the individual CIs.

4.4 Negative controls

Systems without critical dynamics show γ clearly separated from unity (Table 2), confirming falsifiability.

5. Discussion

5.1 Universality of gamma

Three independent biological substrates — zebrafish morphogenesis, human cardiac rhythm, and human EEG during motor imagery — all yield γ values within the metastable band ($|\gamma - 1| < 0.15$). Each substrate independently passes IAAFT surrogate testing ($p < 0.05$), confirming that the observed γ values are not artifacts of autocorrelation or spectral structure. The cross-substrate mean $\bar{\gamma} = 1.003$ with $t(2) = 0.06$ ($p > 0.9$ for $H_0 : \gamma = 1.0$) is statistically indistinguishable from unity.

The cross-substrate mean has a 95% CI containing unity. Two simulation substrates (Gray–Scott, Kuramoto) further corroborate the $\gamma \approx 1.0$ prediction at criticality. Critically, the BN-Syn spiking network ($N = 200$, $k = 10$, $\sigma = 1$) yields $\gamma \approx 0.49$ — an honest finite-size deviation from the mean-field prediction.

5.2 Speculative implications for architecture design

We speculate that the $\gamma \approx 1.0$ signature may distinguish phase-dynamic systems (which encode information in spike timing and phase coherence) from rate-based architectures (which encode in activation magnitudes). Transformer architectures achieve remarkable performance through parameter scaling [6], but operate within a rate-based



Figure 1. Topo-cost scaling across six substrates. *Row 1* (Tier 1, evidential): zebrafish morphogenesis, HRV PhysioNet, EEG PhysioNet. *Row 2* (Tier 2, simulation): Gray–Scott, Kuramoto, BN-Syn. Red lines: Theil–Sen robust regression. Each panel shows γ and 95% CI.

Table 1. Gamma-scaling across all substrates. Tier 1 (evidential, real external data) and Tier 2 (simulation-validated). Cross-substrate mean from Tier 1: $\bar{\gamma} = 1.003 \pm 0.083$, 95% CI containing unity. †EEG R^2 is not applicable: γ is computed as the per-subject mean aperiodic spectral exponent via specparam, not from log-log regression of topo-cost pairs.

Tier	Substrate	γ	95% CI	n	R^2	IAAFT p	$\ln C$ range
1	Zebrafish (McGuirl 2020)	1.055	[0.89, 1.20]	45	0.76	0.005	[−8.1, −7.1]
1	HRV PhysioNet (NSR2DB)	0.885	[0.83, 1.08]	10	0.93	<0.05	[−5.5, −3.2]
1	EEG PhysioNet (EEGBCI)	∼1.07	[0.88, 1.25]	20	n/a†	0.020	[0.8, 3.6]
2	Gray–Scott reaction-diffusion	0.938	[0.93, 0.95]	200	0.99	—	—
2	Kuramoto oscillators (K_c)	0.980	[0.93, 1.01]	300	0.42	—	—
2	BN-Syn spiking criticality	∼0.49	—	1990	—	—	—

Table 2. Statistical power (Tier 1 substrates) and negative controls. Left: all substrates have Cohen's $d > 11$. Right: all controls show γ clearly separated from unity.

Effect sizes

Substrate	γ	SE	$\lvert\gamma-1\rvert$	MDE	d
Zebrafish	1.055	0.078	0.055	0.219	13.5
HRV	0.885	0.063	0.115	0.176	14.1
EEG	1.068	0.094	0.068	0.264	11.3

Negative controls

Control	γ	R^2	Verdict	Δ
White noise	−0.06	−0.11	LOW_R2	>1.0
Random walk	0.18	−0.02	LOW_R2	>0.8
Supercritical	2.00	1.00	COLLAPSE	>0.9



Figure 2. Cross-substrate γ convergence by tier. Green: Tier 1 (evidential). Blue: Tier 2 (simulation). Dashed line: $\gamma = 1.0$ reference. Error bars: 95% bootstrap CI. Note: overlapping CIs across Tier 1 substrates are consistent with the hypothesis that all share a common $\gamma \approx 1.0$; the wide individual CIs reflect small per-substrate n.



Figure 3. Negative controls. Shaded band: metastable zone $[0.85, 1.15]$. All controls fall outside, confirming falsifiability of the $\gamma \approx 1.0$ claim.

paradigm. Whether rate-based systems can endogenously achieve $\gamma \approx 1.0$ remains an open empirical question not addressed by the data in this paper.

5.3 Mechanistic basis for cross-domain Hurst convergence

Why should the Hurst exponent H in zebrafish morphogenesis and H in financial markets produce the same $\gamma \approx 1.0$? The answer lies in universality class:

1. **Scale-free fluctuations at criticality.** $S(f) \propto f^{-\beta}$ where $\beta = 2H+1$ holds regardless of substrate.
2. **SOC.** Systems that self-tune to the edge of instability generically produce $1/f$ noise ($\beta \approx 1$, $H \approx 0$, $\gamma \approx 1$).
3. **RG universality.** Critical exponents depend only on symmetry and spatial dimension, not microscopic details.
4. **Fluctuation-dissipation connection.** $\gamma = -d(\log K)/d(\log C)$ is constrained by the FDT at criticality.

5.4 The cognitive loop as measurable system (illustrative)

The human–AI cognitive loop data extends the scaling relation into cognition, with critical caveats. The aggregate $\gamma_{\text{all}} = 1.059$ (CI containing 1.0) is suggestive but not evidential: the productivity classification was performed by the measured subject, introducing self-report bias, and per-session $R^2 = 0.12$.

5.5 Interpretation through extended mind (speculative)

Following Clark and Chalmers [4], one may speculatively interpret human–AI cognitive coupling as a dynamical system amenable to γ-scaling analysis. Whether such coupling exhibits measurable criticality signatures remains an open question requiring independent replication with blind labeling (see Limitations).

6. Limitations

1. **Self-measurement bias in cognitive substrate (critical)**: Productivity classification was performed by the measured operator ($n = 1$). CNS-AI data is presented as illustrative only.
2. **Low R^2 for cognitive substrate**: $R^2 = 0.12$

for productive sessions; $R^2 = -0.10$ for non-productive.

3. **Pseudo-replication removed**: The "neosynaptex cross-domain" substrate is an aggregate of other substrates, reclassified as illustrative.
4. **Statistical power at small** n: Zebrafish ($n = 47$) has CI width $\sim$0.41 with MDE of $\Delta\gamma = 0.35$ at 80% power.
5. **Single operator**: All CNS-AI data comes from one researcher.
6. **Proxy metrics**: Shannon entropy and TTR are proxies for cognitive complexity, not direct neural measurements.
7. **Non-stationarity**: The three-year archive may reflect learning effects.

7. Conclusion

We present empirical evidence that a scaling exponent $\gamma \approx 1.0$ appears across three independent biological substrates: zebrafish morphogenesis ($\gamma = 1.055$), human cardiac rhythm ($\gamma = 0.885$), and human EEG during motor imagery ($\gamma \approx 1.07$). All three evidential substrates fall within the metastable band ($|\gamma - 1| < 0.15$), with cross-substrate 95% CI containing unity. All pass surrogate testing ($p < 0.05$). Three simulation substrates provide theoretical validation: Gray–Scott ($\gamma = 0.938$) and Kuramoto ($\gamma = 0.980$) confirm $\gamma \approx 1.0$ at criticality, while BN-Syn ($\gamma \approx 0.49$, $N = 200$) demonstrates the expected finite-size deviation.

We propose that $\gamma \approx 1.0$ is a scaling signature of metastability itself — a substrate-independent condition for coherent computation at the edge of chaos. If confirmed by independent replication, this finding has implications for AI alignment (productive coupling has a measurable signature), cognitive enhancement (γ as a real-time diagnostic), and the physics of intelligence (metastability as a necessary condition for computation in any substrate).

Data Availability

All code, data processing pipelines, and proof bundles are available at `github.com/neuron7xLab`. The gamma probe pipeline and proof bundle are included in the neosynaptex repository.

Acknowledgments

This work was conducted independently during wartime in Poltava region, Ukraine, without institutional support or funding. The author acknowledges the use of AI language models (GPT-4o, Claude) as cognitive tools during the research process.

References

[1] P. Bak, C. Tang, K. Wiesenfeld. Self-organized criticality: An explanation of $1/f$ noise. *Phys. Rev. Lett.* **59**, 381 (1987).

[2] J. M. Beggs, D. Plenz. Neuronal avalanches in neocortical circuits. *J. Neurosci.* **23**, 11167 (2003).

[3] W. Maass, T. Natschläger, H. Markram. Real-time computing without stable states. *Neural Comput.* **14**, 2531 (2002).

[4] A. Clark, D. Chalmers. The extended mind. *Analysis* **58**, 7 (1998).

[5] H. Theil. A rank-invariant method of linear and polynomial regression analysis. *Proc. R. Neth. Acad. Sci.* **53**, 386 (1950).

[6] A. Vaswani *et al.* Attention is all you need. *NeurIPS* (2017).

[7] T. E. Harris. *The Theory of Branching Processes.* Springer (1963).

[8] M. A. Muñoz *et al.* Avalanche and spreading exponents in systems with absorbing states. *Phys. Rev. E* **59**, 6175 (1999).

[9] A. Clauset, C. R. Shalizi, M. E. J. Newman. Power-law distributions in empirical data. *SIAM Rev.* **51**, 661 (2009).