
Evolution of the AGN Population in the Universe

Neven Caplar

Simon Lilly, Benny Trakhtenbrot



Noeske+ 07



Reverse engineering approach

Basic idea: Identify key features of the observed galaxy and AGN <u>populations</u> and then explore *analytically* what they tell us, via the most basic continuity equations or via other simple relationships

$$\phi_{\text{red}}(t) = \phi_{\text{blue}}(t_0) \frac{1}{-(1+\alpha_s+\beta)} \frac{m}{M^*}$$

$$\times \int_{t_0}^t -(1+\alpha_s+\beta) \text{sSFR}(t') e^{\int_{t_0}^t -(1+\alpha_s+\beta) \text{sSFR}(t') dt'} dt'$$

$$= \phi_{\text{blue}}(t_0) \frac{1}{-(1+\alpha_s+\beta)} \frac{m}{M^*} e^{\int_{t_0}^t -(1+\alpha_s+\beta) \text{sSFR}(t') dt'}$$

$$= \phi_{\text{blue}}(t) \frac{1}{-(1+\alpha_s+\beta)} \frac{m}{M^*}.$$
(B4)

Putting Equations (B2) and (B4) together gives

$$\frac{1}{\phi_{\text{red}}(t)} \frac{d\phi_{\text{red}}(t)}{dt} = -\text{sSFR}(t)(1 + \alpha_s + \beta)$$

$$= \frac{1}{\phi_{\text{blue}}(t)} \frac{d\phi_{\text{blue}}(t)}{dt}$$

$$\frac{d \ln \phi_{\text{red}}(t)}{dt} = \frac{d \ln \phi_{\text{blue}}(t)}{dt}.$$
(B5)

Peng+ 2012

Peng+ 2010

Reverse engineering approach

Basic idea: Identify key features of the observed galaxy and AGN <u>populations</u> and then explore *analytically* what they tell us, via the most basic continuity equations or via other simple relationships

Observationally self-consistent

Constrains possible physical processes


Reveals connections between different aspects of the problem


Highlights difficulties of establishing uniqueness and causality

Topics for which this approach has been used

- Quenching of star-formation in galaxies (Peng+ 10, Peng+ 12, Knobel+ 14)
- Gas-regulator control of star-formation (Lilly+ 13)
- Size and internal structure of galaxies (Lilly & Carollo 16)

Simplest observables in the Universe






- Galaxy mass function number of galaxies per unit of mass
- Quasar luminosity function number of quasars (AGN) per unit of luminosity

Ansätze - coexistance scenario

- Radiatively efficient AGN are in star forming systems
- Distribution of Eddington ratio does not depend on the mass of the central black hole
- Mass of central black hole proportional to stellar mass
- To make quasar luminosity function use
 - AGN mass function & Eddington ratio function


Ansätze - coexistance scenario

- Radiatively efficient AGN are in star forming systems
- Distribution of Eddington ratio does not depend on the mass of the central black hole
- Mass of central black hole proportional to stellar mass
- To make quasar luminosity function use
 - AGN mass function & Eddington ratio function



Ansätze - coexistance scenario

- Radiatively efficient AGN are in star forming systems
- Distribution of Eddington ratio does not depend on the mass of the central black hole $\frac{L^* \propto M^* m_{hh}/m_* \lambda^*}{L^*}$
- Mass of central black hole proportional to stellar mass
- To make quasar luminosity function use
 - AGN mass function & Eddington ratio function




Ansätze - coexistance scenario


- Radiatively efficient AGN are in star forming systems
- Distribution of Eddington ratio does not depend on the mass of the central black hole $\frac{L^* \propto M^* m_{hh} / m_* \lambda^*}{L^*}$

 $\phi_{QLF}^* \propto \phi_{SF}^* \, \xi_{\lambda}^*$

- Mass of central black hole proportional to stellar mass
- To make quasar luminosity function use
 - AGN mass function & Eddington ratio function




Two interesting results from quasar luminosity function




Hopkins+ 07, but also Aird+ 14, Ueda+ 15, Aird+ 18

Two interesting results from quasar luminosity function




- Normalization of quasar luminosity function changes as normalization of star forming galaxies
- $L^* = (1+z)^4$, z < 2
- $L^* \propto M^* m_{bh} / m_* \lambda^*$


Results from simulating mass-luminosity plane



Results from simulating mass-luminosity plane



Results from simulating mass-luminosity plane











Consequences of size evolution


- Virial relation, Faber-Jackson and size evolution in galaxies implies evolution in either m_{bh}/m_{*} or m_{bh} sigma relation
- redshift evolution in m_{bh}/m_{*} of (1+z)² leads to constant and tighter m_{bh} sigma relation

$$r\Big|_{m} \propto (1+z)^{-1} \Leftrightarrow \sigma^{2}\Big|_{m} \propto (1+z)$$

$$m_{star} \propto \sigma^4 (1+z)^2$$

$$\frac{m_{BH}}{m_{star}} \propto (1+z)^2 \quad \Leftrightarrow \quad \frac{m_{BH}}{\sigma^4} = \text{constant}$$


Consequences of size evolution






 Mass growth of galaxies and AGN does not support such a mass ratio evolution in the co-existence scenario considered until now

Quenching scenario



- Now AGN activity connected with mass-quenching; black hole accretion happens in a single, almost delta-like burst
 - Naturally overcomes "the problem"- mass ratio responds instantly to the changes of the mass accretion rate

Evolution of AGN population in quenching scenario




- Mass ratio evolution is now an output of the model
- Mass ratio = BHARD/SFRD $\propto (1+z)^{1.5}$

Evolution of AGN population in quenching scenario

- Eddington ratio = $\lambda^* = (SFRD/BHARD) \cdot (L^\square/M^\square)$
- Eddington ratio $\propto (1+z)^{2.5}$
- sSFR $\propto (1+z)^{2.5}$




Evolution of AGN population in quenching scenario





- The main difference = which galaxies host AGN
- Co-existence w All SF galaxies have equal chance to host an AGN
- Quenching w More massive SF galaxies have a great chance to host an AGN

- The main difference = which galaxies host AGN
- Co-existence w All SF galaxies have equal chance to host an AGN
- Quenching w More massive SF galaxies have a great chance to host an AGN



- The main difference = which galaxies host AGN
- Co-existence w All SF galaxies have equal chance to host an AGN
- Quenching w More massive SF galaxies have a great chance to host an AGN






Anglés-Alcázar+ 17


- The main difference = which galaxies host AGN
- Co-existence w All SF galaxies have equal chance to host an AGN

 Quenching w More massive SF galaxies have a great chance to host an AGN



- The main difference = which galaxies host AGN
- Co-existence w All SF galaxies have equal chance to host an AGN

 Quenching w More massive SF galaxies have a great chance to host an AGN



Summary

- Simple global model combining galaxy mass function and quasar luminosity function leads to following conclusions
 - Evolution in the m_{bh}/m_{*} relation in star-forming galaxies
 - Non-evolving m_{bh}/m_{*} disfavored by mass-luminosity plane

Rewrite

 Scenario in which AGN growth is connected with quenching of starformation self-consistently produces m_{bh}/m_{*} evolution, Edd. ratio evolution and normalization evolution as seen in observations.