

- CARTER, H. B., and HARDY, MARGARET H. (1947).—Studies in the biology of the skin and fleece of sheep. 4. The hair follicle group and its topographical variations in the skin of the Merino foetus. Coun. Sci. Industr. Res. Aust. Bull. No. 215.
- CLOETE, J. H. L. (1939).—Prenatal growth in the Merino sheep. *Onderstepoort J. Vet. Sci.* **13**: 417–558.
- DUERDEN, J. E. (1939).—The arrangement of fibre follicles in some mammals, with special reference to the Ovidae. *Trans. Roy. Soc. Edin.* **59**: 763–71.
- DUERDEN, J. E., and RITCHIE, M. I. F. (1924).—Development of the Merino wool fibre. *S. Afr. J. Sci.* **21**: 480–97.
- FRASER, A. S. (1951).—Competition between skin follicles in sheep. *Nature* **167**: 202–3.
- FRASER, A. S. (1952a).—Growth of wool fibres in sheep. *Aust. J. Agric. Res.* **3**: 419–34.
- FRASER, A. S. (1952b).—Growth of the N-type fleece. *Aust. J. Agric. Res.* **3**: 435–44.
- FRASER, A. S. (1953).—Factors in the genetic determination of fleece structure in sheep. *J. Genet.* **51**: 222–36.
- FRASER, A. S. (1954).—Development of the skin follicle population in Merino sheep. *Aust. J. Agric. Res.* **5**: 737–44.
- FRASER, A. S., and SHORT, B. F. (1952).—Competition between skin follicles in sheep. *Aust. J. Agric. Res.* **3**: 445–52.
- FRÖLICH, G., SPÖTTEL, W., and TÄNZER, E. (1929).—“Wollkunde.” (Julius Springer: Berlin.)
- HARDY, MARGARET H. (1949).—The development of mouse hair *in vitro* with some observations on pigmentation. *J. Anat., Lond.* **83**: 364–84.
- HARDY, MARGARET H., and LYNE, A. G. (1956a).—The histological development of the skin and wool in the Merino foetus. *Proc. Int. Wool Text. Res. Conf. Aust.* **F**: 26–31.
- HARDY, MARGARET H., and LYNE, A. G. (1956b).—The development of wool follicles from foetal sheep skin in tissue culture. *Proc. Int. Wool Text. Res. Conf. Aust.* **F**: 32–5.
- HARDY, MARGARET H., and LYNE, A. G. (1956c).—Proposed terminology for wool follicles in sheep. *Nature* **177**: 705–6.
- MARGOLENA, L. A. (1954).—Sequence and growth of primary and secondary fiber follicles in Karakul sheep. *J. Anim. Sci.* **13**: 765–80.
- SCHINCKEL, P. G. (1953).—Follicle development in the Australian Merino. *Nature* **171**: 310.
- SCHINCKEL, P. G. (1955a).—The post-natal development of the skin follicle population in a strain of Merino sheep. *Aust. J. Agric. Res.* **6**: 68–76.
- SCHINCKEL, P. G. (1955b).—The relationship of skin follicle development to growth rate in sheep. *Aust. J. Agric. Res.* **6**: 308–23.
- SHORT, B. F. (1955a).—Development of the secondary follicle population in sheep. *Aust. J. Agric. Res.* **6**: 62–7.
- SHORT, B. F. (1955b).—Developmental modification of fleece structure by adverse maternal nutrition. *Aust. J. Agric. Res.* **6**: 863–72.
- SPÖTTEL, W., and TÄNZER, E. (1923).—Rassenanalytische Untersuchungen an Schafen unter besonderer Berücksichtigung von Haut und Haar. *Arch. Naturgesch.* **89**: 1–242.
- TÄNZER, E. (1926).—“Haut und Haar beim Karakul in rassenanalytischen Vergleich.” (Otto Thiele: Halle (Saale).)
- WILDMAN, A. B. (1932).—Coat and fibre development in some British sheep. *Proc. Zool. Soc. Lond.* **2**: 257–85.
- WILDMAN, A. B., and CARTER, H. B. (1939).—Fibre-follicle terminology in the Mammalia. *Nature* **144**: 783–4.

EXPLANATION OF PLATES 1 AND 2

All figures are of sections through skin from the trunk of Merino fetuses. The staining is with haemalum, eosin, and picric acid, except Plate 1, Fig. 4, which is without picric acid. DP, dermal papilla; G, stratum germinativum; H, hair; HN, hair cone; P, periderm; PP, pre-papilla; S, stratum spinosum; SD, derived secondary follicle; SG, sebaceous gland; SO, original secondary follicle.