

Follicle depth, straight length, and curved length  
in plain and wrinkly sheep

Jim Watts and Neville Jackson

27 Aug 2018

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Materials and Methods</b>	<b>2</b>
2.1	Sheep studied . . . . .	2
2.2	Measurements . . . . .	2
2.3	Statistical analysis . . . . .	2
<b>3</b>	<b>Results</b>	<b>3</b>
3.1	Means . . . . .	3
3.2	Variance components . . . . .	3
3.3	Correlations . . . . .	4
<b>4</b>	<b>Discussion</b>	<b>9</b>
<b>5</b>	<b>Conclusion</b>	<b>10</b>
<b>A</b>	<b>Appendix A</b>	<b>11</b>

# 1 Introduction

## 2 Materials and Methods

### 2.1 Sheep studied

This is a small study of 36 sheep from two Flocks, 18 from each Flock. Within each Flock there were 9 sheep chosen to represent the "loose" Skintype, and 9 sheep chosen to represent the "wrinkly" Skintype.

On each sheep 25 follicles chosen at random were measured on a vertical skin section taken from a skin biopsy sample obtained by the technique of Maddocks and Jackson (1988) [8]. The measurements were of follicle depth,  $t$ , and  $h$ , as defined below.

### 2.2 Measurements

The following measurements and scores were available

**SkinType** visual scores for sheep skin type. Two grades "loose" and "wrinkly".

**Follicle depth** perpendicular distance from skin surface to bottom of follicle bulb

**Follicle straight length( $t$ )** distance from skin surface to bottom of follicle bulb, measured at the angle of the follicle to the surface.

**Follicle sagitta( $h$ )** If the straight follicle length is considered to be the chord of a circle, the sagitta is the height of the arc above the centre of the chord.

From these measurements it was possible to calculate

**Radius of curvature of follicle** given by  $R = h/2 + t^2/(8h)$

**Angle subtended by the arc** given by  $\theta = 2\text{asin}(2/(2R))$

**Follicle curved length** given by  $S = R\theta$

The measurement of fibre curvature used by the wool industry is the reciprocal of radius of curvature and is in degrees per mm. The reciprocal of our  $R$  would be in radians per mm.

### 2.3 Statistical analysis

Data were imported into the R statistical program [12] and analysed using the *aov()* function for analysis of variance.

Table 1: Means for follicle measurements separately for each Flock and each Skintype

Flock	Skin.type	Follddepth	Straightlen	Curvlen	Radcurv
1	loose	91.95	91.37	91.95	348.5
2	loose	92.68	92.32	92.68	449.5
1	wrinkly	111.10	106.80	111.10	146.2
2	wrinkly	101.74	96.23	101.74	107.5

### 3 Results

#### 3.1 Means

Means for each Skintype within each Flock are given in Table 1

All four traits ( Follicle depth, straight follicle lenfgh, curved follicle length, and radius of curvature) would appear to differ substantially between loose and wrinkly Skintypes, but the two Flocks would seem to be similar.

We do not present standard deviations or standard errors here. The experimental design has multiple error levels. There are two random effects, Sheep, and Follicles within Sheep. Each of these contributes to the standard error of a mean. We need to estimate variance components for Sheep and Follicles to do the standard errors properly.

However we can test the significance of the Skintype difference and the Flock difference with an appropriate analysis of variance. We show these analyses of variance in Table 3.1 The first thing to note is that there are multiple error levels - corresponding to variation between sheep and variation between follicles within a sheep. The appropriate error level for testing Flock and Skintype differences is Error:Sheep - the line labelled Residuals. We can see that the analyses have used this error level for the F tests. These show that the Skintype difference was significant for all four traits, although the probability level for Follicle straight length was only .027 ( ie between 1 percent and five percent significance). None of the four traits showed a significant Flock difference.

This result is , of course, what we expected to find. Based on our understanding of the histology of wrinkle formation, we expect that wrinkled sheep will have a layer of hard collagen in the dermis which interferes with the downward growth of follicles from the dermis ( ie reduces their length) and which also causes the downgrowing follicles to curve sideways in an attempt to complete their length growth ( ie increases Radius of curvature). So these results support our hypothesis regarding the role of collagen in wrinkle formation and its effect on developing follicles.

#### 3.2 Variance components

With two Error levels ( Sheep and Follicles within sheep) we need to estimate variance components for each level, in order to calculate the standard error of

a mean. In Table 3.1 the mean square for Residual1 is the Follicles within sheep variance component ( $\sigma_F^2$ ). To get the Sheep variance component  $\sigma_S^2$  we calculate

$$\sigma_S^2 = (MS(\text{sheep}) - MS(\text{follicles within sheep}))/N_F$$

where  $N_F$  is the number of follicles per sheep.

We can then calculate the variance of a mean as

$$Var(\text{mean}) = \sigma_S^2/N_S + \sigma_F^2/(N_S N_F)$$

where  $N_S$  is number of sheep in the mean and  $N_F$  is number of follicles per sheep. This gives the variance of a mean. To get a standard error we take its square root.

Table 3 gives the variance component estimates calculated as above for each of the four traits, plus a standard error calculated for a mean of 9 sheep with 25 follicles per sheep.

These are the standard errors which apply to the means given in Table 1 The significance tests in Table 3.1 indicate a higher level of significance because they are for the Skintype difference averaged across both Flocks - ie they correspond to means of 18 Sheep rather than 9. We have not presented the means for Skintype differences averaged across both Flocks, so we present these and their standard errors in Table 4

We see that these means are indeed significantly different, as indicated in the analyses of variance in Table 3.1.

### 3.3 Correlations

Figure 1 shows a scatterplot of curved length of follicles against follicle depth.

It suggests that follicle depth is a rather poor indicator of the curved length of a follicle. The points seem to cluster at or below the 1 to 1 line, indicating that curved length is either equal to or longer than follicle depth, depending on how much the follicle is curved. In this situation a correlation is not a good descriptor.

Figure 2 shows a similar scatterplot of curved length against straight length of follicles.

We see a similar picture to Figure 1 but with much closer agreement. The points again cluster at or below the 1 to 1 line. Curved length is either equal to straight length or slightly longer. Straight length is a better indicator of curved length than is follicle depth, but it misses out if the follicles are curved.

Figure 3 shows a scatterplot of curved length against radius of curvature.

Here we see something radically different. Why the stepped spacing of points, especially those from the loose Skintype with high radius of curvature. Well, the issue is that measuring the follicle sagitta (h) gets to be rather imprecise when the follicle is not curved. Values of 1, 2, 3 for the sagitta produce the grouping effect when converted to radius of curvature.

What Figure 3 shows is that wrinkly sheep have curved follicles that may also be long, while loose sheep have straighter follicles that on average are shorter.




Figure 1: Plot of follicle depth against curved length of follicles. The loose and wrinkly skintypes are shown as points labelled 1 ( red) and 2(green) respectively.

This is not quite what we expected. Those very long follicles in the wrinkly group bear further investigation.

The relationship between length and curvature is complicated. The points occupy the bottom left corner of the plot. Clearly there are other factors involved.




Figure 2: Plot of straight length against curved length of follicles. The loose and wrinkly skintypes are shown as points labelled 1 ( red) and 2(green) respectively.

Table 2: Analyses of variance of Follicle depth, Follicle straight length, Follicle curved length, and Follicle radius of curvature

Follicle depth					
Effect	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Error: Sheep					
Flock	1	1.08	1.08	0.00	0.9861
Skin.type	1	35822.56	35822.56	10.16	0.0031
Residuals	33	116320.51	3524.86		
Error: Within					
Residuals1	860	59641.23	69.35		
Follicle straight length					
Effect	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Error: Sheep					
Flock	1	5218.12	5218.12	1.32	0.2591
Skin.type	1	21042.58	21042.58	5.32	0.0275
Residuals	33	130612.06	3957.94		
Error: Within					
Residuals1	860	66226.12	77.01		
Follicle curved length					
Effect	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Error: Sheep					
Flock	1	4251.50	4251.50	1.03	0.3180
Skin.type	1	44711.74	44711.74	10.81	0.0024
Residuals	33	136454.62	4134.99		
Error: Within					
Residuals1	860	75424.61	87.70		
Follicle radius of curvature					
Effect	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Error: Sheep					
Flock	1	239208.35	239208.35	1.31	0.2603
Skin.type	1	16554724.49	16554724.49	90.79	0.0000
Residuals	33	6017368.02	182344.49		
Error: Within					
Residuals1	860	20509483.53	23848.24		

Table 3: Sheep and follicle within sheep variance components for four traits. Standard error of a mean of 9 sheep with 25 follicles per sheep

	Follddepth	Straightlen	Curvlen	Radcurv
Sheep component	138.22	155.24	161.89	6339
Follicles component	69.35	77.01	87.70	23848
Standard error of mean	3.95	4.19	4.28	28.46



Table 4: Means and standard errors for follicle measurements separately for each Skintype

Skin.type	Follddepth	Straightlen	Curvlen	Radcurv
Mean loose	82.5	91.86	92.32	399.0
Mean wrinkly	95.14	101.6	106.5	127.1
Standard error	2.79	2.96	3.03	20.12




Figure 3: Plot of radius of curvature against curved length of follicles. The loose and wrinkly skintypes are shown as points labelled 1 ( red) and 2(green) respectively.

## 4 Discussion

We wanted to show that wrinkled sheep have curved follicles, and we wanted to use something other than a subjective score for follicle curvature. We have achieved that - the radius of curvature differs significantly between loose and wrinkly sheep.

There are also differences in follicle length, regardless of how it is measured. The wrinkled sheep have longer follicles. This begs an explanation. If the presence of collagen in wrinkled sheep interferes with follicle development causing them to grow curved, that is one thing. But how would presence of collagen cause follicles to grow longer? Perhaps the folding of the epidermal and dermal layers which occurs as wrinkles form 'stretches' the follicles. Perhaps the collagen contracts and pulls on the follicles. Perhaps it is not a simple physical effect at all. Perhaps the collagen moves the papilla cells toward which the follicle downgrowth develops, so that the intital infolding of the epidermis and the papilla cells get out of line, causing the follicle to both curve sideways and grow longer in order to reach the papilla cells. We do not know.

So what needs to be done from here? Well we need to sort out whether the length of follicles is at all important in relation to fibre growth. There has been one experiment selecting for follicle depth (Jackson (2017) [4]) and it was a failure in relation to improving wool production. It changed follicle depth and fibre diameter and variance of fibre diameter. The experiment was started when there was a feeling that 'deep' follicles had a better blood supply and a more vigorous wool growth. That sort of thinking has since disappeared. 'Deep' follicles are just long follicles, not anything special. Having an elongted shaft is not going to help a follicle grow more wool.

There also needs to be some consideration as to whether follicle curvature is the same thing as intrinsic fibre curvature, and whether the current commercial curvature measurement technology ( Laserscan or OFDA) measures intrinsic fibre curvature. It comes down to whether follicle curvature is the only factor determining intrinsic fibre curvature, or whether there are other factors perhaps related to the fibre cross sectional distribution of cortical cell types. Understanding the effect of collagen on follicle curvature is a half-way house. We need to know the effect of collagen on intrinsic fibre curvature, not the 'set' we observe in staple crimp, not some 'set' we measure under Laserscan od OFDA standardised measuring conditions, but the real intrincic curvature which is wahat a fibre can always be relaxed to and which is determined by different growth rates of the ortho and para cortical cells in a fibre with a bilaterally segmented cortex.

That being said, this study has progressed our understanding of why sheep with wrinkle and hard collagen differ from plain bodied sheep in wool growth. They differ because the presence of a layer of hard collagen in the dermis affects the size and shape of wool follicles. In our discussion we have explored possible mechanisms.

## 5 Conclusion

Of all the aspects discussed above, one stands out. Here is what we think is actually happening to follicle development in wrinkly sheep.

Follicles develop in the foetus between 60 and 150 days. Papilla cells migrate to or agglomerate at sites. Sites are thought to be determined by the collagen network. Above each site, at the skin surface, a downgrowth of epidermal tissue occurs, directed toward the site. The papilla cells at the site act as an attractor. Now, if the site moves sideways after the downgrowth commences, the partially completed downgrowth will curve toward the new site location, resulting in a curved follicle. The downgrowth will always be toward the attractor. The curved follicle will also need to be longer in order for the bulb to reach the papilla cells at the new site. What would cause the site to move? Well it would seem that development of hard collagen causes the sites to move. This would have to happen at the critical time while the follicle downgrowths were part complete.

Only P and So follicles have sites. So only P and So follicles would be affected by the 'moving site effect'. So look at Figure 3. For the wrinkly sheep (which are the green points) there are the P and So follicles with their curved length  $l$  140 and tiny radius of curvature, at the top left corner of the scatterplot. All the other green points are SD follicles; they do not have defined sites; they curve because the papilla cells aggregate around other follicles and they have to curve to belong to a common follicle opening. They are not long because the papilla cells aggregate close to other follicles.

So what happens in loose or non-wrinkly sheep? Well, the sites don't move, so the P and So follicles are not forced to curve to follow the papilla cells. So in Figure 3 there are the P and So follicles at the red points on the lower right corner of the scatterplot; they are quite straight (high radius) and normal length. The rest of the red points are loose sheep Sd follicles, and they are similar to the wrinkly sheep Sd follicles, because they form in the same way, and the collagen hardening does not interfere with Sd's.

So there we have it. Our 'moving site hypothesis' explains follicle size and shape, and how hard collagen affects follicle development. It does not explain why a curved follicle grows a curved fibre. That, however is quite simple - in order to form inside a curved follicle a fibre must have either more cortical cells or longer cortical cells on the side of the fibre cross section which corresponds to the outer side of the curve. This is simple material science, the outer side of a curved object has to be longer than the inner side. Once a fibre forms in that way, it retains the intrinsic curvature permanently. That is what we observe. Fibre curvature is a complicated phenomenon because, apart from this intrinsic curvature, fibres can be 'set' into other degrees of curvature, for example in a staple where the crimp observed is not just intrinsic curvature but a complicated phenomenon involving stretching or unfolding of the intrinsic curvature combined with various setting conditions. Staple crimp is semi-permanent. Fibre intrinsic curvature is permanent (ie is always recoverable) and is the same thing as follicle curvature.

So why all this concern about crimp and curvature? Well, wrinkly sheep (and low density sheep with hard collagen that are not wrinkly, such as Downs Wool Breeds) all have characteristically highly crimped fleeces. This is one of the outstanding features of wrinkled sheep, so we must explain crimp and curvature, or we have not grasped the wrinkle issue.

## A Appendix A

Detective stories have almost universal appeal, because they are about the discovery of truth and the triumph of good over evil. There is even an interpretation of the Bible as a detective story. We need to 'play detective' and embark on a search for clues which might support the 'moving sites hypothesis'.

Lets start with the appearance of follicles in vertical sections from adult sheep. Figure 4 is a copy of the Nay follicle curvature score drawings. Dr Nay was an artist; these are 3 dimensional drawings; we need to bear that in mind when looking at the curvature of follicles, some are curved in the plane of the page, some are curved in and out of the page.

The papilla cells ( and therefore the sites in the case of P and So follicles) have to be where the follicle bulbs are. For score 1, the bulbs are all at the same level and directly below the emergence point of the fibre at the epidermal surface. No evidence of movement here. For scores 2 and 3, the bulbs are still all at the same level, but are slightly out of line with the emergence point, so the follicles have a curve, and the sites/papilla cells have moved slightly sideways. For scores 4, 5, 6, and 7 the bulbs are not all at the same level. So do the sites move vertically as well as sideways? It would appear so. The bulbs are also increasingly out of line with the emergence point, so the follicles curve increasingly, as the scores increase. The sites move sideways as well as vertically.

What are these follicles in scores 6 and 7 for which the bulb is above the lowest point of the follicle curve and facing upwards? They would have to be follicles for which the site ( or aggregate of papilla cells) has moved sideways and probably upward after the downgrowth of epidermal tissue has commenced. They start out straight, then have to curve 'late' in their development, producing the typical 'J' curve with a straight upper shaft and a late curve at the lower end.

We have to remember that not all follicles have a site. The Sd follicles do have a papilla cell aggregate. We do not know what if anything controls where these Sd aggregates form, but they seem to cluster alongside existing follicles. We do not know if these Sd papilla cell clusters move, or even if they all form at the same level vertically. So it is difficult to say which might be the Sd follicles in Figure 4. There are no distinguishing features - the follicles all look the same. We just know that most of the follicles are Sd's. So what does this suggest. Maybe the Sd follicles are subject to the same papilla cell aggregate movement as the other follicles . Maybe they simply form all over the place initially.

That is about all we can get out of vertical section from adults. If we




Figure 4: The Nay follicle curvature score drawings. Reproduced from Maddocks and Jackson(1988) [8]

had vertical sections of foetal skin we might actually be able to see directly whether follicles curve as they grow. The alternative is that they are bent around afterwards - that still involves moving the 'site', but it also involves dragging the whole fully formed follicle around, which seems biologically unlikely.

We can also have a look at what effects site movement might have on the arrangement of follicles in horizontal sections. We have to bear in mind that horizontal sections are conventionally taken at sebaceous gland level, which is well above the bulb region, so any effect of site movement will be minimal. We have to also bear in mind that follicles and follicle groups are also moved around by expansion of the skin as the animal grows.

There are some known differences in follicle group arrangement between SRS Merinos and normal wrinkly Merinos.

## References

- [1] Bogolyubsky S.N. (1940) cited by Fraser A.S and Short B.F. (1960) The Biology of the Fleece. Animal Research Laboratories Technical Paper No 3. CSIRO Melbourne 1960.
- [2] Carter H.B. (1943) Studies in the biology of the skin and fleece of sheep. 1. The development and general histology of the follicle group in the skin of the Merino. 2. The use of tanned sheepskin in the study of follicle population density. 3. Notes on the arrangement, nomenclature, and variation of skin folds and wrinkles in the Merino. C.S.I.R. Bulletin No 164, Melbourne, 1943
- [3] Fraser A.S and Short B.F. (1960) The Biology of the Fleece. Animal Research Laboratories Technical Paper No 3. CSIRO Melbourne 1960.
- [4] Jackson, N. (2017) Genetic relationship between skin and wool traits in Merino sheep. Part I Responses to selection and estimates of genetic parameters. URL <https://github.com/nevillejackson/Fleece-genetics/tree/master/skinandfleeceparameters/ab3220/skinwool1.pdf>
- [5] Jackson, N. and Watts, J.E. (2018) Does follicle development affect the spatial layout of sheep skin? URL <https://github.com/nevillejackson/Fleece-biology/tree/master/skinspace/skinspace.pdf>
- [6] Jackson, N. and Watts, J.E. (2017) What is known about the genetics of wrinkle score in Merino sheep? URL <https://github.com/nevillejackson/Fleece-genetics/wrinkle/wrinkle.pdf>
- [7] Knight, K.R., Lepore, D.A., Horne, R.S., Ritz, M., Kumta, S. and O'Brian, B.M. (1993) Collagen content of uninjured skin and scar tissue in foetal and adult sheep. *Int. J. Exp. Pathol.* 74(6):583-591
- [8] Maddocks, I.G. and Jackson, N. (1988) Structural studies of sheep, cattle, and goat skin. CSIRO, Division of Animal Production, Sydney.
- [9] Menton, D.N. and Hess, R.A. (1980) The ultrastructure of collagen in the dermis of tight-skin (Tsk) mutant mice. *The Journal of Investigative Dermatology* 74:139-147
- [10] Mitchell, T.W. et al (1984) Some physical and mechanical properties of sheep skin with a comparison of "thick" and "thin" skins. *Wool Technology and Sheep Breeding*, Vol XXXII, No IV, 200-206
- [11] Nay, T. (1966) Wool follicle arrangement and vascular pattern in the Australian Merino. *Aust. J. Agric. Res.* 17:797-805
- [12] R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.

- [13] Watts, J.E., Jackson, N., and Ferguson, K.A. (2017) Improvements in fleece weight weight and wool quality of Merino sheep selected visually for high fibre density and length. URL [https://github.com/nevillejackson/SRS-Merino/Paper\\_2\\_Revised\\_10\\_November\\_2017.docx](https://github.com/nevillejackson/SRS-Merino/Paper_2_Revised_10_November_2017.docx)