Newborough Warren Weather & Water Watch: October 2026

Weather Summary

Metric	Local Gauge (ILLANF24)	RAF Valley	Typical October*
Total Rainfall	N/A	101.8 mm	~101.6 mm
Average Highs		15.1 C	14.3 C
Average Lows		10.2 C	9.2 C

**Based on 1991-2020 Met Office climate averages (RAF Valley).*

October saw 101.8 mm at RAF Valley, close to the 101.6 mm average (100%). Temperatures were milder than usual with lows of 10.2 C (average 9.2 C).

Rainfall over the twelve months to October 2026 (from November 2025) stands at 864 mm, which is 101% of the 1991–2020 average for the same twelve-month window (857 mm).


Data quality note: ILLANF24: Could not find embedded JSON in WU page (after 4 attempts)

Local Water Levels

The latest well round on 03/11/2026 covered 68 monitoring dipwells across the reserve. Water levels are predominantly rising: 55 wells showed increases, 12 showed decreases, and 1 were stable. The mean change was +0.07 m.

Dry this round (4): D7 (1.06 m), ceh22 (0.94 m), nw3 (1.10 m), nw6 (0.85 m). These wells were visited and found dry, the figure being the depth below ground at which the dipper reached the bottom; they have no water level to compare, which is why they do not appear in the tables above.

Month-on-Month Change: Sep 2026 to Oct 2026




Water level change (m) from Sep 2026 to Oct 2026. Blue = rising, red = falling.

By area — C1 (Lake Edge) +0.12 m (7); C2 (Dune) +0.13 m (17); C3 (Western Residual) +0.10 m (24); C4 (Main Forest) -0.08 m (11); C5 (Coastal Forest) +0.03 m (9). C2 (Dune) rose most (+0.13 m), and C4 (Main Forest) was the only area to fall (-0.08 m); the figures in brackets are the wells compared in each. The outlines on the map are these five areas (nearest-well partitions of the cluster membership, a rendering of membership rather than a surveyed boundary).


Against the last 9 years: the lowest annual minimum of the period was August 2019. October's level is 0.13 m above it (across 62 wells read on both occasions), and lower than 4 of the 9 annual minima.

Annual Minima of Previous Years




Each summer's annual minimum relative to October 2026, compared well by well. Below the line, that year's low was lower than this month; the darker bar is the lowest of the period.

Difference from the Lowest Minimum: Aug 2019 to Oct 2026



Water level change (m) from Aug 2019, the lowest annual minimum of the period, to October 2026. Blue = higher now, red = lower now.

Every October So Far



Each earlier October relative to October 2026, compared well by well. Below the line, that year was lower than now. This October is higher than 1 of the 10 compared and lower than 9.

Flood Outlook for the Coming Winter


Each part of the warren has a measured rainfall threshold: the amount that would have to fall over the coming months to bring the water table up to the ground surface, given how deep it fell this summer. **C1 (Lake Edge) is closest**, needing 495 mm against a typical 400 mm — about 124% of an average season — and 102 mm has fallen so far.

Area	Summer low (m)	Rain needed	Typical	Needed / typical
C1 (Lake Edge)	1.05	495 mm	400 mm	1.24
C2 (Dune)	1.17	597 mm	461 mm	1.30
C3 (Western Residual)	1.31	657 mm	461 mm	1.43
C4 (Main Forest)	1.79	955 mm	461 mm	2.07
C5 (Coastal Forest)	1.75	1037 mm	461 mm	2.25

A ratio above 1 means a wetter-than-average season would be needed for the water table to reach the surface there; below 1, an average season would do it. These are rainfall thresholds, not probabilities.

Year-on-Year Comparison: Oct 2025 vs Oct 2026

Compared to this time last year, 13 out of 67 wells are showing higher water levels, with a mean change of -0.05 m (range -0.25 to +0.12 m).



Year-on-year water level change from Oct 2025 to Oct 2026.

By area — C1 (Lake Edge) -0.17 m (7); C2 (Dune) -0.08 m (17); C3 (Western Residual) -0.01 m (22); C4 (Main Forest) -0.05 m (11); C5 (Coastal Forest) +0.05 m (10). C5 (Coastal Forest) rose most (+0.05 m) while 4 areas fell, C1 (Lake Edge) furthest (-0.17 m); the figures in brackets are the wells compared in each. The outlines on the map are these five areas (nearest-well partitions of the cluster membership, a rendering of membership rather than a surveyed boundary).

Well Measurement Summary

Largest Risers and Fallers (month-on-month)

Well	Change (m)
wmc4	+0.19
D10	+0.19
T41a	+0.19
nw13	+0.18
ceh10	+0.18

Well	Change (m)
ceh34	-0.08
FE1	-0.09
ceh13	-0.12
ceh14	-0.13
nw10	-0.14

Full data: 68 wells. Mean: +0.07 m. Range: -0.14 to +0.19 m.