Question 1

$$\frac{4x^4 + 5x^2 - 7x}{x} = \frac{x(4x^3 + 5x - 7)}{x}$$
$$= 4x^3 + 5x - 7$$
 (1a)

$$\frac{7x^5 - 5x^5 + 9x^3 + x^2}{x} = \frac{x(2x^4 + 9x^2 + x)}{x}$$

$$= 2x^4 + 9x^2 + x$$
(1b)

$$\frac{-x^4 + 4x^2 + 6}{x} = \frac{x(4x - x^3) + 6}{x}$$

$$= \frac{x(4x - x^3)}{x} + \frac{6}{x}$$

$$= 4x - x^3 + \frac{6}{x}$$
(1c)

$$\frac{7x^5 - x^3 - 4}{x} = \frac{x(7x^4 - x^2) - 4}{x}$$

$$= \frac{x(7x^4 - x^2)}{x} - \frac{4}{x}$$

$$= 7x^4 - x^2 - \frac{4}{x}$$
(1d)

$$\frac{8x^4 - 4x^3 + 6x}{2x} = \frac{2x(4x^3 - 2x^2 + 3)}{2x}$$

$$= 4x^3 - 2x^2 + 3$$
(1e)

$$\frac{9x^2 - 12x^3 - 3x}{3x} = \frac{3x(3x - 4x^2 - 1)}{3x}$$
$$= -4x^2 + 3x - 1$$
 (1f)

$$\frac{7x^3 - x^4 - 2}{5x} = \frac{x(7x^2 - x^3)}{x(5)} - \frac{2}{5x}$$

$$= -\frac{1}{5}x^3 + \frac{7}{5}x^2 - \frac{2}{5x}$$
(1g)

$$\frac{-4x^2 + 6x^4 - 2x}{-2x} = \frac{-2x(2x - 3x^3 + 1)}{-2x}$$

$$= -3x^3 + 2x + 1$$
(1h)

$$\frac{-x^8 + 9x^4 - 4x^3 + 6}{-2x} = \frac{-x(x^7 - 9x^3 + 4x^2)}{-x(2)} + \frac{-2(-3)}{-2x}$$

$$= \frac{1}{2}x^7 - \frac{9}{2}x^3 + 2x^2 - \frac{3}{x}$$
(1i)

$$\frac{-9x^9 - 6x^6 + 4x^4 - 2}{-3x} = \frac{-x(9x^8 + 6x^5 - 4x^3)}{-x(3)} - \frac{2}{-3x}$$

$$= 3x^8 + 2x^5 - \frac{4}{3}x^3 + \frac{2}{3x}$$
(1j)

Question 2

$$\frac{(x+3)(x-2)}{(x-2)} = x+3 \tag{2a}$$

$$\frac{(x+4)(3x-1)}{(3x-1)} = x+4 \tag{2b}$$

$$\frac{(x+3)^2}{(x+3)} = x+3\tag{2c}$$

$$\frac{x^2 + 10x + 21}{(x+3)} = \frac{(x+3)(x+7)}{(x+3)}$$

$$= x+7$$
(2d)

$$\frac{x^2 + 9x + 20}{(x+4)} = \frac{(x+4)(x+5)}{(x+4)}$$

$$= x+5$$
(2e)

$$\frac{x^2 + x - 12}{(x - 3)} = \frac{(x - 3)(x + 4)}{(x - 3)}$$

$$= x + 4$$
(2f)

$$\frac{x^2 + x - 20}{x^2 + 2x - 15} = \frac{(x - 4)(x + 5)}{(x - 3)(x + 5)}$$
$$= \frac{x - 4}{x - 3}$$
 (2g)

$$\frac{x^2 + 3x + 2}{x^2 + 5x + 4} = \frac{(x+1)(x+2)}{(x+1)(x+4)}$$

$$= \frac{x+2}{x+4}$$
(2h)

$$\frac{x^2 + x - 12}{x^2 - 9x + 18} = \frac{(x - 3)(x + 4)}{(x - 3)(x - 6)}$$

$$= \frac{x + 4}{x - 6}$$
(2i)

$$\frac{2x^2 + 7x + 6}{(x - 5)(x + 2)} = \frac{(2x + 3)(x + 2)}{(x - 5)(x + 2)}$$

$$= \frac{2x + 3}{x - 5}$$
(2j)

$$\frac{2x^2 + 9x - 18}{(x+6)(x+1)} = \frac{(2x-3)(x+6)}{(x+6)(x+1)}$$

$$= \frac{2x-3}{x+1}$$
(2k)

$$\frac{3x^2 - 7x + 2}{(3x - 1)(x + 2)} = \frac{(3x - 1)(x - 2)}{(3x - 1)(x + 2)}$$
$$= \frac{x - 2}{x + 2}$$
(2l)

$$\frac{2x^2 + 3x + 1}{x^2 - x - 2} = \frac{(2x+1)(x+1)}{(x-2)(x+1)}$$

$$= \frac{2x+1}{x-2}$$
(2m)

$$\frac{x^2 + 6x + 8}{3x^2 + 7x + 2} = \frac{(x+4)(x+2)}{(3x+1)(x+2)}$$
$$= \frac{x+4}{3x+1}$$
 (2n)

$$\frac{2x^2 - 5x - 3}{2x^2 - 9x + 9} = \frac{(2x+1)(x-3)}{(2x-3)(x-3)}$$

$$= \frac{2x+1}{2x-3}$$
(20)

Question 3

Find a, b, and c:

$$\frac{6x^3 + 3x^2 - 84x}{6x^2 - 33x + 42} = \frac{ax(x+b)}{x+c} \tag{3}$$

$$\frac{6x^3 + 3x^2 - 84x}{6x^2 - 33x + 42} = \frac{3x(2x^2 + x - 28)}{3(2x^2 - 11x + 14)}$$
$$= \frac{x(2x - 7)(x + 4)}{(2x - 7)(x - 2)}$$
$$= \frac{x(x + 4)}{x - 2}$$

$$\therefore a = 1, b = 4, c = -2$$