Differentiable Probabilistic Models

William Watson

nextbillyonair@gmail.com

Abstract

d

Contents

1	Intro	oduction	6					
	1.1	Philosophy	5					
2	Preliminary 6							
	2.1	Kronecker Product	5					
	2.2	Gradients	5					
	2.3	Jacobian	5					
	2.4	Hessian	5					
	2.5	Newton Optimization	5					
3	Disti	ributions	6					
	3.1	Distribution	5					
	3.2	Arcsine	5					
	3.3	Bernoulli	5					
	3.4	Beta	5					
	3.5	Categorical	5					
	3.6	Cauchy	5					
	3.7	Chi Square	5					
	3.8	Conditional Model	5					
	3.9	Convolution	5					
	3.10	Data	5					
	3.11	Dirac Delta	5					
	3.12	Dirichlet	5					
	3.13	Exponential	5					
	3.14	Fisher-Snedcor (F-Distribution)	5					
	3.15	Gamma	5					
	3.16	Generator	5					
	3.17	Gumbel Softmax	5					
	3.18	Gumbel	5					
	3.19	Half Cauchy	5					
	3.20	Half Normal	5					
	3.21	Hyperbolic Secant	5					
	3.22	Langevin	5					
	3.23	Laplace	5					
	3.24	Log Cauchy	5					
	3.25	Log Laplace	5					
	3.26	Log Normal	5					
	3.27	Logistic	5					
	3.28	Normal (Multivariate)	5					

	3.29	Rayleigh	6
	3.30	Relaxed Bernoulli	6
	3.31	Student T	6
	3.32	Transformed Distribution	6
	3.33	Uniform	6
4	Mixt	ure Models	6
	4.1	Mixture Model	6
	4.2	Gumbel Mixture Model	6
	4.3	Infinite Mixture Model	6
5	Tran	sforms	6
	5.1	Transform	7
	5.2	Inverse Transform	7
	5.3	Affine	7
	5.4	Exp	7
	5.5	Expm1	7
	5.6	Gumbel	7
	5.7	Log	8
	5.8	Logit	8
	5.9	Power	8
	5.10	Reciprocal	8
	5.11	Sigmoid	9
	5.12	SinhArcsinh	9
	5.13	Softplus	9
	5.14	Softsign	9
	5.15	Square	9
	5.16	Tanh	9
6	Crite	erion and Divergences	9
	6.1	Cross-Entropy	9
	6.2	Perplexity	9
	6.3	Forward KL Divergence	9
	6.4	Reverse KL Divergence	10
	6.5		12
	6.6	Earth Mover's Distance	12
7	ELB	0	12
8	Adve	ersarial Loss	12
	8.1	Adversarial Loss	12

	8.2	GAN Loss	12
	8.3	MMGAN Loss	12
	8.4	WGAN Loss	12
	8.5	LSGAN Loss	12
	8.6	Gradient Penalty	12
	8.7	Spectral Norm	12
9	Mod	lels	12
	9.1	Base Models	12
		9.1.1 Model	12
	9.2	Regression	12
		9.2.1 Linear Regression (Normal)	12
		9.2.2 L1 Regression (Laplace)	12
		9.2.3 Ridge Regression (Normal + Normal Prior on Weights)	12
		9.2.4 Lasso Regression (Normal + Laplace Prior on Weights)	12
	9.3	Classification	12
		9.3.1 Logistic Regression (Bernoulli)	12
		9.3.2 Bayesian Logistic Regression (Bernoulli)	12
		9.3.3 Softmax Regression (Categorical)	12
	9.4	Generative Adversarial Networks	12
		9.4.1 Generative Adversarial Networks	12
		9.4.2 GAN Model	12
		9.4.3 MMGAN Model	12
		9.4.4 WGAN Model	12
		9.4.5 LSGAN Model	12
1Λ	Mon	ate Carlo	12
10		Monte Carlo Approximation	12
		Linear Congruential Generator	12
		Inverse Transform Sampling	12
		Box-Muller	12
			12
		Marsaglia-Bray	12
	10.0	Rejection Sampling	12
11	Mar	kov Chain Monte Carlo (MCMC)	12
		Metropolis	12
	11.2	Metropolis-Hastings	12
	11.3	Metropolis-Adjusted Langevin Algorithm (MALA)	12
	11.4	Hamiltonian Monte Carlo	12

1 Introduction

- 1.1 Philosophy
- 2 Preliminary
- 2.1 Kronecker Product
- 2.2 Gradients
- 2.3 Jacobian
- 2.4 Hessian
- 2.5 Newton Optimization
- 3 Distributions
- 3.1 Distribution
- 3.2 Arcsine
- 3.3 Bernoulli
- 3.4 Beta
- 3.5 Categorical
- 3.6 Cauchy
- 3.7 Chi Square
- 3.8 Conditional Model
- 3.9 Convolution
- 3.10 Data
- 3.11 Dirac Delta
- 3.12 Dirichlet
- 3.13 Exponential
- 3.14 Fisher-Snedcor (F-Distribution)
- **3.15** Gamma
- 3.16 Generator
- 3.17 Gumbel Softmax
- 3.18 Gumbel
- 3.19 Half Cauchy
- 3.20 Half Normal
- 3.21 Hyperbolic Secant
- 3.22 Langevin
- 3.23 Laplace
- 3.24 Log Cauchy
- 3.25 Log Laplace
- 3.26 Log Normal
- 3.27 Logistic

5.1 Transform

5.2 Inverse Transform

5.3 Affine

- Parameters
 - Location $\mu \in \mathbb{R}^n$
 - Scale $\sigma > 0$
- Forward

$$f(x) = \mu + \sigma \cdot x \tag{1}$$

• Inverse

$$f^{-1}(y) = \frac{y - \mu}{\sigma} \tag{2}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = \log|\sigma| \tag{3}$$

5.4 Exp

- Parameters
 - None
- Forward

$$f(x) = e^x (4)$$

Inverse

$$f^{-1}(y) = \log y \tag{5}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = x \tag{6}$$

5.5 Expm1

- Parameters
 - None
- Forward

$$f(x) = e^x - 1 \tag{7}$$

Inverse

$$f^{-1}(y) = \log(1+y) \tag{8}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = x\tag{9}$$

5.6 Gumbel

- Parameters
 - Location $\mu \in \mathbb{R}^n$
 - Scale $\sigma > 0$
- Forward

$$f(x) = \exp\left(-\exp\left(-\frac{x-\mu}{\sigma}\right)\right) \tag{10}$$

Inverse

$$f^{-1}(y) = \mu - \sigma \cdot \log\left(-\log\left(y\right)\right) \tag{11}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = -\log\left(\frac{\sigma}{-\log(y)\cdot y}\right) \tag{12}$$

5.7 Log

- Parameters
 - None
- Forward

$$f(x) = \log x \tag{13}$$

Inverse

$$f^{-1}(y) = \exp y \tag{14}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = -y\tag{15}$$

5.8 Logit

- Parameters
 - None
- Forward

$$f(x) = \log\left(\frac{x}{1-x}\right) \tag{16}$$

• Inverse

$$f^{-1}(y) = \frac{1}{1 + e^{-y}} \tag{17}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = \log(1 + e^{-y}) + \log(1 + e^{y})$$
(18)

5.9 Power

- Parameters
 - Power p
- Forward

$$f(x) = \begin{cases} e^x & p = 0\\ (1 + x \cdot p)^{1/p} & \text{otherwise} \end{cases}$$
 (19)

• Inverse

$$f^{-1}(y) = \begin{cases} \log y & p = 0\\ y^{p-1}/p & \text{otherwise} \end{cases}$$
 (20)

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = \begin{cases} x & p = 0\\ \left(\frac{1}{p} - 1\right) \cdot \log(x \cdot p + 1) & \text{otherwise} \end{cases}$$
 (21)

5.10 Reciprocal

- Parameters
 - None
- Forward

$$f(x) = 1/x \tag{22}$$

• Inverse

$$f^{-1}(y) = 1/y (23)$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = -2 \cdot \log|x| \tag{24}$$

5.11 Sigmoid

- Parameters
 - None
- Forward

$$f(x) = \frac{1}{1 + e^{-x}} \tag{25}$$

• Inverse

$$f^{-1}(y) = \log\left(\frac{y}{1-y}\right) \tag{26}$$

• Log Absolute Determinant Jacobian

$$\log|\det \mathbf{J}|(x,y) = -\log(1 + e^{-x}) - \log(1 + e^{x}) \tag{27}$$

- 5.12 SinhArcsinh
- 5.13 Softplus
- 5.14 Softsign
- 5.15 Square
- 5.16 Tanh

6 Criterion and Divergences

The criterion and divergences listed here can be used to quantify the "distance" between two distributions. Hence, in conjunction with torch optimizers, one can minimize said difference to learn the paramters of a distribution. For sake of notation clarity, p is the true distribution and q is the learned distribution. Hence we "fit" q to match p. In addition, we provide the Monte Carlo approximation.

6.1 Cross-Entropy

$$H(p,q) = -\int p(x) \log q(x) dx$$

$$= -\frac{1}{n} \sum_{x \sim p} \log q(x)$$
(28)

6.2 Perplexity

$$H(p,q) = \exp\left(-\int p(x)\log q(x)dx\right)$$

$$= \exp\left(-\frac{1}{n}\sum_{x\sim p}\log q(x)\right)$$
(29)

6.3 Forward KL Divergence

$$H(p,q) = \int p(x) \log \frac{p(x)}{q(x)} dx$$

$$= \frac{1}{n} \sum_{x \sim p} \log \frac{p(x)}{q(x)}$$
(30)

6.4 Reverse KL Divergence

$$H(p,q) = \int q(x) \log \frac{q(x)}{p(x)} dx$$

$$= \frac{1}{n} \sum_{x \sim q} \log \frac{q(x)}{p(x)}$$
(31)

- 6.5 Jensen-Shannon Divergence
- 6.6 Earth Mover's Distance
- 7 ELBO
- 8 Adversarial Loss
- 8.1 Adversarial Loss
- 8.2 GAN Loss
- 8.3 MMGAN Loss
- 8.4 WGAN Loss
- 8.5 LSGAN Loss
- 8.6 Gradient Penalty
- 8.7 Spectral Norm
- 9 Models
- 9.1 Base Models
- 9.1.1 Model
- 9.2 Regression
- 9.2.1 Linear Regression (Normal)
- 9.2.2 L1 Regression (Laplace)
- 9.2.3 Ridge Regression (Normal + Normal Prior on Weights)
- 9.2.4 Lasso Regression (Normal + Laplace Prior on Weights)
- 9.3 Classification
- 9.3.1 Logistic Regression (Bernoulli)
- 9.3.2 Bayesian Logistic Regression (Bernoulli)
- 9.3.3 Softmax Regression (Categorical)
- 9.4 Generative Adversarial Networks
- 9.4.1 Generative Adversarial Networks
- 9.4.2 GAN Model
- 9.4.3 MMGAN Model
- 9.4.4 WGAN Model
- 9.4.5 LSGAN Model
- 10 Monte Carlo
- 10.1 Monte Carlo Approximation
- **10.2** Linear Congruential Generator
- 10.3 Inverse Transform Sampling
- 10.4 Box-Muller
- 10.5 Marsaglia-Bray
- 10.6 Rejection Sampling