2026-11-04

BDBV (Ebolaviridae) updates, November 2026

The Nextstrain Team, James Hadfield[1,2], Cornelius, JT McCrone[1,2]

[1] Nextstrain

[2] Vaccine & Infectious disease division, Fred Hutchinson Cancer Center, Seattle, WA

Abstract

TKTK

Auspice map through time

See inferred transmissions using Health Zone (epicenter combined) across different periods of the outbreak (Figure 1).

> *Todo: make animation work in* `gazette`

Entire outbreak , Prior to June , July - Sept , Oct onwards




Figure 1. Auspice map with temporal controls.

ADAR

- What is ADAR
- How they were detected
- Detail masking
- Onwards transmission
- Toggle `on` `off` focus on ADAR samples in tree.




Figure 2. Samples with ADAR mutations detected are coloured according to the specific set (cluster) of ADAR mutations identified.

Chains are persisting in Health Zones for long times

> *This section relies a lot on the DTA and is probably overstating the truth*

While we currently infer the outbreak origin to be in the (combined) Health Zones of 'Bunia|Mongbwalu|Rwampara', the longest chains of continuous transmission within individual Health Zones are from Nizi (Figure 3). We infer two separate jumps from Bunia|Mongbwalu|Rwampara into Nizi in around April/May, and each of those chains continues to be present within Nizi.




Figure 3. The duration of within-Health-Zone transmission chains Circle size scales with the number of tips in a chain. The vast majority of chains persist for less than 1 month and involve only one sampled genome.




Figure 4. The duration a chain has been transmitting in a Health Zone vs the date of the chain's most recent sample

Limited transmission outside of DRC

Following JSON string doesn't work!

```
:trigger[Filter to non-DRC]{target=fig:auspice-countries update='{"filters":{"country":["France","Germany","Kenya","Nigeria
```




Figure 5. TKTK

Some mutations are artifactual

Nucleotide 5246 (corresponding to residue 262 in VP40) is `extremely homoplasic on the tree`. When `comparing this mutation against the sequencing protocol employed` it becomes clear that the `5246G` mutation is only present in two sequencing protocols, "Amplicon_Artic_pan-ebola/1000/v2.0.0" and "Amplicon_Artic_bdbv/400".




Figure 6. Homoplasic mutations at nuc 5246 are due to sequencing protocol used