Deep linear networks are mildly singular at every exact fit

Niels Feld*

~~Draft of 28~~ 29 October ~~2026.~~ 2026, 00:19:56 (Paris, UTC+01:00). Not ~~submitted; not~~ refereed.

Visual comparison with the draft of 28 October 2026.
~~Deletions~~ Additions.
Source-provenance identifiers were removed from the comparison baseline;
its mathematical text was not altered.

Abstract

Let $P(A) = A_{D-1} \cdots A_0$ be the product of a chain of real matrices of arbitrary sizes, and let Φ be a matrix. At every point A of the fibre $P^{-1}(\Phi)$ we determine the volume of the sublevel sets $\{B : \frac{1}{2}\|P(B) - \Phi\|_{\mathrm{F}}^2 < \varepsilon\}$ inside a small fixed ball around A: it is comparable to $\varepsilon^{\lambda}(\log 1/\varepsilon)^{m-1}$ as $\varepsilon \to 0$. The exponent λ is half the local codimension of the fibre at A, and the integer $m \geq 1$ is computed by a finite combinatorial procedure from the ranks of the partial products $A_{j-1} \cdots A_i$, equivalently from the barcode of A as a representation of an equioriented quiver of type A. ~~In the language of singular learning theory~~For a nonzero loss germ, λ and m are the local real log canonical threshold of the squared loss and its multiplicity. The equality $\lambda = \frac{1}{2}$ codim at every point was conjectured by Pepin Lehalleur and Rimányi, who proved it for the minimum of λ over the fibre. The multiplicity m can be strictly smaller than the number of top-dimensional strata of the fibre through A. The main statements have been formalized in the Lean 4 proof assistant; we describe the formal statement and exactly which parts of the mathematical statements it covers.

Contents

1 Introduction **3**
1.1 The question . . . 3
1.2 Main results . . . 3
1.3 Context . . . 4
1.4 Method . . . 5
1.5 Formalization . . . 5
1.6 Organization . . . 5

2 Notation and conventions **6**
2.1 Numbers, intervals and matrices . . . 6
2.2 Comparison of functions . . . 6

*~~See the AI use statement, Section 12.~~ See the disclosure of AI use in Section 12.

2.3 Hausdorff measures and Hausdorff dimension . . . 7
2.4 Measures and integrals . . . 8
2.5 Chains . . . 8

3 Definitions and statements **9**
3.1 Local volume order . . . 9
3.2 The first main theorem . . . 10
3.3 Barcodes . . . 10
3.4 The second main theorem . . . 14
3.5 Examples . . . 14

4 Local volume orders **16**

5 Orbits of chains **20**
5.1 Orbits are submanifolds . . . 21
5.2 Linked pairs and the normal space . . . 21
5.3 The slice chart . . . 22
5.4 Switches and the closure order . . . 23
5.5 Strata of a fibre . . . 24

6 Reduction to the zero target **25**
6.1 A Schur complement estimate . . . 25
6.2 The reduction . . . 26

7 The zero target: a recursion over orbits **28**
7.1 The shear . . . 28
7.2 Potentials and a dilation . . . 29
7.3 The recursion . . . 30

8 The exponent and the codimension **32**
8.1 Stationarity . . . 32
8.2 The exponent . . . 33
8.3 The codimension . . . 33

9 The multiplicity **34**

10 Proofs of the main theorems **35**

11 The formalization **36**
11.1 What is formalized . . . 36
11.2 What a reader has to trust . . . 37
11.3 Deceptive formal statements . . . 37
11.4 ~~Compilation records~~The public proof file and its checks . . . 38
11.5 ~~A single file and a local recheck~~ . . . 39

11.5 Reproducing the check 39

12 ~~AI use statement~~ 41

12 Disclosure of AI use 41

A The formal statement and its translation 43

A.1 Reading conventions 43

A.2 The statement 45

A.3 The program 48

B The rank order is generated by switches 53

1 Introduction

1.1 The question

Fix an integer $D \geq 1$ (the *depth*) and integers $n_0, \dots, n_D \geq 0$ (the *widths*). A *chain* is a tuple $A = (A_0, \dots, A_{D-1})$ of real matrices with $A_k \in \mathbb{R}^{n_{k+1} \times n_k}$, and its *product* is

$$P(A) = A_{D-1} A_{D-2} \cdots A_0 \in \mathbb{R}^{n_D \times n_0}.$$

In machine learning, $x \mapsto P(A)x$ is the function computed by a *deep linear network* with weights A. Given a matrix $\Phi \in \mathbb{R}^{n_D \times n_0}$, the squared loss

$$L_\Phi(B) = \tfrac{1}{2} \|P(B) - \Phi\|_{\mathrm{F}}^2$$

uses the Frobenius norm $\|M\|_{\mathrm{F}} = (\sum_{i,j} M_{ij}^2)^{1/2}$ and vanishes exactly on the *fibre* $F_\Phi = P^{-1}(\Phi)$; its points are the *exact fits* of Φ. This paper describes, at every exact fit A, how ~~fast~~ the volume

$$\mathrm{vol}\{B : \ \|B - A\| < \delta, \ L_\Phi(B) < \varepsilon\}$$

~~tends to 0~~ behaves as $\varepsilon \to 0$, for a small fixed radius δ. Here $\|\cdot\|$ is the Euclidean norm in all matrix entries and vol is Lebesgue measure.

1.2 Main results

The precise statements are Theorems A and B in Section 3. In words:

- **Theorem A.** For every exact fit A there are unique $\lambda \in [0, +\infty[$ and $m \in \mathbb{N}^*$ such that, for every sufficiently small fixed $\delta > 0$, the volume above is bounded above and below by constant multiples of $\varepsilon^\lambda (\log 1/\varepsilon)^{m-1}$ for all small ε. Moreover

  $$\lambda = \tfrac{1}{2} \operatorname{codim}_A F_\Phi,$$

  where $\operatorname{codim}_A F_\Phi$ is the dimension of the space of chains minus the local Hausdorff dimension of F_Φ at A. In particular $2\lambda \in \mathbb{N}$. The pair (λ, m) does not change if Lebesgue measure is multiplied by a function that is continuous and positive at A.

- **Theorem B.** The integers 2λ and m are given by an explicit finite procedure whose only inputs are the widths and the ranks of the partial products $A_{j-1} \cdots A_i$ $(0 \leq i \leq j \leq D)$. The procedure is phrased in terms of the *barcode* of A: the decomposition of A, viewed as a representation of the quiver $0 \to 1 \to \cdots \to D$, into interval representations.

The ~~multiplicity m has a geometric description (~~precise formula for the multiplicity is Theorem 9.2~~): $m-1$ is the largest number of *critical* strata, not counting the last one, along a chain of strata of the fibre that degenerate to A and along which the local codimension stays constant. In small cases m equals the number of top-dimensional strata of the fibre through A, but this fails in general~~; the finite chains and the integers maximized along them are defined in Section 3. The multiplicity does not, in general, count the components of maximal dimension meeting at the chosen point: Example 3.18 has $m = 3$ and four ~~top-dimensional strata~~such components.

1.3 Context

In Watanabe's singular learning theory [37, 38], the numbers λ and m attached to a nonnegative analytic loss at a zero, when the loss does not vanish identically near that point, are the *local real log canonical threshold* and its *multiplicity*; they govern, for instance, the asymptotics of Laplace integrals $\int e^{-tL}$ and of Bayesian free energies. We use only the volume definition (Definition 3.1) and none of these consequences. That definition also covers losses identically zero near the point, for which the pair is $(0, 1)$. Real-analytic and ideal-theoretic treatments of these thresholds are also given by Saito [33] and Lin [23]. For a sum of squares of real-analytic functions one has $\lambda \leq \frac{1}{2}$ codim at every zero, where codim is the local codimension of the zero set (Remark 3.4, which uses the stratification of real-analytic sets)~~; singularities with equality are informally called *mildly singular*~~.

For ~~deep linear networks~~ positive widths, the *global* threshold of a deep linear network, the minimum of λ over the whole fibre, was computed by Aoyagi and Watanabe for $D = 2$ [2] and by Aoyagi for all D [3]. Pepin Lehalleur and Rimányi [24] computed the codimension of the fibres and the number of their top-dimensional components using quiver representations. By comparing with Aoyagi's formula, they showed that the global threshold equals half the codimension of the fibre, for every target Φ with nonempty fibre, and they conjectured that the equality holds at every point of the fibre ~~[24, Section 8]~~[24, Remark 8.7]. Theorem A proves this conjecture; our proof does not use Aoyagi's formula. For equal widths, the formula for λ given by Theorems A and B is stated without proof in [16, Appendix B, Theorem B.5], where it serves as ground truth for a benchmark of posterior samplers. The stratification of the fibres of P over $\mathbb{R}$, the dimensions of the strata and their closure relations are studied in detail by Shewchuk and Bhattacharya [34], and the top-dimensional components of the zero fibre are described by Koncki and Rimányi [17]. We have not found the multiplicity formula of Theorem B in the literature for $D \geq 3$ (for $D = 2$ see below); ~~our searchcovered the papers above and their citations, the work of Aoyagi, Lin and Kurumadani on learning coefficients, and work on log canonical thresholds of determinantal varieties, but it was not exhaustive~~this statement reflects a bounded literature search.

For $D = 2$, the pointwise problem is part (a) of the question MAIS-O77 [29] of the *Math for AI Safety* program [25, 30], and also its question MAIS-O70 [28] on reduced-rank regression. Candidate solutions giving the depth-two formula for (λ, m), and the depth-two case of Theorem A, were posted in August 2026 by Sneiderman [26] and by Novikov with OpenAI Codex [27]. A Lean-checked proof is described in [9], and Example 3.16 explains how Theorem B recovers the formula. In concurrent work motivated by low-rank adaptation, Ferry, Monod and Yano [11] study the two-factor case with the Frobenius norm replaced by a positive semidefinite quadratic form on matrices. When that form is positive definite, their loss lies between two positive multiples of L_Φ, so by Lemma 4.6 its local invariants at exact fits are those of Theorem B with $D = 2$. When the form is degenerate, the zero set is the preimage of an affine subspace of matrices rather than a fibre of P; they analyse this case locally, and our methods do not cover it (see Remark 10.3). Estimators of local

learning coefficients ~~[21, 12, 16] are often calibrated on deep linear networks;~~ are studied in [21, 16]; the earlier work of Furman and Lau [12] was merged into [21]. Theorem B ~~gives the exact values at every exact fit , which in general differ from the global value at the minimizer that an optimizer finds. These test cases all~~ determines the local pair at an exact fit from its rank table. This pair need not coincide with the global pair; an application to the output of an optimizer therefore requires the ranks at that output. Hitchcock and Hoogland [16, Remark B.6] already emphasize that deep-linear examples satisfy $\lambda = \frac{1}{2}\operatorname{codim}$ ~~, so they do not exercise the strictly smaller values that occur for other losses~~and consequently do not represent losses with a strictly smaller threshold, such as x^4 at 0.

Related pointwise computations in other statistical models appear in Lin [23, Section 5] ; formulas at nonsingular points of the set of realizable parameters and upper bounds at certain singular points are developed by Kurumadani [19, 20]. Combinatorial formulas for a threshold and its multiplicity, including a maximization over chains of subspaces, also occur for real hyperplane arrangements in Kosta and Windisch [18, Theorem 4.2 and Corollary 4.4] . That work builds on the resolution constructions of De Concini and Procesi [7] and the threshold calculations of Mustaţă [32] and Teitler [35]. These are related precedents, rather than inputs to the proofs below.

1.4 Method

The proof uses four ingredients.

(i) The orbits of $\prod_v \mathrm{GL}_{n_v}(\mathbb{R})$ on chains, classified by barcodes. A transverse slice to the orbit of an interval representation has explicit coordinates, one for each *linked pair* of intervals (Section 5). The interval classification is the type-A case of Gabriel's theorem [13] and, in persistence-module language, is proved by Crawley-Boevey [6]; the normal-space calculation is the interval case of Voigt's lemma [36].

(ii) A Schur-complement change of coordinates, which splits off the rank of Φ and reduces the problem to the zero target (Section 6).

(iii) For the zero target, a dilation of the slice coordinates under which the loss is homogeneous. Weighted polar coordinates then give a recursion for (λ, m) over the finite poset of orbits (Section 7).

(iv) A stationarity argument that identifies 2λ with a combinatorial minimum, and an explicit family of submanifolds of the fibre that identifies the same minimum with the local codimension (Sections 8 and 9).

1.5 Formalization

The public repository [10] contains a Lean 4 ~~proofs of the main statements exist,~~ formalization built on the Mathlib library. ~~They were~~ It was produced with AI assistance; see Section 12. Section 11 and Appendix A state the formal theorem and translate every definition it depends on, so that a reader can check, without reading the formal proofs, how it implies Theorem A and the formulas of Theorem B; Proposition A.1 and the remark after Proposition A.2 list the steps that remain in the text. The mathematical text (Sections 1–10 and Appendix B) does not depend on the formalization.

1.6 Organization

- Section 2 fixes notation.

- Section 3 gives the definitions and states the results, with examples.
- Sections 4–9 contain the proofs, which are assembled in Section 10.
- Section 11 discusses the formalization, and Section 12 is the AI use statement.
- Appendix A contains the formal statement and its translation. Appendix B proves a combinatorial lemma on barcodes.

2 Notation and conventions

2.1 Numbers, intervals and matrices

$\mathbb{N} = \{0, 1, 2, \dots\}$ and $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$. We use the French notation for intervals of real numbers: for real numbers $a \le b$,

$$[a,b] = \{x \in \mathbb{R} :\ a \le x \le b\}, \qquad]a,b[= \{x \in \mathbb{R} :\ a < x < b\},$$
$$[a,b[= \{x \in \mathbb{R} :\ a \le x < b\}, \qquad]a,b] = \{x \in \mathbb{R} :\ a < x \le b\},$$

and similarly $]a,+\infty[$ and $[a,+\infty[$. The extended half-line is $[0,+\infty] = [0,+\infty[\cup \{+\infty\}$, with its usual order, addition and multiplication (and $0 \cdot (+\infty) = 0$). For integers a, b, the *integer interval* is $[\![a,b]\!] = \{k \in \mathbb{Z} :\ a \le k \le b\}$; it is empty if $a > b$. We write $\#X$ for the cardinality of a finite set X.

For $p, q \in \mathbb{N}$, $\mathbb{R}^{p\times q}$ is the space of real $p \times q$ matrices, identified with the linear maps $\mathbb{R}^q \to \mathbb{R}^p$; if $p = 0$ or $q = 0$ it is the zero space. We write I_p for the identity matrix of size p, $\operatorname{rk} M$ for the rank of a matrix M,

~~$\|M\|_{\mathrm{F}} = \left(\sum_{i,j} M_{ij}^2\right)^{1/2}$ and $\|M\|_{\mathrm{op}} = \sup\{\|Mx\| :\ x \in \mathbb{R}^q,\ \|x\| = 1\}$~~

$$\|M\|_{\mathrm{F}} = \left(\sum_{i,j} M_{ij}^2\right)^{1/2} \quad \text{and} \quad \|M\|_{\mathrm{op}} = \sup\{\|Mx\| :\ x \in \mathbb{R}^q,\ \|x\| \le 1\}$$

for its Frobenius norm and its operator norm, where $\|x\|$ is the Euclidean norm of a vector. This definition gives operator norm 0 when $q = 0$, since the closed unit ball of $\mathbb{R}^0$ consists of its zero vector. Recall that $\|MN\|_{\mathrm{F}} \le \|M\|_{\mathrm{op}}\|N\|_{\mathrm{F}}$ and $\|M\|_{\mathrm{op}} \le \|M\|_{\mathrm{F}}$. For a square matrix M, $e^M = \exp(M) = \sum_{k\ge 0} M^k/k!$.

2.2 Comparison of functions

Let $\varepsilon_1 \in]0,+\infty]$ and let $f, g :]0,\varepsilon_1[\to [0,+\infty]$ be two functions. For $\varepsilon_0 \in]0,\varepsilon_1]$ we write $f \asymp g$ on $]0,\varepsilon_0[$ if there are real numbers c, C with $0 < c \le C$ and $c\,g(\varepsilon) \le f(\varepsilon) \le C\,g(\varepsilon)$ for every $\varepsilon \in]0,\varepsilon_0[$. We write $f(\varepsilon) \asymp g(\varepsilon)$ as $\varepsilon \to 0$ if $f \asymp g$ on $]0,\varepsilon_0[$ for some $\varepsilon_0 \in]0,\varepsilon_1]$. Throughout, log is the natural logarithm, and

$$L :]0,1[\to]0,+\infty[, \qquad L(\varepsilon) = \log(1/\varepsilon) = -\log\varepsilon. \tag{2.1}$$

The function L is decreasing, and $L(\varepsilon) > 1$ if and only if $\varepsilon < e^{-1}$. For $\kappa > 0$ and $\varepsilon \in]0, \min(1,\kappa)[$ we have $L(\varepsilon/\kappa) = L(\varepsilon) + \log\kappa$; hence

$$\tfrac{1}{2}L(\varepsilon) \le L(\varepsilon/\kappa) \le 2L(\varepsilon) \qquad \text{whenever } \varepsilon < \min(1,\kappa) \text{ and } L(\varepsilon) \ge 2|\log\kappa|. \tag{2.2}$$

2.3 Hausdorff measures and Hausdorff dimension

Let $(E, \|\cdot\|)$ be a finite-dimensional real normed vector space. The *diameter* of a nonempty subset $U \subseteq E$ is $\operatorname{diam} U = \sup\{\|x-y\| : x, y \in U\} \in [0, +\infty]$, and $\operatorname{diam} \emptyset = 0$. For $s \in [0, +\infty[$ put $\gamma_s(U) = (\operatorname{diam} U)^s$ if $U \neq \emptyset$, with the convention $0^0 = 1$, and $\gamma_s(\emptyset) = 0$. For $S \subseteq E$, $s \in [0, +\infty[$ and $\eta \in]0, +\infty[$ put

$$\mathcal{H}^s_\eta(S) = \inf\Big\{\sum_{i\in\mathbb{N}} \gamma_s(U_i) : \ (U_i)_{i\in\mathbb{N}} \in \mathcal{C}_\eta(S)\Big\} \in [0, +\infty],$$

where $\mathcal{C}_\eta(S)$ is the set of sequences $(U_i)_{i\in\mathbb{N}}$ of subsets of E with $S \subseteq \bigcup_{i\in\mathbb{N}} U_i$ and $\operatorname{diam} U_i \leq \eta$ for all i (η-*covers* of S). The function $\eta \mapsto \mathcal{H}^s_\eta(S)$ is nonincreasing. The *s-dimensional Hausdorff measure* of S (for the norm $\|\cdot\|$) is

$$\mathcal{H}^s(S) = \sup_{\eta>0} \mathcal{H}^s_\eta(S) = \lim_{\eta\to 0} \mathcal{H}^s_\eta(S) \in [0, +\infty],$$

and the *Hausdorff dimension* of S is

$$\dim_{\mathrm{H}} S = \sup\big\{s \in [0, +\infty[: \ \mathcal{H}^s(S) = +\infty\big\}, \qquad \text{with } \sup \emptyset = 0.$$

We use the following standard facts [31, Chapter 4], [8].

(H0) If $0 \leq s < t$ and $\mathcal{H}^s(S) < +\infty$, then $\mathcal{H}^t(S) = 0$. Consequently $\dim_{\mathrm{H}} S = \inf\{t \in [0, +\infty[: \ \mathcal{H}^t(S) = 0\}$. Moreover $\mathcal{H}^t(S) = 0$ for every $t > \dim E$, so $\dim_{\mathrm{H}} S \leq \dim E$.

(H1) If $S \subseteq S'$ then $\dim_{\mathrm{H}} S \leq \dim_{\mathrm{H}} S'$. If $S = \bigcup_{i\in\mathbb{N}} S_i$ then $\dim_{\mathrm{H}} S = \sup_{i\in\mathbb{N}} \dim_{\mathrm{H}} S_i$.

(H2) If E' is another finite-dimensional real normed vector space, $S \subseteq E$, and $f : S \to E'$ is Lipschitz, then $\dim_{\mathrm{H}} f(S) \leq \dim_{\mathrm{H}} S$. Consequently $\dim_{\mathrm{H}} S$ does not depend on the choice of the norm on E (all norms are equivalent, so the identity map is bi-Lipschitz), and $\dim_{\mathrm{H}}$ is invariant under bi-Lipschitz maps, for instance under the restriction of a C^1 diffeomorphism between open sets to a compact subset (a C^1 map is Lipschitz on every compact subset of its open domain).

(H3) A nonempty open subset of a C^1 embedded submanifold of E of dimension k has Hausdorff dimension k.

Definition 2.1 (Local codimension)**.** Let $(E, \|\cdot\|)$ be as above, $S \subseteq E$ and $a \in E$. For $\delta \in]0, +\infty[$ let $B_\delta(a) = \{x \in E : \ \|x - a\| < \delta\}$ be the open ball. The *local Hausdorff dimension* of S at a and the *local codimension* of S at a are

$$d_a(S) = \inf_{\delta>0} \dim_{\mathrm{H}}\big(S \cap B_\delta(a)\big) \in [0, \dim E], \qquad \operatorname{codim}_a S = \dim E - d_a(S) \in [0, \dim E].$$

The function $\delta \mapsto \dim_{\mathrm{H}}(S \cap B_\delta(a))$ is nondecreasing by (H1), so the infimum is a limit as $\delta \to 0$. If $\|\cdot\|'$ is another norm on E, with balls $B'_\delta(a)$, there is $K \geq 1$ with $B'_{\delta/K}(a) \subseteq B_\delta(a) \subseteq B'_{K\delta}(a)$ for all $\delta > 0$; with (H1) and (H2) this shows that $d_a(S)$ does not depend on the norm. If S is a finite union of C^1 embedded submanifolds $S_1, \dots, S_N$ of E and $a \in \overline{S}$, then

$$d_a(S) = \max\{\dim S_i : \ i \in [\![1, N]\!], \ a \in \overline{S_i}\}. \tag{2.3}$$

Indeed, if $a \notin \overline{S_i}$, then $S_i \cap B_\delta(a) = \emptyset$ for small δ; if $a \in \overline{S_i}$, then $S_i \cap B_\delta(a)$ is a nonempty open subset of S_i for every $\delta > 0$. Apply (H1) and (H3).

2.4 Measures and integrals

Let E be a finite-dimensional real vector space, with its Borel σ-algebra. A *Haar measure* on E is a positive multiple of the image of Lebesgue measure under a linear isomorphism $\mathbb{R}^{\dim E} \to E$; this does not depend on the isomorphism, and any two Haar measures differ by a positive constant factor. When $E = \mathbb{R}^N$ we write vol for Lebesgue measure. For a function $\varphi : E \to \mathbb{R}$ we write $\varphi^+ = \max(\varphi, 0)$. For a Borel set $S \subseteq E$, a Haar measure μ and a function $g : E \to [0, +\infty]$, the *lower integral*

$$\int_S^- g\, d\mu = \sup\Big\{ \int_S g'\, d\mu :\ g' : E \to [0,+\infty] \text{ Borel measurable},\ g' \le g \Big\} \in [0,+\infty]$$

equals the Lebesgue integral when g is measurable. It is used only so that no measurability hypothesis on the functions φ below is needed. It is nondecreasing in S and in g, and if $c_1, c_2 \in [0, +\infty[$ satisfy $c_1 \le \varphi^+(x) \le c_2$ for all $x \in S$, then

$$c_1\, \mu(S) \le \int_S^- \varphi^+\, d\mu \le c_2\, \mu(S). \tag{2.4}$$

(For the lower bound take ~~$g' = c_1$~~ g' equal to c_1 on S ~~; for~~ and to 0 outside S; this function is Borel measurable and lies below φ^+. For the upper bound note that every admissible g' is at most c_2 on S.)

2.5 Chains

Let $D \in \mathbb{N}$. The *quiver* of type A_{D+1} considered here has the *vertices* $v \in [\![0, D]\!]$ and the *arrows* $k \in [\![0, D-1]\!]$, the arrow k going from the vertex k to the vertex $k+1$:

$$0 \longrightarrow 1 \longrightarrow \cdots \longrightarrow D.$$

A *width vector* is an element $n = (n_0, \dots, n_D) \in \mathbb{N}^{D+1}$; it determines D. For a width vector n put

$$\mathrm{Rep}(n) = \prod_{k=0}^{D-1} \mathbb{R}^{n_{k+1} \times n_k}, \qquad N(n) = \dim \mathrm{Rep}(n) = \sum_{k=0}^{D-1} n_k n_{k+1}.$$

The elements of $\mathrm{Rep}(n)$ are the *chains* $A = (A_0, \dots, A_{D-1})$; the matrix A_k is attached to the arrow k. For $D = 0$, $\mathrm{Rep}(n)$ is the zero space. Unless stated otherwise, $\mathrm{Rep}(n)$ carries the Euclidean norm $\|A\| = (\sum_k \|A_k\|_{\mathrm{F}}^2)^{1/2}$, and $B_\delta(A)$ denotes its open ball of radius δ around A. For $0 \le i \le j \le D$ the *partial products* of a chain A are

$$P_{ij}(A) = A_{j-1} A_{j-2} \cdots A_i \in \mathbb{R}^{n_j \times n_i} \quad (i < j), \qquad P_{ii}(A) = \mathrm{I}_{n_i},$$

and its *product* is $P(A) = P_{0D}(A)$. The *rank table* of A is the family $R(A) = (R_{ij}(A))_{0 \le i \le j \le D}$ of integers $R_{ij}(A) = \mathrm{rk}\, P_{ij}(A)$; note that $R_{vv}(A) = n_v$ for every vertex v.

For $\Phi \in \mathbb{R}^{n_D \times n_0}$ we define the *fibre* $F_{n,\Phi}$ and the *loss* $L_{n,\Phi} : \mathrm{Rep}(n) \to \mathbb{R}$ by

$$F_{n,\Phi} = \{A \in \mathrm{Rep}(n) :\ P(A) = \Phi\}, \qquad L_{n,\Phi}(A) = \tfrac{1}{2}\|P(A) - \Phi\|_{\mathrm{F}}^2.$$

The *zero fibre* is $Z(n) = F_{n,0}$, and we put $K_n(A) = \|P(A)\|_{\mathrm{F}}^2 = 2L_{n,0}(A)$. When only one width vector n is involved, we write F_Φ and L_Φ for $F_{n,\Phi}$ and $L_{n,\Phi}$. The points of F_Φ are the *exact fits* of Φ.

The group $G(n) = \prod_{v=0}^{D} \mathrm{GL}_{n_v}(\mathbb{R})$ acts linearly on $\mathrm{Rep}(n)$ by

$$(g \cdot A)_k = g_{k+1} A_k g_k^{-1} \qquad (g = (g_v)_{v \in [\![0,D]\!]} \in G(n),\ k \in [\![0, D-1]\!]).$$

Then $P_{ij}(g \cdot A) = g_j P_{ij}(A) g_i^{-1}$, so the rank table is constant on the orbits of $G(n)$. The group $G(n)$ is a Lie group, open in the real vector space

$$\mathfrak{g}(n) = \prod_{v=0}^{D} \mathbb{R}^{n_v \times n_v},$$

which is its Lie algebra. Its exponential map is $\exp : \mathfrak{g}(n) \to G(n)$, $\xi = (\xi_v)_v \mapsto (e^{\xi_v})_v$. We use only that $t \mapsto \exp(t\xi)$ is a one-parameter subgroup of $G(n)$ with derivative ξ at $t = 0$, and that exp restricts to a diffeomorphism from a neighbourhood of 0 in $\mathfrak{g}(n)$ onto a neighbourhood of the identity in $G(n)$.

3 Definitions and statements

3.1 Local volume order

Definition 3.1 (Local volume order)**.** Let $(E, \|\cdot\|)$ be a finite-dimensional real normed vector space, $U \subseteq E$ an open subset, $a \in U$, and $f : U \to \mathbb{R}$ a continuous function. Let μ be a Haar measure on E, and let $\varphi : E \to \mathbb{R}$ be a function that is continuous at a and satisfies $\varphi(a) > 0$ (a *prior*). For $\delta \in]0, +\infty[$ with $B_\delta(a) \subseteq U$ and for $\varepsilon \in]0, +\infty[$ put

$$S_{f,a}(\delta, \varepsilon) = \{x \in B_\delta(a) : \ |f(x) - f(a)| < \varepsilon\}, \qquad V_{f,a}^{\mu,\varphi}(\delta, \varepsilon) = \int^{-}_{S_{f,a}(\delta,\varepsilon)} \varphi^+ \, d\mu \ \in [0, +\infty].$$

Let $\lambda \in \mathbb{R}$ and $m \in \mathbb{N}$. We say that f has *local volume order* (λ, m) *at* a (for μ and φ) if $\lambda \geq 0$, $m \geq 1$, and there is $\delta_0 \in]0, +\infty[$ with $B_{\delta_0}(a) \subseteq U$ such that for every $\delta \in]0, \delta_0[$ there are real numbers c, C, ε_0 with $0 < c \leq C$ and $0 < \varepsilon_0 < e^{-1}$ such that

$$c\,\varepsilon^\lambda L(\varepsilon)^{m-1} \ \leq \ V_{f,a}^{\mu,\varphi}(\delta, \varepsilon) \ \leq \ C\,\varepsilon^\lambda L(\varepsilon)^{m-1} \qquad \text{for all } \varepsilon \in]0, \varepsilon_0[, \tag{3.1}$$

where $L(\varepsilon) = \log(1/\varepsilon)$ is the function (2.1) of Section 2.2.

The set $S_{f,a}(\delta, \varepsilon)$ is open, since f is continuous on the open set U; in particular it is a Borel set. The order of the quantifiers is: first δ_0; then every fixed radius $\delta \in]0, \delta_0[$; then constants c, C, ε_0, which may depend on δ; then all $\varepsilon \in]0, \varepsilon_0[$. Nothing uniform in δ is required. If $f \geq 0$ and $f(a) = 0$, then $S_{f,a}(\delta, \varepsilon)$ is the sublevel set $\{x \in B_\delta(a) : \ f(x) < \varepsilon\}$.

Lemma 3.2. *Keep the notation and hypotheses of Definition 3.1.*

(a) *There is at most one pair* $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$ *such that* f *has local volume order* (λ, m) *at* a *for* μ *and* φ*.*

(b) *Let* μ' *be a Haar measure on* E*,* $\varphi' : E \to \mathbb{R}$ *a function continuous at* a *with* $\varphi'(a) > 0$*, and* $\|\cdot\|'$ *a norm on* E*. If* f *has local volume order* (λ, m) *at* a *for* μ*,* φ *and the balls of* $\|\cdot\|$*, then it has local volume order* (λ, m) *at* a *for* μ'*,* φ' *and the balls of* $\|\cdot\|'$*.*

(c) *If* f *is constant on a neighbourhood of* a*, then* f *has local volume order* $(0, 1)$ *at* a*.*

The proof is in Section 4. By (a) and (b), when f has a local volume order at a for some choice of $(\mu, \varphi, \|\cdot\|)$, it has the same one for every choice; we then denote it by $(\lambda_a(f), m_a(f))$.

Remark 3.3 (Terminology). Let $f \geq 0$ be real-analytic on a neighbourhood of a with $f(a) = 0$. ~~By~~ If f is identically zero on a neighbourhood of a, its local volume order is $(0, 1)$ by Lemma 3.2(c). Otherwise, Hironaka's resolution of singularities~~, f then has~~ [15] gives a

local volume order at a~~, and~~; in this case $\lambda_a(f)$ and $m_a(f)$ are the local real log canonical threshold of f at a and its multiplicity ([37, Chapters 2 and 6]). We do not use ~~this~~resolution of singularities: all existence statements below are proved directly~~, without resolution of singularities~~. The losses L_Φ are polynomials, so this remark applies to them; the content of the theorems below is the value of the pair, not its existence.

Remark 3.4 (An upper bound). Let $f = \sum_{j=1}^{J} f_j^2$, where the functions $f_j : U \to \mathbb{R}$ are of class C^1 and vanish at a. Suppose that the zero set $f^{-1}(0)$ contains a C^1 embedded submanifold S of E of codimension c_S with $a \in \overline{S}$, and that f has a local volume order (λ, m) at a. Then $\lambda \leq c_S/2$. Indeed, identify E with $\mathbb{R}^{\dim E}$ by a linear isomorphism. By Lemma 3.2(b), f has local volume order (λ, m) at a for Lebesgue measure vol, the prior 1 and the Euclidean norm, with some radius δ_0 as in Definition 3.1. Let $\delta \in]0, \delta_0[$, choose $x_0 \in S \cap B_{\delta/2}(a)$, and choose a C^1 diffeomorphism Ψ from an open neighbourhood Q' of 0 in $\mathbb{R}^{\dim E - c_S} \times \mathbb{R}^{c_S}$ onto an open neighbourhood $Q \subseteq B_\delta(a)$ of x_0 with $\Psi(0) = x_0$ and $\Psi^{-1}(S \cap Q) = Q' \cap (\mathbb{R}^{\dim E - c_S} \times \{0\})$. Let $Q_0 = [-t, t]^{\dim E} \subseteq Q'$ be a cube, and write its points (y, w) with $y \in \mathbb{R}^{\dim E - c_S}$ and $w \in \mathbb{R}^{c_S}$. Each $f_j \circ \Psi$ is of class C^1 and vanishes on $Q_0 \cap \{w = 0\}$, so by the mean value inequality on the segments $\{(y, \sigma w) : \ \sigma \in [0,1]\} \subseteq Q_0$ there is $K > 0$ with $f(\Psi(y,w)) \leq K\|w\|^2$ for $(y,w) \in Q_0$. Let $k_0 > 0$ be a lower bound of $|\det d\Psi|$ on Q_0. Since $f \geq 0$ and $f(a) = 0$, the set $\Psi(\{(y,w) \in Q_0 : \ K\|w\|^2 < \varepsilon\})$ is contained in $S_{f,a}(\delta, \varepsilon)$, so

$$\begin{aligned} V_{f,a}^{\mathrm{vol},1}(\delta, \varepsilon) &\geq \operatorname{vol} \Psi\big(\{(y,w) \in Q_0 : \ K\|w\|^2 < \varepsilon\}\big) \\ &\geq k_0 \, (2t)^{\dim E - c_S} \ \operatorname{vol}\{w \in [-t,t]^{c_S} : \ \|w\|^2 < \varepsilon/K\}, \end{aligned}$$

which is at least a positive constant times $\varepsilon^{c_S/2}$ for small ε. Since $\varepsilon^\lambda L(\varepsilon)^{m-1}/\varepsilon^{c_S/2} \to 0$ as $\varepsilon \to 0$ when $\lambda > c_S/2$, the upper bound in (3.1) forces $\lambda \leq c_S/2$.

If f is real-analytic, its zero set is, near a, a finite union of real-analytic embedded submanifolds of E [4]. By (2.3), one of them, S, satisfies $a \in \overline{S}$ and has codimension $c_S = \operatorname{codim}_a f^{-1}(0)$. Hence $\lambda_a(f) \leq \frac{1}{2} \operatorname{codim}_a f^{-1}(0)$ whenever f has a local volume order at a. For the losses L_Φ, the submanifolds are given directly by Lemma 5.9. Theorem A below ~~therefore says that the zero sets of the losses~~ asserts equality in this upper bound for L_Φ ~~are "as mildly singular as possible"~~at every exact fit.

3.2 The first main theorem

Theorem A. *Let $D \in \mathbb{N}$, let $n \in \mathbb{N}^{D+1}$ be a width vector, let $\Phi \in \mathbb{R}^{n_D \times n_0}$, and let $A \in F_\Phi$. Then the loss $L_\Phi : \mathrm{Rep}(n) \to \mathbb{R}$ has a local volume order $(\lambda_A(L_\Phi), m_A(L_\Phi))$ at A, and*

$$\lambda_A(L_\Phi) = \tfrac{1}{2} \operatorname{codim}_A F_\Phi.$$

In particular $2\lambda_A(L_\Phi) \in \mathbb{N}$, and the pair does not depend on the Haar measure, on the prior or on the norm used in Definition 3.1.

For $D = 0$ the space $\mathrm{Rep}(n)$ is a point and P is the constant map with value I_{n_0}. The statement is then trivial: if $A \in F_\Phi$, then L_Φ is identically 0, its local volume order is $(0,1)$ by Lemma 3.2(c), and the codimension of a point in a point is 0. The content is the case $D \geq 1$.

3.3 Barcodes

In this subsection $D \in \mathbb{N}^*$ is fixed.

Definition 3.5 (Intervals, barcodes, slots). (a) An *interval* (of length D) is a subset of $[\![0, D]\!]$ of the form $[\![b, e]\!]$ with $0 \leq b \leq e \leq D$. The integers b and e are determined by the interval I; we write $b(I) = b$ and $e(I) = e$. The set of intervals is denoted Int_D.

(b) A *barcode* (of length D) is a map $h : \mathrm{Int}_D \to \mathbb{N}$. For $I \in \mathrm{Int}_D$, $\mathbf{1}_I$ is the barcode with $\mathbf{1}_I(I) = 1$ and $\mathbf{1}_I(I') = 0$ for $I' \neq I$. Barcodes are added, subtracted (when the result has values in $\mathbb{N}$) and compared pointwise, and $h = \sum_{I \in \mathrm{Int}_D} h(I)\,\mathbf{1}_I$ for every barcode h. In examples we write a barcode as the formal sum $\sum_I h(I)\,I$ of its intervals; for instance $2[\![0,0]\!] + [\![0,2]\!]$ is the barcode $2 \cdot \mathbf{1}_{[\![0,0]\!]} + \mathbf{1}_{[\![0,2]\!]}$.

(c) The set of *slots* of a barcode h is the finite set

$$\mathcal{S}(h) = \{(I, j) \in \mathrm{Int}_D \times \mathbb{N}^* : \ j \leq h(I)\}.$$

For a slot $s = (I, j)$ we put $I_s = I$, $b_s = b(I)$ and $e_s = e(I)$, so that $I_s = [\![b_s, e_s]\!]$. Thus $h(I)$ is the number of slots s with $I_s = I$, and distinct slots may have the same interval.

Definition 3.6 (Dimension vector and rank table of a barcode)**.** Let h be a barcode of length D. For a vertex $v \in [\![0, D]\!]$ put

$$\mathcal{S}_v(h) = \{s \in \mathcal{S}(h) : \ v \in I_s\}, \qquad \dim_v(h) = \#\mathcal{S}_v(h) = \sum_{I \ni v} h(I).$$

The *dimension vector* of h is $\dim(h) = (\dim_0(h), \ldots, \dim_D(h)) \in \mathbb{N}^{D+1}$. For $0 \leq i \leq j \leq D$ put

$$\rho_{ij}(h) = \#\{s \in \mathcal{S}(h) : \ [\![i, j]\!] \subseteq I_s\} = \sum_{I \supseteq [\![i,j]\!]} h(I).$$

The family $\rho(h) = (\rho_{ij}(h))_{0 \leq i \leq j \leq D}$ is the *rank table* of h. Note that $\rho_{vv}(h) = \dim_v(h)$ for every vertex v.

A slot s of h is *full-span* if $I_s = [\![0, D]\!]$, an *input slot* if $b_s = 0$, an *output slot* if $e_s = D$, and *interior* if $0 < b_s$ and $e_s < D$. A full-span slot is both an input and an output slot; every other slot is of exactly one of the three kinds input, output, interior. The barcode h is *zero-output* if $h([\![0, D]\!]) = 0$, i.e. if it has no full-span slot. Its *zero-output part* is the barcode

$$h^\circ = h - h([\![0, D]\!])\,\mathbf{1}_{[\![0,D]\!]},$$

which agrees with h except that $h^\circ([\![0, D]\!]) = 0$; thus $\mathcal{S}(h^\circ)$ is the set of slots of h that are not full-span. Since every vertex lies in $[\![0, D]\!]$,

$$\dim_v(h^\circ) = \dim_v(h) - h([\![0, D]\!]), \qquad \rho_{ij}(h^\circ) = \rho_{ij}(h) - h([\![0, D]\!]) \tag{3.2}$$

for every vertex v and all $0 \leq i \leq j \leq D$. For a width vector $n \in \mathbb{N}^{D+1}$ we denote by $\mathcal{Z}(n)$ the set of zero-output barcodes h of length D with $\dim(h) = n$, i.e. with $\dim_v(h) = n_v$ for every vertex $v \in [\![0, D]\!]$. It is a finite set, since $h(I) \leq n_{b(I)}$ for every $h \in \mathcal{Z}(n)$ and $I \in \mathrm{Int}_D$. For an integer $r \in \mathbb{N}$ with $r \leq n_v$ for all v, we write $n - r = (n_0 - r, \ldots, n_D - r)$.

Definition 3.7 (Interval chain)**.** Let h be a barcode of length D, put $n = \dim(h)$, and fix a total order on the finite set $\mathcal{S}(h)$. For each vertex v the set $\mathcal{S}_v(h)$ has exactly n_v elements (Definition 3.6). Let $\sigma_v : [\![1, n_v]\!] \to \mathcal{S}_v(h)$ be the unique increasing bijection, and for $s \in \mathcal{S}_v(h)$ let ϵ_s^v be the $\sigma_v^{-1}(s)$-th vector of the standard basis of $\mathbb{R}^{n_v}$. Thus $(\epsilon_s^v)_{s \in \mathcal{S}_v(h)}$ is the standard basis of $\mathbb{R}^{n_v}$, indexed by the slots present at v. The *interval chain* $\mathbb{I}(h) \in \mathrm{Rep}(n)$ is defined, for each arrow $k \in [\![0, D-1]\!]$, by

$$\mathbb{I}(h)_k\,\epsilon_s^k = \begin{cases} \epsilon_s^{k+1} & \text{if } k+1 \in I_s, \\ 0 & \text{if } k+1 \notin I_s, \end{cases} \qquad s \in \mathcal{S}_k(h).$$

So each matrix $\mathbb{I}(h)_k$ is a partial permutation matrix: a slot is carried from one vertex to the next as long as its interval lasts. Changing the total order on $\mathcal{S}(h)$ replaces $\mathbb{I}(h)$ by $g \cdot \mathbb{I}(h)$, where $g \in G(n)$ consists of permutation matrices; whenever $\mathbb{I}(h)$ is used below, a total order is fixed, and no statement depends on it.

Fact 3.8 (Interval decomposition)**.** (a) *For every width vector $n \in \mathbb{N}^{D+1}$ and every chain $A \in \mathrm{Rep}(n)$ there are a barcode h of length D with $\dim(h) = n$ and an element $g \in G(n)$ with $g \cdot A = \mathbb{I}(h)$.*

(b) *For every barcode h of length D, the rank table of the chain $\mathbb{I}(h) \in \mathrm{Rep}(\dim(h))$ (Section 2.5) is the rank table of the barcode h: $R(\mathbb{I}(h)) = \rho(h)$.*

(c) *For every barcode h of length D and all $0 \le b \le e \le D$,*

$$h(\llbracket b, e \rrbracket) = \rho_{be}(h) - \rho_{b-1,e}(h) - \rho_{b,e+1}(h) + \rho_{b-1,e+1}(h),$$

where $\rho_{ij}(h)$ is read as 0 when $i < 0$ or $j > D$. In particular the rank table $\rho(h)$ determines h.

Part (a) is Gabriel's theorem for the quiver $0 \to 1 \to \cdots \to D$ of type A_{D+1} [13]; for this quiver it is also the structure theorem for persistence modules [6]. Part (b) is a direct computation: for $s \in \mathcal{S}_i(h)$, $P_{ij}(\mathbb{I}(h))$ maps ϵ^i_s to ϵ^j_s if $\llbracket i, j \rrbracket \subseteq I_s$ and to 0 otherwise, so its rank is $\rho_{ij}(h)$. For (c), $\rho_{be}(h) - \rho_{b-1,e}(h)$ is the number of slots s with $b_s = b$ and $e_s \ge e$, and $\rho_{b,e+1}(h) - \rho_{b-1,e+1}(h)$ is the number of slots with $b_s = b$ and $e_s \ge e+1$.

Lemma 3.9. *Let $n \in \mathbb{N}^{D+1}$ and $A \in \mathrm{Rep}(n)$.*

(i) *There is exactly one barcode h of length D with $\rho(h) = R(A)$. It satisfies $\dim(h) = n$ and $h(\llbracket 0, D \rrbracket) = \operatorname{rk} P(A)$, and there is $g \in G(n)$ with $g \cdot A = \mathbb{I}(h)$.*

(ii) *Two chains $A, A' \in \mathrm{Rep}(n)$ lie in the same $G(n)$-orbit if and only if $R(A) = R(A')$.*

Proof. (i) By Fact 3.8(a) there are a barcode h of length D with $\dim(h) = n$ and $g \in G(n)$ with $g \cdot A = \mathbb{I}(h)$. Since the rank table is constant on orbits, $R(A) = R(\mathbb{I}(h)) = \rho(h)$ by Fact 3.8(b). A barcode h' with $\rho(h') = R(A) = \rho(h)$ equals h by Fact 3.8(c). Finally $h(\llbracket 0, D \rrbracket) = \rho_{0D}(h) = R_{0D}(A) = \operatorname{rk} P(A)$, because $\llbracket 0, D \rrbracket$ is the only interval containing $\llbracket 0, D \rrbracket$.

(ii) The rank table is constant on orbits. Conversely, if $R(A) = R(A')$, then (i) gives the same barcode h for A and A', and both lie in the orbit of $\mathbb{I}(h)$. □

Definition 3.10 (Barcode of a chain; residual barcode)**.** Let $n \in \mathbb{N}^{D+1}$ and $A \in \mathrm{Rep}(n)$.

(a) The *barcode of A* is the barcode $\beta(A)$ of Lemma 3.9(i): the unique barcode of length D with $\rho(\beta(A)) = R(A)$.

(b) The *residual barcode* of A is the zero-output part $h_A = \beta(A)^\circ$ of $\beta(A)$.

Put $r = \operatorname{rk} P(A)$. By Lemma 3.9(i), $\beta(A)$ has exactly r full-span slots, and each of them contains every vertex, so $r \le \dim_v(\beta(A)) = n_v$ for every vertex v. By (3.2), $h_A \in \mathcal{Z}(n - r)$. By Lemma 3.9, two chains $A, A' \in \mathrm{Rep}(n)$ lie in the same $G(n)$-orbit if and only if $\beta(A) = \beta(A')$, and A lies in the orbit of $\mathbb{I}(\beta(A))$.

Definition 3.11 (Links)**.** Let h be a barcode of length D. An ordered pair $(s, s') \in \mathcal{S}(h) \times \mathcal{S}(h)$ is *linked*, written $s \rightarrowtail s'$, if

$$b_s < b_{s'} \leq e_s + 1 \leq e_{s'}.$$

We denote by $\mathrm{Lk}(h) \subseteq \mathcal{S}(h) \times \mathcal{S}(h)$ the set of linked pairs. For a slot s let $\mathrm{in}(s) = \#\{s' \in \mathcal{S}(h) : \ s' \rightarrowtail s\}$ and $\mathrm{out}(s) = \#\{s' \in \mathcal{S}(h) : \ s \rightarrowtail s'\}$. The *output count* of h is

$$c_{\mathrm{out}}(h) = \#\{(s, s') \in \mathrm{Lk}(h) : \ s' \text{ is an output slot}\}.$$

The barcode h is *balanced* if $\mathrm{in}(s) = \mathrm{out}(s)$ for every interior slot s.

In words: $s \rightarrowtail s'$ means that $I_{s'}$ begins strictly after I_s begins and at most one vertex after I_s ends, and that $I_{s'}$ ends strictly after I_s ends; equivalently, $I_s \cup I_{s'}$ is an interval, and I_s begins and ends strictly before $I_{s'}$. No pair involving a full-span slot s_0 is linked: $s_0 \rightarrowtail s'$ would require $e_{s'} > D$, and $s \rightarrowtail s_0$ would require $b_s < 0$. If $s \rightarrowtail s'$, then s is not an output slot (since $e_s < e_{s'} \leq D$) and s' is not an input slot (since $b_{s'} > b_s \geq 0$). By Lemma 5.2 below, $\#\,\mathrm{Lk}(h)$ is the codimension of the orbit of $\mathbb{I}(h)$.

Definition 3.12 (Order and combinatorial invariants)**.** For barcodes h, k of length D we write $h \preceq k$ if $\dim(h) = \dim(k)$ and $\rho_{ij}(h) \leq \rho_{ij}(k)$ for all $0 \leq i \leq j \leq D$, and $h \prec k$ if moreover $h \neq k$. Let $n \in \mathbb{N}^{D+1}$ and $h \in \mathcal{Z}(n)$.

(i) $C(h) = \min\{c_{\mathrm{out}}(k) : \ k \in \mathcal{Z}(n), \ h \preceq k, \ k \text{ balanced}\}$. (Lemma 3.13 shows that this set is nonempty.)

(ii) h is *critical* if it is balanced and $c_{\mathrm{out}}(h) = C(h)$.

(iii) $F(h)$ is the maximum of $\#\{i \in [\![0, \ell - 1]\!] : \ h_i \text{ is critical}\}$ over all $\ell \in \mathbb{N}$ and all finite sequences $h = h_0 \prec h_1 \prec \cdots \prec h_\ell$ of elements of $\mathcal{Z}(n)$ with $C(h_i) = C(h)$ for all $i \in [\![0, \ell]\!]$.

The relation $\preceq$ is a partial order on the barcodes of length D; it is antisymmetric by Fact 3.8(c). Because $\mathcal{Z}(n)$ is finite and $\prec$ is a strict order, the maximum in (iii) is taken over a finite nonempty set (the sequence with $\ell = 0$ is allowed), and $F(h) \leq \#\mathcal{Z}(n)$. The last term h_ℓ of a sequence is never counted.

Lemma 3.13. *Let $n \in \mathbb{N}^{D+1}$ and $h \in \mathcal{Z}(n)$, and let k be a maximal element, for the order $\preceq$, of the finite set $\{k' \in \mathcal{Z}(n) : \ h \preceq k'\}$. Then every linked pair of k goes from an input slot to an output slot, and k is balanced. In particular the set minimized in the definition of $C(h)$ is nonempty.*

Proof. The set contains h, so it has maximal elements. Suppose that k has a linked pair $\pi = (s, s')$ such that s is not an input slot or s' is not an output slot, and let k^π be the barcode of Lemma 5.5; its definition, the equality $\dim(k^\pi) = n$ and the formula for its rank table proved there are purely combinatorial. The intervals of k^π are those of k, except that $I_s, I_{s'}$ are replaced by $[\![b_s, e_{s'}]\!]$ and (if $b_{s'} \leq e_s$) $[\![b_{s'}, e_s]\!]$. The first equals $[\![0, D]\!]$ only if $b_s = 0$ and $e_{s'} = D$, which is excluded; the second does not contain 0, since $b_{s'} > b_s \geq 0$. So $k^\pi \in \mathcal{Z}(n)$. By the formula of Lemma 5.5, $\rho_{ij}(k^\pi) \geq \rho_{ij}(k)$ for all $i \leq j$, with strict inequality for $(i, j) = (b_s, e_{s'})$. Hence $h \preceq k \prec k^\pi$, which contradicts the maximality of k. So every linked pair of k goes from an input slot to an output slot. An interior slot is neither, so it belongs to no linked pair, and $\mathrm{in}(s) = \mathrm{out}(s) = 0$ for every interior slot s: k is balanced. It belongs to the set minimized in $C(h)$. □

3.4 The second main theorem

Let $D \in \mathbb{N}^*$ and $n \in \mathbb{N}^{D+1}$. For $r \in \mathbb{N}$ with $r \le \min(n_0, n_D)$ put

$$c_{\mathrm{reg}}(n, r) = r\,(n_0 + n_D - r);$$

it is the dimension of the manifold of $n_D \times n_0$ real matrices of rank r.

Theorem B. *Let $D \in \mathbb{N}^*$, let $n \in \mathbb{N}^{D+1}$ be a width vector, let $\Phi \in \mathbb{R}^{n_D \times n_0}$, let $A \in F_\Phi$, and put $r = \operatorname{rk}\Phi$. Let $h_A \in \mathcal{Z}(n-r)$ be the residual barcode of A (Definition 3.10). The set minimized in the definition of $C(h_A)$ is nonempty, and*

$$\operatorname{codim}_A F_\Phi = 2\lambda_A(L_\Phi) = c_{\mathrm{reg}}(n, r) + C(h_A), \qquad m_A(L_\Phi) = 1 + F(h_A).$$

Moreover $C(h_A) = \min\{\#\operatorname{Lk}(k) :\ k \in \mathcal{Z}(n-r),\ h_A \preceq k\}$.

Since $r = R_{0D}(A)$ and $h_A = \beta(A)^\circ$ with $\rho(\beta(A)) = R(A)$, the right-hand sides depend only on n and on the rank table $R(A)$. The rank table is computable from A, and the sets involved are finite, so Theorem B gives an algorithm. For a width vector n and the rank table $R = R(A)$ of a chain $A \in \mathrm{Rep}(n)$ we put

$$\mathsf{Prog}(n, R) = \{(c_{\mathrm{reg}}(n, r) + C(h_A),\ 1 + F(h_A))\} \ \text{ if } D \ge 1, \qquad \mathsf{Prog}(n, R) = \{(0, 1)\} \ \text{ if } D = 0,$$

a one-element set, which depends only on (n, R); the value for $D = 0$ agrees with the discussion after Theorem A. Appendix A compares Prog with the formal definition.

The geometric meaning of F is given in Theorem 9.2: by Proposition 5.7, the relation $h \preceq k$ means that the orbit of $\mathbb{I}(h)$ lies in the closure of the orbit of $\mathbb{I}(k)$. Thus $F(h_A)$ counts critical strata along chains of strata of the residual zero fibre that degenerate to $\mathbb{I}(h_A)$ and along which the local codimension stays constant.

3.5 Examples

Barcodes are written as formal sums of intervals (Definition 3.5(b)).

Example 3.14 (One matrix)**.** Let $D = 1$, $n = (n_0, n_1)$, and $A = (A_0)$ with $A_0 = \Phi$ of rank r. Then $\beta(A) = r\,[\![0,1]\!] + (n_0 - r)[\![0,0]\!] + (n_1 - r)[\![1,1]\!]$ and $h_A = (n_0 - r)[\![0,0]\!] + (n_1 - r)[\![1,1]\!]$. The linked pairs of h_A are the pairs (s, s') with $I_s = [\![0,0]\!]$ and $I_{s'} = [\![1,1]\!]$, so $c_{\mathrm{out}}(h_A) = \#\operatorname{Lk}(h_A) = (n_0 - r)(n_1 - r)$. There is no interior slot, so h_A is balanced. A barcode of length 1 without full-span slot is determined by its dimension vector, so $\mathcal{Z}(n-r) = \{h_A\}$. Hence $C(h_A) = (n_0 - r)(n_1 - r)$ and $F(h_A) = 0$, and Theorem B gives $2\lambda = r(n_0 + n_1 - r) + (n_0 - r)(n_1 - r) = n_0 n_1$ and $m = 1$. This is consistent with the fact that P is the identity map of $\mathrm{Rep}(n) = \mathbb{R}^{n_1 \times n_0}$, so that $F_\Phi = \{\Phi\}$ is a point.

Example 3.15 (Scalar chains)**.** Let $n_v = 1$ for all $v \in [\![0, D]\!]$. A barcode with dimension vector $(1, \dots, 1)$ is a partition of $[\![0, D]\!]$ into consecutive intervals, and $h \preceq k$ means that every interval of h is contained in an interval of k. If $A_k \ne 0$ for all k, then $r = 1$, $\beta(A) = [\![0, D]\!]$, $h_A = 0$ is the empty barcode, and Theorem B gives $(\lambda, m) = (\frac{1}{2}, 1)$. If exactly $p \ge 1$ of the matrices vanish, say $A_{k_1} = \cdots = A_{k_p} = 0$ with $k_1 < \cdots < k_p$, then $\Phi = 0$, $r = 0$ and

$$h_A = [\![0, k_1]\!] + [\![k_1 + 1, k_2]\!] + \cdots + [\![k_p + 1, D]\!].$$

Consecutive intervals are linked and no others, so $\operatorname{Lk}(h_A)$ consists of p pairs forming a path. Every interior slot has one incoming and one outgoing linked pair, and $c_{\mathrm{out}}(h_A) = 1$. The barcodes $k \in \mathcal{Z}(n)$ with $h_A \preceq k$ are obtained by merging consecutive intervals of h_A while

keeping at least two intervals. They all have the same shape, so they are all balanced with $c_{\mathrm{out}} = 1$, all critical, and $C = 1$ on all of them. A longest sequence $h_A = h_0 \prec \cdots \prec h_\ell$ merges one pair at each step and has $\ell = p - 1$, so $F(h_A) = p - 1$ and $(\lambda, m) = (\frac{1}{2}, p)$. Directly: near A, the loss L_0 is comparable to the square of a product of p independent coordinates.

Example 3.16 (Depth two; the O77 table)**.** Let $D = 2$ and $n = (I, H, O)$; in this example only, the letters I, H, O denote these widths (input, hidden, output), and $a, b, p, q, s, \eta, c(t)$ denote the quantities defined below. Let $a = \operatorname{rk} A_0$, $b = \operatorname{rk} A_1$ and $r = \operatorname{rk} \Phi$. For $I = H = O = 2$ the program gives:

a	b	r	$\beta(A)$	$(2\lambda, m)$
0	0	0	$2[\![0,0]\!] + 2[\![1,1]\!] + 2[\![2,2]\!]$	$(3, 1)$
1	1	0	$[\![0,0]\!] + [\![0,1]\!] + [\![1,2]\!] + [\![2,2]\!]$	$(3, 1)$
2	0	0	$2[\![0,1]\!] + 2[\![2,2]\!]$	$(4, 1)$
1	1	1	$[\![0,0]\!] + [\![0,2]\!] + [\![1,1]\!] + [\![2,2]\!]$	$(4, 2)$
2	1	1	$[\![0,1]\!] + [\![0,2]\!] + [\![2,2]\!]$	$(4, 1)$
2	2	2	$2[\![0,2]\!]$	$(4, 1)$

In depth two, a closed formula for the pair was proposed by Sneiderman [26] (for the problem MAIS-O70) and by Novikov [27] (for MAIS-O77), and proved in Lean in [9]. For $I, O > 0$ and $r < H$ it reads

$$2\lambda = s + \min_{0 \le t \le \min(\eta, q)} c(t), \qquad m = \#\{t \in [\![0, \min(\eta, q)]\!] : \ c(t) = \min c\},$$

where $p = O - b$, $q = I - a$, $\eta = H + r - a - b$, $s = Oa + b(I - a)$ and $c(t) = (\eta - t)(q - t) + pt$. Since a local volume order is unique (Lemma 3.2), Theorem B and this formula agree under these hypotheses. The last row of the table, with $r = H$, is outside them: there P is a submersion at A, so F_Φ is a submanifold of codimension $O \cdot I = 4$ near A.

Example 3.17 (A squared logarithm in depth three)**.** Let $n = (2, 2, 2, 2)$ and $\beta(A) = [\![0,0]\!] + [\![0,3]\!] + [\![1,1]\!] + [\![2,2]\!] + [\![3,3]\!]$, so that $r = 1$. The residual barcode is the barcode $h_A = [\![0,0]\!] + [\![1,1]\!] + [\![2,2]\!] + [\![3,3]\!] \in \mathcal{Z}(1,1,1,1)$ of Example 3.15 with $p = 3$. Hence $2\lambda = c_{\mathrm{reg}}(n, 1) + 1 = 3 + 1 = 4$ and $m = 3$.

Example 3.18 (Multiplicity smaller than the number of top strata)**.** Let $n = (2, 2, 2, 2, 2)$, $\Phi = 0$, and let $h \in \mathcal{Z}(n)$ be the barcode with six slots $s_1, \dots, s_6$:

$$h = \underbrace{[\![0,0]\!]}_{s_1} + \underbrace{[\![0,2]\!]}_{s_2} + \underbrace{[\![1,1]\!]}_{s_3} + \underbrace{[\![2,4]\!]}_{s_4} + \underbrace{[\![3,3]\!]}_{s_5} + \underbrace{[\![4,4]\!]}_{s_6}.$$

The linked pairs are $s_1 \rightarrowtail s_3$, $s_2 \rightarrowtail s_4$, $s_2 \rightarrowtail s_5$, $s_3 \rightarrowtail s_4$ and $s_5 \rightarrowtail s_6$. Give them the coordinates x_1, x_2, x_3, x_4, x_5 respectively. With the bases ordered (s_1, s_2), (s_2, s_3), (s_2, s_4), (s_4, s_5), (s_4, s_6) at the vertices $0, \dots, 4$, the chains $\mathbb{I}(h) + \sum_\pi x_\pi J_\pi$ of the slice chart of Lemma 5.3 are

$$A_0 = \begin{pmatrix} 0 & 1 \\ x_1 & 0 \end{pmatrix}, \quad A_1 = \begin{pmatrix} 1 & 0 \\ x_2 & x_4 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & 1 \\ x_3 & 0 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 1 & 0 \\ 0 & x_5 \end{pmatrix},$$

$$\text{so that} \quad A_3 A_2 A_1 A_0 = \begin{pmatrix} x_1 x_4 & x_2 \\ 0 & x_3 x_5 \end{pmatrix}.$$

By Section 7, near $\mathbb{I}(h)$ the loss is comparable to $x_2^2 + (x_1 x_4)^2 + (x_3 x_5)^2$ in suitable coordinates, up to spectator variables. A direct computation gives volume $\asymp \varepsilon^{3/2} L(\varepsilon)^2$: the estimate $\mathrm{vol}\{(x, x') \in [-1, 1]^2 : \ |xx'| < \eta\} \asymp \eta L(\eta)$, applied to $x_1 x_4$ and to $x_3 x_5$, gives the upper

bound, and requiring each of the three terms to be $< \varepsilon/3$ gives the lower bound. So $(\lambda, m) = (\frac{3}{2}, 3)$, as Theorem B predicts.

The zero set of (x_2, x_1x_4, x_3x_5) has four irreducible components $\{x_2 = 0\} \cap \{x_1 = 0 \text{ or } x_4 = 0\} \cap \{x_3 = 0 \text{ or } x_5 = 0\}$, all of codimension 3. They are the closures of the four orbits of the barcodes $[\![0,0]\!]+[\![0,2]\!]+[\![1,4]\!]+[\![3,4]\!]$, $[\![0,0]\!]+[\![0,3]\!]+[\![1,4]\!]+[\![4,4]\!]$, $[\![0,1]\!]+[\![0,2]\!]+[\![2,4]\!]+[\![3,4]\!]$ and $[\![0,1]\!] + [\![0,3]\!] + [\![2,4]\!] + [\![4,4]\!]$. Each of these barcodes has exactly three linked pairs, all from an input slot to an output slot. So four top-dimensional strata pass through $\mathbb{I}(h)$, while $m = 3$. The logarithmic exponent $m - 1 = 2$ counts the two independent "product crossings" x_1x_4 and x_3x_5, whereas the number of components counts the 2×2 ways of resolving them.

Corollary 3.19 (Global minimum)**.** *Let $D \in \mathbb{N}$, $n \in \mathbb{N}^{D+1}$ and $\Phi \in \mathbb{R}^{n_D \times n_0}$ with $F_\Phi \neq \emptyset$. Then*

$$\min_{A \in F_\Phi} \lambda_A(L_\Phi) = \tfrac{1}{2}\big(N(n) - \dim_{\mathrm{H}} F_\Phi\big).$$

Proof. For $D = 0$ both sides are 0. For $D \geq 1$, F_Φ is a finite union of embedded submanifolds (Lemma 5.9), so by (H1) and (2.3) $\dim_{\mathrm{H}} F_\Phi = \max_{A \in F_\Phi} d_A(F_\Phi)$. Apply Theorem A at each point. □

Pepin Lehalleur and Rimányi [24] proved this global equality for the real log canonical threshold defined through zeta functions, by comparing their codimension formula with Aoyagi's computation [3].

4 Local volume orders

Throughout this section $(E, \|\cdot\|)$ is a finite-dimensional real normed vector space, $U \subseteq E$ is an open subset, $a \in U$, and $f : U \to \mathbb{R}$ is a continuous function. A *prior* is a function $\varphi : E \to \mathbb{R}$ that is continuous at a with $\varphi(a) > 0$. We put $\delta_U = \sup\{\delta > 0 : \ B_\delta(a) \subseteq U\} \in]0, +\infty]$. It is convenient to allow more general domains of integration than balls.

Definition 4.1 (Window systems)**.** A *window system at a in U* is a family $\mathcal{W} = (W_\delta)_{\delta \in]0, \delta_{\mathcal{W}}[}$ of Borel subsets of U, where $\delta_{\mathcal{W}} \in]0, +\infty]$, which *shrinks to a*: for every $\eta > 0$ there is $\theta > 0$ such that $W_\delta \subseteq B_\eta(a)$ for every $\delta \in]0, \min(\theta, \delta_{\mathcal{W}})[$. For a Haar measure μ on E, a prior φ, $\delta \in]0, \delta_{\mathcal{W}}[$ and $\varepsilon \in]0, +\infty[$ we put

$$V_{f,a}^{\mathcal{W},\mu,\varphi}(\delta, \varepsilon) = \int^{-}_{\{x \in W_\delta :\ |f(x) - f(a)| < \varepsilon\}} \varphi^+ \, d\mu \ \in [0, +\infty].$$

We say that f has *local volume order (λ, m) at a with respect to $\mathcal{W}$* (for μ and φ) if the condition of Definition 3.1 holds with $V_{f,a}^{\mu,\varphi}$ replaced by $V_{f,a}^{\mathcal{W},\mu,\varphi}$ and the condition $B_{\delta_0}(a) \subseteq U$ replaced by $\delta_0 \leq \delta_{\mathcal{W}}$. Two window systems $\mathcal{W}$ and $\mathcal{W}'$ at a in U are *equivalent* if for every $\eta > 0$ there is $\theta > 0$ such that for every $\delta \in]0, \min(\theta, \delta_{\mathcal{W}})[$ there are $\delta_1, \delta_2 \in]0, \min(\eta, \delta_{\mathcal{W}'})[$ with $W'_{\delta_1} \subseteq W_\delta \subseteq W'_{\delta_2}$, and if the same holds with $\mathcal{W}$ and $\mathcal{W}'$ exchanged.

The *ball system* $\mathcal{B} = (B_\delta(a))_{\delta \in]0, \delta_U[}$ is a window system at a in U, and $V_{f,a}^{\mathcal{B},\mu,\varphi} = V_{f,a}^{\mu,\varphi}$, so local volume orders with respect to $\mathcal{B}$ are those of Definition 3.1. The set $\{x \in W_\delta : |f(x) - f(a)| < \varepsilon\}$ is the intersection of the Borel set W_δ with an open set, hence a Borel set. The function $V_{f,a}^{\mathcal{W},\mu,\varphi}(\delta, \varepsilon)$ is nondecreasing in ε, and nondecreasing in the window: if $\mathcal{W}'$ is another window system and $W'_{\delta'} \subseteq W_\delta$, then $V_{f,a}^{\mathcal{W}',\mu,\varphi}(\delta', \varepsilon) \leq V_{f,a}^{\mathcal{W},\mu,\varphi}(\delta, \varepsilon)$ for all ε.

Lemma 4.2 (Sandwich)**.** *Let $\mathcal{W}$ and $\mathcal{W}'$ be equivalent window systems at a in U, let μ be a Haar measure on E and φ a prior. For all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, f has local volume order (λ, m) at a with respect to $\mathcal{W}$ if and only if it does with respect to $\mathcal{W}'$ (both for μ and φ).*

Proof. In this proof write $V = V_{f,a}^{\mathcal{W},\mu,\varphi}$ and $V' = V_{f,a}^{\mathcal{W}',\mu,\varphi}$. Suppose that f has local volume order (λ, m) with respect to $\mathcal{W}'$, with radius δ_0'. Apply the definition of equivalence with $\eta = \delta_0'$, and let θ be the corresponding number; put $\delta_0 = \min(\theta, \delta_{\mathcal{W}})$. Let $\delta \in]0, \delta_0[$, and choose $\delta_1, \delta_2 \in]0, \delta_0'[$ with $W'_{\delta_1} \subseteq W_\delta \subseteq W'_{\delta_2}$. Then $V'(\delta_1, \varepsilon) \le V(\delta, \varepsilon) \le V'(\delta_2, \varepsilon)$ for all $\varepsilon > 0$. Let $(c_i, C_i, \varepsilon_i)$ be the constants of (3.1) for V' at the radius δ_i $(i = 1, 2)$. Then (3.1) holds for V at the radius δ with the constants c_1, $\max(C_2, c_1)$ and $\min(\varepsilon_1, \varepsilon_2)$. The converse is the same argument with $\mathcal{W}$ and $\mathcal{W}'$ exchanged. □

Example 4.3. The following window systems at a in U are equivalent to the ball system $\mathcal{B}$ of $\|\cdot\|$.

(a) The ball system of another norm $\|\cdot\|'$ on E. Indeed, there is $K \ge 1$ with $B'_{\delta/K}(a) \subseteq B_\delta(a) \subseteq B'_{K\delta}(a)$ for all $\delta > 0$, where B' denotes the balls of $\|\cdot\|'$.

(b) When $E = E_1 \times E_2$ is the product of two finite-dimensional real normed vector spaces $(E_1, \|\cdot\|_1)$ and $(E_2, \|\cdot\|_2)$ and $a = (a_1, a_2)$: the family of products $B_\delta^{E_1}(a_1) \times B_\delta^{E_2}(a_2)$ (for small δ), since these are the balls of the norm $\max(\|x_1\|_1, \|x_2\|_2)$; then apply (a).

(c) Let $(E', \|\cdot\|')$ be a finite-dimensional real normed vector space, $U' \subseteq E'$ open, $a' \in U'$, and $\Psi : U' \to U$ a C^1 diffeomorphism onto U with $\Psi(a') = a$. Choose $r_0 > 0$ such that the closed ball $\overline{B'}_{r_0}(a')$ is contained in U'. Then the family $(\Psi(B'_\delta(a')))_{\delta \in]0, r_0[}$ is a window system at a in U equivalent to $\mathcal{B}$. Indeed, its members are open, since Ψ is a homeomorphism onto the open set U. The map Ψ is Lipschitz on the compact convex set $\overline{B'}_{r_0}(a')$, by the mean value inequality, and Ψ^{-1} is Lipschitz on a closed ball $\overline{B}_{r_1}(a) \subseteq U$; let $K \ge 1$ be a common Lipschitz constant. Then $B_{\delta/K}(a) \subseteq \Psi(B'_\delta(a')) \subseteq B_{K\delta}(a)$ for every $\delta \in]0, \min(r_0, r_1)[$.

Lemma 4.4 (Uniqueness)**.** *Let $\mathcal{W}$ be a window system at a in U, μ a Haar measure on E and φ a prior. There is at most one pair $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$ such that f has local volume order (λ, m) at a with respect to $\mathcal{W}$, for μ and φ.*

Proof. Write $V = V_{f,a}^{\mathcal{W},\mu,\varphi}$. Suppose (λ, m) and (λ', m') are both local volume orders, and fix δ smaller than both radii δ_0. Let (c, C, ε_0) and (c', C', ε_0') be the corresponding constants. For $\varepsilon < \min(\varepsilon_0, \varepsilon_0')$, writing $L = L(\varepsilon) > 1$,

$$c\,\varepsilon^\lambda L^{m-1} \le V(\delta, \varepsilon) \le C'\varepsilon^{\lambda'} L^{m'-1}.$$

Taking logarithms, $(\lambda' - \lambda)\, L \le \log(C'/c) + (m' - m)\log L$. Dividing by L and letting $\varepsilon \to 0$, so that $L \to +\infty$, gives $\lambda' \le \lambda$; by symmetry $\lambda = \lambda'$. Then $c\,L^{m-1} \le C' L^{m'-1}$ for all small ε, so $m \le m'$, and by symmetry $m = m'$. □

Lemma 4.5 (Measure and prior)**.** *Let $\mathcal{W}$ be a window system at a in U, let μ, μ' be Haar measures on E, and let φ, φ' be priors. For all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, f has local volume order (λ, m) at a with respect to $\mathcal{W}$ for (μ, φ) if and only if it does for (μ', φ').*

Proof. Write $V = V_{f,a}^{\mathcal{W},\mu,\varphi}$ and $V' = V_{f,a}^{\mathcal{W},\mu',\varphi'}$.

Step 1: the measures. There is a real number $\kappa > 0$ with $\mu' = \kappa\mu$ (Section 2.4).

Step 2: the priors near a. Since φ is continuous at a and $\varphi(a) > 0$, there is $\eta > 0$ with $\frac{1}{2}\varphi(a) \le \varphi(x) \le 2\varphi(a)$ for all $x \in B_\eta(a)$; in particular $\varphi^+ = \varphi$ on $B_\eta(a)$. Likewise there is $\eta' > 0$ with $\frac{1}{2}\varphi'(a) \le \varphi'(x) \le 2\varphi'(a)$ for all $x \in B_{\eta'}(a)$. Put $\eta'' = \min(\eta, \eta')$.

Step 3: comparison of the volume functions. Since $\mathcal{W}$ shrinks to a, there is $\theta > 0$ with $W_\delta \subseteq B_{\eta''}(a)$ for all $\delta \in]0, \min(\theta, \delta_{\mathcal{W}})[$. For such δ and every $\varepsilon > 0$, the set $S = \{x \in W_\delta : |f(x) - f(a)| < \varepsilon\}$ is contained in $B_{\eta''}(a)$, so by (2.4)

$$\tfrac{1}{2}\varphi(a)\,\mu(S) \le V(\delta,\varepsilon) \le 2\varphi(a)\,\mu(S), \qquad \tfrac{1}{2}\varphi'(a)\,\kappa\,\mu(S) \le V'(\delta,\varepsilon) \le 2\varphi'(a)\,\kappa\,\mu(S).$$

Hence, with $k_1 = \kappa\varphi'(a)/(4\varphi(a))$ and $k_2 = 4\kappa\varphi'(a)/\varphi(a)$,

$$k_1\,V(\delta,\varepsilon) \le V'(\delta,\varepsilon) \le k_2\,V(\delta,\varepsilon) \qquad \text{for all } \delta \in]0, \min(\theta, \delta_{\mathcal{W}})[\text{ and } \varepsilon > 0. \tag{4.1}$$

Step 4: the quantifiers. Suppose that f has local volume order (λ, m) for (μ, φ), with radius δ_0. Put $\delta_0' = \min(\delta_0, \theta)$. Let $\delta \in]0, \delta_0'[$, and let (c, C, ε_0) be the constants of (3.1) for V at the radius δ. By (4.1), V' satisfies (3.1) at the radius δ with the constants $(k_1 c,\ k_2 C,\ \varepsilon_0)$; note $k_1 c \le k_2 C$. So f has local volume order (λ, m) for (μ', φ'), with radius δ_0'. The converse follows by exchanging the roles of (μ, φ) and (μ', φ'). □

Lemma 4.6 (Comparison)**.** *Let $g_1, g_2 : U \to [0, +\infty[$ be continuous with $g_1(a) = g_2(a) = 0$. Suppose that there are $\eta > 0$ and real numbers $0 < \kappa_1 \le \kappa_2$ such that $B_\eta(a) \subseteq U$ and*

$$\kappa_1\,g_1(x) \le g_2(x) \le \kappa_2\,g_1(x) \qquad \text{for all } x \in B_\eta(a).$$

Let $\mathcal{W}$ be a window system at a in U, μ a Haar measure and φ a prior. For all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, g_1 has local volume order (λ, m) at a with respect to $\mathcal{W}$ if and only if g_2 does.

Proof. Write $V_i = V_{g_i,a}^{\mathcal{W},\mu,\varphi}$ for $i = 1, 2$. Let $\theta > 0$ be such that $W_\delta \subseteq B_\eta(a)$ for $\delta < \min(\theta, \delta_{\mathcal{W}})$. For such δ and every $\varepsilon > 0$,

$$\{x \in W_\delta :\ g_1(x) < \varepsilon/\kappa_2\} \subseteq \{x \in W_\delta :\ g_2(x) < \varepsilon\} \subseteq \{x \in W_\delta :\ g_1(x) < \varepsilon/\kappa_1\},$$

hence $V_1(\delta, \varepsilon/\kappa_2) \le V_2(\delta, \varepsilon) \le V_1(\delta, \varepsilon/\kappa_1)$. Suppose that g_1 has local volume order (λ, m) with radius δ_0, and put $\delta_0' = \min(\delta_0, \theta)$. Let $\delta \in]0, \delta_0'[$, let (c, C, ε_0) be the constants of (3.1) for V_1 at the radius δ, and put

$$\varepsilon_0' = \min\big(\kappa_1\varepsilon_0,\ \varepsilon_0,\ e^{-2|\log\kappa_1|},\ e^{-2|\log\kappa_2|}\big) \in]0, e^{-1}[.$$

Let $\varepsilon \in]0, \varepsilon_0'[$. Then $\varepsilon < \kappa_1\varepsilon_0 < \kappa_1 \le \kappa_2$ and $\varepsilon < 1$, and $L(\varepsilon) > 2|\log\kappa_i|$ because $\varepsilon < e^{-2|\log\kappa_i|}$ $(i = 1, 2)$; so the hypotheses of (2.2) hold for $\kappa = \kappa_1$ and $\kappa = \kappa_2$; moreover $\varepsilon/\kappa_2 \le \varepsilon/\kappa_1 < \varepsilon_0$, so the bounds (3.1) for V_1 apply at ε/κ_1 and ε/κ_2, and $\frac{1}{2}L(\varepsilon) \le L(\varepsilon/\kappa_i) \le 2L(\varepsilon)$ for $i = 1, 2$ by (2.2). Therefore

$$V_2(\delta,\varepsilon) \le C\Big(\frac{\varepsilon}{\kappa_1}\Big)^\lambda L(\varepsilon/\kappa_1)^{m-1} \le C\kappa_1^{-\lambda}2^{m-1}\,\varepsilon^\lambda L(\varepsilon)^{m-1},$$
$$V_2(\delta,\varepsilon) \ge c\Big(\frac{\varepsilon}{\kappa_2}\Big)^\lambda L(\varepsilon/\kappa_2)^{m-1} \ge c\,\kappa_2^{-\lambda}2^{1-m}\,\varepsilon^\lambda L(\varepsilon)^{m-1},$$

and $c\kappa_2^{-\lambda}2^{1-m} \le C\kappa_1^{-\lambda}2^{m-1}$. So g_2 has local volume order (λ, m), with radius δ_0'. For the converse apply the same argument to $\kappa_2^{-1}g_2 \le g_1 \le \kappa_1^{-1}g_2$. □

Lemma 4.7 (Change of variables)**.** *Let $(E', \|\cdot\|')$ be a finite-dimensional real normed vector space, $U' \subseteq E'$ an open subset, $a' \in U'$, and $\Psi : U' \to U$ a C^1 diffeomorphism from U' onto U with $\Psi(a') = a$. For all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, f has local volume order (λ, m) at a if and only if $f \circ \Psi : U' \to \mathbb{R}$ has local volume order (λ, m) at a'.*

Proof. Since Ψ is a diffeomorphism, $\dim E' = \dim E =: N$. Choose linear isomorphisms $\mathbb{R}^N \to E$ and $\mathbb{R}^N \to E'$, and let μ and μ' be the images of Lebesgue measure; we use the constant prior 1 on E and on E'. By Lemma 3.2(b) (whose proof below uses only Lemmas 4.2, 4.4 and 4.5), it suffices to prove the claim for these measures and priors. Write $J(x) = |\det d\Psi(x)|$, the absolute value of the Jacobian determinant of Ψ at $x \in U'$, computed in the chosen coordinates.

Choose $r_0 > 0$ with $\overline{B'}_{r_0}(a') \subseteq U'$. The function J is continuous and positive on this compact ball, so there is $k \geq 1$ with $k^{-1} \leq J \leq k$ on it. Let $\mathcal{W} = (\Psi(B'_\delta(a')))_{\delta \in]0,r_0[}$; by Example 4.3(c) it is a window system at a in U, equivalent to the ball system. For $\delta \in]0, r_0[$ and $\varepsilon > 0$ put $S' = \{x \in B'_\delta(a') : \ |f(\Psi(x)) - f(a)| < \varepsilon\}$, an open set; then $\Psi(S') = \{y \in \Psi(B'_\delta(a')) : \ |f(y) - f(a)| < \varepsilon\}$. The change of variables formula gives $\mu(\Psi(S')) = \int_{S'} J \, d\mu'$, so

$$k^{-1} \, V^{\mu',1}_{f\circ\Psi,a'}(\delta,\varepsilon) \ \leq \ V^{\mathcal{W},\mu,1}_{f,a}(\delta,\varepsilon) \ \leq \ k \, V^{\mu',1}_{f\circ\Psi,a'}(\delta,\varepsilon) \qquad (\delta \in]0, r_0[, \ \varepsilon > 0).$$

As in Step 4 of the proof of Lemma 4.5, the constant k is absorbed into the constants c and C: $f \circ \Psi$ has local volume order (λ, m) at a' if and only if f has local volume order (λ, m) at a with respect to $\mathcal{W}$. By Lemma 4.2, the latter holds if and only if f has local volume order (λ, m) at a. □

Lemma 4.8 (Spectator variables)**.** *Let $(E_1, \|\cdot\|_1)$ and $(E_2, \|\cdot\|_2)$ be finite-dimensional real normed vector spaces, let $U_1 \subseteq E_1$ be open, $a_1 \in U_1$, $a_2 \in E_2$, and $f_1 : U_1 \to \mathbb{R}$ continuous. Define $\tilde{f} : U_1 \times E_2 \to \mathbb{R}$ by $\tilde{f}(x_1, x_2) = f_1(x_1)$. For all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, $\tilde{f}$ has local volume order (λ, m) at (a_1, a_2) if and only if f_1 has local volume order (λ, m) at a_1.*

Proof. By Lemma 3.2(b) we may choose the norm, the measures and the priors. Give $E_1 \times E_2$ the norm $\|(x_1, x_2)\| = \max(\|x_1\|_1, \|x_2\|_2)$, whose balls are $B_\delta((a_1, a_2)) = B^{E_1}_\delta(a_1) \times B^{E_2}_\delta(a_2)$. Let μ_1, μ_2 be Haar measures on E_1, E_2; then $\mu_1 \otimes \mu_2$ is a Haar measure on $E_1 \times E_2$. Use the priors 1. For $\delta > 0$ with $B^{E_1}_\delta(a_1) \subseteq U_1$ (equivalently, $B_\delta((a_1, a_2)) \subseteq U_1 \times E_2$) and $\varepsilon > 0$,

$$\{x \in B_\delta((a_1, a_2)) : \ |\tilde{f}(x) - \tilde{f}(a_1, a_2)| < \varepsilon\} = S_{f_1,a_1}(\delta, \varepsilon) \times B^{E_2}_\delta(a_2),$$

so by the definition of the product measure

$$V^{\mu_1\otimes\mu_2,1}_{\tilde{f},(a_1,a_2)}(\delta,\varepsilon) = V^{\mu_1,1}_{f_1,a_1}(\delta,\varepsilon) \cdot \mu_2\big(B^{E_2}_\delta(a_2)\big).$$

The factor $\mu_2(B^{E_2}_\delta(a_2))$ is a positive real number that depends on δ but not on ε. Since the constants c and C in (3.1) may depend on δ, the bounds (3.1) hold for $\tilde{f}$ at the radius δ if and only if they hold for f_1 at the radius δ (with the constants multiplied, respectively divided, by this factor), and the radii δ_0 correspond. □

Lemma 4.9 (A nondegenerate quadratic block)**.** *Let $p, k \in \mathbb{N}$, let $U \subseteq \mathbb{R}^p$ be open, $a \in U$, and $g : U \to [0, +\infty[$ continuous with $g(a) = 0$. Define $G : U \times \mathbb{R}^k \to [0, +\infty[$ by $G(x, y) = g(x) + \|y\|^2$. For all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, g has local volume order (λ, m) at a if and only if G has local volume order $(\lambda + k/2, m)$ at $(a, 0)$.*

Proof. If $k = 0$ there is nothing to prove; let $k \geq 1$. We use Lebesgue measures, the priors 1, and on $\mathbb{R}^p \times \mathbb{R}^k$ the norm $\max(\|x\|, \|y\|)$, whose balls are $W_\delta = B_\delta(a) \times \{y : \ \|y\| < \delta\}$ (Lemma 3.2(b)). Let $\delta \in]0, \delta_U[$, write $v_\delta(t) = V^{\mathrm{vol},1}_{g,a}(\delta, t) = \mathrm{vol}\{x \in B_\delta(a) : \ g(x) < t\}$ for $t > 0$ (a nondecreasing function), and let ω_k be the volume of the Euclidean unit ball of $\mathbb{R}^k$. For $0 < \varepsilon < \delta^2$, the condition $\|y\|^2 < \varepsilon$ implies $\|y\| < \delta$, so Fubini's theorem gives

$$V^{\mathrm{vol},1}_{G,(a,0)}(\delta,\varepsilon) = \int_{\{y\in\mathbb{R}^k : \ \|y\|^2<\varepsilon\}} v_\delta\big(\varepsilon - \|y\|^2\big) \, dy.$$

Bounding the integrand above by $v_\delta(\varepsilon)$, and below by $v_\delta(\varepsilon/2)$ on $\{\|y\|^2 < \varepsilon/2\}$, we get, for $0 < \varepsilon < \delta^2$,

$$\omega_k(\varepsilon/2)^{k/2}\, v_\delta(\varepsilon/2) \;\le\; V^{\mathrm{vol},1}_{G,(a,0)}(\delta,\varepsilon) \;\le\; \omega_k\, \varepsilon^{k/2}\, v_\delta(\varepsilon). \tag{4.2}$$

Suppose $c\,\varepsilon^\lambda L(\varepsilon)^{m-1} \le v_\delta(\varepsilon) \le C\,\varepsilon^\lambda L(\varepsilon)^{m-1}$ for $\varepsilon \in]0,\varepsilon_0[$. For $\varepsilon < \min(\varepsilon_0, \delta^2, \frac14)$ we have $\frac12 L(\varepsilon) \le L(\varepsilon/2) \le 2L(\varepsilon)$ by (2.2), and (4.2) gives

$$\omega_k 2^{-k/2-\lambda-m+1} c\; \varepsilon^{\lambda+k/2} L(\varepsilon)^{m-1} \le V^{\mathrm{vol},1}_{G,(a,0)}(\delta,\varepsilon) \le \omega_k C\; \varepsilon^{\lambda+k/2} L(\varepsilon)^{m-1}.$$

Conversely, suppose $c\,\varepsilon^{\lambda+k/2} L(\varepsilon)^{m-1} \le V^{\mathrm{vol},1}_{G,(a,0)}(\delta,\varepsilon) \le C\,\varepsilon^{\lambda+k/2} L(\varepsilon)^{m-1}$ for $\varepsilon \in]0,\varepsilon_0[$. The right inequality of (4.2) gives $v_\delta(\varepsilon) \ge \omega_k^{-1} c\,\varepsilon^\lambda L(\varepsilon)^{m-1}$; since $v_\delta(\varepsilon) \le \operatorname{vol} B_\delta(a) < +\infty$ for all ε, this forces $\lambda \ge 0$. Next, the left inequality of (4.2) at 2ε gives $v_\delta(\varepsilon) \le \omega_k^{-1}\varepsilon^{-k/2} V^{\mathrm{vol},1}_{G,(a,0)}(\delta, 2\varepsilon) \le \omega_k^{-1} 2^{\lambda+k/2} C\,\varepsilon^\lambda L(\varepsilon)^{m-1}$, for $\varepsilon < \min(\varepsilon_0/2, \delta^2/2, \frac14)$, using $L(2\varepsilon) \le L(\varepsilon)$. In both directions the radii δ correspond, since $W_\delta \subseteq U \times \mathbb{R}^k$ if and only if $B_\delta(a) \subseteq U$. □

Lemma 4.10 (Homogeneity)**.** *Let $\Lambda : E \to E$ be a linear isomorphism, $\kappa > 0$ a real number, and $H : E \to \mathbb{R}$ a continuous function with $H(\Lambda z) = \kappa H(z)$ for all $z \in E$. Then for every $z \in E$ and all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$, H has local volume order (λ, m) at z if and only if it has local volume order (λ, m) at Λz.*

Proof. By Lemma 4.7 applied to $\Psi = \Lambda$, H has local volume order (λ, m) at Λz if and only if $H \circ \Lambda = \kappa H$ has local volume order (λ, m) at z. By Lemma 4.6, applied to $g_1 = |H - H(z)|$ and $g_2 = \kappa|H - H(z)|$, this holds if and only if H has local volume order (λ, m) at z; note that $|H(x) - H(z)| < \varepsilon$ is the condition defining the volume functions of H and of $|H - H(z)|$ alike. □

Proof of Lemma 3.2. (a) is Lemma 4.4 for the ball system. (b) By Lemma 4.5 we may replace (μ, φ) by (μ', φ'); by Example 4.3(a) and Lemma 4.2 we may replace the balls of $\|\cdot\|$ by those of $\|\cdot\|'$. (c) Suppose f is constant on $B_{\delta_1}(a) \subseteq U$. For $\delta \in]0, \delta_1[$ and every $\varepsilon > 0$, $S_{f,a}(\delta,\varepsilon) = B_\delta(a)$, so $V^{\mu,\varphi}_{f,a}(\delta,\varepsilon) = \int^-_{B_\delta(a)} \varphi^+\, d\mu$. For δ small this is a real number in $[\frac12\varphi(a)\mu(B_\delta(a)),\, 2\varphi(a)\mu(B_\delta(a))] \subseteq]0, +\infty[$ (Step 2 of the proof of Lemma 4.5), so (3.1) holds with $(\lambda, m) = (0, 1)$. □

The volume functions of f at a coincide with those of $|f - f(a)|$; this is how Lemmas 4.6 and 4.10 apply to functions that do not vanish at the base point. Below, all losses vanish at the base point.

5 Orbits of chains

Throughout this section $D \in \mathbb{N}^*$, $n \in \mathbb{N}^{D+1}$ is a width vector, and h is a barcode of length D with $\dim(h) = n$. A total order on $\mathcal{S}(h)$ is fixed, so that the interval chain $\mathbb{I}(h) \in \mathrm{Rep}(n)$ and the bases $(\epsilon^v_s)_{s \in \mathcal{S}_v(h)}$ of the spaces $\mathbb{R}^{n_v}$ are defined (Definition 3.7). We denote by

$$O_h = G(n) \cdot \mathbb{I}(h) = \{g \cdot \mathbb{I}(h) :\; g \in G(n)\}$$

the orbit of the interval chain $\mathbb{I}(h)$ under the action of the group $G(n)$. By Lemma 3.9 and Definition 3.10, O_h does not depend on the total order on $\mathcal{S}(h)$, and

$$O_h = \{A \in \mathrm{Rep}(n) :\; R(A) = \rho(h)\} = \{A \in \mathrm{Rep}(n) :\; \beta(A) = h\}. \tag{5.1}$$

5.1 Orbits are submanifolds

Let $\partial_h : \mathfrak{g}(n) \to \mathrm{Rep}(n)$ be the derivative at the identity of the orbit map $G(n) \to \mathrm{Rep}(n)$, $g \mapsto g \cdot \mathbb{I}(h)$, where $\mathfrak{g}(n)$ is the Lie algebra of $G(n)$ (Section 2.5). It is the linear map

$$(\partial_h \xi)_k = \xi_{k+1}\, \mathbb{I}(h)_k - \mathbb{I}(h)_k\, \xi_k \qquad (\xi = (\xi_v)_v \in \mathfrak{g}(n),\ k \in [\![0, D-1]\!]).$$

Its kernel is the Lie algebra of the stabilizer $\{g \in G(n) :\ g \cdot \mathbb{I}(h) = \mathbb{I}(h)\}$ of $\mathbb{I}(h)$.

Lemma 5.1. *The orbit O_h is an embedded real-analytic submanifold of* $\mathrm{Rep}(n)$*, with tangent space $\partial_h(\mathfrak{g}(n))$ at $\mathbb{I}(h)$, and* $\dim O_h = \dim \mathfrak{g}(n) - \dim \ker \partial_h$*. If $\mathfrak{g}' \subseteq \mathfrak{g}(n)$ is a linear complement of* $\ker \partial_h$*, then $\xi \mapsto (\exp \xi) \cdot \mathbb{I}(h)$ maps a neighbourhood of 0 in $\mathfrak{g}'$ diffeomorphically onto a neighbourhood of $\mathbb{I}(h)$ in O_h.*

Proof. By (5.1), O_h is the finite intersection of the sets $\{A :\ \mathrm{rk}\, P_{ij}(A) = \rho_{ij}(h)\}$. Each of them is the intersection of the open set $\{\mathrm{rk}\, P_{ij} \geq \rho_{ij}(h)\}$ with the closed set $\{\mathrm{rk}\, P_{ij} \leq \rho_{ij}(h)\}$, so O_h is locally closed in $\mathrm{Rep}(n)$, hence locally compact. The Lie group $G(n)$ has finitely many connected components and acts transitively on O_h. By [14, Chapter II, Theorem 3.2], the orbit map induces a homeomorphism from the homogeneous space of $G(n)$ modulo the stabilizer of $\mathbb{I}(h)$ onto O_h. This map is also an injective immersion [22, Chapter 21]. Hence it is a diffeomorphism onto an embedded submanifold, with the stated tangent space and dimension; the last assertion follows from the inverse function theorem. The submanifold is real-analytic because the action is algebraic. □

5.2 Linked pairs and the normal space

For a linked pair $\pi = (s, s') \in \mathrm{Lk}(h)$, let $J_\pi \in \mathrm{Rep}(n)$ be the chain whose only nonzero entry is a 1 in the matrix of the arrow $b_{s'} - 1$, in the column of $\epsilon_s^{b_{s'}-1}$ and the row of $\epsilon_{s'}^{b_{s'}}$. This makes sense: $b_s \leq b_{s'} - 1 \leq e_s$, so $s \in \mathcal{S}_{b_{s'}-1}(h)$, and $s' \in \mathcal{S}_{b_{s'}}(h)$. For $x = (x_\pi)_{\pi \in \mathrm{Lk}(h)} \in \mathbb{R}^{\mathrm{Lk}(h)}$ put $Jx = \sum_{\pi \in \mathrm{Lk}(h)} x_\pi J_\pi \in \mathrm{Rep}(n)$.

Lemma 5.2. $\mathrm{Rep}(n) = \partial_h(\mathfrak{g}(n)) \oplus \bigoplus_{\pi \in \mathrm{Lk}(h)} \mathbb{R} J_\pi$*. Consequently* $\mathrm{codim}\, O_h = \#\,\mathrm{Lk}(h)$*.*

Proof. Write elements of $\mathfrak{g}(n)$ and of $\mathrm{Rep}(n)$ in the bases (ϵ_s^v). An entry of ξ_v is indexed by a vertex v and an ordered pair of slots (s, s') with $v \in I_s \cap I_{s'}$ (column s, row s'). An entry of $Y \in \mathrm{Rep}(n)$ is indexed by an arrow k and an ordered pair of slots (s, s') with $k \in I_s$ and $k+1 \in I_{s'}$. Since every matrix $\mathbb{I}(h)_k$ maps basis vectors to basis vectors of the same slot or to 0, the map ∂_h respects the decomposition according to the pair (s, s'). On the summand of (s, s') it is the map

$$\partial_{s,s'} : \mathbb{R}^{I_s \cap I_{s'}} \to \mathbb{R}^{\mathrm{Ar}(s,s')}, \qquad (\partial_{s,s'} u)_k = u_{k+1} - u_k,$$

where $\mathrm{Ar}(s, s') = \{k \in [\![0, D-1]\!] :\ k \in I_s,\ k+1 \in I_{s'}\}$, and u_v is read as 0 for $v \notin I_s \cap I_{s'}$. Write $I_s = [\![b, e]\!]$ and $I_{s'} = [\![b', e']\!]$.

Linked pairs. If $b < b' \leq e + 1 \leq e'$, then $I_s \cap I_{s'} = [\![b', e]\!]$ (empty if $b' = e + 1$) and $\mathrm{Ar}(s, s') = [\![b' - 1, e]\!]$. In this case $\partial_{s,s'}$ is injective: the coordinate $k = b' - 1$ gives $u_{b'} = 0$, and inductively $u = 0$. Its image lies in the hyperplane $\{y :\ \sum_k y_k = 0\}$, by telescoping, and has dimension $e - b' + 1 = \#\mathrm{Ar}(s, s') - 1$. So the image is exactly this hyperplane, and J_π, whose coordinate sum is 1, spans a complement.

Non-linked pairs. We show that $\partial_{s,s'}$ is surjective. If $\mathrm{Ar}(s, s') = \emptyset$ there is nothing to prove; otherwise $b \leq e' - 1$ and $b' \leq e + 1$.

- If $b' \le b$ and $e' \le e$, then $I_s \cap I_{s'} = [\![b, e']\!]$ and $\mathrm{Ar}(s, s') = [\![b, e' - 1]\!]$. ~~The map $\partial_{s,s'}$ is the coboundary map of a path with $e' - b + 1$ vertices, which~~ For any $y \in \mathbb{R}^{\mathrm{Ar}(s,s')}$, set $u_b = 0$ and $u_{v+1} = u_v + y_v$ for $v \in [\![b, e' - 1]\!]$. Then $\partial_{s,s'} u = y$, so the map is surjective.
- If $b' \le b$ and $e' > e$, then both index sets equal $[\![b, e]\!]$, and $\partial_{s,s'}$ is triangular with diagonal entries -1, since $(\partial_{s,s'} u)_e = -u_e$.
- If $b' > b$, then non-linkedness and $b' \le e + 1$ force $e' \le e$. So $I_s \cap I_{s'} = [\![b', e']\!]$ and $\mathrm{Ar}(s, s') = [\![b' - 1, e' - 1]\!]$. Then $(\partial_{s,s'} u)_{b'-1} = u_{b'}$, and $\partial_{s,s'}$ is triangular with diagonal entries 1.

Summing over all pairs gives the decomposition, and $\operatorname{codim} O_h = \dim \mathrm{Rep}(n) - \dim \partial_h(\mathfrak{g}(n)) = \#\,\mathrm{Lk}(h)$ by Lemma 5.1. □

~~In the language of representation theory, $\partial_{s,s'}$ is the standard two-term complex computing Hom and Ext^1 from the interval representation of I_s to that of $I_{s'}$. The lemma says that Ext^1 is one-dimensional exactly for linked pairs. The equality $\operatorname{codim} O_h = \dim \mathrm{Ext}^1(\mathbb{I}(h), \mathbb{I}(h))$ is~~ The normal-space calculation above is the interval-chain instance of Voigt's lemma [36]. Its proof here uses only the displayed difference maps.

5.3 The slice chart

Fix a linear complement $\mathfrak{g}'_h$ of $\ker \partial_h$ in $\mathfrak{g}(n)$ and define

$$\Psi_h : \mathfrak{g}'_h \times \mathbb{R}^{\mathrm{Lk}(h)} \to \mathrm{Rep}(n), \qquad \Psi_h(\xi, x) = (\exp \xi) \cdot \big(\mathbb{I}(h) + Jx\big).$$

Lemma 5.3 (Slice chart). *There is an open neighbourhood U_h of $(0, 0)$ in $\mathfrak{g}'_h \times \mathbb{R}^{\mathrm{Lk}(h)}$ such that Ψ_h maps U_h real-analytically and diffeomorphically onto an open neighbourhood of $\mathbb{I}(h)$ in $\mathrm{Rep}(n)$, and $\Psi_h(U_h) \cap O_h = \Psi_h\big(U_h \cap (\mathfrak{g}'_h \times \{0\})\big)$. In particular $\mathbb{I}(h) + Jx \notin O_h$ for every $x \neq 0$ with $(0, x) \in U_h$.*

Proof. The differential $d\Psi_h$ at $(0, 0)$ is $(\xi, x) \mapsto \partial_h \xi + Jx$, which is bijective by Lemma 5.2, since ∂_h is injective on $\mathfrak{g}'_h$. The inverse function theorem gives a neighbourhood U_1 of $(0, 0)$ on which Ψ_h is a diffeomorphism onto an open set. By Lemma 5.1 there are an open neighbourhood $\mathcal{V}$ of $\mathbb{I}(h)$ in $\mathrm{Rep}(n)$ and a neighbourhood $\mathcal{U}$ of 0 in $\mathfrak{g}'_h$ with $\mathcal{U} \times \{0\} \subseteq U_1$ and $O_h \cap \mathcal{V} = \{(\exp \xi) \cdot \mathbb{I}(h) : \ \xi \in \mathcal{U}\}$. Let $U_h \subseteq U_1$ be an open neighbourhood of $(0, 0)$ with $\Psi_h(U_h) \subseteq \mathcal{V}$ and $U_h \cap (\mathfrak{g}'_h \times \{0\}) \subseteq \mathcal{U} \times \{0\}$. If $\Psi_h(\xi, x) \in O_h$ with $(\xi, x) \in U_h$, then $\Psi_h(\xi, x) = \Psi_h(\xi', 0)$ for some $\xi' \in \mathcal{U}$, and injectivity of Ψ_h on U_1 gives $x = 0$. □

Lemma 5.4 (Path expansion). *Let $x \in \mathbb{R}^{\mathrm{Lk}(h)}$, let s be an input slot and s' an output slot of h. The entry of $P(\mathbb{I}(h) + Jx)$ in the column of ϵ^0_s and the row of $\epsilon^D_{s'}$ equals*

$$\sum_{s = s_0 \rightarrowtail s_1 \rightarrowtail \cdots \rightarrowtail s_p = s'} \ \prod_{l=1}^{p} x_{(s_{l-1}, s_l)},$$

the sum over all $p \in \mathbb{N}$ and all sequences $(s_0, \ldots, s_p)$ of slots with $s_0 = s$, $s_p = s'$ and $(s_{l-1}, s_l) \in \mathrm{Lk}(h)$ for $l \in [\![1, p]\!]$. The sequence with $p = 0$ occurs only if $s = s'$, which is then a full-span slot.

Proof. At the arrow k, the matrix $(\mathbb{I}(h) + Jx)_k$ sends ϵ^k_u to ϵ^{k+1}_u if $k + 1 \in I_u$, and to 0 otherwise, plus $\sum x_{(u,u')} \epsilon^{k+1}_{u'}$ over the linked pairs (u, u') with $b_{u'} = k + 1$. Expanding the product $(\mathbb{I}(h) + Jx)_{D-1} \cdots (\mathbb{I}(h) + Jx)_0$, the entry in question is a sum over the sequences

$(u_0, \dots, u_D)$ of slots with $u_0 = s$, $u_D = s'$ and $k \in I_{u_k}$ for all k. At each arrow k, either $u_{k+1} = u_k$ (weight 1, possible only if $k+1 \in I_{u_k}$), or $u_k \rightarrowtail u_{k+1}$ with $b_{u_{k+1}} = k+1$ (weight $x_{(u_k, u_{k+1})}$). The distinct slots visited form a sequence $s = s_0 \rightarrowtail \cdots \rightarrowtail s_p = s'$ of linked pairs with $b_{s_1} < \cdots < b_{s_p}$. Conversely, such a sequence determines (u_k) uniquely: $u_k = s_l$ for $b_{s_l} \le k < b_{s_{l+1}}$, with the conventions $b_{s_0} = 0$ and $b_{s_{p+1}} = D+1$. This sequence is admissible, since $b_{s_{l+1}} \le e_{s_l} + 1$ means that s_l is present until the jump, and $e_{s'} = D$. So each sequence of linked pairs contributes exactly the product of its coordinates. □

Full-span slots. Let $r = h(\llbracket 0, D \rrbracket)$ be the number of full-span slots of h, and let h° be the zero-output part of h, so that $\mathcal{S}(h^\circ) \subseteq \mathcal{S}(h)$ consists of the slots that are not full-span and $\dim(h^\circ) = n - r$ by (3.2). Since no linked pair involves a full-span slot, $\mathrm{Lk}(h) = \mathrm{Lk}(h^\circ)$. Suppose that the full-span slots come first in the total order on $\mathcal{S}(h)$, and give $\mathcal{S}(h^\circ)$ the induced order. Then at each vertex v the first r basis vectors ϵ_s^v belong to the full-span slots, and in the decomposition $\mathbb{R}^{n_v} = \mathbb{R}^r \oplus \mathbb{R}^{n_v - r}$ given by this splitting of the basis,

$$\mathbb{I}(h) + Jx = (\mathrm{I}_r, \dots, \mathrm{I}_r) \oplus \big(\mathbb{I}(h^\circ) + J^\circ x\big), \qquad P(\mathbb{I}(h) + Jx) = \mathrm{I}_r \oplus P\big(\mathbb{I}(h^\circ) + J^\circ x\big) \tag{5.2}$$

for all $x \in \mathbb{R}^{\mathrm{Lk}(h)}$, where J° is the map J defined for the barcode h°, and $(\mathrm{I}_r, \dots, \mathrm{I}_r) \in \mathrm{Rep}(r, \dots, r)$.

5.4 Switches and the closure order

Lemma 5.5 (Switch). *Let $\pi = (s, s') \in \mathrm{Lk}(h)$, and define the barcode*

$$h^\pi = h - \mathbf{1}_{I_s} - \mathbf{1}_{I_{s'}} + \mathbf{1}_{\llbracket b_s, e_{s'} \rrbracket} + \mathbf{1}_{\llbracket b_{s'}, e_s \rrbracket},$$

where the last term is omitted if $b_{s'} = e_s + 1$. Then $\dim(h^\pi) = n$, and $\mathbb{I}(h) + \tau J_\pi \in O_{h^\pi}$ for every real number $\tau \neq 0$. Hence $O_h \subseteq \overline{O_{h^\pi}}$. Moreover, for $0 \le i \le j \le D$, $\rho_{ij}(h^\pi) = \rho_{ij}(h) + 1$ if $b_s \le i < b_{s'}$ and $e_s < j \le e_{s'}$, and $\rho_{ij}(h^\pi) = \rho_{ij}(h)$ otherwise.

Proof. Since $b_s < b_{s'}$, the intervals I_s and $I_{s'}$ are distinct and $h(I_s), h(I_{s'}) \ge 1$, so h^π has values in $\mathbb{N}$. A vertex v lies in as many of the intervals $\llbracket b_s, e_{s'} \rrbracket$, $\llbracket b_{s'}, e_s \rrbracket$ as of the intervals I_s, $I_{s'}$, because $b_s < b_{s'} \le e_s + 1 \le e_{s'}$; hence $\dim(h^\pi) = \dim(h)$. Fix $\tau \neq 0$ and consider the vectors

$$w^v = \begin{cases} \epsilon_s^v & \text{if } v \in \llbracket b_s, b_{s'} - 1 \rrbracket, \\ \epsilon_s^v + \tau \epsilon_{s'}^v & \text{if } v \in \llbracket b_{s'}, e_s \rrbracket, \\ \tau \epsilon_{s'}^v & \text{if } v \in \llbracket e_s + 1, e_{s'} \rrbracket. \end{cases}$$

Put $Y = \mathbb{I}(h) + \tau J_\pi$. We group vectors into *strings*: a string with interval $\llbracket b, c \rrbracket \in \mathrm{Int}_D$ is a family $(z^v)_{v \in \llbracket b, c \rrbracket}$ with $z^v \in \mathbb{R}^{n_v}$, such that $Y_v z^v = z^{v+1}$ for $b \le v < c$ and $Y_c z^c = 0$ if $c < D$. The following are strings:

- $(w^v)_{v \in \llbracket b_s, e_{s'} \rrbracket}$, with interval $\llbracket b_s, e_{s'} \rrbracket$: at the arrow $b_{s'} - 1$ the term τJ_π adds $\tau \epsilon_{s'}^{b_{s'}}$, and at the other arrows Y acts as $\mathbb{I}(h)$;
- $(\epsilon_s^v)_{v \in \llbracket b_{s'}, e_s \rrbracket}$, with interval $\llbracket b_{s'}, e_s \rrbracket$ (if $b_{s'} \le e_s$), since Y acts as $\mathbb{I}(h)$ at the arrows $v \ge b_{s'}$;
- $(\epsilon_t^v)_{v \in I_t}$ for every slot $t \notin \{s, s'\}$, with interval I_t.

At each vertex v, the vectors of these strings present at v form a basis of $\mathbb{R}^{n_v}$, since $\tau \neq 0$. For $i \le j$, the map $P_{ij}(Y)$ therefore sends the basis vectors at i of the strings whose interval contains $\llbracket i, j \rrbracket$ to distinct basis vectors at j, and the other basis vectors at i to 0. Hence $\mathrm{rk}\, P_{ij}(Y)$ is the number of strings whose interval contains $\llbracket i, j \rrbracket$. The intervals of the strings

are those of h^π, counted with multiplicity, so $R(Y) = \rho(h^\pi)$ and $Y \in O_{h^\pi}$ by (5.1). Letting $\tau \to 0$ gives $\mathbb{I}(h) \in \overline{O_{h^\pi}}$, hence $O_h \subseteq \overline{O_{h^\pi}}$, because $\overline{O_{h^\pi}}$ is $G(n)$-invariant. For the ranks: the interval $[\![i, j]\!]$ is contained in $[\![b_s, e_{s'}]\!]$ or $[\![b_{s'}, e_s]\!]$ as often as in I_s or $I_{s'}$, except when $b_s \le i < b_{s'}$ and $e_s < j \le e_{s'}$, in which case it is contained in exactly one of the new intervals and in neither of the old ones. □

Definition 5.6. For barcodes h, k of length D with $\dim(h) = \dim(k) = n$, we write $h \sqsubseteq k$ if $O_h \subseteq \overline{O_k}$. Equivalently, $\mathbb{I}(h) \in \overline{O_k}$, since $\overline{O_k}$ is $G(n)$-invariant.

Proposition 5.7 (Closure order)**.** *Let h, k be barcodes of length D with $\dim(h) = \dim(k) = n$.*

(a) *If $h \sqsubseteq k$, then $h \preceq k$. Hence $\sqsubseteq$ is a partial order on the barcodes of length D with dimension vector n.*

(b) *If $h \preceq k$, then $h \sqsubseteq k$.*

Proof. (a) The set $\{A \in \mathrm{Rep}(n) : \operatorname{rk} P_{ij}(A) \le \rho_{ij}(k) \text{ for all } 0 \le i \le j \le D\}$ is closed and contains O_k, hence it contains $\overline{O_k} \ni \mathbb{I}(h)$; so $\rho(h) = R(\mathbb{I}(h)) \le \rho(k)$. Reflexivity and transitivity of $\sqsubseteq$ are clear, and antisymmetry follows from that of $\preceq$ (Fact 3.8(c)).

(b) By Proposition B.1 (Appendix B), if $h \preceq k$ there is a sequence $h = h_0, h_1, \dots, h_\ell = k$ in which each h_{t+1} is obtained from h_t by a switch along a linked pair of h_t (Lemma 5.5). By Lemma 5.5, $h_t \sqsubseteq h_{t+1}$ for each t, and $\sqsubseteq$ is transitive. □

Part (b) is classical over algebraically closed fields [1]; over $\mathbb{R}$ we give a complete argument in Appendix B. The stratification of the fibres of P over $\mathbb{R}$ by the ranks of partial products, the dimensions of its strata and their closure relations are studied in detail by Shewchuk and Bhattacharya [34]. Only part (a) is used in the proof of Theorem A. Part (b) is used to express the multiplicity and the codimension through the computable order $\preceq$ in Theorem B.

Lemma 5.8 (Strict monotonicity of dimension)**.** *Let k be a barcode of length D with $\dim(k) = n$. If $h \sqsubseteq k$ and $h \ne k$, then $\dim O_h < \dim O_k$, i.e. $\#\,\mathrm{Lk}(k) < \#\,\mathrm{Lk}(h)$.*

Proof. For a real number $t > 0$ let $\Gamma(t) \in G(n)$ act at each vertex v by $\epsilon^v_s \mapsto t^{b_s}\epsilon^v_s$ $(s \in \mathcal{S}_v(h))$, and let Δ_t be the linear automorphism of $\mathbb{R}^{\mathrm{Lk}(h)}$ given by $(\Delta_t x)_\pi = t^{b_{s'} - b_s} x_\pi$ for $\pi = (s, s')$. The matrices $\mathbb{I}(h)_j$, $j \in [\![0, D-1]\!]$, join two basis vectors of the same slot, so $\Gamma(t)\cdot\mathbb{I}(h) = \mathbb{I}(h)$, and $\Gamma(t) \cdot J_\pi = t^{b_{s'} - b_s} J_\pi$. Hence $\Gamma(t) \cdot (\mathbb{I}(h) + Jx) = \mathbb{I}(h) + J\Delta_t x$, with $b_{s'} - b_s \ge 1$ for every $\pi \in \mathrm{Lk}(h)$. Since $\mathbb{I}(h) \in \overline{O_k}$ and $\Psi_h(U_h)$ is a neighbourhood of $\mathbb{I}(h)$, there is $(\xi, x) \in U_h$ with $\Psi_h(\xi, x) \in O_k$, hence $\mathbb{I}(h) + Jx = (\exp(-\xi)) \cdot \Psi_h(\xi, x) \in O_k$. Here $x \ne 0$: otherwise $\Psi_h(\xi, x) \in O_h$, and $O_h \cap O_k = \emptyset$ since $h \ne k$. Replacing x by $\Delta_t x$ for a small t (which keeps $\mathbb{I}(h) + Jx$ in O_k), we may assume that $\{(\xi', \Delta_t x) : \ \xi' \in \mathcal{U}',\ t \in]0, 1[\} \subseteq U_h$ for a small open ball $\mathcal{U}' \ni 0$ of $\mathfrak{g}'_h$. The map $(\xi', t) \mapsto \Psi_h(\xi', \Delta_t x) = (\exp \xi') \cdot \Gamma(t) \cdot (\mathbb{I}(h) + Jx)$ then takes values in O_k. It is injective and immersive, because Ψ_h is a diffeomorphism on U_h and $t \mapsto \Delta_t x$ is an injective immersion (its derivative has the coordinates $(b_{s'} - b_s) t^{b_{s'} - b_s - 1} x_\pi$, not all zero). Since O_k is an embedded submanifold (Lemma 5.1), this is an injective immersion of a manifold of dimension $\dim \mathfrak{g}'_h + 1 = \dim O_h + 1$ into O_k, so $\dim O_k \ge \dim O_h + 1$. The last assertion follows from Lemma 5.2. □

5.5 Strata of a fibre

Lemma 5.9. *Let $\Phi \in \mathbb{R}^{n_D \times n_0}$ be of rank r. Then F_Φ is the disjoint union, over the barcodes k of length D with $\dim(k) = n$ and $k([\![0, D]\!]) = r$, of the sets $F_\Phi \cap O_k$. Each of them is empty or an embedded submanifold of $\mathrm{Rep}(n)$ of dimension $\dim O_k - c_{\mathrm{reg}}(n, r)$.*

Proof. The orbits O_k are pairwise disjoint and cover $\mathrm{Rep}(n)$ by (5.1), and a point $A \in O_k$ has $\operatorname{rk} P(A) = R_{0D}(A) = k(\llbracket 0, D \rrbracket)$; this gives the decomposition. Let $M_r \subseteq \mathbb{R}^{n_D \times n_0}$ be the manifold of matrices of rank r, of dimension $c_{\mathrm{reg}}(n, r)$. The map $P|_{O_k} : O_k \to M_r$ satisfies $P(g \cdot A) = g_D P(A) g_0^{-1}$. Since $D \geq 1$, the vertices 0 and D are distinct, so $g \mapsto (g_D, g_0)$ maps $G(n)$ onto $\mathrm{GL}_{n_D}(\mathbb{R}) \times \mathrm{GL}_{n_0}(\mathbb{R})$, which acts transitively on M_r. Thus $P|_{O_k}$ is an equivariant map between homogeneous spaces of $G(n)$, so its rank is constant. It is surjective onto M_r (when $O_k \neq \emptyset$), so by Sard's theorem its rank is $\dim M_r$: it is a submersion. Its fibres are therefore embedded submanifolds of O_k of codimension $c_{\mathrm{reg}}(n, r)$. □

For $D = 0$ the lemma fails as stated: the action on the target is then by conjugation $\Phi \mapsto g_0 \Phi g_0^{-1}$, which is not transitive on M_r.

6 Reduction to the zero target

6.1 A Schur complement estimate

Let $p, q, r \in \mathbb{N}$ with $r \leq \min(p, q)$, and let $\Phi_0 = \begin{pmatrix} \mathrm{I}_r & 0 \\ 0 & 0 \end{pmatrix} \in \mathbb{R}^{p \times q}$. For $M \in \mathbb{R}^{p \times q}$ write $M = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix}$ with $M_{11} \in \mathbb{R}^{r \times r}$, $M_{12} \in \mathbb{R}^{r \times (q-r)}$, $M_{21} \in \mathbb{R}^{(p-r) \times r}$ and $M_{22} \in \mathbb{R}^{(p-r) \times (q-r)}$. On the open set of matrices M with M_{11} invertible put

$$\omega(M) = (M_{11} - \mathrm{I}_r,\ M_{12},\ M_{21}) \in \mathbb{R}^{r \times r} \times \mathbb{R}^{r \times (q-r)} \times \mathbb{R}^{(p-r) \times r},$$
$$\mathrm{Sch}(M) = M_{22} - M_{21} M_{11}^{-1} M_{12} \in \mathbb{R}^{(p-r) \times (q-r)}.$$

The space of values of ω has dimension $r^2 + r(q-r) + r(p-r) = r(p+q-r)$; we give it the Euclidean norm $\|\omega(M)\|_{\mathrm{F}} = (\|M_{11} - \mathrm{I}_r\|_{\mathrm{F}}^2 + \|M_{12}\|_{\mathrm{F}}^2 + \|M_{21}\|_{\mathrm{F}}^2)^{1/2}$.

Lemma 6.1. (a) *On a neighbourhood of Φ_0 in $\mathbb{R}^{p \times q}$,*

$$\tfrac{1}{3}\big(\|\omega(M)\|_{\mathrm{F}}^2 + \|\mathrm{Sch}(M)\|_{\mathrm{F}}^2\big) \leq \|M - \Phi_0\|_{\mathrm{F}}^2 \leq 3\big(\|\omega(M)\|_{\mathrm{F}}^2 + \|\mathrm{Sch}(M)\|_{\mathrm{F}}^2\big),$$

and $M = \Phi_0$ if and only if $\omega(M) = 0$ and $\mathrm{Sch}(M) = 0$.

(b) *There is a neighbourhood of $(\mathrm{I}_p, \mathrm{I}_q, 0)$ in $\mathrm{GL}_p(\mathbb{R}) \times \mathrm{GL}_q(\mathbb{R}) \times \mathbb{R}^{(p-r) \times (q-r)}$ on which*

$$\tfrac{1}{2}\|X\|_{\mathrm{F}} \leq \|\mathrm{Sch}\big(U(\mathrm{I}_r \oplus X)V\big)\|_{\mathrm{F}} \leq 2\|X\|_{\mathrm{F}}.$$

Proof. (a) Restrict to the matrices M with $\|M_{11} - \mathrm{I}_r\|_{\mathrm{op}} < \frac{1}{2}$ and $\|\omega(M)\|_{\mathrm{F}} \leq \frac{1}{2}$. Then $\|M_{11}^{-1}\|_{\mathrm{op}} \leq 2$, and

$$\|M_{21} M_{11}^{-1} M_{12}\|_{\mathrm{F}} \leq 2\|M_{21}\|_{\mathrm{F}} \|M_{12}\|_{\mathrm{F}} \leq \|\omega(M)\|_{\mathrm{F}}^2 \leq \tfrac{1}{2}\|\omega(M)\|_{\mathrm{F}}.$$

Since $\|M - \Phi_0\|_{\mathrm{F}}^2 = \|\omega(M)\|_{\mathrm{F}}^2 + \|M_{22}\|_{\mathrm{F}}^2$ and $M_{22} = \mathrm{Sch}(M) + M_{21} M_{11}^{-1} M_{12}$, the inequality $(\alpha + \beta)^2 \leq 2\alpha^2 + 2\beta^2$, applied to $\|M_{22}\|_{\mathrm{F}} \leq \|\mathrm{Sch}(M)\|_{\mathrm{F}} + \frac{1}{2}\|\omega(M)\|_{\mathrm{F}}$ and to $\|\mathrm{Sch}(M)\|_{\mathrm{F}} \leq \|M_{22}\|_{\mathrm{F}} + \frac{1}{2}\|\omega(M)\|_{\mathrm{F}}$, gives both bounds. The last assertion follows.

(b) For U near I_p and V near I_q we have the block factorizations

$$U = \begin{pmatrix} \mathrm{I} & 0 \\ U_{21} U_{11}^{-1} & \mathrm{I} \end{pmatrix} \begin{pmatrix} U_{11} & 0 \\ 0 & S_U \end{pmatrix} \begin{pmatrix} \mathrm{I} & y_U \\ 0 & \mathrm{I} \end{pmatrix}, \qquad V = \begin{pmatrix} \mathrm{I} & 0 \\ y_V & \mathrm{I} \end{pmatrix} \begin{pmatrix} V_{11} & 0 \\ 0 & S_V \end{pmatrix} \begin{pmatrix} \mathrm{I} & V_{11}^{-1} V_{12} \\ 0 & \mathrm{I} \end{pmatrix},$$

with $S_U = \mathrm{Sch}(U)$, $S_V = \mathrm{Sch}(V)$ (for the block sizes $(r, p-r)$ and $(r, q-r)$), $y_U = U_{11}^{-1} U_{12}$ and $y_V = V_{21} V_{11}^{-1}$. If $\mathcal{L}$ is block lower triangular and $\mathcal{R}$ is block upper triangular with

invertible diagonal blocks, and $\mathcal{M}_{11}$ is invertible, a direct computation gives $\operatorname{Sch}(\mathcal{L}\mathcal{M}\mathcal{R}) = \mathcal{L}_{22}\operatorname{Sch}(\mathcal{M})\mathcal{R}_{22}$. Applying this,

$$\operatorname{Sch}\big(U(\mathrm{I}_r \oplus X)V\big) = S_U \operatorname{Sch}(\mathcal{M})\, S_V,$$

where

$$\mathcal{M} = \begin{pmatrix} \mathrm{I} & y_U \\ 0 & \mathrm{I} \end{pmatrix}\begin{pmatrix} \mathrm{I} & 0 \\ 0 & X \end{pmatrix}\begin{pmatrix} \mathrm{I} & 0 \\ y_V & \mathrm{I} \end{pmatrix} = \begin{pmatrix} \mathrm{I} + y_U X y_V & y_U X \\ X y_V & X \end{pmatrix}.$$

By the identity $(\mathrm{I} + y_V y_U X)^{-1} = \mathrm{I} - y_V(\mathrm{I} + y_U X y_V)^{-1} y_U X$, valid for small X,

$$\operatorname{Sch}(\mathcal{M}) = X - X y_V(\mathrm{I} + y_U X y_V)^{-1} y_U X = X(\mathrm{I} + y_V y_U X)^{-1}.$$

Hence $\operatorname{Sch}(U(\mathrm{I}_r \oplus X)V) = S_U\, X\,(\mathrm{I} + y_V y_U X)^{-1} S_V$. As $(U, V, X) \to (\mathrm{I}_p, \mathrm{I}_q, 0)$, the three matrices S_U, $(\mathrm{I}+y_V y_U X)^{-1}$, S_V and their inverses tend to identity matrices; so near $(\mathrm{I}_p, \mathrm{I}_q, 0)$ they all have operator norm at most $2^{1/3}$. Since $\|MXN\|_{\mathrm{F}} \le \|M\|_{\mathrm{op}}\|X\|_{\mathrm{F}}\|N\|_{\mathrm{op}}$, this gives $\|\operatorname{Sch}(U(\mathrm{I}_r \oplus X)V)\|_{\mathrm{F}} \le 2\|X\|_{\mathrm{F}}$, and, writing $X = S_U^{-1}\operatorname{Sch}(U(\mathrm{I}_r \oplus X)V)\, S_V^{-1}(\mathrm{I}+y_V y_U X)$, also $\|X\|_{\mathrm{F}} \le 2\|\operatorname{Sch}(U(\mathrm{I}_r \oplus X)V)\|_{\mathrm{F}}$. □

6.2 The reduction

In this subsection $D \in \mathbb{N}^*$, $n \in \mathbb{N}^{D+1}$, $\Phi \in \mathbb{R}^{n_D \times n_0}$, $A \in F_\Phi$, and $r = \operatorname{rk}\Phi$. By Lemma 3.9(i), $\beta(A)$ has exactly r full-span slots, its zero-output part is the residual barcode $h_A \in \mathcal{Z}(n-r)$, and $\operatorname{Lk}(\beta(A)) = \operatorname{Lk}(h_A)$ (Section 5). We fix a total order on $\mathcal{S}(\beta(A))$ in which the r full-span slots come first, and give $\mathcal{S}(h_A) \subseteq \mathcal{S}(\beta(A))$ the induced order. This defines the interval chains $\mathbb{I}(\beta(A)) \in \operatorname{Rep}(n)$ and $\mathbb{I}(h_A) \in \operatorname{Rep}(n-r)$.

Proposition 6.2. *For all $(\lambda', m') \in \mathbb{R} \times \mathbb{N}$: if $L_{n-r,0}$ has local volume order (λ', m') at $\mathbb{I}(h_A)$, then $L_{n,\Phi}$ has local volume order $(\frac{1}{2}c_{\mathrm{reg}}(n,r) + \lambda',\, m')$ at A. Moreover*

$$\operatorname{codim}_A F_{n,\Phi} = c_{\mathrm{reg}}(n,r) + \operatorname{codim}_{\mathbb{I}(h_A)} Z(n-r).$$

Proof. Step 1: normal form. By Lemma 3.9(i) there is $g \in G(n)$ with $g \cdot A = \mathbb{I}(\beta(A))$. Put $\Phi_0 = P(\mathbb{I}(\beta(A))) = g_D \Phi g_0^{-1}$; by (5.2) (with $x = 0$), $\Phi_0 = \mathrm{I}_r \oplus 0$ in the chosen bases. The linear automorphism $T : Y \mapsto g^{-1} \cdot Y$ of $\operatorname{Rep}(n)$ maps $\mathbb{I}(\beta(A))$ to A and F_{n,Φ_0} onto $F_{n,\Phi}$, and

$$L_{n,\Phi}(T(Y)) = \tfrac{1}{2}\|g_D^{-1}(P(Y) - \Phi_0)g_0\|_{\mathrm{F}}^2 \qquad (Y \in \operatorname{Rep}(n)),$$

which lies between two positive multiples of $L_{n,\Phi_0}(Y)$, since $\|MXN\|_{\mathrm{F}} \le \|M\|_{\mathrm{op}}\|X\|_{\mathrm{F}}\|N\|_{\mathrm{op}}$. By Lemmas 4.7 and 4.6, $L_{n,\Phi}$ at A and L_{n,Φ_0} at $\mathbb{I}(\beta(A))$ have the same local volume orders, and by (H2) $\operatorname{codim}_A F_{n,\Phi} = \operatorname{codim}_{\mathbb{I}(\beta(A))} F_{n,\Phi_0}$. So it suffices to prove the proposition for the chain $\mathbb{I}(\beta(A))$ and the target Φ_0.

Step 2: slices. Let $\Psi_{\beta(A)}$ and Ψ_{h_A} be the slice charts of $\mathbb{I}(\beta(A))$ and $\mathbb{I}(h_A)$ (Lemma 5.3), with complements $\mathfrak{g}'_{\beta(A)} \subseteq \mathfrak{g}(n)$ and $\mathfrak{g}'_{h_A} \subseteq \mathfrak{g}(n-r)$ and domains $U_{\beta(A)}$ and U_{h_A}. Both have the same second factor $\mathbb{R}^{\operatorname{Lk}(h_A)}$. Put

$$X(x) = P\big(\mathbb{I}(h_A) + J^\circ x\big) \in \mathbb{R}^{(n_D-r)\times(n_0-r)} \qquad (x \in \mathbb{R}^{\operatorname{Lk}(h_A)}),$$

a polynomial map with $X(0) = 0$, where J° is the map J of h_A. By (5.2),

$$M(\xi, x) := P\big(\Psi_{\beta(A)}(\xi, x)\big) = e^{\xi_D}\big(\mathrm{I}_r \oplus X(x)\big)e^{-\xi_0} \qquad \big((\xi, x) \in \mathfrak{g}'_{\beta(A)} \times \mathbb{R}^{\operatorname{Lk}(h_A)}\big). \tag{6.1}$$

For the chain $\mathbb{I}(h_A)$ (the case $r = 0$ of (6.1)), $L_{n-r,0}(\Psi_{h_A}(\xi, x)) = \frac{1}{2}\|e^{\xi_D}X(x)e^{-\xi_0}\|_{\mathrm{F}}^2$ lies between two positive multiples of $\|X(x)\|_{\mathrm{F}}^2$ near $(0,0)$. By Lemmas 4.7, 4.6 and 4.8, $L_{n-r,0}$ at $\mathbb{I}(h_A)$ has the same local volume orders as the function

$$x \mapsto \|X(x)\|_{\mathrm{F}}^2 \quad \text{at } 0 \in \mathbb{R}^{\operatorname{Lk}(h_A)},$$

and $Z(n-r)\cap\Psi_{h_A}(U_{h_A})=\Psi_{h_A}\big((\mathfrak{g}'_{h_A}\times Y_A)\cap U_{h_A}\big)$, where $Y_A=\{x\in\mathbb{R}^{\mathrm{Lk}(h_A)}:\ X(x)=0\}$.

Step 3: Schur coordinates. Let ω be as in Lemma 6.1 for $(p,q)=(n_D,n_0)$; we identify its space of values, of dimension $c_{\mathrm{reg}}(n,r)$, with $\mathbb{R}^{c_{\mathrm{reg}}(n,r)}$ by listing the entries of the three blocks. This is an isometry for the Euclidean norm given to it before Lemma 6.1, so that $\|u\|=\|\omega(M)\|_{\mathrm{F}}$ below. Consider the map $\theta(\xi)=\omega(e^{\xi_D}\Phi_0e^{-\xi_0})$, defined on the neighbourhood of 0 in $\mathfrak{g}'_{\beta(A)}$ where the upper left $r\times r$ block of $e^{\xi_D}\Phi_0e^{-\xi_0}$ is invertible, with values in $\mathbb{R}^{c_{\mathrm{reg}}(n,r)}$. Its differential at 0 is $\xi\mapsto\omega'(\xi_D\Phi_0-\Phi_0\xi_0)$, where $\omega'(M)=(M_{11},M_{12},M_{21})$ is the linear part of ω. Writing $\xi_D=\begin{pmatrix}\alpha_1&\alpha_2\\ \alpha_3&\alpha_4\end{pmatrix}$ and $\xi_0=\begin{pmatrix}\gamma_1&\gamma_2\\ \gamma_3&\gamma_4\end{pmatrix}$ in blocks of sizes (r,n_D-r) and (r,n_0-r), we get $\omega'(\xi_D\Phi_0-\Phi_0\xi_0)=(\alpha_1-\gamma_1,\,-\gamma_2,\,\alpha_3)$. On all of $\mathfrak{g}(n)$ this linear map is onto $\mathbb{R}^{c_{\mathrm{reg}}(n,r)}$. It vanishes on $\ker\partial_{\beta(A)}$: indeed $\xi_D\Phi_0-\Phi_0\xi_0$ is the derivative at $t=0$ of $t\mapsto P(\exp(t\xi)\cdot\mathbb{I}(\beta(A)))$, so it equals $dP(\partial_{\beta(A)}\xi)$, where dP is the differential of P at $\mathbb{I}(\beta(A))$. Since $\mathfrak{g}(n)=\mathfrak{g}'_{\beta(A)}\oplus\ker\partial_{\beta(A)}$, the map is onto already on $\mathfrak{g}'_{\beta(A)}$. Choose a decomposition $\mathfrak{g}'_{\beta(A)}=\mathfrak{g}'_1\oplus\mathfrak{g}'_2$ such that $d\theta_0$ maps $\mathfrak{g}'_1$ isomorphically onto $\mathbb{R}^{c_{\mathrm{reg}}(n,r)}$ and vanishes on $\mathfrak{g}'_2$, and define

$$\Theta(\xi,x)=\big(\omega(M(\xi,x)),\ \xi_2,\ x\big),\qquad \xi=\xi_1+\xi_2\text{ with }\xi_1\in\mathfrak{g}'_1,\ \xi_2\in\mathfrak{g}'_2.$$

We have $\omega(M(0,x))=\omega(\mathrm{I}_r\oplus X(x))=0$ for all x, and $\omega(M(\xi,0))=\theta(\xi)$. Hence $d\Theta_{(0,0)}(\xi,x)=(d\theta_0(\xi_1),\xi_2,x)$ is bijective, and Θ is a diffeomorphism from a neighbourhood of $(0,0)$ onto a neighbourhood of 0. In the coordinates $(u,\xi_2,x)=\Theta(\xi,x)$, Lemma 6.1 and (6.1) give, near 0,

$$\tfrac{1}{24}\big(\|u\|^2+\|X(x)\|_{\mathrm{F}}^2\big)\le L_{n,\Phi_0}\big(\Psi_{\beta(A)}(\Theta^{-1}(u,\xi_2,x))\big)\le 6\big(\|u\|^2+\|X(x)\|_{\mathrm{F}}^2\big):\tag{6.2}$$

by Lemma 6.1(a), $L_{n,\Phi_0}=\frac12\|M-\Phi_0\|_{\mathrm{F}}^2$ is between $\frac16$ and $\frac32$ times $\|\omega(M)\|_{\mathrm{F}}^2+\|\mathrm{Sch}(M)\|_{\mathrm{F}}^2$, and by Lemma 6.1(b) with $U=e^{\xi_D}$, $V=e^{-\xi_0}$, $\frac14\|X(x)\|_{\mathrm{F}}^2\le\|\mathrm{Sch}(M)\|_{\mathrm{F}}^2\le4\|X(x)\|_{\mathrm{F}}^2$. Moreover, for a small open neighbourhood U_* of $(0,0)$,

$$F_{n,\Phi_0}\cap\Psi_{\beta(A)}(U_*)=\Psi_{\beta(A)}\Theta^{-1}\big(\{0\}\times\mathfrak{g}'_2\times Y_A\big)\cap\Psi_{\beta(A)}(U_*).$$

Step 4: conclusion. By (6.2) and Lemmas 4.7, 4.6, 4.8 and 4.9 (applied with $k=c_{\mathrm{reg}}(n,r)$), L_{n,Φ_0} has local volume order $(\frac12c_{\mathrm{reg}}(n,r)+\lambda',m')$ at $\mathbb{I}(\beta(A))$ whenever $x\mapsto\|X(x)\|_{\mathrm{F}}^2$ has local volume order (λ',m') at 0. By Step 2, this is the case if $L_{n-r,0}$ has local volume order (λ',m') at $\mathbb{I}(h_A)$. For the codimension, by (H2) and the product formula (H4) below,

$$d_{\mathbb{I}(\beta(A))}(F_{n,\Phi_0})=\dim\mathfrak{g}'_2+d_0(Y_A),\qquad d_{\mathbb{I}(h_A)}(Z(n-r))=\dim\mathfrak{g}'_{h_A}+d_0(Y_A).$$

By Lemma 5.2, $N(n)=\dim\mathfrak{g}'_{\beta(A)}+\#\,\mathrm{Lk}(h_A)=c_{\mathrm{reg}}(n,r)+\dim\mathfrak{g}'_2+\#\,\mathrm{Lk}(h_A)$ and $N(n-r)=\dim\mathfrak{g}'_{h_A}+\#\,\mathrm{Lk}(h_A)$. Hence

$$\mathrm{codim}_{\mathbb{I}(\beta(A))}F_{n,\Phi_0}=c_{\mathrm{reg}}(n,r)+\#\,\mathrm{Lk}(h_A)-d_0(Y_A)=c_{\mathrm{reg}}(n,r)+\mathrm{codim}_{\mathbb{I}(h_A)}Z(n-r).\qquad\square$$

The product formula used above is the following standard fact [8, Product formulae 7.2 and 7.3].

(H4) For every nonempty Borel set $S\subseteq\mathbb{R}^b$ and every nonempty open ball (or open cube) $Q\subseteq\mathbb{R}^a$, $\dim_{\mathrm{H}}(Q\times S)=a+\dim_{\mathrm{H}}S$.

Consequently, for a closed set $S\subseteq\mathbb{R}^b$ and a point $y\in S$, the local Hausdorff dimension at $(0,y)$ of $\mathbb{R}^a\times S$ is $a+d_y(S)$ (the sets $S\cap B_\delta(y)$ are nonempty Borel sets), and the local Hausdorff dimension at $(0,y)$ of $\{0\}\times S$ is $d_y(S)$. In Step 4 these facts are applied to the closed set $S=Y_A$ and the point $y=0\in Y_A$, after the changes of coordinates given by the charts, which are bi-Lipschitz on compact neighbourhoods (see (H2)).

7 The zero target: a recursion over orbits

Throughout this section $D \in \mathbb{N}^*$ and $n \in \mathbb{N}^{D+1}$ is a width vector. Recall the function $K_n = \|P\|_{\mathrm{F}}^2 = 2L_{n,0}$ on $\mathrm{Rep}(n)$ (Section 2.5); by Lemma 4.6, K_n and $L_{n,0}$ have the same local volume orders. The zero fibre $Z(n)$ is the disjoint union of the orbits O_h, $h \in \mathcal{Z}(n)$, by (5.1) and Lemma 3.9(i): a chain A with $P(A) = 0$ has a barcode $\beta(A)$ without full-span slot. Since $K_n(g \cdot Y) = \|g_D P(Y) g_0^{-1}\|_{\mathrm{F}}^2$ lies between two positive multiples of $K_n(Y)$, Lemmas 4.7 and 4.6 show that K_n has the same local volume orders at all points of an orbit. For $h \in \mathcal{Z}(n)$ we write (λ_h, m_h) for the local volume order of K_n at the points of O_h, when it exists.

In the rest of this section we fix $h \in \mathcal{Z}(n)$, a total order on $\mathcal{S}(h)$, and the slice chart Ψ_h of Lemma 5.3, with domain U_h. Since h is zero-output, every slot of h is of exactly one of the kinds input, output, interior. A linked pair $(s, s') \in \mathrm{Lk}(h)$ is *direct* if s is an input slot and s' an output slot, and *residual* otherwise. We write $\mathrm{Lk}(h) = \mathrm{Lk}_{\mathrm{dir}}(h) \sqcup \mathrm{Lk}_{\mathrm{res}}(h)$ accordingly, and $x = (y, z)$ for the corresponding decomposition of $x \in \mathbb{R}^{\mathrm{Lk}(h)} = \mathbb{R}^{\mathrm{Lk}_{\mathrm{dir}}(h)} \times \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}$. Put $q_h = \frac{1}{2}\#\,\mathrm{Lk}_{\mathrm{dir}}(h)$.

7.1 The shear

The rows of $P(\mathbb{I}(h) + Jx)$ are indexed by the output slots (the slots present at the vertex D) and its columns by the input slots (the slots present at the vertex 0).

Lemma 7.1. *For an input slot s and an output slot s' of h,*

$$P(\mathbb{I}(h) + Jx)_{s',s} = [s \rightarrowtail s']\, y_{(s,s')} + Q_{s',s}(z) \qquad \big(x = (y, z) \in \mathbb{R}^{\mathrm{Lk}(h)}\big),$$

where $[s \rightarrowtail s']$ is 1 if $(s, s') \in \mathrm{Lk}(h)$ and 0 otherwise, and $Q_{s',s}$ is a polynomial in the residual coordinates z only: the sum in Lemma 5.4 over the sequences of length $p \geq 2$. Consequently the map $\mathrm{Sh}_h(y, z) = (y + Q_{\mathrm{dir}}(z), z)$, where $Q_{\mathrm{dir}}(z) = (Q_{s',s}(z))_{(s,s') \in \mathrm{Lk}_{\mathrm{dir}}(h)}$, is a polynomial automorphism of $\mathbb{R}^{\mathrm{Lk}(h)}$ with polynomial inverse and Jacobian determinant 1, and

$$K_n(\mathbb{I}(h) + Jx) = \|y + Q_{\mathrm{dir}}(z)\|^2 + H_h(z), \qquad H_h(z) = \sum_{(s,s')} Q_{s',s}(z)^2,$$

the last sum being over the pairs (input slot s, output slot s') that are not linked.

Proof. There are no full-span slots, so the term $p = 0$ of Lemma 5.4 does not occur, and the term $p = 1$ is the coordinate of the linked pair (s, s') if there is one. In a sequence $s = s_0 \rightarrowtail s_1 \rightarrowtail \cdots \rightarrowtail s_p = s'$ with $p \geq 2$, the slots $s_1, \ldots, s_{p-1}$ are interior: they are not input slots because $b_{s_l} > b_s \geq 0$, and not output slots because an output slot is never the first slot of a linked pair. So every pair of such a sequence starts or ends at an interior slot, i.e. is residual. The rest follows by expanding $\|P(\mathbb{I}(h) + Jx)\|_{\mathrm{F}}^2$ entry by entry. □

By Lemma 5.3, $K_n(\Psi_h(\xi, x)) = \|e^{\xi_D} P(\mathbb{I}(h) + Jx) e^{-\xi_0}\|_{\mathrm{F}}^2$, which lies between two positive multiples of $K_n(\mathbb{I}(h) + Jx)$ for (ξ, x) near $(0, 0)$. Applying Lemmas 4.7, 4.6, 4.8 and 4.9 at a point $(0, \mathrm{Sh}_h^{-1}(0, z_0))$ of the chart, we obtain:

Corollary 7.2. *Let $z_0 \in \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}$ with $H_h(z_0) = 0$, and put*

$$p_h(z_0) = \mathbb{I}(h) + J\,\mathrm{Sh}_h^{-1}(0, z_0) = \mathbb{I}(h) + J\big(-Q_{\mathrm{dir}}(z_0), z_0\big).$$

Then $p_h(z_0) \in Z(n)$. If z_0 is small enough that $(0, \mathrm{Sh}_h^{-1}(0, z_0)) \in U_h$, then for all $(\lambda, m) \in \mathbb{R} \times \mathbb{N}$: K_n has local volume order (λ, m) at $p_h(z_0)$ if and only if H_h has local volume order $(\lambda - q_h, m)$ at z_0. This includes $z_0 = 0$, for which $p_h(0) = \mathbb{I}(h)$.

Here the case $\mathrm{Lk}_{\mathrm{res}}(h) = \emptyset$ is included: then H_h is the zero function on the point $\mathbb{R}^0$, with local volume order $(0, 1)$.

7.2 Potentials and a dilation

Definition 7.3. A (rational) *admissible potential* for h is a function $u : \mathcal{S}(h) \to \mathbb{Q}$ with $u_s = 0$ for the input slots s, $u_s = 1$ for the output slots s, and $u_s < u_{s'}$ for every linked pair $(s, s') \in \mathrm{Lk}(h)$. Its *threshold* is

$$T(u) = \tfrac{1}{2} \sum_{(s,s') \in \mathrm{Lk}(h)} (u_{s'} - u_s).$$

Let $\nu(b, e) = b/(b + D - e)$ for integers $0 \le b \le e \le D$ with $(b, e) \neq (0, D)$. The *canonical potential* is $u_s = \nu(b_s, e_s)$; its denominator is positive because h has no full-span slot. It is admissible: it is 0 on input slots and 1 on output slots, and the function ν is nondecreasing in b, and nondecreasing in e, strictly when $b > 0$. If $(s, s') \in \mathrm{Lk}(h)$ and $b_s > 0$, then $u_s = \nu(b_s, e_s) < \nu(b_s, e_{s'}) \le \nu(b_{s'}, e_{s'}) = u_{s'}$, since $e_s < e_{s'}$ and $b_s < b_{s'}$; if $b_s = 0$, then $u_s = 0 < u_{s'}$ because $b_{s'} > 0$. The canonical potential is the one used in the formal definition of the program (Appendix A).

Lemma 7.4 (Dilation)**.** *Let u be an admissible potential for h, and $M \in \mathbb{N}^*$ with $Mu_s \in \mathbb{Z}$ for all $s \in \mathcal{S}(h)$. For $\pi = (s, s') \in \mathrm{Lk}(h)$ put $w_\pi = M(u_{s'} - u_s) \in \mathbb{N}^*$. For a real number $r > 0$, let $\Gamma(r) \in G(n)$ act at each vertex v by $\epsilon^v_s \mapsto r^{Mu_s} \epsilon^v_s$ ($s \in \mathcal{S}_v(h)$), and let Δ_r be the linear automorphism of $\mathbb{R}^{\mathrm{Lk}(h)}$ with $(\Delta_r x)_\pi = r^{w_\pi} x_\pi$. Then for all $x \in \mathbb{R}^{\mathrm{Lk}(h)}$ and $z \in \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}$:*

$$\Gamma(r) \cdot (\mathbb{I}(h) + Jx) = \mathbb{I}(h) + J\Delta_r x, \qquad P(\mathbb{I}(h) + J\Delta_r x) = r^M P(\mathbb{I}(h) + Jx),$$
$$Q_{\mathrm{dir}}(\Delta_r z) = r^M Q_{\mathrm{dir}}(z), \qquad H_h(\Delta_r z) = r^{2M} H_h(z).$$

Here Δ_r also denotes its restriction to the coordinate subspace $\mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}$, which it preserves.

Proof. The first identity is checked arrow by arrow, as in the proof of Lemma 5.8: $\Gamma(r) \cdot \mathbb{I}(h) = \mathbb{I}(h)$, and $\Gamma(r) \cdot J_\pi = r^{w_\pi} J_\pi$. Every slot present at the vertex 0 is an input slot and every slot present at the vertex D is an output slot, so $\Gamma(r)_0 = \mathrm{I}$ and $\Gamma(r)_D = r^M \mathrm{I}$. Hence $P(\Gamma(r) \cdot Y) = r^M P(Y)$ for every chain Y, which gives the second identity. Direct pairs have weight $w_\pi = M$. Comparing the entries of Lemma 7.1 on both sides of the second identity, as polynomial identities in x, gives the other two. □

Lemma 7.5 (Zeros of H_h and higher orbits)**.** (a) *For every $z \neq 0$ with $H_h(z) = 0$, the point $p_h(z)$ of Corollary 7.2 lies in an orbit O_k with $k \in \mathcal{Z}(n)$, $h \sqsubseteq k$ and $k \neq h$, and so do the points $p_h(\Delta_r z)$ for all $r > 0$.*

(b) *Conversely, if $k \in \mathcal{Z}(n)$, $h \sqsubseteq k$ and $k \neq h$, then for every $\eta > 0$ there is $z \in \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}$ with $0 < \|z\| < \eta$, $H_h(z) = 0$ and $p_h(z) \in O_k$.*

Proof. (a) $p_h(z) \in Z(n)$ by Corollary 7.2, so $p_h(z) \in O_k$ for some $k \in \mathcal{Z}(n)$. By Lemma 7.4, $\Gamma(r) \cdot p_h(z) = p_h(\Delta_r z)$, so all the points $p_h(\Delta_r z)$ lie in O_k, and $p_h(\Delta_r z) \to \mathbb{I}(h)$ as $r \to 0$. Hence $\mathbb{I}(h) \in \overline{O_k}$, i.e. $h \sqsubseteq k$. Finally $k \neq h$: for small r, the point $p_h(\Delta_r z) = \mathbb{I}(h) + Jx$ has $x \neq 0$ small, so it is not in O_h by Lemma 5.3.

(b) Since $\mathbb{I}(h) \in \overline{O_k}$ and $\Psi_h(U_h)$ is a neighbourhood of $\mathbb{I}(h)$, there are points $\Psi_h(\xi, x) \in O_k$ with $(\xi, x) \in U_h$ arbitrarily small. Then $\mathbb{I}(h) + Jx = (\exp(-\xi)) \cdot \Psi_h(\xi, x) \in O_k \subseteq Z(n)$. By Lemma 7.1, $K_n(\mathbb{I}(h) + Jx) = 0$ forces $y = -Q_{\mathrm{dir}}(z)$ and $H_h(z) = 0$, so $\mathbb{I}(h) + Jx = p_h(z)$. If $z = 0$, then $x = 0$ and $p_h(z) = \mathbb{I}(h) \notin O_k$; hence $z \neq 0$. □

7.3 The recursion

Proposition 7.6. *Suppose that K_n has a local volume order (λ_k, m_k) along every orbit O_k with $k \in \mathcal{Z}(n)$, $h \sqsubseteq k$ and $k \neq h$. Put*

$$a_* = \min\{\lambda_k : \ k \in \mathcal{Z}(n),\ h \sqsubseteq k,\ k \neq h\},$$
$$m_* = \max\{m_k : \ k \in \mathcal{Z}(n),\ h \sqsubseteq k,\ k \neq h,\ \lambda_k = a_*\},$$

with $a_* = +\infty$ (and m_* undefined) if there is no such k. Then for every admissible potential u for h, K_n has at $\mathbb{I}(h)$ the local volume order (λ_h, m_h) given by

$$\lambda_h = \min\big(T(u), a_*\big), \qquad m_h = \begin{cases} 1 & \text{if } T(u) < a_*, \\ m_* & \text{if } T(u) > a_*, \\ m_* + 1 & \text{if } T(u) = a_*. \end{cases}$$

Proof. Fix u, an integer M as in Lemma 7.4, the weights w_π and the dilations Δ_r. Each direct pair contributes $\frac{1}{2}$ to $T(u)$, so

$$T(u) = q_h + T_H, \qquad T_H = \tfrac{1}{2} \sum_{(s,s') \in \mathrm{Lk}_{\mathrm{res}}(h)} (u_{s'} - u_s) = \frac{|w|}{2M}, \quad \text{where } |w| = \sum_{\pi \in \mathrm{Lk}_{\mathrm{res}}(h)} w_\pi.$$

By Corollary 7.2 with $z_0 = 0$, it suffices to prove that H_h has at $0 \in \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}$ the local volume order $(\min(T_H, a_* - q_h), m_h)$, with m_h as in the statement. If $\mathrm{Lk}_{\mathrm{res}}(h) = \emptyset$, then H_h is the zero function on a point, with order $(0, 1)$; in that case Lemma 7.5(b) shows that no k as in the statement exists, so $a_* = +\infty$, and the claim reads $(T_H, 1) = (0, 1)$. Assume from now on that $d := \#\,\mathrm{Lk}_{\mathrm{res}}(h) \geq 1$, and put $a = a_* - q_h \in]-\infty, +\infty]$.

Step 1: weighted polar coordinates. Let $M_0 \in \mathbb{N}^*$ be a common multiple of the weights w_π, $\pi \in \mathrm{Lk}_{\mathrm{res}}(h)$, and

$$\varrho(z) = \sum_{\pi \in \mathrm{Lk}_{\mathrm{res}}(h)} z_\pi^{2M_0/w_\pi} \qquad \big(z \in \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)}\big).$$

This is a polynomial, positive on $\mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)} \setminus \{0\}$, with $\varrho(\Delta_r z) = r^{2M_0} \varrho(z)$. The set $\mathbb{S} = \varrho^{-1}(1)$ is compact. The vector field $\mathcal{E}(z) = (w_\pi z_\pi)_\pi$ generates the dilations ($\frac{d}{dr} \Delta_r z = r^{-1} \mathcal{E}(\Delta_r z)$), and Euler's identity gives $d\varrho_z(\mathcal{E}(z)) = 2M_0 \varrho(z)$. Hence $\mathbb{S}$ is a smooth hypersurface transverse to $\mathcal{E}$, and

$$\Pi :]0, +\infty[\times \mathbb{S} \to \mathbb{R}^{\mathrm{Lk}_{\mathrm{res}}(h)} \setminus \{0\}, \qquad (r, \vartheta) \mapsto \Delta_r \vartheta,$$

is a diffeomorphism, with inverse $z \mapsto (\varrho(z)^{1/2M_0}, \Delta_{\varrho(z)^{-1/2M_0}} z)$. For a Borel set $S \subseteq \mathbb{S}$ put $\sigma(S) = \mathrm{vol}\, \Pi(]0, 1[\times S)$. Since Δ_r is linear with determinant $r^{|w|}$, we get $\mathrm{vol}\, \Pi(]0, r_0[\times S) = r_0^{|w|} \sigma(S)$ for $r_0 > 0$. Hence Π^{-1} carries Lebesgue measure to the product of the measure $|w|\, r^{|w|-1} dr$ on $]0, +\infty[$ with σ, by uniqueness of measures that agree on products. In a chart $\chi : V_\chi \to \mathbb{S}$ ($V_\chi \subseteq \mathbb{R}^{d-1}$ open), the map $(r, v) \mapsto \Delta_r \chi(v)$ has Jacobian determinant $r^{|w|-1} j(v)$, with $j(v) = \det(\mathcal{E}(\chi(v)), d\chi(v))$ continuous and nowhere zero (by transversality). Therefore $\sigma = |w|^{-1} |j(v)|\, dv$ on $\chi(V_\chi)$: the measure σ is finite and has a continuous positive density in every chart.

Step 2: the angular function. For $\eta > 0$ put $\Xi(\eta) = \sigma\{\vartheta \in \mathbb{S} : \ H_h(\vartheta) < \eta\}$. For $r_0 > 0$ consider the window $W_{r_0} = \{z : \ \varrho(z) < r_0^{2M_0}\} = \Pi(]0, r_0[\times \mathbb{S}) \cup \{0\}$. The polar decomposition and $H_h(\Delta_r \vartheta) = r^{2M} H_h(\vartheta)$ give

$$V_{r_0}(\varepsilon) := \mathrm{vol}\{z \in W_{r_0} : \ H_h(z) < \varepsilon\} = |w| \int_0^{r_0} r^{|w|-1}\, \Xi(\varepsilon r^{-2M})\, dr. \tag{7.1}$$

Step 3: angular asymptotics. Let $Z_{\mathbb{S}} = \{\vartheta \in \mathbb{S} :\ H_h(\vartheta) = 0\}$, a compact set.

Case $Z_{\mathbb{S}} = \emptyset$. Then $0 < \eta_1 \le H_h \le \eta_2$ on $\mathbb{S}$ for some real numbers η_1, η_2, so $\Xi(\eta) = 0$ for $\eta \le \eta_1$ and $\Xi(\eta) = \sigma(\mathbb{S})$ for $\eta > \eta_2$. No k as in the statement exists, so $a_* = +\infty$: otherwise Lemma 7.5(b) would give a zero $z \neq 0$ of H_h, and then $\Delta_{\varrho(z)^{-1/2M_0} z} \in Z_{\mathbb{S}}$ by Lemma 7.4.

Case $Z_{\mathbb{S}} \neq \emptyset$. Let $\vartheta \in Z_{\mathbb{S}}$. For small $r > 0$, $\Delta_r \vartheta$ is a small zero of H_h, and $p_h(\Delta_r \vartheta) \in O_k$ for some $k \in \mathcal{Z}(n)$ with $h \sqsubseteq k \neq h$, independent of r (Lemma 7.5(a)). By Corollary 7.2 and Lemma 4.10 (applied to the linear map Δ_r and the function H_h, with the constant r^{2M}), H_h has local volume order $(\lambda_k - q_h, m_k)$ at $\Delta_r \vartheta$, hence at ϑ. Near ϑ, use the coordinates $(r, v) \mapsto \Delta_r \chi(v)$ around $(1, v_\vartheta)$, where $\chi(v_\vartheta) = \vartheta$. In them $H_h(\Delta_r \chi(v)) = r^{2M} H_h(\chi(v))$, which lies between two positive multiples of $H_h(\chi(v))$ for $r \in]\frac{1}{2}, 2[$. By Lemmas 4.7, 4.6 and 4.8, the function $v \mapsto H_h(\chi(v))$ has local volume order $(\lambda_k - q_h, m_k)$ at v_ϑ. By Lemma 4.5, this holds for the measure σ (transported by χ) as well, since σ has a continuous positive density.

By compactness, finitely many chart balls $U_1, \dots, U_N$ cover $Z_{\mathbb{S}}$, each centred at a zero ϑ_i with associated barcode k_i, and each of radius below the radius δ_0 of the corresponding local volume order. On the compact set $\mathbb{S} \setminus \bigcup_i U_i$ we have $H_h \ge \eta_3$ for some $\eta_3 > 0$. Hence for $\eta < \eta_3$

$$\Xi(\eta) \le \sum_{i=1}^{N} \sigma\{\vartheta \in U_i :\ H_h(\vartheta) < \eta\} \le C\, \eta^a L(\eta)^{m_*-1} \qquad (\eta \text{ small}),$$

~~because~~ Indeed, $\lambda_{k_i} - q_h \ge a$~~, and~~. If $\lambda_{k_i} - q_h = a$, then $m_{k_i} \le m_*$ ~~when $\lambda_{k_i} - q_h = a.$~~ and $L(\eta) > 1$ for $\eta < e^{-1}$. If $\lambda_{k_i} - q_h > a$, then

$$\frac{\eta^{\lambda_{k_i} - q_h} L(\eta)^{m_{k_i}-1}}{\eta^a L(\eta)^{m_*-1}} = \eta^{\lambda_{k_i} - q_h - a} L(\eta)^{m_{k_i} - m_*} \longrightarrow 0 \qquad (\eta \to 0).$$

Since the cover is finite, a single positive constant and a single positive cutoff suffice for the displayed upper bound. Conversely, by Lemma 7.5(b) and Lemma 7.4, every orbit O_k with $k \in \mathcal{Z}(n)$, $h \sqsubseteq k \neq h$, contains $p_h(\vartheta)$ for some $\vartheta \in Z_{\mathbb{S}}$: take z from Lemma 7.5(b) and $\vartheta = \Delta_{\varrho(z)^{-1/2M_0} z}$. The local volume order at that ϑ has exponent $\lambda_k - q_h \ge 0$, by Definition 3.1. This applies to every k in the minimum defining a_*; in particular $a = a_* - q_h \ge 0$. Choosing k with $(\lambda_k, m_k) = (a_*, m_*)$ and a small chart ball around the corresponding ϑ gives the matching lower bound. Thus there is $\eta_0 \in]0, e^{-1}[$ with

$$\Xi(\eta) \asymp \eta^a L(\eta)^{m_*-1} \ \text{ on }]0, \eta_0[, \qquad \Xi(\eta_0) \le \Xi(\eta) \le \sigma(\mathbb{S}) \ \text{ for } \eta \ge \eta_0,$$

and $\Xi(\eta_0) > 0$.

Step 4: radial integration. First suppose $Z_{\mathbb{S}} \neq \emptyset$. Fix $r_0 > 0$ and split (7.1) at $r_\varepsilon = (\varepsilon/\eta_0)^{1/2M}$, which is $< r_0$ for small ε. For $r < r_\varepsilon$ we have $\varepsilon r^{-2M} > \eta_0$, so the inner part satisfies

$$|w| \int_0^{r_\varepsilon} r^{|w|-1}\, \Xi(\varepsilon r^{-2M})\, dr \asymp r_\varepsilon^{|w|} \asymp \varepsilon^{T_H}.$$

In the outer part substitute $r = (\varepsilon e^\tau)^{1/2M}$, so that $\varepsilon r^{-2M} = e^{-\tau}$ and $dr = r\, d\tau / 2M$:

$$|w| \int_{r_\varepsilon}^{r_0} r^{|w|-1}\, \Xi(\varepsilon r^{-2M})\, dr \asymp \varepsilon^{T_H} \int_{\tau_0}^{\tau_\varepsilon} e^{(T_H - a)\tau}\, \tau^{m_*-1}\, d\tau,$$

with ~~$\tau_0 = \log(1/\eta_0)$ and $\tau_\varepsilon = \log(r_0^{2M}/\varepsilon)$. For a fixed $\tau_0 \ge 1$ and $\tau_\varepsilon \to +\infty$, the last integral is~~ $\tau_0 = \log(1/\eta_0) > 1$ and $\tau_\varepsilon = \log(r_0^{2M}/\varepsilon) \to +\infty$. If $T_H > a$ and $\tau_\varepsilon \ge \max(\tau_0 + 1, 2)$, then

$$\begin{aligned} e^{(T_H - a)(\tau_\varepsilon - 1)} (\tau_\varepsilon - 1)^{m_*-1} &\le \int_{\tau_0}^{\tau_\varepsilon} e^{(T_H - a)\tau} \tau^{m_*-1}\, d\tau \\ &\le \frac{e^{(T_H - a)\tau_\varepsilon} \tau_\varepsilon^{m_*-1}}{T_H - a}. \end{aligned}$$

The lower bound uses the interval $[\tau_\varepsilon - 1, \tau_\varepsilon]$; the upper bound uses $\tau^{m_*-1} \le \tau_\varepsilon^{m_*-1}$. Since $\tau_\varepsilon - 1 \ge \tau_\varepsilon/2$, these inequalities give an integral $\asymp e^{(T_H-a)\tau_\varepsilon}\tau_\varepsilon^{m_*-1}$~~if $T_H > a$, $\asymp \tau_\varepsilon^{m_*}$ if~~. If $T_H = a$, ~~and $\asymp 1$ if $T_H < a$. (In the first case~~the integral equals $(\tau_\varepsilon^{m_*} - \tau_0^{m_*})/m_*$ and is $\asymp \tau_\varepsilon^{m_*}$. If $T_H < a$, the integral ~~over $[\tau_\varepsilon - 1, \tau_\varepsilon]$ gives the lower bound.)~~ converges to a strictly positive finite number: it is bounded below by its integral over $[\tau_0, \tau_0 + 1]$, and for sufficiently large τ its integrand is at most $e^{(T_H-a)\tau/2}$, because $\tau^{m_*-1} \le e^{(a-T_H)\tau/2}$. Thus it is $\asymp 1$. Since $\tau_\varepsilon = L(\varepsilon) + 2M\log r_0$:

- if $T_H > a$: $V_{r_0}(\varepsilon) \asymp \varepsilon^{T_H} + \varepsilon^a L(\varepsilon)^{m_*-1} \asymp \varepsilon^a L(\varepsilon)^{m_*-1}$;
- if $T_H = a$: $V_{r_0}(\varepsilon) \asymp \varepsilon^{T_H} + \varepsilon^a L(\varepsilon)^{m_*} \asymp \varepsilon^a L(\varepsilon)^{m_*}$;
- if $T_H < a$: $V_{r_0}(\varepsilon) \asymp \varepsilon^{T_H}$.

In the case $Z_{\mathbb{S}} = \emptyset$, $\Xi(\varepsilon r^{-2M})$ vanishes for $r > (\varepsilon/\eta_1)^{1/2M}$ and equals $\sigma(\mathbb{S})$ for $r < (\varepsilon/\eta_2)^{1/2M}$, so again $V_{r_0}(\varepsilon) \asymp \varepsilon^{T_H}$. In all cases H_h has the claimed local volume order with respect to the window system $(W_{r_0})_{r_0>0}$, for Lebesgue measure and the prior 1. This window system is equivalent to the ball system of the Euclidean norm: the sets $W_{r_0} = \Delta_{r_0}(W_1)$ are bounded open neighbourhoods of 0 that shrink to 0 as $r_0 \to 0$, and $B_\delta(0) \subseteq W_{r_0}$ as soon as $r_0^{2M_0} > \max\{\varrho(z) : \|z\| \le \delta\}$. Lemma 4.2 concludes. Adding q_h (Corollary 7.2) gives $\lambda_h = \min(q_h + T_H,\, q_h + a) = \min(T(u), a_*)$ and the stated m_h. □

Corollary 7.7 (Existence)**.** *For every $h \in \mathcal{Z}(n)$, K_n has a local volume order (λ_h, m_h) at the points of O_h. Moreover $\lambda_h \le T(u)$ for every admissible potential u for h, and $\lambda_h \le \lambda_k$ for every $k \in \mathcal{Z}(n)$ with $h \sqsubseteq k$.*

Proof. $(\mathcal{Z}(n), \sqsubseteq)$ is a finite partially ordered set (Proposition 5.7(a)). We argue by descending induction: for a maximal element, Proposition 7.6 applies with $a_* = +\infty$; in general it applies once all strictly larger elements have been treated. The inequalities follow from $\lambda_h = \min(T(u), a_*)$. □

8 The exponent and the codimension

We keep the notation of Section 7: $D \in \mathbb{N}^*$, $n \in \mathbb{N}^{D+1}$, and for $h \in \mathcal{Z}(n)$, (λ_h, m_h) is the local volume order of $K_n = \|P\|_{\mathrm{F}}^2$ at the points of O_h, which exists by Corollary 7.7. In this section the order $\sqsubseteq$ of Definition 5.6 is used, and for $h \in \mathcal{Z}(n)$ we put

$$C^{\sqsubseteq}(h) = \min\{c_{\mathrm{out}}(k) : \ k \in \mathcal{Z}(n),\ h \sqsubseteq k,\ k \text{ balanced}\} \in \mathbb{N} \cup \{+\infty\},$$

with $\min\emptyset = +\infty$. By Proposition 5.7, $C^{\sqsubseteq} = C$, but the proof of Theorem A does not need this.

8.1 Stationarity

For a slot s of h let $\mathrm{in}(s)$ and $\mathrm{out}(s)$ be as in Definition 3.11. Input slots are never the second slot of a linked pair, and output slots never the first one. Hence, for every admissible potential u for h (Definition 7.3),

$$2T(u) = \sum_{(s,s')\in \mathrm{Lk}(h)} (u_{s'} - u_s) = \sum_{s\in\mathcal{S}(h)} \big(\mathrm{in}(s) - \mathrm{out}(s)\big)u_s = c_{\mathrm{out}}(h) + \sum_{s \text{ interior}} \big(\mathrm{in}(s) - \mathrm{out}(s)\big)u_s, \tag{8.1}$$

using $u_s = 0$ on input slots and $u_s = 1$ on output slots. In particular, if h is balanced then $2T(u) = c_{\mathrm{out}}(h)$ for every admissible potential u.

Lemma 8.1 (Stationarity)**.** *Let $h \in \mathcal{Z}(n)$. If $\lambda_h = T(u_0)$ for some admissible potential u_0 for h, then h is balanced and $2\lambda_h = c_{\mathrm{out}}(h)$. Equivalently: if h is not balanced, then $\lambda_h < T(u)$ for every admissible potential u for h.*

Proof. By Corollary 7.7, $\lambda_h \le T(u)$ for every admissible potential u. Suppose $\lambda_h = T(u_0)$. Let Ω be the set of real functions u on $\mathcal{S}(h)$ with $u_s = 0$ on input slots, $u_s = 1$ on output slots, and $u_s < u_{s'}$ for every $(s, s') \in \mathrm{Lk}(h)$. It is an open subset of an affine space (the values on interior slots are free), the admissible potentials are its rational points, and $u_0 \in \Omega$. If some interior slot s had $\mathrm{in}(s) \neq \mathrm{out}(s)$, then by (8.1) a small rational change of u_0 at s alone, in the appropriate direction, would give an admissible potential u with $T(u) < T(u_0) = \lambda_h$, a contradiction. Hence h is balanced, and $2\lambda_h = 2T(u_0) = c_{\mathrm{out}}(h)$ by (8.1). □

8.2 The exponent

Theorem 8.2. *For every $h \in \mathcal{Z}(n)$, the set minimized in $C^{\sqsubseteq}(h)$ is nonempty, and $2\lambda_h = C^{\sqsubseteq}(h)$.*

Proof. Descending induction on the finite partially ordered set $(\mathcal{Z}(n), \sqsubseteq)$. Fix h and assume the claim for all $k \in \mathcal{Z}(n)$ with $h \sqsubseteq k \neq h$. Let a_* be as in Proposition 7.6. The balanced elements k'' with $h \sqsubseteq k'' \neq h$ are exactly the balanced elements above some k with $h \sqsubseteq k \neq h$ (take $k = k''$). So, by the induction hypothesis,

$$\min\{c_{\mathrm{out}}(k'') : \ k'' \in \mathcal{Z}(n),\ h \sqsubseteq k'' \neq h,\ k'' \text{ balanced}\} = \min_{h \sqsubseteq k \neq h} C^{\sqsubseteq}(k) = \min_{h \sqsubseteq k \neq h} 2\lambda_k = 2a_*,$$

with the convention $\min \emptyset = +\infty$ on both sides. Fix an admissible potential u for h; by Proposition 7.6, $\lambda_h = \min(T(u), a_*)$.

If $\lambda_h < a_*$, then $\lambda_h = T(u)$, so h is balanced with $c_{\mathrm{out}}(h) = 2\lambda_h$ (Lemma 8.1). Hence h itself belongs to the minimized set, and $C^{\sqsubseteq}(h) = \min(2\lambda_h, 2a_*) = 2\lambda_h$.

If $\lambda_h = a_*$, then $a_* < +\infty$, and either h is not balanced, or it is balanced with $c_{\mathrm{out}}(h) = 2T(u) \ge 2\lambda_h = 2a_*$. In both cases the minimized set is nonempty and $C^{\sqsubseteq}(h) = 2a_* = 2\lambda_h$.

Maximal elements fall under the first case, since $a_* = +\infty$ for them. □

8.3 The codimension

Proposition 8.3 (Output cut)**.** *For every $h \in \mathcal{Z}(n)$ and every $X \in O_h$,*

$$\mathrm{codim}_X Z(n) = \min\{\#\,\mathrm{Lk}(k) : \ k \in \mathcal{Z}(n),\ h \sqsubseteq k\} = C^{\sqsubseteq}(h).$$

Proof. The first equality. $Z(n)$ is the finite disjoint union of the embedded submanifolds O_k, $k \in \mathcal{Z}(n)$ (Lemma 5.1). We have $X \in \overline{O_k}$ if and only if $O_h \subseteq \overline{O_k}$, i.e. $h \sqsubseteq k$, since $\overline{O_k}$ is $G(n)$-invariant. By (2.3), $d_X(Z(n)) = \max\{\dim O_k : \ k \in \mathcal{Z}(n),\ h \sqsubseteq k\}$, and Lemma 5.2 gives the first equality. In particular $\mathrm{codim}_X Z(n) \le \mathrm{codim}_{X'} Z(n)$ for $X' \in O_k$ with $h \sqsubseteq k$.

Upper bound by $C^{\sqsubseteq}(h)$. Let $k \in \mathcal{Z}(n)$, and let $\mathrm{Lk}_{\to\mathrm{out}}(k) \subseteq \mathrm{Lk}(k)$ be the set of linked pairs (s, s') with s' an output slot; $\#\,\mathrm{Lk}_{\to\mathrm{out}}(k) = c_{\mathrm{out}}(k)$. Every sequence of linked pairs from an input slot to an output slot ends with a pair in $\mathrm{Lk}_{\to\mathrm{out}}(k)$. By Lemma 5.4 (applied to k, which has no full-span slot), $P(\mathbb{I}(k) + Jx) = 0$ whenever $x_\pi = 0$ for all $\pi \in \mathrm{Lk}_{\to\mathrm{out}}(k)$. Since $Z(n)$ is $G(n)$-invariant, the slice chart Ψ_k maps the intersection of its domain with the subspace $\{(\xi, x) : \ x_\pi = 0 \text{ for all } \pi \in \mathrm{Lk}_{\to\mathrm{out}}(k)\}$ into $Z(n)$. The image is an embedded submanifold of codimension $c_{\mathrm{out}}(k)$ through $\mathbb{I}(k)$. By (H1) and (H3), $\mathrm{codim}_{\mathbb{I}(k)} Z(n) \le c_{\mathrm{out}}(k)$. With the

monotonicity above, $\operatorname{codim}_X Z(n) \le c_{\mathrm{out}}(k)$ for every $k \in \mathcal{Z}(n)$ with $h \sqsubseteq k$, in particular for the balanced ones: $\operatorname{codim}_X Z(n) \le C^{\sqsubseteq}(h)$.

Lower bound. Choose $k^* \in \mathcal{Z}(n)$ with $h \sqsubseteq k^*$ and $\#\operatorname{Lk}(k^*)$ minimal, so that $\operatorname{codim}_X Z(n) = \#\operatorname{Lk}(k^*)$. Then k^* is maximal in $(\mathcal{Z}(n), \sqsubseteq)$: if $k^* \sqsubseteq k'' \neq k^*$, Lemma 5.8 would give $\#\operatorname{Lk}(k'') < \#\operatorname{Lk}(k^*)$ with $h \sqsubseteq k''$. Suppose k^* had a residual linked pair π. By Lemma 5.4 (applied to k^*, which has no full-span slot, so that sequences with $p = 0$ do not occur), the only sequence of linked pairs that uses no pair other than π is π itself, and it does not go from an input slot to an output slot. Hence $P(\mathbb{I}(k^*) + \tau J_\pi) = 0$, i.e. $\mathbb{I}(k^*) + \tau J_\pi \in Z(n)$, for every real τ. By Lemma 5.5, $\mathbb{I}(k^*) + \tau J_\pi \in O_{k''}$ for every $\tau \neq 0$, where $k'' = (k^*)^\pi$; so $O_{k''} \subseteq Z(n)$, i.e. $k'' \in \mathcal{Z}(n)$, and $k^* \sqsubseteq k''$ (Lemma 5.5). Moreover $k'' \neq k^*$, since $\rho_{b_s,e_{s'}}(k'') = \rho_{b_s,e_{s'}}(k^*) + 1$ for $\pi = (s, s')$. This contradicts maximality. So every linked pair of k^* is direct. Then no interior slot of k^* belongs to a linked pair, so k^* is balanced, and $c_{\mathrm{out}}(k^*) = \#\operatorname{Lk}(k^*)$. Therefore $\operatorname{codim}_X Z(n) = c_{\mathrm{out}}(k^*) \ge C^{\sqsubseteq}(h)$. □

Combining Theorem 8.2 and Proposition 8.3:

Corollary 8.4. *For every $h \in \mathcal{Z}(n)$ and every $X \in O_h$, K_n has local volume order (λ_h, m_h) at X, and $2\lambda_h = \operatorname{codim}_X Z(n) = C^{\sqsubseteq}(h)$.*

9 The multiplicity

We keep the notation of Sections 7 and 8. Call $h \in \mathcal{Z}(n)$ *$\sqsubseteq$-critical* if it is balanced and $c_{\mathrm{out}}(h) = C^{\sqsubseteq}(h)$. Let $F^{\sqsubseteq}(h)$ be the maximum of $\#\{i \in [\![0, \ell-1]\!] : \ h_i \text{ is } \sqsubseteq\text{-critical}\}$ over all $\ell \in \mathbb{N}$ and all finite sequences $h = h_0 \sqsubseteq h_1 \sqsubseteq \cdots \sqsubseteq h_\ell$ in $\mathcal{Z}(n)$ with $h_i \neq h_{i+1}$ and $C^{\sqsubseteq}(h_i) = C^{\sqsubseteq}(h)$ for all i. By Proposition 5.7, these notions agree with "critical" and F of Definition 3.12.

Lemma 9.1. *Let $h \in \mathcal{Z}(n)$ and let u be an admissible potential for h. Then h is $\sqsubseteq$-critical if and only if $\lambda_h = T(u)$.*

Proof. If $\lambda_h = T(u)$, then h is balanced and $c_{\mathrm{out}}(h) = 2\lambda_h$ by Lemma 8.1, and $2\lambda_h = C^{\sqsubseteq}(h)$ by Theorem 8.2. Conversely, if h is balanced, then $2T(u) = c_{\mathrm{out}}(h)$ by (8.1); if moreover $c_{\mathrm{out}}(h) = C^{\sqsubseteq}(h) = 2\lambda_h$, then $T(u) = \lambda_h$. □

Theorem 9.2. *For every $h \in \mathcal{Z}(n)$, $m_h = 1 + F^{\sqsubseteq}(h)$.*

Proof. Let $\operatorname{Succ}(h) = \{k \in \mathcal{Z}(n) : \ h \sqsubseteq k \neq h, \ C^{\sqsubseteq}(k) = C^{\sqsubseteq}(h)\}$. Splitting off the first step of a sequence gives

$$F^{\sqsubseteq}(h) = \max\Big(0, \ \max_{k \in \operatorname{Succ}(h)} \big([h \text{ is } \sqsubseteq\text{-critical}] + F^{\sqsubseteq}(k)\big)\Big), \tag{9.1}$$

where $\max \emptyset = 0$ and $[\cdot] \in \{0, 1\}$ is the truth value. We argue by descending induction on $(\mathcal{Z}(n), \sqsubseteq)$, and compare (9.1) with Proposition 7.6 for a fixed admissible potential u. By Theorem 8.2, $k \in \operatorname{Succ}(h)$ if and only if $h \sqsubseteq k \neq h$ and $\lambda_k = \lambda_h$. Since $\lambda_h = \min(T(u), a_*) \le a_* \le \lambda_k$ for all such k, the set $\operatorname{Succ}(h)$ is nonempty exactly when $\lambda_h = a_*$, and then it consists of the k with $h \sqsubseteq k \neq h$ and $\lambda_k = a_*$.

- If $T(u) < a_*$: h is $\sqsubseteq$-critical (Lemma 9.1) and $\operatorname{Succ}(h) = \emptyset$, so $F^{\sqsubseteq}(h) = 0$ and $m_h = 1$.
- If $T(u) > a_*$: h is not $\sqsubseteq$-critical, and by the induction hypothesis

$$m_h = m_* = \max_{k \in \operatorname{Succ}(h)} m_k = 1 + \max_{k \in \operatorname{Succ}(h)} F^{\sqsubseteq}(k) = 1 + F^{\sqsubseteq}(h).$$

- If $T(u) = a_*$: h is $\sqsubseteq$-critical, $\mathrm{Succ}(h) \neq \emptyset$, and

$$m_h = m_* + 1 = 1 + \max_{k \in \mathrm{Succ}(h)} \big(1 + F^{\sqsubseteq}(k)\big) = 1 + F^{\sqsubseteq}(h). \qquad \square$$

Remark 9.3 (Critical strata). By Corollary 8.4 and Proposition 8.3, $C^{\sqsubseteq}(k)$ is the local codimension of $Z(n)$ at the points of O_k. The sequences in the definition of $F^{\sqsubseteq}(h)$ are therefore chains of orbits $O_h \subseteq \overline{O_{h_1}}$, $O_{h_1} \subseteq \overline{O_{h_2}}, \dots$ along which the local codimension of $Z(n)$ is constant. An orbit O_k is critical if and only if the polynomial H_k of Section 7 has local volume exponent exactly T_H, the exponent "expected" from the weights; equivalently, the radial integral in the proof of Proposition 7.6 contributes at the leading order. Each critical orbit along a chain contributes one factor $L(\varepsilon)$. Combinatorial formulas of a similar kind, for the real log canonical threshold and its multiplicity, are known for real hyperplane arrangements [18].

Remark 9.4 (Multiplicity and the number of top strata). For $h \in \mathcal{Z}(n)$~~let $N_{\mathrm{top}}(h)$ be the number of $k \in \mathcal{Z}(n)$~~, the orbits $O_k \subseteq Z(n)$ whose closures contain $\mathbb{I}(h)$ and whose dimension is largest are precisely those with $h \sqsubseteq k$ and $\#\,\mathrm{Lk}(k) = \mathrm{codim}_{\mathbb{I}(h)} Z(n)$~~; the orbits O_k of these k are the top-dimensional strata of $Z(n)$ through $\mathbb{I}(h)$. (~~, by Proposition 8.3. The top-dimensional components of $Z(n)$ itself are described in [24, 17]. ~~)~~ In Example 3.18 ~~, $m_h = 3 < 4 = N_{\mathrm{top}}(h)$. In the companion research notes we show that $m_h = N_{\mathrm{top}}(h)$ for every h when $D = 2$, using the depth-two formula of Example 3.16. A computer search found $m_h = N_{\mathrm{top}}(h)$ for all barcodes with $D = 3$ and all widths at most 16, and $m_h \leq N_{\mathrm{top}}(h)$ in all computed cases, including depths four and five. We do not use these facts here, and the~~ there are four such orbits, whereas $m_h = 3$. Thus the multiplicity is not, in general, the number of these orbits. A general relation between ~~m_h and $N_{\mathrm{top}}(h)$ is~~ the two numbers remains open.

10 Proofs of the main theorems

Proof of Theorem A. The case $D = 0$ was treated after the statement. Let $D \geq 1$, $n \in \mathbb{N}^{D+1}$, $\Phi \in \mathbb{R}^{n_D \times n_0}$, $A \in F_\Phi$, $r = \mathrm{rk}\,\Phi$, and let $h_A \in \mathcal{Z}(n-r)$ be the residual barcode of A. By Corollary 8.4 applied to the width vector $n-r$ and the barcode h_A, the function K_{n-r} has a local volume order (λ_{h_A}, m_{h_A}) at $\mathbb{I}(h_A)$, with $2\lambda_{h_A} = \mathrm{codim}_{\mathbb{I}(h_A)} Z(n-r)$. By Lemma 4.6, $L_{n-r,0} = \frac{1}{2} K_{n-r}$ has the same local volume order. By Proposition 6.2, L_Φ has local volume order $(\frac{1}{2} c_{\mathrm{reg}}(n,r) + \lambda_{h_A},\ m_{h_A})$ at A, and

$$2\lambda_A(L_\Phi) = c_{\mathrm{reg}}(n,r) + 2\lambda_{h_A} = c_{\mathrm{reg}}(n,r) + \mathrm{codim}_{\mathbb{I}(h_A)} Z(n-r) = \mathrm{codim}_A F_\Phi.$$

By Lemma 3.2(b) the same pair is the local volume order for every Haar measure, every prior and every norm, and by Lemma 3.2(a) it is unique. Finally $\mathrm{codim}_A F_\Phi = N(n) - d_A(F_\Phi)$ is an integer, since $d_A(F_\Phi)$ is the dimension of a stratum, by (2.3) and Lemma 5.9. $\square$

Proof of Theorem B. With the notation of the previous proof, Corollary 8.4, Proposition 8.3 and Theorem 9.2, applied to $h_A \in \mathcal{Z}(n-r)$, give

$$2\lambda_{h_A} = \mathrm{codim}_{\mathbb{I}(h_A)} Z(n-r) = C^{\sqsubseteq}(h_A) = \min\{\#\,\mathrm{Lk}(k) :\ k \in \mathcal{Z}(n-r),\ h_A \sqsubseteq k\},$$
$$m_{h_A} = 1 + F^{\sqsubseteq}(h_A),$$

and the set defining $C^{\sqsubseteq}(h_A)$ is nonempty (Theorem 8.2). By Proposition 5.7, $\sqsubseteq$ coincides with $\preceq$ on $\mathcal{Z}(n-r)$. Hence $C^{\sqsubseteq}(h_A) = C(h_A)$, "$\sqsubseteq$-critical" means "critical", and $F^{\sqsubseteq}(h_A) = F(h_A)$. The theorem follows from Proposition 6.2, as in the proof of Theorem A. $\square$

Remark 10.1 (What depends on what). Theorem A uses Sections 4–8; of Proposition 5.7 it uses only part (a), the easy direction. Theorem B additionally uses Proposition 5.7(b), proved

in Appendix B, and Section 9. Besides standard results of real analysis (the inverse function theorem, the change of variables formula, the Fubini–Tonelli theorem), the external inputs are: Fact 3.8(a) (Gabriel's theorem for A_{D+1}); the facts (H0)–(H4) on Hausdorff dimension; the homeomorphism theorem for transitive actions [14] and the fact that an injective immersion which is a homeomorphism onto its image is an embedding [22], both used in Lemma 5.1; Sard's theorem, used in Lemma 5.9; and, in Appendix B, the rank-matrix criterion for the Bruhat order [5].

Remark 10.2 (Dependence on the data). That (λ, m) depends only on the barcode of A also follows directly: if $\beta(A) = \beta(A')$ and A, A' are exact fits of Φ, Φ', then $A' = g \cdot A$ for some $g \in G(n)$, and Step 1 of the proof of Proposition 6.2 transports one problem to the other. The content of the theorems is the value of (λ, m).

Remark 10.3 (Data on a subspace). Assume $D \geq 1$. Let $\Sigma \in \mathbb{R}^{n_0 \times n_0}$ be symmetric positive semidefinite of rank n_0'. The population loss of linear regression with input covariance Σ is $B \mapsto \frac{1}{2}\|(P(B) - \Phi)\Sigma^{1/2}\|_{\mathrm{F}}^2$. Choose an orthonormal basis of $\mathbb{R}^{n_0}$ whose first n_0' vectors span the image of Σ, let Σ' be the (invertible) restriction of Σ to its image, and for a chain $B = (B_0, \dots, B_{D-1}) \in \mathrm{Rep}(n)$ split $B_0 = (B_0' \mid B_0'')$ accordingly, with $B_0' \in \mathbb{R}^{n_1 \times n_0'}$. Then the loss of B equals $\frac{1}{2}\|(P(B') - \Phi')\Sigma'^{1/2}\|_{\mathrm{F}}^2$, where $B' = (B_0', B_1, \dots, B_{D-1}) \in \mathrm{Rep}(n')$ with $n' = (n_0', n_1, \dots, n_D)$ and Φ' is the restriction of Φ to the image of Σ; it lies between two positive multiples of $L_{n',\Phi'}(B')$, and the $(n_0 - n_0')n_1$ entries of B_0'' are spectator variables. By Lemmas 4.6 and 4.8, at a zero A of this loss the local volume order is that of $L_{n',\Phi'}$ at A', given by Theorems A and B for the width vector n'. A positive semidefinite weight on the output side is handled in the same way, with the rows of ~~A_{D-1}~~ B_{D-1} in place of the columns of ~~A_0~~B_0. A degenerate quadratic form on $\mathbb{R}^{n_D \times n_0}$ that is not of this product form is not covered; for $D = 2$ it is studied in [11].

11 The formalization

11.1 What is formalized

The Lean theorem `MAIS.ExactFitStatement.exact_fit_classification`, stated and translated in Appendix A, asserts the following for all D, n, every Haar measure μ, every exact fit A of Φ, and every function φ continuous at A with $\varphi(A) > 0$. There are $c, m \in \mathbb{N}$ such that:

(F1) the formal version of the program Prog returns exactly $\{(c, m)\}$ on $(n, R(A))$;

(F2) c is the local Hausdorff codimension of F_Φ at A;

(F3) for every (λ, m'), L_Φ has local volume order (λ, m') at A if and only if $(\lambda, m') = (c/2, m)$.

By Proposition A.1, this implies Theorem A (with Lemma 3.2(b) for the independence of the norm) and, with Proposition A.2 and Lemma 3.13, Theorem B except its last sentence, which is proved only in the text. The proof of Proposition A.2 uses three results of the text (Proposition 5.7(a), Theorem 8.2 and Lemma 8.1) at one point, as explained after it. The formal proof of this statement only composes three theorems of the formal development:

- `MAIS.Run2ArithmeticPositive.exactFitPairs_nonempty_singleton` (the program returns one pair);
- `MAIS.Run2ArithmeticPositive.entry_half_prior_pair_iff` (local volume orders are characterized by the program);

- `MAIS.Run2ArithmeticPositiveGeometry.exactFitPair_literal_hausdorff_codimension` (the first coordinate is the local codimension);

together with the definitional equivalence `MAIS.DirectBandOrder.measuredBandPair_iff`, whose proof is `rfl`. The ~~import closure of the statementconsists of the Mathlib library at commit and 282 files of the repository (the root file of an earlier formalization of the depth-two case and 281 modules of the library)~~public file `ExactFitFull_2026_10_28_UTC.lean` in [10] concatenates 283 source modules, including this statement. Its only external import is Mathlib at commit `664cbbe52927ce6acaa39ca35498336116e123b8`. The Lean version is `4.34.0-rc2`. Module names in this section identify declarations within that file; separate module files are not required to check it.

The formal development follows the same strategy as Sections 5–9, with some technical differences. For instance, the radial integration of Proposition 7.6 is done with dyadic annuli instead of weighted polar coordinates, and Proposition B.1 is proved with a different reduction to permutations.

11.2 What a reader has to trust

A Lean proof is accepted by Lean's kernel, a small type checker. Besides the kernel, the trusted base consists of the axioms used, the meaning of the Mathlib definitions that occur in the statement, and the translation of the statement into mathematics.

- *Axioms.* The command `#print axioms` reports that the theorem depends only on `propext`, `Classical.choice` and `Quot.sound`, the standard axioms of classical mathematics in Lean. A `sorry` or a newly declared axiom would appear in this list. The file checks this list at compile time.
- *Definitions.* Appendix A quotes ~~every repository definition used by the statement and every Mathlib definition that controls its meaning,~~ the controlling project definitions and the Mathlib definitions that fix the norms, measure and Hausdorff dimension, with a translation~~. The script~~ ; the path product is characterized by the three displayed proved identities. Other standard Mathlib notions, such as rank and finite dimension, retain their usual meanings. The audit in `exact-fit-local/` computes the *definitional closure* of the statement: the constants reachable from its type through types and bodies of definitions, but not through proofs. It prints every repository constant in this closure. This list is what a reader has to check against Appendix A and the public file.
- *Compilation.* See Sections 11.4 and 11.5.

11.3 Deceptive formal statements

A kernel-checked theorem can still fail to mean what its reader thinks. Lean allows user-defined notation, local instances (which can change, for example, which norm $\|\cdot\|$ denotes), shadowing of names, metaprograms run at compile time, and identifiers that look alike, such as `rn` and `m`, or Latin and Greek letters. The kernel checks the elaborated proposition, not the reader's interpretation of the source text. ~~We therefore ran a static scan of the 282 repository files in the import closure. It~~ The constituent sources and their concatenation were inspected for these mechanisms. The source checks found:

- no `axiom`, `sorry`/`admit`, `native_decide`, `implemented_by`, `extern`, `unsafe`, or `debug.skipKernelTC`;

- no metaprogramming commands (`macro`, `syntax`, `elab`, `run_cmd`, `#eval`, `initialize`);
- no global notation (only `local notation` abbreviations inside proofs of the root depth-two file), and no reopening of Mathlib namespaces (in particular none of `Matrix`, `Real`, `MeasureTheory`, `Metric`);
- no instance of a norm, metric, measure or measurable-space class (the instances declared are decidability and finiteness instances, Prop-valued instances such as `BorelSpace`, and the group action on chains);
- no project declaration whose name could shadow a Mathlib name occurring in the statement;
- no non-ASCII characters other than standard mathematical symbols and Greek letters, and no invisible or bidirectional characters.

Lean does not accept unquoted identifiers in the Cyrillic alphabet, so the most common look-alike letters are excluded automatically. In the ~~statement file itself, every~~ final statement, every project name is either fully qualified or defined ~~in the file.~~ immediately before the statement. These source checks apply to the released bytes, not to arbitrary later Lean files.

Many files raise the option `maxHeartbeats` (up to 3.2×10^6 instead of the default 2×10^5). This option bounds the time spent by the *elaborator*, which searches for proofs and fills in implicit arguments. It plays no role in the kernel check, so it cannot make a false statement provable; it indicates slow proofs, which may need maintenance when the dependencies change. The option `backward.isDefEq.respectTransparency false`, also used in many files, likewise only affects elaboration.

11.4 ~~Compilation records~~The public proof file and its checks

The public repository [10] contains `ExactFitFull_2026_10_28_UTC.lean` (63 475 lines). Its SHA-256 is

`b362a3ee35df68adad7aa24a1ddd288e63e4e59d074445aaafef4093e8462a2c`.

Its header lists the constituent modules and their source hashes. Their bodies are placed inside sections, with internal imports removed. Concatenation changes which declarations are visible during elaboration; equality of source fragments alone is therefore not a proof that the elaborated statement is unchanged. The ~~workflow of the repository~~was run on 27 fingerprint comparison described below checks the statement and its controlling definitions in addition to compiling the proof.

The author ran the released file through the local harness on 28 October ~~2026 on a hosted GitHub runner (2 cores, 7 GB of memory), at the source commit . The logs are kept on the branch (commit , folder). The run did the following. It installed Lean (commit) from the file , and the precompiled Mathlib at the commit fixed by . It compiled with the modules , and with all their imports (among them), and then the module : 9130 build jobs in total, all successful, with no warning. Compilation passes every declaration through the kernel. It also evaluates the #check of~~ 2026. The published receipt in `exact-fit-local/logs/` records a successful target build, ordinary kernel replay, the ~~axiom list written in the statement file, which would make the build fail if the list changed. It ran the audit script . The script printed the statement as elaborated, once with notation and once with notation disabled and fully qualified names. Both agree with the source quoted in Appendix~~ A~~. It printed the axiom list . Finally it printed the definitional closure of the statement: 92 repository constants and~~

270 constants of Mathlib and the Lean core. The repository constants are 77 definitions, one inductive type with its constructor, axiom list `[propext, Classical.choice, Quot.sound ]`, and 13 auxiliary theorems that the elaborator generates for proof terms occurring inside definitions; there is no axiom and no opaque constant among them. They all lie in the 12 files quoted in Appendix A or in the two files defining the composable-arrows construction behind , which the appendix pins down by three proved lemmas instead of quoting. It ran on the five modules , , , and . Each run exited with status 0. Without the option , re-checks with the kernel the declarations of the named module, taking its imports as given. The other modules of the import closure were checked by the kernel when they were compiled in (ii). A replay of the whole closure from an empty environment () was not run. It recorded the SHA-256 hashes of , , , the statement file, and a fingerprint match. The target build reused cached compiled output in that run; the replay was executed anew. The published replay, audit and fingerprint logs have the hashes listed in the receipt. The scan log was redacted after the run and consequently no longer has the receipt's scan-log hash; the accompanying notice records this limitation. The receipt is unsigned. It is execution evidence, not a certification of the audit script. The two Lean files quoted in Appendix A for the statement are byte-identical to the compiled ones (the hashes are in). The other quoted files are taken from the same commit. The run took 33 minutes, of which 25 were spent compiling the import closuretranslation into mathematics.

11.5 A single file and a local recheck

For readers who want to recheck the proof themselves, the import closure of the statement (the 282 files above and the statement file) has been concatenated into a single Lean file, at the root of the repository (63 475 lines, SHA-256 beginning with), which imports only Mathlib. The script copies each module byte for byte inside its own , removes only the imports internal to the repository, and closes the scopes that the module leaves open. The header of the file lists every module with the SHA-256 of its source, and the option of the script regenerates the file and compares it The logs in `verification/2026-10-28-agent-check/` record a separate check on 28 October that compiled the exact same file from source with the pinned toolchain and cached Mathlib, then replayed its compiled module and recomputed the fingerprint. Compilation exited with status 0 after about 14 minutes, with one tactic-style warning; replay exited with status 0, and the fingerprint matched the reference byte for byte. No statement or proof was changed. In a single file, the declarations of a module become visible to modules that did not import it; the fingerprint below shows that this changes neither the statement nor the definitions it depends on. This was an agent-operated execution of Lean, not an independent implementation of its kernel. No fresh replay of all imported Mathlib and Lean-core declarations is claimed for either check.

11.5 Reproducing the check

The folder `exact-fit-local/` of the repository contains a container harnessadapted from the one published with [9]provides a Podman/Docker harness. Its image contains only Ubuntu and the Lean version manager. It downloads the pinned versions of Setup obtains the pinned Lean and Mathlibonce, and then runs every check ; subsequent checks run without network access , as an unprivileged container user, with the source file input source mounted read-only. Its The command `all` does the following performs the following operations.

(i) It scans the file, outside comments and strings, for the constructs listed in Section 11.3A source scan reports proof placeholders, added axioms, native-evaluation escapes,

metaprogramming commands, notation declarations and invisible control characters. A lexical scan is a reading aid, not a proof that an arbitrary file cannot execute code.

(ii) ~~It~~ A target build compiles the file~~as a module. This also evaluates~~, or reuses an unchanged compiled target. Source compilation checks the final `#guard_msgs` ~~check of the~~ assertion of the theorem's axiom list.

(iii) ~~It replays every declaration of this module with . Since the module contains the whole repository part of the proof , this replay covers the entire import closure except Mathlib and the Lean core , whereas the replay of Section 11.4(iv) covered only five modules~~`lean checker` replays the declarations of the compiled module with Lean's kernel. The entire project proof is in that module. Ordinary replay takes Mathlib and Lean core as given; the option `FRESH=1` also replays those imports with the same kernel.

(iv) ~~It~~ The audit prints the statement, ~~its axioms, and~~ once with notation and once with all implicit arguments and instances displayed; it prints the axiom list and a fingerprint of ~~its definitional closure: for each definition, inductive type and constructor of the closure, a 128-bit structural hash of its type and of its body,~~ the definitions that determine the statement's meaning.

(v) The fingerprint is compared with `exact-fit-local/reference/fingerprint-multi.txt`, and ~~for the theorems occurring in these terms,~~ the ~~hashes of their statements. The hashes are computed after normalizations that do not change the meaning of the statement: the module part of the names of private declarations is removed, bound variables are not named, metadata annotations (which the kernel ignores) are dropped, and a theorem occurring inside a term is identified by its statement rather than by its name (by proof irrelevance ; the elaborator shares identical auxiliary lemmas between the declarations of one module, so their names differ between the two builds). The harness compares this fingerprint with the one printed by the same script in the original multi-file development.~~

(vi) ~~It writes a receipt with the SHA-256 hashes of the input file, of the container image and of the logs, which the person who ran the check can sign.~~ harness writes a receipt and full logs.

~~*Record.* The workflow ran this harness twice with Docker on 28 October 2026, on hosted runners with 2 cores and 7 GB of memory, at the source commits and . The single file was the same in both runs; between them, only the fingerprint script and one line of the receipt writer changed. In each run, the script reproduced the single file byte for byte, the scan found no forbidden construct, the file compiled (17 and 12 minutes), the replay of (iii) exited with status 0 (less than 2 minutes), and the axioms printed were . In each run the fingerprint was also computed in the multi-file development, and the statement printed with was identical in both builds. The first run used a first version of the fingerprint, which still took into account the names of auxiliary lemmas and of bound variables; it reported differences. The second run used the version described in (iv): the two fingerprints were identical, byte for byte (~~The reference fingerprint has SHA-256 ~~beginning with). They list~~

`34e2fa7f3635b4f82eb985509eb7abad5e820413bc0eb34cd06241be172169d6`.

It contains the printed statement, 79 ~~constants of the repository, namely 77 definitions, one inductive type and its constructor, and~~ project definitions, inductive types and constructors, 17 ~~distinct statementsof theorems occurring in them. The 79 constants are exactly the repository constants of the closure listed in Section 11.4(iii), minus its 13 auxiliary theorems.~~

~~The logs of the two runs are kept on the branch , in the folders and ; the reference fingerprint used by the harness is the one printed in the second run.~~ distinct theorem statements used inside them, and the names of the external constants. For project definitions it records structural hashes of types and bodies. Bound-variable names and metadata are ignored, private declaration names lose their module prefix, and proofs are identified by their propositions. Proof irrelevance justifies the last normalization. The structural hash is not cryptographic and the comparison is not a formally proved equivalence or an adversarial security mechanism. It supplies evidence against accidental changes during extraction; the mathematical interpretation still requires reading the definitions.

12 ~~AI use statement~~

The harness README gives the commands and dependency-reuse options. A shared O77 dependency volume remains writable and must not be deleted by the harness's `clean` command. The source file should remain unchanged throughout a run. Preserve the logs together with the receipt, and require an explicit fingerprint match rather than the mere presence of a receipt file.

12 Disclosure of AI use

~~This section describes how the results and this text were produced. *Drafted by an AI assistant from the author's instructions, last revised on 28 October 2026 from his comments; to be checked by the author before any circulation.*~~ This work was directed by Niels Feld and produced with substantial assistance from AI systems. The following distinctions concern the mathematical text, the formal statements, the formal proofs and their verification.

~~Who did what~~Origin and contributions

- ~~**Origin**~~**Research direction.** ~~The project began in August 2026 with~~ Following Lionel Levine's invitation ~~to mathematicians [25] and the open problems of the~~ and the Math for AI Safety ~~repository [30]. The author first directed the formalization in Lean 4 of an answer to~~ problems [25, 30], the author directed the Lean formalization of the depth-two problem MAIS-O77 ~~[29]; this development, with its blueprint, is public [9] (). He then asked for the generalization to arbitrary depth, which is the subject of this paper, and directed that project~~[29, 9], then requested the arbitrary-depth exact-fit generalization studied here.

- **Formal proofs.** ~~The Lean 4/Mathlib development that proves Theorems A and B was written by AI agents (~~AI agents using OpenAI GPT models ~~, run through OpenAI's Codex tooling) during October 2026,~~ through Codex wrote the Lean development under the author's direction. The author did not write ~~the formal proofs.~~ these proofs. The checked formal endpoint is (F1)–(F3) of Appendix A. Proposition A.1 and the discussion after Proposition A.2 specify the additional mathematical arguments connecting it to Theorems A and B; in particular, the selected endpoint does not itself assert the last sentence of Theorem B.

- ~~**Mathematical text.** The~~ **Mathematical exposition.** Claude (Anthropic) wrote the natural-language ~~proofs in Sections 4–10 and Appendix B were written by an AI assistant (Claude, Anthropic), working~~ arguments from the formal development and its documentation, ~~and revised following~~ then revised them after the author's reading.

~~Earlier drafts were reviewed by other AI agents (Claude and GPT), whose findings were incorporated. Example 3.18 and the explanation of the discrepancy in it were first written up by~~ OpenAI GPT/Codex and Claude agents reviewed successive versions and proposed corrections. GPT/Codex ~~. The proof of Proposition B.1 given~~ first wrote the depth-four example and its explanation; Claude wrote the proof in Appendix B, which differs from the formal ~~one and was written for this paper~~proof. GPT/Codex prepared the present revision, including explicit estimates and the public source references.

- ~~**Formal statement**~~ **Statement and** ~~**translation**~~**checking tools.** ~~The single formal statement of Appendix A~~ Claude wrote the single-statement wrapper and its translation~~were written by Claude~~, the single-file extractor and the checking harness, the latter adapted from [9]. The formal ~~statement~~ wrapper and its proof ~~are checked by Lean's kernel; the~~ have been accepted by Lean; its translation into mathematics ~~is not machine-checked.~~ remains a reading task. Source quotations are copied mechanically and checked against the released Lean file or pinned Mathlib sources.

- ~~**Compilation.** The formal development was compiled many times by the AI agents while they wrote it, and by the continuous-integration runs recorded in Section 11.4. A single-file version and a container that rechecks it offline are described in Section 11.5; any reader, including the author, can rerun this check.~~

- ~~**Literature.** References were collected with AI assistance. The attributions to [24] were checked against the arXiv text. The bibliographic details of the classical references should be checked before submission.~~ **Literature.** AI agents assisted with the search and attribution of prior work. References distinguish results used in proofs from related precedents. No claim that the literature search is exhaustive is made.

~~What the author has checked~~Human review and execution

The author has read the ~~whole paper and found no mistake, but he~~ mathematical exposition and the formal statement with its controlling definitions. He found no error in those readings, but has not checked ~~all the algebraic computations by hand . He has read the formal statement and the definitions it depends on (Appendix A) and found no discrepancy with Theorems A and B; he~~ every algebraic calculation by hand and does not claim a complete certification of ~~this correspondence~~the translation. He has not read all the formal proofs~~: they are checked by Lean's kernel. The author is a mathematician working~~ . His expertise is in algebraic geometry~~and homotopy theory , not in~~ , motivic homotopy theory and K-theory, rather than singular learning theory.

~~What the machines have checked~~

~~The formal proofs are checked by the kernel of Lean 4; the trusted base is~~ He ran the local Lean check described in Section ~~11. Independent computer programs, also written~~ 11.4; this is distinct from the source compilations and checks performed by AI agents~~, recomputed the examples of Section 3.5 and tested the definitions on many small cases. These computations are not used in any proof.~~ .

The present version incorporates further AI-proposed revisions. The author's readings of earlier versions should not be interpreted as approval of every later edit. The red-and-blue comparison file accompanying this version identifies the mathematical and editorial changes for review.

~~Responsibility~~Scope of machine verification and responsibility

~~Errors may remain in the natural-language text even where the formal statementis correct; conversely,~~ Lean's kernel checks proof terms for the elaborated formal statement, subject to the axioms and imported declarations described in Section 11. It does not check whether the statement expresses the ~~formal statement could fail to express the~~ intended mathematics~~in ways that a reader of Appendix A should check. The author takes responsibility for the decision to circulate this text~~, whether the prose proof is correct, or whether a bibliographic attribution is complete. Kernel replay uses the same kernel; agreement among AI reviewers is not independent human refereeing. No referee report is claimed. The author is responsible for the version he chooses to circulate and for its mathematical claims.

Acknowledgements

~~**To be completed by the author.** The problems MAIS-O70 and MAIS-O77 were posed in~~ The author acknowledges Lionel Levine and the Math for AI Safety ~~program led by Lionel Levine [30, 25]. Rob Sneiderman [26]~~ contributors for the questions that initiated this work; Rob Sneiderman and Svyatoslav Novikov~~(~~, with OpenAI Codex~~) [27] made public, in August 2026, candidate solutionsof these problems that state the depth-two case of the pointwise formula. The geometric viewpoint on the fibres of the multiplication map, and the global identification of the threshold with half the codimension, are due to Simon Pepin Lehalleur and~~ , for the posted depth-two candidate solutions [26, 27]; Simon Pepin Lehalleur and Richárd Rimányi for the geometric study of the fibres, the global half-codimension theorem and the pointwise conjecture [24]; ~~the global learning coefficients of deep linear networks are due to Miki Aoyagi [3].~~ and Miki Aoyagi and Sumio Watanabe for the global learning-coefficient calculations [2, 3]. Related contributions, including precedents not used in the proof, are identified in Section 1 and where they enter the argument. The formal development depends on Lean and the work of the Mathlib contributors. The substantial contributions of the AI systems are described above rather than presented as personal communications or human refereeing.

A The formal statement and its translation

This appendix is for a reader who wants to check that the Lean theorem says what Theorems A and B say. ~~All Lean text~~ Each project quotation below is copied mechanically from the ~~repository files, at the commit given~~ public file `ExactFitFull_2026_10_28_UTC.lean` identified in Section ~~11. Each block is followed by its file, line numbers and the beginning of the SHA-256 hash of the~~ 11.4; its displayed line numbers refer to that file. Mathlib ~~files~~ quotations are from the pinned ~~commit .~~ public Mathlib source. Each block gives its source and the beginning of its SHA-256 hash.

A.1 Reading conventions

- `ℕ`, `ℝ`, `ℚ` are $\mathbb{N}$, $\mathbb{R}$, $\mathbb{Q}$. `ℝ≥0∞` (also `ENNReal`) is $[0, +\infty]$. `Fin k` is $\{0, \dots, k-1\}$. For e in `Fin D`, `e.castSucc` and `e.succ` are the vertices e and $e+1$, and `Fin.last D` is D.
- Subtraction on $\mathbb{N}$ is truncated at 0; so is subtraction on $[0, +\infty]$, with $(+\infty) - x = +\infty$ for finite x. `ENNReal.ofReal x` is $\max(x, 0) \in [0, +\infty[$.
- `Matrix (Fin p) (Fin q) ℝ` is $\mathbb{R}^{p \times q}$, as functions of (row, column). `M.rank` is the rank.

- `Real.rpow ε λ` is ε^λ (used only for $\varepsilon > 0$), `Real.log` is the natural logarithm, `Real.exp` the exponential, `Real.sqrt` the nonnegative square root.
- `∃ c m : ℕ, P` means "there exist $c, m \in \mathbb{N}$ with P". `{(c, m)}` is a one-element finite set, and `s = {(c, m)}` says that the finite set s is exactly $\{(c, m)\}$.

The following Mathlib definitions control the meaning of the statement.

Norms. After `open scoped Matrix.Norms.Frobenius`, the norm of a matrix is the Frobenius norm:

```
theorem frobenius_norm_def (A : Matrix m n α) :
    ‖A‖ = (∑ i, ∑ j, ‖A i j‖ ^ (2 : ℝ)) ^ (1 / 2 : ℝ) :=
  (congr_arg ((↑) : ℝ≥0 → ℝ) (frobenius_nnnorm_def A)).trans <| by simp [NNReal.coe_sum]
```

[Q1] `Mathlib 664cbbe5, Mathlib/Analysis/Matrix/Normed.lean`, lines 561–563, file SHA-256 `e430f9176d36…`

```
namespace Norms.Frobenius
attribute [scoped instance]
  Matrix.frobeniusSeminormedAddCommGroup
  Matrix.frobeniusNormedAddCommGroup
  Matrix.frobeniusNormedSpace
  Matrix.frobeniusNormedRing
  Matrix.frobeniusNormedAlgebra
  Matrix.frobeniusIsBoundedSMul
  Matrix.frobeniusNormSMulClass
end Norms.Frobenius
```

[Q2] `Mathlib 664cbbe5, Mathlib/Analysis/Matrix/Normed.lean`, lines 681–690, file SHA-256 `e430f9176d36…`

Translation. $\|A\| = \|A\|_{\mathrm{F}}$ for every real matrix A. A chain is an element of a finite product, which Mathlib normalizes with the sup norm:

```
/-- Finite product of seminormed groups, using the sup norm. -/
@[to_additive /-- Finite product of seminormed groups, using the sup norm. -/]
instance Pi.seminormedGroup : SeminormedGroup (∀ i, G i) where
  norm f := ↑(Finset.univ.sup fun b => ‖f b‖₊)
```

[Q3] `Mathlib 664cbbe5, Mathlib/Analysis/Normed/Group/Constructions.lean`, lines 301–304, file SHA-256 `ab6a98350718…`

Translation. For a chain B, $\|B\| = \max_e \|B_e\|_{\mathrm{F}}$, so `Metric.ball A r`$= B_r^\infty(A) := \{B : \max_e \|B_e - A_e\|_{\mathrm{F}} < r\}$. This is *not* the Euclidean ball; it is used only in the codimension conjunct, where the choice of norm does not matter (Definition 2.1).

Measures. A measure on the chain space is defined on its Borel σ-algebra. An additive Haar measure is:

```
/-- A measure on an additive group is an additive Haar measure if it is left-invariant, and
gives finite mass to compact sets and positive mass to open sets.

Textbooks generally require an additional regularity assumption to ensure nice behavior on
arbitrary locally compact groups. Use `[IsAddHaarMeasure μ] [Regular μ]` or
`[IsAddHaarMeasure μ] [InnerRegular μ]` in these situations. Note that a Haar measure in our
sense is automatically regular and inner regular on second countable locally compact groups, as
checked just below this definition. -/
class IsAddHaarMeasure {G : Type*} [AddGroup G] [TopologicalSpace G] [MeasurableSpace G]
    (μ : Measure G) : Prop
    extends IsFiniteMeasureOnCompacts μ, IsAddLeftInvariant μ, IsOpenPosMeasure μ
```

[Q4] `Mathlib 664cbbe5, Mathlib/MeasureTheory/Group/Measure.lean`, lines 764–774, file SHA-256 `cba62e69e14a…`

Translation. A translation-invariant measure, finite on compact sets and positive on nonempty open sets. On a finite-dimensional real vector space these are exactly the positive multiples of Lebesgue measure.

```
/-- Given a measure `μ : Measure α` and a function `f : α → ℝ≥0∞`, `μ.withDensity f` is the
measure such that for a measurable set `s` we have `μ.withDensity f s = ∫⁻ a in s, f a ∂μ`. -/
noncomputable
def Measure.withDensity {m : MeasurableSpace α} (μ : Measure α) (f : α → ℝ≥0∞) : Measure α :=
  Measure.ofMeasurable (fun s _ => ∫⁻ a in s, f a ∂μ) (by simp) fun _ hs hd =>
    lintegral_iUnion hs hd _
```

[Q5] Mathlib 664cbbe5, Mathlib/MeasureTheory/Measure/WithDensity.lean, lines 36–41, file SHA-256 975237c8f778…

Translation. $(\mu.\text{withDensity}\, f)(S) = \int_S^- f\, d\mu$ for measurable S, where $\int^-$ (Mathlib's `lintegral`) is the lower integral of Section 2.4. The sets S to which it is applied below are open, hence measurable.

Hausdorff dimension.

```
/-- Hausdorff dimension of a set in an (e)metric space. -/
@[irreducible] noncomputable def dimH (s : Set X) : ℝ≥0∞ := by
  borelize X; exact ⨆ (d : ℝ≥0) (_ : @hausdorffMeasure X _ _ (rfl) d s = ∞), d
```

[Q6] Mathlib 664cbbe5, Mathlib/Topology/MetricSpace/HausdorffDimension.lean, lines 96–98, file SHA-256 d525b021e2f4…

Translation. $\dim_{\mathrm{H}} S = \sup\{d \geq 0 :\ \mathcal{H}^d(S) = \infty\}$, computed with the metric of the ambient normed space. This agrees with Section 2.3.

A.2 The statement

The theorem and the six definitions it introduces ~~are in the file~~ occur near the end of the public file:

```
open MeasureTheory
open scoped ENNReal Matrix.Norms.Frobenius

namespace MAIS.ExactFitStatement

/-- Coordinate index family: the vertex `v` carries the coordinate set `Fin (n v)`. -/
abbrev idx {D : ℕ} (n : Fin (D + 1) → ℕ) : Fin (D + 1) → Type :=
  fun v => Fin (n v)

/-- A chain `A = (A_0, …, A_{D-1})`, where `A e` is a real `n (e+1) × n e` matrix.
It unfolds to `(e : Fin D) → Matrix (Fin (n e.succ)) (Fin (n e.castSucc)) ℝ`. -/
abbrev Chain {D : ℕ} (n : Fin (D + 1) → ℕ) : Type :=
  MAIS.QuiverRealOrbit.Rep (idx n)

/-- The product `A_{D-1} ⋯ A_0` (the identity matrix when `D = 0`). -/
def product {D : ℕ} (n : Fin (D + 1) → ℕ) (A : Chain n) :
    Matrix (Fin (n (Fin.last D))) (Fin (n 0)) ℝ :=
  MAIS.MatrixChainIsomorphism.pathMatrix (ι := idx n) A 0 (Fin.last D) (Nat.zero_le D)

/-- `rankTable n A i j` is the rank of `A_{j-1} ⋯ A_i` when `i ≤ j`, and `0` otherwise. -/
def rankTable {D : ℕ} (n : Fin (D + 1) → ℕ) (A : Chain n) (i j : Fin (D + 1)) : ℕ :=
  MAIS.Run2BarcodeEnumeration.pathRankInput (idx n) A i j

/-- The loss `B ↦ ½ ‖P(B) - Φ‖²`, with the Frobenius norm. -/
def halfLoss {D : ℕ} (n : Fin (D + 1) → ℕ)
    (Φ : Matrix (Fin (n (Fin.last D))) (Fin (n 0)) ℝ) (B : Chain n) : ℝ :=
  (1 / 2 : ℝ) * ‖product n B - Φ‖ ^ 2

/-- The Euclidean ball of radius `δ` around `A`, in all matrix entries. -/
def entryBall {D : ℕ} (n : Fin (D + 1) → ℕ) (A : Chain n) (δ : ℝ) : Set (Chain n) :=
  {B | Real.sqrt (∑ e : Fin D, ∑ r : Fin (n e.succ), ∑ s : Fin (n e.castSucc),
      (B e r s - A e r s) ^ 2) < δ}
```

[Q7] ExactFitFull_2026_10_28_UTC.lean, lines 63365–63396, file SHA-256 b362a3ee35df…

Translations.

- `Chain n` is $\mathrm{Rep}(n)$. It unfolds to the type of tuples whose component e is a real $n_{e+1} \times n_e$ matrix, by the definition

```
abbrev Rep (ι : Fin (D + 1) → Type*) :=
  (e : Fin D) → Matrix (ι e.succ) (ι e.castSucc) ℝ
```

[Q8] ExactFitFull_2026_10_28_UTC.lean, lines 788–789, file SHA-256 b362a3ee35df…

- `product n A` is $P(A) = A_{D-1} \cdots A_0$. The function `pathMatrix A i j` is defined through Mathlib's category of composable arrows, but it is pinned down by the three proved lemmas

```
noncomputable def pathMatrix (A : Rep ι) (i j : Fin (d + 1)) (hij : i ≤ j) :
    Matrix (ι j) (ι i) ℝ := LinearMap.toMatrix' ((matrixChain A).map i j hij)

lemma pathMatrix_self (A : Rep ι) (i : Fin (d + 1)) : pathMatrix A i i le_rfl = 1 := by
  rw [pathMatrix, (matrixChain A).map_id]
  exact LinearMap.toMatrix'_id

lemma pathMatrix_comp (A : Rep ι) (i j k : Fin (d + 1)) (hij : i ≤ j) (hjk : j ≤ k) :
    pathMatrix A j k hjk * pathMatrix A i j hij = pathMatrix A i k (hij.trans hjk) := by
  rw [pathMatrix, pathMatrix, ← LinearMap.toMatrix'_comp, (matrixChain A).map_comp]
  rfl

lemma pathMatrix_arrow (A : Rep ι) (e : Fin d) :
    pathMatrix A e.castSucc e.succ (by exact Nat.le_succ e.val) = A e := by
  rw [pathMatrix, matrixChain, ofArrows_map_succ]
  exact LinearMap.toMatrix'_toLin' _
```

[Q9] ExactFitFull_2026_10_28_UTC.lean, lines 4551–4566, file SHA-256 b362a3ee35df…

  which say $M_{ii} = \mathrm{I}$, $M_{jk}M_{ij} = M_{ik}$ and $M_{e,e+1} = A_e$ for $M_{ij} =$ `pathMatrix A i j`. By induction on $j - i$, every family with these three properties equals $(A_{j-1} \cdots A_i)_{i \le j}$. Hence `pathMatrix A i j`$= P_{ij}(A)$ and `product n A`$= P(A)$; for $D = 0$ this is I_{n_0}.

- `rankTable n A i j` is $R_{ij}(A)$ for $i \le j$ and 0 for $i > j$, by

```
def pathRankInput (A : Rep ι) (i j : Fin (D + 1)) : ℕ :=
  if hij : i ≤ j then (pathMatrix A i j hij).rank else 0
```

[Q10] ExactFitFull_2026_10_28_UTC.lean, lines 56724–56725, file SHA-256 b362a3ee35df…

- `halfLoss n Φ B` is $L_\Phi(B) = \frac{1}{2}\|P(B) - \Phi\|_{\mathrm{F}}^2$, and `entryBall n A δ` is the Euclidean ball $B_\delta(A)$ in all entries.

The theorem:

```
/-- **Exact-fit classification.**  Let `A` be a chain with product `Φ`, let `μ` be an
additive Haar measure and `φ` a real function, continuous at `A`, with `φ A > 0`.
Then there are natural numbers `c` and `m` such that

1. the finite program returns exactly `{(c, m)}`;
2. `c` is the local Hausdorff codimension of the fibre `{B | P(B) = Φ}` at `A`
   (ambient dimension minus the infimum over all radii `r > 0` of the Hausdorff
   dimension of the fibre inside the open ball of radius `r`; the balls and the
   Hausdorff dimension are those of the norm `max_e ‖B e‖_F`);
3. for every real `lam` and natural `m'`, the band volume
   `(δ, ε) ↦ ∫_{B ∈ entryBall A δ, |L(B) - L(A)| < ε} φ⁺ dμ` has the two-sided
   power-log order `(lam, m')` (`HasOrder`) if and only if `lam = c / 2` and `m' = m`. -/
theorem exact_fit_classification
    {D : ℕ} (n : Fin (D + 1) → ℕ)
    (μ : Measure (Chain n)) [μ.IsAddHaarMeasure]
    (A : Chain n) (Φ : Matrix (Fin (n (Fin.last D))) (Fin (n 0)) ℝ)
    (hfit : product n A = Φ)
    (φ : Chain n → ℝ) (hφ : ContinuousAt φ A) (hφA : 0 < φ A) :
```

```
    ∃ c m : ℕ,
      MAIS.Run2ArithmeticPositiveData.exactFitPairs n (rankTable n A) = {(c, m)} ∧
      (Module.finrank ℝ (Chain n) : ℝ≥0∞) -
          (⨅ (r : ℝ) (_ : 0 < r),
            dimH ({B : Chain n | product n B = Φ} ∩ Metric.ball A r)) =
        (c : ℝ≥0∞) ∧
      ∀ (lam : ℝ) (m' : ℕ),
        MAIS.DirectBandOrder.HasOrder
            (fun δ ε => μ.withDensity (fun B => ENNReal.ofReal (φ B))
              {B | B ∈ entryBall n A δ ∧ |halfLoss n Φ B - halfLoss n Φ A| < ε})
            lam m' ↔
          (lam = (c : ℝ) / 2 ∧ m' = m) := by
```

[Q11] ExactFitFull_2026_10_28_UTC.lean, lines 63398–63427, file SHA-256 b362a3ee35df…

The predicate `HasOrder` is defined in one block:

```
/-- Two-sided small-loss order at every sufficiently small fixed radius. -/
def HasOrder (volume : _root_.Real → _root_.Real → ENNReal)
    (exponent : _root_.Real) (multiplicity : Nat) : Prop :=
  0 ≤ exponent ∧ 1 ≤ multiplicity ∧
  ∃ radius : _root_.Real, 0 < radius ∧
    ∀ delta : _root_.Real, 0 < delta → delta < radius →
      ∃ lower upper cutoff : _root_.Real,
        0 < lower ∧ lower ≤ upper ∧ 0 < cutoff ∧ cutoff < Real.exp (-1) ∧
        ∀ epsilon : _root_.Real, 0 < epsilon → epsilon < cutoff →
          ENNReal.ofReal lower *
            ENNReal.ofReal (Real.rpow epsilon exponent *
              (Real.log (1 / epsilon)) ^ (multiplicity - 1)) ≤ volume delta epsilon ∧
          volume delta epsilon ≤ ENNReal.ofReal upper *
            ENNReal.ofReal (Real.rpow epsilon exponent *
              (Real.log (1 / epsilon)) ^ (multiplicity - 1))
```

[Q12] ExactFitFull_2026_10_28_UTC.lean, lines 63297–63311, file SHA-256 b362a3ee35df…

Translation of `HasOrder V λ m`. For a function $V : \mathbb{R} \times \mathbb{R} \to [0, +\infty]$: $\lambda \geq 0$, $m \geq 1$, and there is $\delta_0 > 0$ such that for every $\delta \in]0, \delta_0[$ there are real numbers c, C, ε_0 with $0 < c \leq C$ and $\varepsilon_0 \in]0, e^{-1}[$ such that

$$c\,\varepsilon^{\lambda} L(\varepsilon)^{m-1} \leq V(\delta, \varepsilon) \leq C\,\varepsilon^{\lambda} L(\varepsilon)^{m-1} \qquad \text{for all } \varepsilon \in]0, \varepsilon_0[.$$

Here $L(\varepsilon) = \log(1/\varepsilon)$ as in (2.1). Indeed, for $\varepsilon \in]0, e^{-1}[$ the real number $\varepsilon^{\lambda} L(\varepsilon)^{m-1}$ is positive, the natural subtraction $m - 1$ is ordinary because $m \geq 1$, and $\mathrm{ofReal}(c) \cdot \mathrm{ofReal}(t) = ct$ for $c, t \geq 0$.

Translation of the theorem. For all $D \in \mathbb{N}$ and $n \in \mathbb{N}^{D+1}$, every additive Haar measure μ on $\mathrm{Rep}(n)$, every $A \in \mathrm{Rep}(n)$ and $\Phi \in \mathbb{R}^{n_D \times n_0}$ with $P(A) = \Phi$, and every $\varphi : \mathrm{Rep}(n) \to \mathbb{R}$ continuous at A with $\varphi(A) > 0$, there are $c, m \in \mathbb{N}$ such that:

(F1) the formal program returns $\{(c, m)\}$ on the input $(n, R(A))$;

(F2) $N(n) \mathbin{\dot{-}} \inf_{r>0} \dim_{\mathrm{H}}\big(F_\Phi \cap B_r^\infty(A)\big) = c$ in $[0, +\infty]$, where $\dot{-}$ is truncated subtraction and $\dim_{\mathrm{H}}$ uses the norm $\max_e \|\cdot\|_{\mathrm{F}}$;

(F3) for all $\lambda \in \mathbb{R}$ and $m' \in \mathbb{N}$: L_Φ has local volume order (λ, m') at A, for the measure μ, the prior φ and the Euclidean balls, if and only if $\lambda = c/2$ and $m' = m$.

Here (F3) is Definition 3.1 verbatim, with $V^{\mu,\varphi}_{L_\Phi, A}(\delta, \varepsilon) = (\mu.\mathrm{withDensity}\,\varphi^+)(S_{L_\Phi, A}(\delta, \varepsilon))$ and $S_{L_\Phi, A}(\delta, \varepsilon) = \{B \in B_\delta(A) : \ |L_\Phi(B) - L_\Phi(A)| < \varepsilon\}$.

Proposition A.1. *The statement (F1)–(F3), together with Lemma 3.2(b), implies Theorem A. Together moreover with Proposition A.2 below and Lemma 3.13, it implies Theorem B except its last sentence.*

Proof. By (F3) with $(\lambda, m') = (c/2, m)$, L_Φ has a local volume order at A for (μ, φ), and by the converse direction it is unique. In (F2), every subset of $\mathrm{Rep}(n)$ has Hausdorff dimension at most $N(n)$, so the truncated subtraction is ordinary. The infimum is $d_A(F_\Phi)$ (Definition 2.1, which allows any norm). Hence $c = \operatorname{codim}_A F_\Phi$ and $\lambda = c/2 = \frac{1}{2}\operatorname{codim}_A F_\Phi$; since $c \in \mathbb{N}$, $2\lambda \in \mathbb{N}$. Independence of (μ, φ) is part of the quantification. Independence of the norm is not: (F3) uses Euclidean balls, and the other norms are covered by Lemma 3.2(b). For Theorem B, Proposition A.2 identifies the formal program with Prog, and the nonemptiness asserted in Theorem B is Lemma 3.13. The last sentence of Theorem B, $C(h_A) = \min\{\#\,\mathrm{Lk}(k) : \ k \in \mathcal{Z}(n-r),\ h_A \preceq k\}$, does not follow from (F1)–(F3); it is proved in Section 10 from Propositions 8.3 and 5.7. □

The reader who only wants Theorem A can stop here: (F2) and (F3) determine c and m without reference to the program.

A.3 The program

The formal program is `exactFitPairs`. We quote its definitions in dependency order, then translate.

```
def residualDimensions (dims : Fin (D + 1) → ℕ) (r : ℕ) : Fin (D + 1) → ℕ :=
  fun v => dims v - r

def residualRanks (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) (r : ℕ) :
    Fin (D + 1) → Fin (D + 1) → ℕ := fun i j => ranks i j - r

abbrev ResidualCarrier (dims : Fin (D + 1) → ℕ) (r : ℕ) (v : Fin (D + 1)) :=
  Fin (residualDimensions dims r v)

/-- Cardinalities of the literal canonical finite residual carriers. These
are proved equal to the supplied residual dimensions below. -/
def residualArchitecture (dims : Fin (D + 1) → ℕ) (r : ℕ) : Fin (D + 1) → ℕ :=
  fun v => Fintype.card (ResidualCarrier dims r v)

def residualBudget (dims : Fin (D + 1) → ℕ) (r : ℕ) : ℕ :=
  ∑ v, residualArchitecture dims r v

def regularCoefficient (dims : Fin (D + 1) → ℕ) (r : ℕ) : ℕ :=
  r * (dims 0 + dims (Fin.last D) - r)

def residualNodes (dims : Fin (D + 1) → ℕ) (r : ℕ) :
    Finset (Multiplicities D (residualBudget dims r)) :=
  fastCandidates (residualArchitecture dims r) (fun _ _ => 0)

theorem residualArchitecture_eq (dims : Fin (D + 1) → ℕ) (r : ℕ) :
    residualArchitecture dims r = residualDimensions dims r := by
  funext v
  exact Fintype.card_fin _

theorem residualBudget_eq (dims : Fin (D + 1) → ℕ) (r : ℕ) :
    residualBudget dims r = ∑ v, (dims v - r) := by
  simp only [residualBudget, residualArchitecture_eq, residualDimensions]

def residualPairs (dims : Fin (D + 1) → ℕ)
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) (r : ℕ) : Finset (ℕ × ℕ) :=
  computedPairs (residualNodes dims r) (residualRanks ranks r)

/-- The first returned integer is twice the actual exponent, and the second
is its logarithmic multiplicity. Actual correctness and singleton output
are proved separately; no desired measured value enters this program. -/
def exactFitPairs (dims : Fin (D + 1) → ℕ)
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) : Finset (ℕ × ℕ) :=
  if D = 0 then {(0, 1)} else
    (residualPairs dims ranks (ranks 0 (Fin.last D))).image
```

```
      (fun p => (regularCoefficient dims (ranks 0 (Fin.last D)) + p.1, p.2))
```

[Q13] ExactFitFull_2026_10_28_UTC.lean, lines 5560–5604, file SHA-256 b362a3ee35df…

Translation. For $D = 0$ the output is $\{(0, 1)\}$. Otherwise, with $r = R_{0D}$, the output is the image of the *residual program* under $(c_0, m) \mapsto (c_{\mathrm{reg}}(n, r) + c_0, m)$, where c_{reg} is computed with truncated subtraction. On actual inputs $r \le n_0, n_D$, so it is ordinary. The residual program runs on the widths $n \dot{-} r = (n_v \dot{-} r)_v$ and the ranks $(R_{ij} \dot{-} r)_{i \le j}$ (exact on actual inputs, since $r \le n_v$ and $r \le R_{ij}$ for $i \le j$), with the slot bound $M_{\mathrm{slot}} = \sum_v (n_v \dot{-} r)$. Its candidate set is `fastCandidates` for the widths $n \dot{-} r$ with zero lower bounds on the ranks. The theorems `residualArchitecture_eq` and `residualBudget_eq` only record that the cardinalities are the intended numbers.

Barcodes. A barcode is encoded by multiplicities of pairs of endpoints:

```
abbrev Endpoint (D : ℕ) := Fin (D + 2)
abbrev IntervalKey (D : ℕ) := Endpoint D × Endpoint D
abbrev Multiplicities (D M : ℕ) := IntervalKey D → Fin (M + 1)
abbrev Slot {D M : ℕ} (h : Multiplicities D M) := Σ p : IntervalKey D, Fin (h p).val

def birth {D M : ℕ} {h : Multiplicities D M} (s : Slot h) : ℕ := s.1.1.val
def death {D M : ℕ} {h : Multiplicities D M} (s : Slot h) : ℕ := s.1.2.val

def Valid {D M : ℕ} (h : Multiplicities D M) : Prop :=
  ∀ s : Slot h, birth s < death s

def vertexCount {D M : ℕ} (h : Multiplicities D M) (v : Fin (D + 1)) : ℕ :=
  Fintype.card {s : Slot h // birth s ≤ v.val ∧ v.val < death s}

def pathCount {D M : ℕ} (h : Multiplicities D M) (i j : Fin (D + 1)) : ℕ :=
  Fintype.card {s : Slot h // birth s ≤ i.val ∧ j.val < death s}

def Linked {D M : ℕ} {h : Multiplicities D M} (r s : Slot h) : Prop :=
  birth s < birth r ∧ birth r ≤ death s ∧ death s < death r
instance linkedDecidable {D M : ℕ} {h : Multiplicities D M} (r s : Slot h) :
    Decidable (Linked r s) := inferInstanceAs (Decidable (_ ∧ _ ∧ _))

def incoming {D M : ℕ} {h : Multiplicities D M} (r : Slot h) : ℕ :=
  Fintype.card {s : Slot h // Linked r s}
def outgoing {D M : ℕ} {h : Multiplicities D M} (s : Slot h) : ℕ :=
  Fintype.card {r : Slot h // Linked r s}
def Balanced {D M : ℕ} (h : Multiplicities D M) : Prop :=
  ∀ s : Slot h, birth s ≠ 0 → death s ≠ D + 1 → incoming s = outgoing s

def outputCount {D M : ℕ} (h : Multiplicities D M) : ℕ :=
  Fintype.card {p : Slot h × Slot h // Linked p.1 p.2 ∧ death p.1 = D + 1}

/-- Actual finite arithmetic feasibility for a prescribed architecture and
prescribed integer path-rank input. -/
def Feasible {D M : ℕ} (dims : Fin (D + 1) → ℕ)
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) (h : Multiplicities D M) : Prop :=
  Valid h ∧ (∀ v, vertexCount h v = dims v) ∧
  pathCount h 0 (Fin.last D) = 0 ∧
  ∀ i j, i ≤ j → ranks i j ≤ pathCount h i j
```

[Q14] ExactFitFull_2026_10_28_UTC.lean, lines 307–345, file SHA-256 b362a3ee35df…

Dictionary. A key `(b, d)` with $b < d$ (`Valid`) is the interval $[\![b, d-1]\!]$, i.e. `birth`$= b_s$ and `death`$= e_s + 1$. A `Slot h` is a pair (key, copy index), i.e. an element of $\mathcal{S}(h)$. Then:

- `vertexCount h v` is $\dim_v(h)$ and `pathCount h i j` is $\rho_{ij}(h)$ (Definition 3.6), since $b \le i$ and $j < d$ means $[\![i, j]\!] \subseteq [\![b, d-1]\!]$.
- `Linked r s` is $s \rightarrowtail r$, with the first argument the *target*: $b_s < b_r \le e_s + 1 \le e_r$.

- `incoming r` is $\mathrm{in}(r)$ and `outgoing s` is $\mathrm{out}(s)$.
- `Balanced` quantifies over interior slots (`birth`$\neq 0$ and `death`$\neq D+1$, i.e. $b_s \neq 0$ and $e_s \neq D$).
- `outputCount h` counts pairs (t, s) with $s \rightarrowtail t$ and $e_t = D$, i.e. $c_{\mathrm{out}}(h)$.
- `Feasible dims ranks h` says: valid, dimension vector `dims`, no full-span slot ($\rho_{0D}(h) = 0$), and $\texttt{ranks}_{ij} \leq \rho_{ij}(h)$ for $i \leq j$.

Multiplicities are bounded by M_{slot} (`Fin (M + 1)`, where `M` is M_{slot}). This loses nothing: a barcode with dimension vector $n \dot{-} r$ has at most $\sum_v (n_v \dot{-} r) = M_{\mathrm{slot}}$ slots.

Candidates.

```
abbrev ValidKey (D : ℕ) := {p : IntervalKey D // p.1.val < p.2.val}
abbrev Assignment (D M : ℕ) := Fin M → Option (ValidKey D)

def histogram {D M : ℕ} (f : Assignment D M) : Multiplicities D M := fun p =>
  (Fintype.card {i : Fin M // Option.map Subtype.val (f i) = some p}, by
    have h := Fintype.card_le_of_injective
      (Subtype.val : {i : Fin M // Option.map Subtype.val (f i) = some p} → Fin M)
      Subtype.val_injective
    simpa only [Fintype.card_fin] using Nat.lt_succ_of_le h)

def generated {D M : ℕ} : Finset (Multiplicities D M) :=
  Finset.univ.image histogram

def fastCandidates {D M : ℕ} (dims : Fin (D + 1) → ℕ)
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) : Finset (Multiplicities D M) :=
  (Finset.univ.filter (fun f : Assignment D M => Feasible dims ranks (histogram f))).image histogram
```

[Q15] `ExactFitFull_2026_10_28_UTC.lean`, lines 5478–5493, file SHA-256 `b362a3ee35df…`

Translation. The candidate set is the set of histograms of maps $[\![0, M_{\mathrm{slot}} - 1]\!] \to \{\text{valid keys}\} \cup \{\text{none}\}$ that are feasible. Every barcode with at most M_{slot} slots is such a histogram. Hence, with zero lower bounds, the candidate set is exactly $\mathcal{Z}(n - r)$.

The coefficient, the threshold, the weight and the edges.

```
def rankInput (h : Multiplicities D M) (i j : Fin (D + 1)) : ℕ :=
  if i ≤ j then pathCount h i j else 0

def Matches (h : Multiplicities D M)
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) : Prop :=
  ∀ i j : Fin (D + 1), i ≤ j → pathCount h i j = ranks i j

instance matchesDecidable (h : Multiplicities D M)
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) : Decidable (Matches h ranks) := by
  unfold Matches
  infer_instance

def RankLE (h k : Multiplicities D M) : Prop :=
  ∀ i j : Fin (D + 1), i ≤ j → pathCount h i j ≤ pathCount k i j

def RankLT (h k : Multiplicities D M) : Prop :=
  RankLE h k ∧ ∃ i j : Fin (D + 1), i ≤ j ∧ pathCount h i j < pathCount k i j

instance rankLEDecidable (h k : Multiplicities D M) : Decidable (RankLE h k) := by
  unfold RankLE
  infer_instance
instance rankLTDecidable (h k : Multiplicities D M) : Decidable (RankLT h k) := by
  unfold RankLT
  infer_instance

/-- Independently enumerated balanced-output objective above a given array.
```

```
The supplied list may be replaced by any proved-equal faster enumeration. -/
def higherOutputValues (ν : Finset (Multiplicities D M)) (h : Multiplicities D M) : Finset ℕ :=
  (ν.filter (fun k => RankLE h k ∧ Balanced k)).image outputCount

/-- Total arithmetic coefficient. Off the feasible domain an empty list
returns zero; the actual measured adapter proves nonemptiness on-domain. -/
def coefficient (ν : Finset (Multiplicities D M)) (h : Multiplicities D M) : ℕ :=
  (higherOutputValues ν h).min.untopD 0

/-- Exact rational canonical potential threshold, before any real cast. -/
def threshold (h : Multiplicities D M) : ℚ :=
  (∑ p : MAIS.IntervalCochainProduct.NormalIndex
      (@birth D M h) (@death D M h),
    MAIS.IntervalPositivePotentials.edgeWeight D (@birth D M h) (@death D M h) p) / 2

/-- A critical source contributes one precisely at exact rational equality.
This equality test never compares arbitrary real numbers. -/
def weight (ν : Finset (Multiplicities D M)) (h : Multiplicities D M) : ℕ :=
  if (coefficient ν h : ℚ) / 2 = threshold h then 1 else 0

def Edge (ν : Finset (Multiplicities D M)) (h k : Multiplicities D M) : Prop :=
  RankLT h k ∧ coefficient ν h = coefficient ν k
```

[Q16] ExactFitFull_2026_10_28_UTC.lean, lines 5328–5375, file SHA-256 b362a3ee35df…

```
def Linked (r s : S) : Prop :=
  birth s < birth r ∧ birth r ≤ death s ∧ death s < death r

instance linked_decidable (r s : S) : Decidable (Linked birth death r s) := by
  unfold Linked
  infer_instance

abbrev NormalIndex := {p : S × S // Linked birth death p.1 p.2}
abbrev NormalParams := NormalIndex birth death → ℝ
```

[Q17] ExactFitFull_2026_10_28_UTC.lean, lines 4817–4825, file SHA-256 b362a3ee35df…

```
variable {S : Type*} (D : ℕ) (birth death : S → ℕ)

def denominator (s : S) : ℕ := birth s + (D + 1 - death s)

/-- Rational interval potential, before clearing any finite denominators. -/
def potential (s : S) : ℚ := (birth s : ℚ) / (denominator D birth death s : ℚ)

def edgeWeight (p : NormalIndex birth death) : ℚ :=
  potential D birth death p.val.1 - potential D birth death p.val.2
```

[Q18] ExactFitFull_2026_10_28_UTC.lean, lines 5009–5017, file SHA-256 b362a3ee35df…

Translation. For h, k in the candidate set ν:

- `RankLE h k` is $h \preceq k$ (all candidates have the same dimension vector), and `RankLT h k` is $h \prec k$.
- `coefficient ν h` is $\min\{c_{\text{out}}(k) : k \in \nu,\ h \preceq k,\ k \text{ balanced}\}$, with the value 0 if this set is empty.
- `potential` is the canonical potential $u_s = b_s/(b_s + D - e_s)$, since $D + 1 - \text{death} = D - e_s$. `edgeWeight` of a linked pair $s \rightarrowtail s'$ is $u_{s'} - u_s$, so `threshold h` is $T(u)$ for the canonical potential, as an exact rational.
- `weight ν h` is 1 if coefficient$/2 = T(u)$, and 0 otherwise.
- `Edge ν h k` means $h \prec k$ and equal coefficients.

The longest weighted path and the output.

```
abbrev Node (ν : Finset (Multiplicities D M)) := ↥ν

def nodeEdge (ν : Finset (Multiplicities D M)) (h k : Node ν) : Prop := Edge ν h.val k.val
instance nodeEdgeDecidable (ν : Finset (Multiplicities D M)) : DecidableRel (nodeEdge ν) := by
  intro h k
  exact edgeDecidable ν h.val k.val

def nodeWeight (ν : Finset (Multiplicities D M)) (h : Node ν) : ℕ := weight ν h.val

abbrev PathIndex (D : ℕ) := {p : Fin (D + 1) × Fin (D + 1) // p.1 ≤ p.2}

/-- Strict rank-order edges strictly increase this literal finite sum. -/
def height (h : Multiplicities D M) : ℕ :=
  ∑ p : PathIndex D, pathCount h p.val.1 p.val.2

def capacity (D M : ℕ) : ℕ := Fintype.card (PathIndex D) * M

def defect (h : Multiplicities D M) : ℕ := capacity D M - height h

/-- A fully executable finite-arithmetic longest-path candidate. Its fuel
is an architecture-independent finite bound for all feasible arrays. -/
def criticalDepth (ν : Finset (Multiplicities D M)) (h : Node ν) : ℕ :=
  MAIS.Run2ArithmeticCriticalPathDynamic.depthFuel (nodeEdge ν) (nodeWeight ν)
    (capacity D M) h

/-- Complete proof-free algorithm output from a supplied integer rank
table: all coefficient/multiplicity pairs at matching finite sources.
Actual correctness later proves this output is a nonempty singleton. -/
def computedPairs (ν : Finset (Multiplicities D M))
    (ranks : Fin (D + 1) → Fin (D + 1) → ℕ) : Finset (ℕ × ℕ) :=
  (ν.attach.filter (fun h => Matches h.val ranks)).image
    (fun h => (coefficient ν h.val, 1 + criticalDepth ν h))
```

[Q19] `ExactFitFull_2026_10_28_UTC.lean`, lines 5384–5415, file SHA-256 `b362a3ee35df…`

```
/-- Maximum source-weight sum over paths using at most the supplied number
of edges. Stopping at the current vertex always has cost zero. -/
def depthFuel (edge : J → J → Prop) [DecidableRel edge]
    (weight : J → ℕ) : ℕ → J → ℕ
  | 0, _ => 0
  | n + 1, i => Finset.univ.sup (fun j =>
      if edge i j then weight i + depthFuel edge weight n j else 0)
```

[Q20] `ExactFitFull_2026_10_28_UTC.lean`, lines 5226–5232, file SHA-256 `b362a3ee35df…`

Translation. `depthFuel edge w k i` is the largest value of $w(i_0) + \cdots + w(i_{l-1})$ over paths $i = i_0 \to i_1 \to \cdots \to i_l$ along edges with $l \le k$. Stopping contributes 0, so the last vertex is not weighted. `computedPairs ν R'` is the set of pairs (coefficient(h), $1 +$ criticalDepth(h)) over the $h \in \nu$ whose rank table is R'.

Proposition A.2. *Let $D \ge 1$ and $A \in \mathrm{Rep}(n)$. The formal program on the input $(n, R(A))$ returns* $\mathsf{Prog}(n, R(A)) = \{(c_{\mathrm{reg}}(n, r) + C(h_A),\ 1 + F(h_A))\}$ *(Section 3).*

Proof. Put $r = R_{0D}(A)$. By the translations above, the candidate set is $\mathcal{Z}(n-r)$, and the only candidate h with $\rho(h) = (R_{ij}(A) - r)_{i \le j}$ is h_A, by (3.2) and Fact 3.8(c). The coefficient is C, because the minimized set is nonempty (Lemma 3.13, a combinatorial statement). The weight of h is 1 if and only if h is critical: if h is balanced then $2T(u) = c_{\mathrm{out}}(h)$ by (8.1); if not, then $C(h) \le C^{\sqsubseteq}(h) = 2\lambda_h < 2T(u)$. Here the inequality holds because the set minimized in $C^{\sqsubseteq}(h)$ is contained in the one minimized in $C(h)$, by Proposition 5.7(a); the equality is Theorem 8.2; and the strict inequality is Lemma 8.1, since the canonical potential is admissible (Section 7). Along an edge, $\sum_{i \le j} \rho_{ij}$ strictly increases and stays at most $\#\{(i,j) :\ 0 \le i \le j \le D\} \cdot M_{\mathrm{slot}}$,

so the fuel bounds the length of every path and `criticalDepth` is the unbounded maximum $F(h_A)$. □

Besides the translations, the proof of Proposition A.2 uses combinatorial facts proved in the text (Lemma 3.13, (3.2), Fact 3.8(c), (8.1) and the bound on the fuel), and, in one place, three geometric results of the text: Proposition 5.7(a), Theorem 8.2 and Lemma 8.1. They serve only to see that the formal weight agrees with "critical" for unbalanced barcodes. A reader who prefers not to rely on them can read Theorem B with "critical" replaced by the formal test $C(h) = 2T(u)$, where u is the canonical potential of h. Then (F1)–(F3), the translations of this appendix and these combinatorial facts certify the displayed formulas of Theorem B.

B The rank order is generated by switches

Proposition B.1. *Let $D \in \mathbb{N}^*$, and let $h \preceq k$ be barcodes of length D. There is a finite sequence $h = h_0, h_1, \dots, h_N = k$ such that each h_{t+1} is obtained from h_t by the switch of Lemma 5.5 along a linked pair of h_t.*

The statement is purely combinatorial; for its use see Proposition 5.7(b). Over algebraically closed fields the equality of the rank order and the degeneration order is due to Abeasis, Del Fra and Kraft [1]. The proof below reduces to the Bruhat order of the symmetric group by "standardization".

Padded matrices

Fix $D \in \mathbb{N}^*$ and $N_0 \in \mathbb{N}$. For a barcode h of length D and $a \in [\![0, D]\!]$ let $\mathrm{st}_a(h) = \#\{s \in \mathcal{S}(h) : \ b_s = a\}$ be the number of slots starting at a, and assume $\mathrm{st}_a(h) \le N_0$ for all a. Define the $(D+2) \times (D+2)$ matrix $X = X(h)$, with rows and columns indexed by $[\![0, D+1]\!]$, by

$$X(b,c) = h([\![b, c-1]\!]) \ (b < c), \qquad X(a,a) = N_0 - \mathrm{st}_a(h) \ (a \le D), \qquad X(D+1, D+1) = N_0,$$

and $X(b,c) = 0$ for $c < b$. So the number of slots with interval $[\![b, c-1]\!]$ is recorded in position (b,c), and the diagonal entries are "padding". Every row sum of X is N_0. Let $n = \dim(h)$ and put $n_{-1} = n_{D+1} = 0$. The column sums are

$$\kappa_c = N_0 - (n_c - n_{c-1}) \quad (c \in [\![0, D]\!]), \qquad \kappa_{D+1} = N_0 + n_D,$$

because $\mathrm{st}_c(h) - \#\{s \in \mathcal{S}(h) : \ e_s = c-1\} = n_c - n_{c-1}$. For $p, q \in [\![0, D+1]\!]$ put

$$\mathrm{NE}_M(p,q) = \sum_{b \le p, \ c \ge q} M(b,c)$$

for any matrix M. Then:

(a) $\mathrm{NE}_X(i, j+1) = \rho_{ij}(h)$ for $0 \le i \le j \le D$. The diagonal entries do not contribute, since $a \le i < j+1 \le a$ is impossible.

(b) $\mathrm{NE}_X(p,q) = \sum_{c \ge q} \kappa_c - \sum_{b > p} N_0$ for $p \ge q$, since every entry with $b > p \ge q$ has $c \ge b > p$.

Hence, for barcodes h, k of length D with the same dimension vector (and $N_0 \ge \mathrm{st}_a(h), \mathrm{st}_a(k)$ for all a), $X(h)$ and $X(k)$ have the same row and column sums, and $h \preceq k$ if and only if $\mathrm{NE}_{X(h)} \le \mathrm{NE}_{X(k)}$ entrywise.

Moves

For $b, c \in [\![0, D+1]\!]$ let E_{bc} be the $(D+2)\times(D+2)$ matrix with a single nonzero entry, equal to 1, at (b, c). For a $(D+2)\times(D+2)$ matrix M with nonnegative integer entries, indices $b_1 < b_2$, $c_1 < c_2$ with $M(b_1, c_1) \geq 1$ and $M(b_2, c_2) \geq 1$, the *move*

$$M' = M - E_{b_1c_1} - E_{b_2c_2} + E_{b_1c_2} + E_{b_2c_1}$$

satisfies $\mathrm{NE}_{M'}(p, q) = \mathrm{NE}_M(p, q) + 1$ if $b_1 \leq p < b_2$ and $c_1 < q \leq c_2$, and $\mathrm{NE}_{M'}(p, q) = \mathrm{NE}_M(p, q)$ otherwise. This follows from $([b_1 \leq p] - [b_2 \leq p])([c_2 \geq q] - [c_1 \geq q]) \geq 0$.

Lemma B.2. *Let $M = X(h)$, and let M' be the move of M with indices $b_1 < b_2$ and $c_1 < c_2$. Suppose that M' is upper triangular ($M'(b, c) = 0$ for $c < b$). Then $b_1 < b_2 \leq c_1 < c_2$, the intervals $I = [\![b_1, c_1 - 1]\!]$ and $I' = [\![b_2, c_2 - 1]\!]$ satisfy $h(I) \geq 1$ and $h(I') \geq 1$, every pair of slots (s, s') with $I_s = I$ and $I_{s'} = I'$ is linked, and $M' = X(h^\pi)$ for $\pi = (s, s')$ and the switched barcode h^π of Lemma 5.5.*

Proof. Since $M'(b_2, c_1) \geq 1$ and M' is upper triangular, $b_2 \leq c_1$. Hence $b_1 < b_2 \leq c_1 < c_2$, so (b_1, c_1) and (b_2, c_2) are strictly above the diagonal, and $h(I) = M(b_1, c_1) \geq 1$, $h(I') = M(b_2, c_2) \geq 1$. For slots s, s' with $I_s = I$ and $I_{s'} = I'$, the linking condition $b_s < b_{s'} \leq e_s + 1 \leq e_{s'}$ reads $b_1 < b_2 \leq c_1 \leq c_2 - 1$, which holds. The move adds 1 at (b_1, c_2), i.e. to the multiplicity of $[\![b_1, c_2 - 1]\!] = [\![b_s, e_{s'}]\!]$, and 1 at (b_2, c_1): this is the multiplicity of $[\![b_2, c_1 - 1]\!] = [\![b_{s'}, e_s]\!]$ if $b_2 < c_1$, and the padding entry at (b_2, b_2) if $b_2 = c_1$, i.e. if $b_{s'} = e_s + 1$. In the latter case st_{b_2} decreases by one, consistently with the padding. Comparing with the definition of h^π in Lemma 5.5 gives $M' = X(h^\pi)$. □

Lemma B.3. *Let $h \preceq k$ be barcodes of length D with dimension vector n, let $N_0 \geq \mathrm{st}_a(h), \mathrm{st}_a(k)$ for all a, and put $X = X(h)$, $Y = X(k)$. If a $(D+2)\times(D+2)$ matrix M has nonnegative integer entries, the same row and column sums as X, and $\mathrm{NE}_X \leq \mathrm{NE}_M \leq \mathrm{NE}_Y$, then $M = X(h'')$ for a barcode h'' with $h \preceq h'' \preceq k$.*

Proof. For $c \in [\![0, D]\!]$,

$$\mathrm{NE}_M(c, c+1) = \sum_{b \leq c} N_0 - \sum_{c' \leq c} \kappa_{c'} + \sum_{b > c,\ c' \leq c} M(b, c').$$

The left side is squeezed between $\mathrm{NE}_X(c, c+1) = n_c = \mathrm{NE}_Y(c, c+1)$, and the last sum vanishes for X. Hence $M(b, c') = 0$ whenever $b > c'$ (take $c = c'$), so M is upper triangular. Define $h''([\![b, c-1]\!]) = M(b, c)$ for $b < c$. Row sums give $M(a, a) = N_0 - \mathrm{st}_a(h'')$, so $M = X(h'')$; the equalities $\mathrm{NE}_M(c, c+1) = n_c$ give $\dim(h'') = n$, and (a) gives $h \preceq h'' \preceq k$. □

Permutations

Let $N_1 = (D+2)N_0$. Index rows by pairs (b, α), $b \in [\![0, D+1]\!]$, $\alpha \in [\![1, N_0]\!]$, and columns by pairs (c, γ), $\gamma \in [\![1, \kappa_c]\!]$, both ordered lexicographically and identified with $[\![1, N_1]\!]$. A permutation w of $[\![1, N_1]\!]$ is identified with the 0/1 matrix having ones at $(a, w(a))$. Its *projection* $\mathrm{pr}(w)$ is the $(D+2)\times(D+2)$ matrix counting the ones in each block (rows of b)$\times$(columns of c). Put $w[p, q] = \#\{a \leq p :\ w(a) \geq q\}$. By [5, Thm. 2.1.5], the Bruhat order satisfies

$$u \leq w \iff u[p, q] \leq w[p, q] \text{ for all } p, q. \tag{B.1}$$

By definition [5, Def. 2.1.1], $u < w$ means that there is a chain $u = u_0, u_1, \ldots, u_\ell = w$ where each u_{t+1} is obtained from u_t by exchanging the values at two positions $a < a'$ with

$u_t(a) < u_t(a')$. The projection is monotone: $\mathrm{NE}_{\mathrm{pr}(w)}(b,c) = w[p_b, q_c]$ with p_b the last row of block b and q_c the first column of block c. Each step $u_t \to u_{t+1}$ projects either to an equality (if a, a' lie in the same row block or $u_t(a), u_t(a')$ in the same column block) or to a move.

Lemma B.4 (Lifting)**.** *Let X, Y have nonnegative integer entries, row sums N_0 and column sums κ, with $\mathrm{NE}_X \le \mathrm{NE}_Y$. Then there are permutations $u \le w$ with $\mathrm{pr}(u) = X$ and $\mathrm{pr}(w) = Y$.*

Proof. Rows. Let $X^{\uparrow}$ be the $N_1 \times (D+2)$ matrix with one unit in each row, obtained by giving the N_0 rows of block b the N_0 units of row b of X in increasing order of column. Let $Y^{\downarrow}$ be defined likewise from Y, in decreasing order. Put $\mathrm{NE}_Z(p,c) = \#\{\text{units at rows} \le p,\ \text{columns} \ge c\}$. Fix c and b, and for $\alpha \in [\![0, N_0]\!]$ let $p = (b, \alpha)$ denote the α-th row of block b ($\alpha = 0$: the end of block $b-1$). With x (resp. y) the number of units in row b of X (resp. Y) in columns $\ge c$,

$$\mathrm{NE}_{X^{\uparrow}}((b,\alpha),c) = \mathrm{NE}_X(b-1,c) + \max(0, \alpha - N_0 + x),$$
$$\mathrm{NE}_{Y^{\downarrow}}((b,\alpha),c) = \mathrm{NE}_Y(b-1,c) + \min(\alpha, y).$$

The first is convex and the second concave in α. At $\alpha = 0$ they equal $\mathrm{NE}_X(b-1,c) \le \mathrm{NE}_Y(b-1,c)$, and at $\alpha = N_0$ they equal $\mathrm{NE}_X(b,c) \le \mathrm{NE}_Y(b,c)$. A convex function lies below its chord and a concave one above it, so the first is $\le$ the second for all $\alpha \in [\![0, N_0]\!]$ (the right-hand sides of the two displays make sense for real $\alpha \in [0, N_0]$, where they are respectively convex and concave).

Columns. Let u be obtained from $X^{\uparrow}$ by giving the κ_c columns of block c the units of column c in increasing order of row, and w from $Y^{\downarrow}$ in decreasing order of row. These are permutation matrices with $\mathrm{pr}(u) = X$ and $\mathrm{pr}(w) = Y$. Fix a row p and a column block c. With z (resp. z') the number of units of column c of $X^{\uparrow}$ (resp. $Y^{\downarrow}$) in rows $\le p$, and $\gamma \in [\![1, \kappa_c + 1]\!]$ (where $(c, \kappa_c + 1)$ stands for the first column of block $c+1$),

$$u[p, (c,\gamma)] = \mathrm{NE}_{X^{\uparrow}}(p, c+1) + \max(0, z - \gamma + 1),$$
$$w[p, (c,\gamma)] = \mathrm{NE}_{Y^{\downarrow}}(p, c+1) + \min(z', \kappa_c - \gamma + 1).$$

The right-hand sides, extended to real $\gamma \in [1, \kappa_c + 1]$, are respectively convex and concave in γ. At $\gamma = 1$ they equal $\mathrm{NE}_{X^{\uparrow}}(p,c)$ and $\mathrm{NE}_{Y^{\downarrow}}(p,c)$, and at $\gamma = \kappa_c + 1$ they equal the values at $c+1$. The row step gives the inequality at both endpoints, hence everywhere. By (B.1), $u \le w$. □

Proof of Proposition B.1. Choose $N_0 \ge \max_a \max(\mathrm{st}_a(h), \mathrm{st}_a(k))$, put $X = X(h)$ and $Y = X(k)$, and take $u \le w$ from Lemma B.4. Let $u = u_0, \dots, u_\ell = w$ be a chain as above, and $M_t = \mathrm{pr}(u_t)$. By monotonicity $X = M_0$, $\mathrm{NE}_X \le \mathrm{NE}_{M_t} \le \mathrm{NE}_Y$, and $M_\ell = Y$. By Lemma B.3, $M_t = X(h_t)$ for barcodes h_t with $h \preceq h_t \preceq k$. Consecutive matrices are equal or differ by a move between upper triangular matrices, which by Lemma B.2 is a switch along a linked pair. Deleting repetitions gives the required sequence. □

The formal development contains an independent Lean proof of the same statement, also by reduction to permutations (`MAIS.LinearChain.Barcode.actual_rank_order_generated`), and of the resulting Euclidean closure order over $\mathbb{R}$ (`MAIS.RealTypeAClosure.mem_closure_orbit_iff_path_rank_le`).

Acknowledgements

To be written by the author. The geometric viewpoint on fibres of the multiplication map, and the global identification of the threshold with half the codimension, are due to Simon Pepin Lehalleur and Richárd Rimányi [24]; the global coefficients of deep linear networks are due to Miki Aoyagi [3]. The question O77 is from the Math for AI Safety agenda [29].

References

[1] S. Abeasis, A. Del Fra, H. Kraft, *The geometry of representations of A_m*, Math. Ann. **256** (1981), 401–418.

[2] M. Aoyagi, S. Watanabe, *Stochastic complexities of reduced rank regression in Bayesian estimation*, Neural Networks **18** (2005), 924–933.

[3] M. Aoyagi, *Consideration on the learning efficiency of multiple-layered neural networks with linear units*, Neural Networks **172** (2024), 106132.

[4] E. Bierstone, P. D. Milman, *Semianalytic and subanalytic sets*, Publ. Math. IHÉS **67** (1988), 5–42.

[5] A. Björner, F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics **231**, Springer, 2005.

[6] W. Crawley-Boevey, *Decomposition of pointwise finite-dimensional persistence modules*, J. Algebra Appl. **14** (2015), 1550066.

[7] C. De Concini, C. Procesi, *Wonderful models of subspace arrangements*, Selecta Math. (N.S.) **1** (1995), 459–494.

[8] K. Falconer, *Fractal Geometry: Mathematical Foundations and Applications*, 3rd ed., Wiley, 2014.

[9] N. Feld, *MAIS-O77: learning coefficients on fibers and saddles*, blueprint, lectures and Lean development, October 2026, `https://github.com/nfeld/mais-o77-lean`.

[10] N. Feld, *Deep linear exact fits: Lean development and verification materials*, standalone proof `ExactFitFull_2026_10_28_UTC.lean`, SHA-256 beginning `b362a3ee35df`, October 2026, `https://github.com/nfeld/deep-linear-exact-fits`.

[11] K. Ferry, A. Monod, K. Yano, *Singular geometry of low-rank adaptation*, arXiv:2610.32854 (2026).

[12] Z. Furman, E. Lau, *Estimating the local learning coefficient at scale*, arXiv:2402.03698v1 (2024); withdrawn and expanded into [21].

[13] P. Gabriel, *Unzerlegbare Darstellungen I*, Manuscripta Math. **6** (1972), 71–103.

[14] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, 1978.

[15] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero: I, II*, Ann. of Math. **79** (1964), 109–203, 205–326.

[16] R. Hitchcock, J. Hoogland, *From global to local: a scalable benchmark for local posterior sampling*, arXiv:2507.21449v2 (2026).

[17] J. Koncki, R. Rimányi, *The main reasons for matrices multiplying to zero*, arXiv:2503.13326 (2025).

[18] D. Kosta, D. Windisch, *Classification of real hyperplane singularities by real log canonical thresholds*, arXiv:2411.13392v2 (2026).

[19] Y. Kurumadani, *Real log canonical thresholds at non-singular points*, arXiv:2408.13030 (2024).

[20] Y. Kurumadani, *Upper bounds for local learning coefficients of three-layer neural networks*, arXiv:2603.12785v2 (2026).

[21] E. Lau, Z. Furman, G. Wang, D. Murfet, S. Wei, *The local learning coefficient: a singularity-aware complexity measure*, arXiv:2308.12108v2 (2024).

[22] J. M. Lee, *Introduction to Smooth Manifolds*, 2nd ed., Graduate Texts in Mathematics **218**, Springer, 2013.

[23] S. Lin, *Ideal-theoretic strategies for asymptotic approximation of marginal likelihood integrals*, J. Algebraic Stat. **8** (2017), no. 1, 22–55; arXiv:1003.5338v3.

[24] S. Pepin Lehalleur, R. Rimányi, *Geometry of fibers of the multiplication map of deep linear neural networks*, arXiv:2411.19920v2 (2024).

[25] L. Levine, *Math for AI safety: an invitation for mathematicians*, arXiv:2609.15289 (2026). An August 2026 revision was distributed in [30] as the paper MAIS-P1.

[26] R. Sneiderman, *MAIS-O70: candidate complete solution, local RLCT table for reduced-rank regression*, issue #3 of the MAIS repository, 3 August 2026, `https://github.com/Math-for-AI-Safety/MAIS/issues/3`.

[27] S. Novikov (with OpenAI Codex), *MAIS-O77: candidate complete solution — fiber rank strata and the universal saddle pair*, issue #12 of the MAIS repository, 4 August 2026, `https://github.com/Math-for-AI-Safety/MAIS/issues/12`.

[28] Math for AI Safety, *Problem MAIS-O70: local learning coefficients of reduced-rank regression*, `https://github.com/Math-for-AI-Safety/MAIS/blob/main/open-problems/MAIS-O70.md`.

[29] Math for AI Safety, *Problem MAIS-O77: learning coefficients on the matrix factorization fiber and saddles*, `https://github.com/Math-for-AI-Safety/MAIS/blob/main/open-problems/MAIS-O77.md`.

[30] Math for AI Safety, *Open problems and agendas*, `https://github.com/Math-for-AI-Safety/MAIS`.

[31] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge University Press, 1995.

[32] M. Mustaţă, *Multiplier ideals of hyperplane arrangements*, Trans. Amer. Math. Soc. **358** (2006), 5015–5023.

[33] M. Saito, *On real log canonical thresholds*, arXiv:0707.2308 (2007).

[34] J. R. Shewchuk, S. Bhattacharya, *The geometry of the set of equivalent linear neural networks*, arXiv:2404.14855v2 (2026); see also *The hierarchy of manifolds in a stratification of the set of equivalent linear neural networks*, in: 42nd International Symposium on Computational Geometry (SoCG 2026), LIPIcs **367** (2026), 91:1–91:19.

[35] Z. Teitler, *A note on Mustaţă's computation of multiplier ideals of hyperplane arrangements*, Proc. Amer. Math. Soc. **136** (2008), 1575–1579.

[36] D. Voigt, *Induzierte Darstellungen in der Theorie der endlichen, algebraischen Gruppen*, Lecture Notes in Mathematics **592**, Springer, 1977.

[37] S. Watanabe, *Algebraic Geometry and Statistical Learning Theory*, Cambridge University Press, 2009.

[38] S. Watanabe, *Mathematical Theory of Bayesian Statistics*, CRC Press, 2018.