

A
Lytic Reactivation Is Intrinsic Across Thaws (70 Pairs, 24 Donors)
Same-batch technical replicates (n = 24 pairs)
Different-batch independent thaws (n = 46 pairs)
Pairwise Replicate Correlation (r)
0.96
0.95
0.97
0.91
0.95
0.81
0.97
0.76
0.95
0.68
0.96
0.64
0.86
0.24
BGLF5 (Shutoff)
log2(IE / Latent)
Lytic IE RPKM
log2(Late / Latent)
Lytic Early RPKM
Lytic Late RPKM
Latent Pol II mRNA
B
Decisive H1 Test: CN Tracks Lytic IE, Not Latent Dosage (n = 717)
2023 MAGE Viral Transcript log2 (RPKM + 0.01)
Latent Pol II mRNA (EBNA/LMP): r = -0.0286 (null)
Lytic Immediate-Early (BZLF1/BRLF1): r = +0.6402 (p = 6.2e-84)
2019 30x WGS log2 (EBV Copy Number)
C
Invariance of Lytic–CN Coupling to Population & Batch
Marginal (n = 717)
Within 26 populations
Within 26 pops × 17 batches
Correlation (r) with WGS log2 (EBV CN)
+0.65
+0.63
+0.62
+0.61
+0.61
-0.06
Spliced BZLF1 (Zta) RPM
Lytic IE RPKM
log2(Late / Latent)
Lytic Late RPKM
Lytic Early RPKM
Latent Pol II mRNA RPKM
D
Exclusively Latency III (Cp/Wp) & Zero Latency I (Qp) Splicing
Wp (W1–W2) IR1 Repeat
7,732.4 reads (59/59 runs)
BART Exons I–VII
4,845.0 reads (59/59 runs)
LMP-1 Introns a/b
4,481.7 reads (59/59 runs)
LMP-2A/2B Exons 2–8
2,003.1 reads (59/59 runs)
Cp (C1–C2–W1) Latency III
1,090.5 reads (59/59 runs)
W2 -> Y1/Y2 (EBNA-2/LP)
465.6 reads (59/59 runs)
W2/Y -> BHRF1 (Latent vBCL2)
142.8 reads (59/59 runs)
Cp/Wp U -> K (Latent III EBNA1)
79.4 reads (59/59 runs)
Qp -> U -> K (Latency I = 0/59)
0.0 reads (0/59 runs)
log10 (Mean Spliced Junction Reads + 1)