



# Mathematical Fundamentals



# Contents

1	Set Theory	5
1.1	Concepts	5
1.1.1	Set	5
1.1.2	Component of a Set	5
1.1.3	Empty Set	5
1.1.4	Cardinality	5
1.1.5	Universal Set	6
2	Linear Algebra	9
2.1	Introduction	9
2.2	Vectors and Vector Spaces	9
2.2.1	Definition of a Vector Space	9
2.2.2	Examples of Vector Spaces	9
2.3	Matrices	10
2.3.1	Operations on Matrices	10
2.3.2	Determinants	10
2.3.3	Inverse of a Matrix	10
2.4	Systems of Linear Equations	10
2.4.1	Solving Systems of Linear Equations	10
2.5	Eigenvalues and Eigenvectors	11
2.5.1	Finding Eigenvalues and Eigenvectors	11
2.6	Linear Transformations	11
3	Differential Equations and Numerical Methods	15
3.1	Introduction to Differential Equations	15
3.1.1	Types of Differential Equations	15
3.2	Methods for Solving Differential Equations	15
3.2.1	Analytical Methods	15
3.2.2	Numerical Methods	16
3.3	Applications of Differential Equations	16
4	Graph Theory	17
4.1	Introduction to Graph Theory	17
4.1.1	Types of Graphs	17
4.2	Fundamental Concepts	17
4.2.1	Degree of a Vertex	17
4.2.2	Paths and Cycles	17
4.2.3	Connected Graphs	18
4.3	Graph Representations	18
4.3.1	Adjacency Matrix	18
4.3.2	Adjacency List	18
4.3.3	Incidence Matrix	18
4.4	Special Classes of Graphs	18
4.4.1	Trees	18
4.4.2	Planar Graphs	18

4.4.3	Graphs and Subgraphs	18
4.5	Algorithms in Graph Theory	18
4.5.1	Breadth-First Search (BFS)	18
4.5.2	Depth-First Search (DFS)	19
4.5.3	Shortest Path Algorithms	19
4.5.4	Minimum Spanning Tree	19
4.6	Applications of Graph Theory	19
5	Lagrangian and Hamiltonian Mechanics	21
5.1	Introduction	21
5.2	Lagrangian Mechanics	21
5.2.1	Euler-Lagrange Equations	21
5.2.2	Generalized Coordinates and Velocities	21
5.2.3	Example: Simple Pendulum	22
5.3	Hamiltonian Mechanics	22
5.3.1	Hamilton's Equations	22
5.3.2	Generalized Coordinates and Momenta	22
5.3.3	Example: Simple Pendulum in Hamiltonian Form	23
5.4	Key Differences Between Lagrangian and Hamiltonian Mechanics	23
5.4.1	Lagrangian Mechanics	23
5.4.2	Hamiltonian Mechanics	23
5.4.3	Applications	23
6	Fourier Analysis	25
6.1	Introduction	25
6.2	Fourier Series	25
6.2.1	Convergence of Fourier Series	25
6.3	Fourier Transform	25
6.3.1	Properties of Fourier Transforms	26
6.4	Discrete Fourier Transform (DFT)	26
6.4.1	Fast Fourier Transform (FFT)	26
6.5	Applications of Fourier Analysis	26
6.5.1	Signal Processing	26
6.5.2	Image Processing	27
6.5.3	Quantum Mechanics	27

# Chapter 1

## Set Theory

Set theory is a fundamental branch of mathematics that studies sets, that is, collections of well-defined objects. These objects, called elements, can be numbers, points, or even other sets. Set theory is essential for formalizing mathematical concepts and the foundations of mathematical logic.

Here are the basic concepts of set theory:

### 1.1 Concepts

#### 1.1.1 Set

a Set is a collection of distinct elements.  $S = \{a_1, a_2, \dots, a_n\}$ . Example :  $A = \{1, 2, 3\}$  is a set which contains 1, 2 and 3.  
 $B = \{\text{banane}, \text{pomme}, \text{orange}, \text{ananas}\}$

**Property 1.** An infinite set is a set that contains an infinite number of elements. There are two main types of infinite sets:

- **Countable:** An infinite set is countable if it can be put into a one-to-one correspondence with the set of integers  $\mathbb{N}$ .  
Example: The set of integers  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$  is infinite and countable
- **Uncountable:** A set is uncountable if it is infinite, but there is no one-to-one correspondence with  $\mathbb{N}$ . Example: The set of real numbers  $\mathbb{R}$  is infinite and uncountable.

#### 1.1.2 Component of a Set

An object is an element of a set if it belongs to that set. We use the symbol  $\in$  to indicate that an element belongs to a set.  
Example :  $1 \in A$  but  $4 \notin A$ .

#### 1.1.3 Empty Set

The Empty set, designated by  $\emptyset$  or  $\{\}$  is a set that doesn't contain any element.

#### 1.1.4 Cardinality

The cardinality of a set is the number of elements in that set. If a set contains a finite number of elements, it is said to be finite, otherwise it is infinite.

Example : Si  $A = \{1, 2, 3\}$ , alors la cardinalité de  $A$  est  $|A| = 3$ .

### 1.1.5 Universal Set

The universal set, denoted  $U$ , is the set that contains all possible objects in a given context. Other sets are considered subsets of the universal set.

Example: If we work with integers, the universal set can be  $U=Z$ , the set of all integers.

**Property 2.** The paradox arises when we consider the set  $R$  of sets that do not contain themselves. The question is: does the set  $RR$  contain itself? If  $R$  contains itself, then even by definition it should not contain itself. If  $R$  does not contain itself, then by definition it should contain itself.

This leads to a logical contradiction.

**Property 3.** It doesn't exist a Set which contains of the Sets and a Set can't contain itself

### Union, Complement, and Intersection

#### Union

The union of two sets  $A$  and  $B$ , denoted  $A \cup B$ , is the set of all elements that are in  $A$ ,  $B$ , or both.

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\}.$$

Example: If  $A = \{1, 2, 3\}$  and  $B = \{3, 4, 5\}$ , then:

$$A \cup B = \{1, 2, 3, 4, 5\}.$$

#### Complement

The complement of a set  $A$ , denoted  $A^c$  or  $\bar{A}$ , is the set of all elements in the universal set  $U$  that are not in  $A$ .

$$A^c = \{x \mid x \in U \text{ and } x \notin A\}.$$

Example: If  $U = \{1, 2, 3, 4, 5\}$  and  $A = \{1, 2, 3\}$ , then:

$$A^c = \{4, 5\}.$$

#### Intersection

The intersection of two sets  $A$  and  $B$ , denoted  $A \cap B$ , is the set of all elements that are in both  $A$  and  $B$ .

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}.$$

Example: If  $A = \{1, 2, 3\}$  and  $B = \{3, 4, 5\}$ , then:

$$A \cap B = \{3\}.$$

### 1.1.7 Cartesian Product

The Cartesian product of two sets  $A$  and  $B$ , denoted  $A \times B$ , is the set of all ordered pairs where the first element is from  $A$  and the second is from  $B$ .

$$A \times B = \{(a, b) \mid a \in A, b \in B\}.$$

Example: If  $A = \{1, 2\}$  and  $B = \{x, y\}$ , then:

$$A \times B = \{(1, x), (1, y), (2, x), (2, y)\}.$$

### 1.1.8 Functions or Relations between Two Sets

#### Relation

A relation between two sets  $A$  and  $B$  is a subset of their Cartesian product  $A \times B$ . It specifies how elements of  $A$  are related to elements of  $B$ . Example: If  $A = \{1, 2\}$  and  $B = \{x, y\}$ , then a relation  $R$  could be:

$$R = \{(1, x), (2, y)\}.$$

## Function

A function  $f : A \rightarrow B$  is a special type of relation where every element of  $A$  is related to exactly one element of  $B$ . Example: If  $A = \{1, 2\}$  and  $B = \{x, y\}$ , then  $f = \{(1, x), (2, y)\}$  is a function. However,  $\{(1, x), (1, y)\}$  is not a function because 1 is related to multiple elements.

### 1.1.9 Cartesian Product (Repeated)

This concept is identical to the explanation in Section 1.1.7.

### 1.1.10 Parts of a Set (Power Set)

The parts of a set, also known as the power set, are the set of all subsets of a given set  $A$ . The power set of  $A$  is denoted  $\mathcal{P}(A)$ . If  $A$  has  $n$  elements, then  $\mathcal{P}(A)$  contains  $2^n$  subsets, including the empty set  $\emptyset$  and  $A$  itself. Example: If  $A = \{1, 2\}$ , then:

$$\mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}.$$

## Zermelo-Fraenkel Set Theory (ZF)

Zermelo-Fraenkel Set Theory is a foundational system for mathematics that defines the rules for constructing sets. It consists of a series of axioms that describe how sets behave and interact, aiming to eliminate paradoxes such as Russell's Paradox. The ZF axioms are often extended with the Axiom of Choice, forming ZFC (Zermelo-Fraenkel with Choice).

### 1. Axiom of Extensionality

Statement: Two sets are equal if and only if they have the same elements.

$$\forall A \forall B (\forall x (x \in A \iff x \in B) \implies A = B).$$

Explanation: This axiom ensures that sets are defined solely by their elements, not by how they are constructed.

### 2. Pair Axiom

Statement: For any two sets  $A$  and  $B$ , there exists a set  $\{A, B\}$  containing exactly  $A$  and  $B$ .

$$\forall A \forall B \exists C \forall x (x \in C \iff (x = A \vee x = B)).$$

Explanation: This axiom guarantees the existence of pairs of sets and allows constructing ordered pairs.

### 3. Reunion and Union Axiom

Statement: For any set  $A$ , there exists a set  $B$  that is the union of all elements of  $A$ .

$$\forall A \exists B \forall x (x \in B \iff \exists C (C \in A \wedge x \in C)).$$

Explanation: This axiom allows forming a set that contains all elements of the sets within a given set  $A$ .

### 4. Empty Set Axiom

Statement: There exists a set  $\emptyset$  that contains no elements.

$$\exists A \forall x (x \notin A).$$

Explanation: This axiom ensures the existence of the empty set, which is the foundation for building other sets.

### 5. Axiom of Understanding (or Separation Axiom)

Statement: *Given any set  $A$  and a property  $P(x)$ , there exists a subset  $B \subseteq A$  containing all elements of  $A$  that satisfy  $P(x)$ .*

$$\forall A \exists B \forall x (x \in B \iff (x \in A \wedge P(x))).$$

Explanation: *This axiom restricts set formation to avoid paradoxes, allowing subsets based on a defining property.*

### 6. Exponentiation Axiom

Statement: *For any two sets  $A$  and  $B$ , there exists a set  $C$  consisting of all functions from  $B$  to  $A$ .*

$$\forall A \forall B \exists C (C = \text{Fun}(B, A)).$$

Explanation: *This axiom allows constructing sets of functions and defines the "power" of one set relative to another.*

### 7. Axiom of Infinity

Statement: *There exists a set  $I$  that contains the empty set and is closed under the operation of adding a single element to a set.*

$$\exists I (\emptyset \in I \wedge \forall x (x \in I \implies x \cup \{x\} \in I)).$$

Explanation: *This axiom guarantees the existence of infinite sets, foundational for defining natural numbers.*

### 8. Replacement Axiom

Statement: *If  $F$  is a function and  $A$  is a set, then the image of  $A$  under  $F$  is also a set.*

$$\forall A \exists B \forall y (y \in B \iff \exists x (x \in A \wedge y = F(x))).$$

Explanation: *This axiom ensures that sets remain well-defined when transformed by a function.*

### 9. Axiom of Foundation (or Regularity)

Statement: *Every non-empty set  $A$  contains an element that is disjoint from  $A$ .*

$$\forall A (A \neq \emptyset \implies \exists x (x \in A \wedge x \cap A = \emptyset)).$$

Explanation: *This axiom eliminates infinitely descending chains of sets (no set can contain itself directly or indirectly).*

### 10. Axiom of Choice

Statement: *For any collection of non-empty, disjoint sets, there exists a set containing exactly one element from each of these sets.*

$$\forall \mathcal{A} (\forall A \in \mathcal{A}, A \neq \emptyset) \implies \exists C (\forall A \in \mathcal{A}, C \cap A \neq \emptyset).$$

Explanation: *This axiom enables selecting elements from sets even when no explicit rule for selection is defined. It's crucial for proving many theorems in modern mathematics but is independent of ZF, forming ZFC when included.*



## Chapter 2

# Linear Algebra

### 2.1 Introduction

*Linear algebra is a branch of mathematics that deals with vector spaces, linear transformations, and systems of linear equations. It is foundational to many areas of mathematics and science, including physics, computer science, economics, and engineering. In linear algebra, we study the properties and operations on vectors and matrices, which are used to represent data, transformations, and systems of equations.*

### 2.2 Vectors and Vector Spaces

*A vector is an ordered list of numbers (or components) that represent a point in space. Vectors can be added together and multiplied by scalars (real numbers). A vector space is a collection of vectors that satisfies certain properties such as closure under addition and scalar multiplication.*

#### 2.2.1 Definition of a Vector Space

*A set  $V$  is a vector space if it satisfies the following properties for all vectors  $u, v, w \in V$  and scalars  $c, d \in \mathbb{R}$ :*

- Closure under addition:  $u + v \in V$
- Closure under scalar multiplication:  $c \cdot u \in V$
- Commutativity of addition:  $u + v = v + u$
- Associativity of addition:  $(u + v) + w = u + (v + w)$
- Existence of an additive identity: *There exists an element  $0 \in V$  such that  $u + 0 = u$*
- Existence of additive inverses: *For every vector  $u \in V$ , there exists a vector  $-u \in V$  such that  $u + (-u) = 0$*
- Distributivity of scalar multiplication over vector addition:  $c \cdot (u + v) = c \cdot u + c \cdot v$
- Distributivity of scalar multiplication over scalar addition:  $(c + d) \cdot u = c \cdot u + d \cdot u$
- Compatibility of scalar multiplication with field multiplication:  $c \cdot (d \cdot u) = (c \cdot d) \cdot u$
- Existence of a multiplicative identity:  $1 \cdot u = u$

#### 2.2.2 Examples of Vector Spaces

*Examples of vector spaces include:*

- *The set of all 2-dimensional vectors  $\mathbb{R}^2$*
- *The set of all polynomials of degree at most  $n$*
- *The set of all matrices of a fixed size*

## 2.3 Matrices

A matrix is a rectangular array of numbers, arranged in rows and columns. Matrices are used to represent linear transformations, systems of linear equations, and data structures. The size of a matrix is defined by its number of rows and columns, denoted  $m \times n$  where  $m$  is the number of rows and  $n$  is the number of columns.

### 2.3.1 Operations on Matrices

Some common matrix operations include:

- Addition: Matrices can be added if they have the same dimensions.
- Scalar multiplication: A matrix can be multiplied by a scalar by multiplying each element of the matrix by the scalar.
- Matrix multiplication: Two matrices  $A$  and  $B$  can be multiplied if the number of columns in  $A$  equals the number of rows in  $B$ .
- Transpose: The transpose of a matrix  $A$ , denoted  $A^T$ , is obtained by swapping its rows and columns.

### 2.3.2 Determinants

The determinant of a square matrix  $A$  is a scalar value that can be computed from its elements and provides important information about the matrix. The determinant is used to determine whether a matrix is invertible, and it is also used in solving systems of linear equations.

$$\det(A) = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det(A_{ij})$$

where  $A_{ij}$  is the submatrix obtained by removing the  $i$ -th row and  $j$ -th column of  $A$ .

### 2.3.3 Inverse of a Matrix

The inverse of a matrix  $A$ , denoted  $A^{-1}$ , is the matrix that satisfies:

$$A \cdot A^{-1} = I$$

where  $I$  is the identity matrix. A matrix has an inverse if and only if its determinant is non-zero.

## 2.4 Systems of Linear Equations

A system of linear equations is a collection of linear equations involving the same set of variables. The general form of a linear equation is:

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$$

where  $a_1, a_2, \dots, a_n$  are constants,  $x_1, x_2, \dots, x_n$  are variables, and  $b$  is a constant.

### 2.4.1 Solving Systems of Linear Equations

Systems of linear equations can be solved using several methods:

- Gaussian elimination: This method transforms the system into an upper triangular form and then solves it using back substitution.
- Matrix methods: The system can be represented as a matrix equation  $A \cdot x = b$  and solved using matrix inversion or other methods.

## 2.5 Eigenvalues and Eigenvectors

Given a square matrix  $A$ , a non-zero vector  $v$  is called an eigenvector of  $A$  if there is a scalar  $\lambda$  such that:

$$A \cdot v = \lambda \cdot v$$

The scalar  $\lambda$  is called the eigenvalue corresponding to the eigenvector  $v$ . The process of finding eigenvalues and eigenvectors is essential in various applications, including principal component analysis (PCA) and quantum mechanics.

### 2.5.1 Finding Eigenvalues and Eigenvectors

To find the eigenvalues of a matrix  $A$ , we solve the characteristic equation:

$$\det(A - \lambda I) = 0$$

where  $I$  is the identity matrix. The solutions to this equation are the eigenvalues  $\lambda$ . Once the eigenvalues are found, the eigenvectors can be determined by solving the system:

$$(A - \lambda I) \cdot v = 0$$

## 2.6 Linear Transformations

A linear transformation is a function  $T : V \rightarrow W$  between two vector spaces  $V$  and  $W$  that satisfies the following properties:

- Additivity:  $T(u + v) = T(u) + T(v)$
- Homogeneity:  $T(c \cdot u) = c \cdot T(u)$

Linear transformations can be represented by matrices. If  $T$  is a linear transformation from  $R^n$  to  $R^m$ , then there exists a matrix  $A$  such that:

$$T(v) = A \cdot v$$

where  $v$  is a vector in  $R^n$ .

## Fundamental Notions

### 1. Scalar

Definition: A scalar is a single numerical value, typically representing magnitude. Scalars are often used in mathematical operations such as addition, multiplication, and scaling vectors. Examples include real numbers ( $R$ ), integers ( $Z$ ), and complex numbers ( $C$ ).

### 2. Vectors

Definition: A vector is an ordered set of numbers (or elements) that can be represented geometrically as a directed line segment in space. Algebraically, a vector in  $n$ -dimensional space is written as:

$$\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}.$$

Vectors have both magnitude and direction.

### 3. Vector Space

Definition: A vector space is a set  $V$  over a field  $F$  (e.g.,  $R$  or  $C$ ) where vector addition and scalar multiplication satisfy the following properties:

- Closure under addition and scalar multiplication.
- Existence of additive identity ( $0$ ).
- Existence of additive inverses.
- Associativity and commutativity of addition.
- Distributive properties of scalar multiplication.

### 4. Linear Combinations

Definition: A linear combination of vectors  $v_1, v_2, \dots, v_n \in V$  is any expression of the form:

$$w = a_1 v_1 + a_2 v_2 + \dots + a_n v_n,$$

where  $a_1, a_2, \dots, a_n \in F$  are scalars.

### 5. Matrices

Definition: A matrix is a rectangular array of numbers arranged in rows and columns. An  $m \times n$  matrix has  $m$  rows and  $n$  columns, typically written as:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

### 6. Determinants

Definition: The determinant of a square matrix  $A$  is a scalar value that provides information about the matrix, such as whether it is invertible. For a  $2 \times 2$  matrix:

$$\det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$

Determinants generalize to higher dimensions using recursive expansions.

### 7. Eigenvalues and Eigenvectors

Definition: For a square matrix  $A$ , an eigenvector  $v$  is a non-zero vector such that:

$$Av = \lambda v,$$

where  $\lambda$  is the corresponding eigenvalue.

### 8. Linear Transformations

Definition: A linear transformation  $T : V \rightarrow W$  between vector spaces preserves vector addition and scalar multiplication:

$$T(u + v) = T(u) + T(v),$$

$$T(cu) = cT(u),$$

for all  $u, v \in V$  and  $c \in F$ .

## 9. Rank

*Definition: The rank of a matrix  $A$  is the dimension of the column space (or equivalently, the row space) of  $A$ . It represents the maximum number of linearly independent columns (or rows) in the matrix.*

### Systems of Linear Equations

*Definition: A system of linear equations is a set of equations in the form:*

$$a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1,$$

$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2,$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m.$$

*Solutions are found using methods such as substitution, elimination, or matrix operations.*

### Gram-Schmidt Process

*Definition: The Gram-Schmidt process is a method for orthonormalizing a set of vectors in a vector space. Given a set of linearly independent vectors  $v_1, v_2, \dots, v_n$ , the process constructs an orthonormal basis  $u_1, u_2, \dots, u_n$  by:*

$$u_1 = \frac{v_1}{\|v_1\|},$$

$$u_k = \frac{v_k - \sum_{i=1}^{k-1} (v_k \cdot u_i) u_i}{\|v_k - \sum_{i=1}^{k-1} (v_k \cdot u_i) u_i\|}, \quad k = 2, 3, \dots, n.$$

## Navigating Between Vector Spaces

### Introduction

*Navigating between vector spaces involves understanding how vectors in one space can be mapped, transformed, or related to vectors in another space. This concept is foundational in linear algebra and has broad applications in areas like computer graphics, quantum mechanics, and data science.*

*The relationship between vector spaces is often established through linear transformations, which preserve the structure of vector spaces, making them predictable and mathematically elegant to analyze.*

### Linear Transformations

*A linear transformation is a mapping  $T : V \rightarrow W$  between two vector spaces  $V$  and  $W$  over the same field  $F$  that satisfies the following properties:*

- *Additivity:  $T(u + v) = T(u) + T(v)$ ,  $\forall u, v \in V$ ,*
- *Homogeneity:  $T(cu) = cT(u)$ ,  $\forall c \in F, u \in V$ .*

*Linear transformations can often be represented by matrices. If  $v \in V$  is expressed in terms of a basis, then  $T(v)$  can be computed using matrix multiplication:*

$$T(v) = Av,$$

*where  $A$  is the transformation matrix corresponding to  $T$ .*

## Change of Basis

To navigate between vector spaces, it is often necessary to change the basis in which vectors are expressed. A basis of a vector space  $V$  is a set of linearly independent vectors that span  $V$ . If  $B = \{b_1, b_2, \dots, b_n\}$  is a basis for  $V$ , any vector  $v \in V$  can be uniquely written as:

$$v = c_1 b_1 + c_2 b_2 + \dots + c_n b_n,$$

where  $c_1, c_2, \dots, c_n \in F$  are the coordinates of  $v$  in the basis  $B$ .

When transitioning between two different bases, a change of basis matrix  $P$  is used. For instance, if  $v$  is expressed in basis  $B_1$  and needs to be expressed in basis  $B_2$ , the coordinates are transformed as:

$$[v]_{B_2} = P[v]_{B_1}.$$

## Kernel and Image

For a linear transformation  $T : V \rightarrow W$ , two important subspaces help in navigating between vector spaces:

- Kernel: The kernel of  $T$ , denoted as  $\ker(T)$ , is the set of vectors in  $V$  that are mapped to the zero vector in  $W$ :

$$\ker(T) = \{v \in V \mid T(v) = 0\}.$$

- Image: The image of  $T$ , denoted as  $\text{Im}(T)$ , is the set of all vectors in  $W$  that can be expressed as  $T(v)$  for some  $v \in V$ :

$$\text{Im}(T) = \{T(v) \mid v \in V\}.$$

The rank-nullity theorem connects these two subspaces:

$$\dim(\ker(T)) + \dim(\text{Im}(T)) = \dim(V).$$

## Applications in Projection and Transformation

One common example of navigating between vector spaces is through projections. A projection maps a vector  $v$  in a higher-dimensional space  $V$  to a subspace  $W \subseteq V$ . The projection is given by:

$$P_W(v) = v_{\parallel} = \frac{v \cdot w}{w \cdot w} w,$$

where  $w$  is a basis vector of  $W$ . This operation is crucial in areas like computer graphics, where 3D points are projected onto 2D planes.

## Chapter 3

# Differential Equations and Numerical Methods

### 3.1 Introduction to Differential Equations

*Differential equations are mathematical equations that involve an unknown function and its derivatives. They are essential for modeling a wide variety of phenomena in physics, engineering, biology, economics, and other sciences. The general form of a differential equation is:*

$$F\left(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots\right) = 0,$$

*where  $y$  is the unknown function, and its derivatives  $\frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots$  are involved.*

#### 3.1.1 Types of Differential Equations

- Ordinary Differential Equations (ODEs): *These involve a single independent variable and its derivatives, such as:*

$$\frac{dy}{dx} + p(x)y = q(x).$$

- Partial Differential Equations (PDEs): *These involve multiple independent variables and partial derivatives, such as:*

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

- Linear vs. Nonlinear Equations: *Linear equations have the unknown function and its derivatives appearing linearly, while nonlinear equations involve nonlinear terms.*

### 3.2 Methods for Solving Differential Equations

#### 3.2.1 Analytical Methods

*Analytical methods involve finding an exact solution to a differential equation. These include:*

- Separation of Variables: *Used when a differential equation can be written as:*

$$\frac{dy}{dx} = g(x)h(y).$$

*By separating variables and integrating:*

$$\int \frac{1}{h(y)} dy = \int g(x) dx.$$

- Integrating Factors: *Used for first-order linear equations of the form:*

$$\frac{dy}{dx} + p(x)y = q(x).$$

*The solution is obtained using the integrating factor:*

$$\mu(x) = e^{\int p(x) dx}.$$

- Characteristic Equations: *Used for linear differential equations with constant coefficients, such as:*

$$a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} + cy = 0.$$

*Solutions are derived from the roots of the characteristic polynomial.*

### 3.2.2 Numerical Methods

*When analytical solutions are not feasible, numerical methods are employed to approximate solutions.*

- Euler's Method: *A simple numerical method for solving first-order ODEs. Given:*

$$\frac{dy}{dx} = f(x, y),$$

*the solution is approximated using:*

$$y_{n+1} = y_n + hf(x_n, y_n),$$

*where  $h$  is the step size.*

- Runge-Kutta Methods: *These are more accurate methods for solving ODEs. The most common is the fourth-order Runge-Kutta method:*

$$y_{n+1} = y_n + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4),$$

*where:*

$$\begin{aligned} k_1 &= f(x_n, y_n), & k_2 &= f\left(x_n + \frac{h}{2}, y_n + \frac{h}{2}k_1\right), \\ k_3 &= f\left(x_n + \frac{h}{2}, y_n + \frac{h}{2}k_2\right), & k_4 &= f(x_n + h, y_n + hk_3). \end{aligned}$$

- Finite Difference Methods: *Used for solving PDEs by approximating derivatives with finite differences. For example, the second derivative  $\frac{\partial^2 u}{\partial x^2}$  is approximated as:*

$$\frac{\partial^2 u}{\partial x^2} \approx \frac{u(x+h) - 2u(x) + u(x-h)}{h^2}.$$

## 3.3 Applications of Differential Equations

*Differential equations are applied in many domains:*

- Physics: *Modeling motion, heat transfer, wave propagation, etc.*
- Biology: *Modeling population dynamics, spread of diseases, etc.*
- Engineering: *Modeling electrical circuits, mechanical vibrations, etc.*
- Economics: *Modeling market dynamics, optimal control problems, etc.*



# Chapter 4

## Graph Theory

### 4.1 Introduction to Graph Theory

*Graph theory is the study of mathematical structures used to model pairwise relations between objects. A graph  $G = (V, E)$  consists of:*

- *A set of vertices (or nodes)  $V$ , which represent entities.*
- *A set of edges  $E$ , which represent relationships between the vertices.*

*Graphs are widely used in computer science, physics, biology, social sciences, and many other fields.*

#### 4.1.1 Types of Graphs

- *Undirected Graph: A graph where edges have no direction.*
- *Directed Graph (Digraph): A graph where edges have a direction, represented as ordered pairs  $(u, v)$ .*
- *Weighted Graph: A graph where each edge is assigned a weight or cost.*
- *Simple Graph: A graph with no loops or multiple edges between the same vertices.*
- *Complete Graph: A graph where every pair of vertices is connected by an edge.*
- *Bipartite Graph: A graph where vertices can be divided into two disjoint sets such that every edge connects a vertex from one set to the other.*

### 4.2 Fundamental Concepts

#### 4.2.1 Degree of a Vertex

*The degree of a vertex in an undirected graph is the number of edges incident to it. In a directed graph:*

- *In-degree: Number of incoming edges to the vertex.*
- *Out-degree: Number of outgoing edges from the vertex.*

#### 4.2.2 Paths and Cycles

- *Path: A sequence of edges connecting a sequence of vertices.*
- *Cycle: A path that starts and ends at the same vertex.*
- *Simple Path: A path with no repeated vertices.*

### 4.2.3 Connected Graphs

A graph is connected if there is a path between every pair of vertices. For directed graphs, the concepts of strong and weak connectivity are used:

- Strongly Connected: There is a directed path between every pair of vertices.
- Weakly Connected: The graph becomes connected if the direction of the edges is ignored.

## 4.3 Graph Representations

Graphs can be represented in various ways:

### 4.3.1 Adjacency Matrix

An  $n \times n$  matrix  $A$  where  $A[i][j] = 1$  if there is an edge between vertices  $i$  and  $j$ , and  $A[i][j] = 0$  otherwise.

$$A[i][j] = \begin{cases} 1 & \text{if } (i, j) \in E, \\ 0 & \text{otherwise.} \end{cases}$$

### 4.3.2 Adjacency List

A collection of lists, where each list corresponds to a vertex and contains all the vertices connected to it.

### 4.3.3 Incidence Matrix

An  $n \times m$  matrix  $B$ , where  $n$  is the number of vertices and  $m$  is the number of edges.  $B[i][j] = 1$  if vertex  $i$  is incident to edge  $j$ , and  $B[i][j] = 0$  otherwise.

## 4.4 Special Classes of Graphs

### 4.4.1 Trees

A tree is a connected graph with no cycles. It has the following properties:

- A tree with  $n$  vertices has  $n - 1$  edges.
- Any two vertices are connected by exactly one path.

### 4.4.2 Planar Graphs

A graph is planar if it can be drawn on a plane without any edges crossing.

### 4.4.3 Graphs and Subgraphs

A subgraph is a subset of a graph's vertices and edges. If a subgraph includes all the edges between its selected vertices, it is called an induced subgraph.

## 4.5 Algorithms in Graph Theory

### 4.5.1 Breadth-First Search (BFS)

An algorithm for traversing or searching graph data structures. It explores all neighbors at the current depth level before moving to nodes at the next depth level.

### 4.5.2 Depth-First Search (DFS)

*An algorithm that starts at a source vertex and explores as far along each branch as possible before backtracking.*

### 4.5.3 Shortest Path Algorithms

- Dijkstra's Algorithm: *Finds the shortest path from a source vertex to all other vertices in a weighted graph with non-negative weights.*
- Bellman-Ford Algorithm: *Solves the single-source shortest path problem in graphs with negative weights.*

### 4.5.4 Minimum Spanning Tree

*A spanning tree of a graph that has the smallest possible total edge weight. Algorithms to compute this include:*

- Kruskal's Algorithm: *Adds edges in increasing order of weight while avoiding cycles.*
- Prim's Algorithm: *Builds the MST incrementally, starting with a single vertex.*

## 4.6 Applications of Graph Theory

- Networking: *Modeling computer and social networks.*
- Transportation: *Optimizing routes and traffic flows.*
- Biology: *Representing relationships in ecosystems and genetic structures.*
- Physics: *Modeling systems of particles and interactions.*



## Chapter 5

# Lagrangian and Hamiltonian Mechanics

### 5.1 Introduction

*Lagrangian and Hamiltonian mechanics are two formulations of classical mechanics that provide powerful tools for analyzing the motion of physical systems. These formalisms are particularly useful in systems with constraints and play a central role in modern physics, including quantum mechanics.*

### 5.2 Lagrangian Mechanics

*Lagrangian mechanics is based on the principle of least action. The action  $S$  is the integral of the Lagrangian  $L$  over time:*

$$S = \int L dt$$

*where  $L$  is the Lagrangian, defined as:*

$$L = T - V$$

*Here,  $T$  is the kinetic energy and  $V$  is the potential energy of the system. The Lagrangian provides a way to derive the equations of motion of a system through the **Euler-Lagrange equations**.*

#### 5.2.1 Euler-Lagrange Equations

*The equations of motion are derived from the principle of least action, which states that the path a system follows is the one that minimizes the action. The Euler-Lagrange equation for a generalized coordinate  $q_i$  is:*

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0$$

*where  $q_i$  are the generalized coordinates, and  $\dot{q}_i$  are the generalized velocities. These equations describe how the generalized coordinates evolve over time.*

#### 5.2.2 Generalized Coordinates and Velocities

*In Lagrangian mechanics, we often use generalized coordinates  $q_i$  that describe the configuration of a system. The generalized velocities  $\dot{q}_i$  are the time derivatives of these coordinates. These variables simplify the description of systems with constraints, where the number of independent variables is reduced.*

### 5.2.3 Example: Simple Pendulum

Consider a simple pendulum with mass  $m$  and length  $l$ . The generalized coordinate is the angle  $\theta$  between the pendulum and the vertical axis. The kinetic energy  $T$  and potential energy  $V$  of the pendulum are given by:

$$T = \frac{1}{2}ml^2\dot{\theta}^2, \quad V = mgl(1 - \cos \theta)$$

The Lagrangian is:

$$L = T - V = \frac{1}{2}ml^2\dot{\theta}^2 - mgl(1 - \cos \theta)$$

The Euler-Lagrange equation for  $\theta$  is:

$$\frac{d}{dt}(ml^2\dot{\theta}) + mgl \sin \theta = 0$$

This gives the equation of motion for the pendulum.

## 5.3 Hamiltonian Mechanics

Hamiltonian mechanics reformulates Lagrangian mechanics using the **Hamiltonian function**, which represents the total energy of the system:

$$H = T + V$$

The Hamiltonian is a function of generalized coordinates  $q_i$  and generalized momenta  $p_i$ , where the generalized momentum is defined as:

$$p_i = \frac{\partial L}{\partial \dot{q}_i}$$

In Hamiltonian mechanics, the evolution of the system is described by **Hamilton's equations**.

### 5.3.1 Hamilton's Equations

Hamilton's equations describe the time evolution of the generalized coordinates and momenta:

$$\dot{q}_i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q_i}$$

These equations are a set of first-order differential equations, which are often easier to solve than the second-order Euler-Lagrange equations.

### 5.3.2 Generalized Coordinates and Momenta

In Hamiltonian mechanics, we use generalized coordinates  $q_i$  and generalized momenta  $p_i$  instead of velocities. The momentum  $p_i$  is defined as the derivative of the Lagrangian with respect to the generalized velocity  $\dot{q}_i$ :

$$p_i = \frac{\partial L}{\partial \dot{q}_i}$$

The Hamiltonian describes the total energy of the system, and Hamilton's equations govern the time evolution of the system.

### 5.3.3 Example: Simple Pendulum in Hamiltonian Form

For the simple pendulum example, we can write the kinetic energy  $T$  and potential energy  $V$  in terms of generalized coordinates and momenta. The generalized coordinate is  $\theta$ , and the generalized momentum is:

$$p_\theta = ml^2\dot{\theta}$$

The Hamiltonian is:

$$H = \frac{p_\theta^2}{2ml^2} + mgl(1 - \cos \theta)$$

Hamilton's equations for this system are:

$$\dot{\theta} = \frac{\partial H}{\partial p_\theta} = \frac{p_\theta}{ml^2}, \quad \dot{p}_\theta = -\frac{\partial H}{\partial \theta} = -mgl \sin \theta$$

These equations describe the motion of the pendulum.

## 5.4 Key Differences Between Lagrangian and Hamiltonian Mechanics

### 5.4.1 Lagrangian Mechanics

Lagrangian mechanics focuses on the difference between kinetic and potential energy. It uses generalized coordinates and velocities and is particularly useful for systems with constraints.

### 5.4.2 Hamiltonian Mechanics

Hamiltonian mechanics uses the total energy of the system and reformulates the equations of motion using generalized coordinates and momenta. It is often preferred in quantum mechanics and statistical mechanics due to its more convenient mathematical structure.

### 5.4.3 Applications

Both Lagrangian and Hamiltonian mechanics are used extensively in various fields such as classical mechanics, quantum mechanics, and field theory. Lagrangian mechanics is often used for systems with constraints, while Hamiltonian mechanics is favored for systems involving statistical mechanics or quantum field theory.





## Chapter 6

# Fourier Analysis

### 6.1 Introduction

*Fourier analysis is a mathematical technique used to analyze and represent functions as sums of sinusoidal functions (sines and cosines). It is widely used in various fields such as signal processing, heat conduction, acoustics, and quantum mechanics. Fourier analysis transforms complex functions into simpler components, making it easier to study their properties.*

### 6.2 Fourier Series

*The Fourier series is a way to represent a periodic function as a sum of sines and cosines. A function  $f(x)$  defined on the interval  $[-\pi, \pi]$  can be written as:*

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx))$$

*where  $a_0$ ,  $a_n$ , and  $b_n$  are the Fourier coefficients. These coefficients are given by the following formulas:*

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx, \quad n \geq 1$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx, \quad n \geq 1$$

*These Fourier coefficients capture the amplitude of the corresponding sine and cosine functions in the decomposition of  $f(x)$ .*

#### 6.2.1 Convergence of Fourier Series

*The Fourier series of a function converges to the function at points where it is continuous. However, at points of discontinuity, the Fourier series converges to the average of the left-hand and right-hand limits of the function at that point.*

### 6.3 Fourier Transform

*The Fourier transform generalizes the Fourier series to non-periodic functions. It transforms a function  $f(t)$  in the time domain into a function  $F(\omega)$  in the frequency domain. The Fourier transform of a function  $f(t)$  is defined as:*

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$$

where  $\omega$  is the angular frequency, and  $e^{-i\omega t}$  is a complex exponential function representing a sinusoidal wave. The inverse Fourier transform is given by:

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{i\omega t} d\omega$$

The Fourier transform provides a way to decompose a time-domain signal into its frequency components.

### 6.3.1 Properties of Fourier Transforms

Some important properties of the Fourier transform include:

- Linearity:  $\mathcal{F}[af(t) + bg(t)] = aF(\omega) + bG(\omega)$
- Shift in Time Domain:  $\mathcal{F}[f(t - t_0)] = e^{-i\omega t_0} F(\omega)$
- Convolution: The Fourier transform of the convolution of two functions is the product of their individual Fourier transforms:  $\mathcal{F}[f * g] = F(\omega)G(\omega)$
- Parseval's Theorem: The total energy of the function in the time domain is equal to the total energy in the frequency domain:

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(\omega)|^2 d\omega$$

## 6.4 Discrete Fourier Transform (DFT)

The discrete Fourier transform (DFT) is used to analyze discrete signals and is an approximation of the continuous Fourier transform. The DFT of a discrete signal  $x_n$  of length  $N$  is given by:

$$X_k = \sum_{n=0}^{N-1} x_n e^{-i\frac{2\pi}{N}kn}, \quad k = 0, 1, 2, \dots, N-1$$

The inverse DFT is given by:

$$x_n = \frac{1}{N} \sum_{k=0}^{N-1} X_k e^{i\frac{2\pi}{N}kn}, \quad n = 0, 1, 2, \dots, N-1$$

The DFT allows us to analyze the frequency content of a finite set of discrete data points.

### 6.4.1 Fast Fourier Transform (FFT)

The fast Fourier transform (FFT) is an efficient algorithm to compute the DFT. It reduces the computational complexity from  $O(N^2)$  to  $O(N \log N)$ , making it feasible to analyze large datasets.

## 6.5 Applications of Fourier Analysis

Fourier analysis has a wide range of applications across various fields:

### 6.5.1 Signal Processing

In signal processing, Fourier analysis is used to analyze and filter signals. It helps in identifying the frequency components of a signal, which is essential for tasks like audio compression, noise reduction, and digital communication.

### 6.5.2 Image Processing

*In image processing, the Fourier transform is used to analyze spatial frequencies in an image. By transforming an image to the frequency domain, one can apply various filters to enhance or remove certain features, such as edges or noise. In Metascript, it's the principal use of the FOurier Analysis*

### 6.5.3 Quantum Mechanics

*Fourier analysis plays a crucial role in quantum mechanics, where wavefunctions are often represented in terms of their frequency components. The Fourier transform connects the position and momentum representations of a quantum state.*