

NOTES ON SIMPLICIAL HOMOTOPY THEORY

NICOLA GAMBINO AND SIMON HENRY

ABSTRACT. Some notes on simplicial homotopy theory

1. Introduction

2. Review of the semi-model structure on simplicial sets

Let \mathcal{E} be a category with an initial object 0 and a terminal object 1. Recall that a semi-model structure on \mathcal{E} consists of three classes of maps **Weq**, **Fib** and **Cof** in \mathcal{E} satisfying some axioms, which we now recall. When stating these axioms, and in the following, we refer to elements of **Weq**, **Fib** and **Cof** as weak equivalences, fibrations and cofibrations, respectively, call *fibrant* the objects X for which the unique map $\top_X: X \rightarrow 1$ is a fibration, and *cofibrant* the objects Y for which the unique map $\perp_Y: 0 \rightarrow Y$ is a cofibration.

The axioms for a semi-model structure can then be stated as follows:

(SM1) A

(SM2)

It is important to note the similarities and differences between semi-model structures and model structures.

- Basics on **SSet** and notation.
- We use Δ^n , $\partial\Delta^n$ and Λ_k^n .
- Recall the semi model structure on **SSet**:
 - Generating cofibrations are the boundary inclusions,

$$\mathcal{I} =_{\text{def}} \{i^n: \partial\Delta^n \rightarrow \Delta^n \mid n \geq 0\}$$

- Trivial Kan fibrations = maps with the right lifting property with respect to \mathcal{I} .
- Cofibrations = Levelwise complemented monomorphisms such that degeneracies are decidable on the complement of the image.
- Cofibrant objects are simplicial sets in which degeneracies are decidable.
- Generating trivial cofibrations are the horn inclusions:

$$\mathcal{J} =_{\text{def}} \{h_n^k: \Lambda_k^n \rightarrow \Delta^n \mid 0 \leq k \leq n\}.$$

- Kan fibrations = maps with the right lifting property with respect to \mathcal{J} .
- Fibrant objects are Kan complexes.
- Interval object is Δ^1 . Endpoint inclusions $\delta^k: \{k\} \rightarrow \Delta^1$ are defined by $\delta^k =_{\text{def}} h_k^1$.
- It would be good to have an explicit definition of the cofibrant replacement functor.
- Recall the pushout product property.

Lemma 2.1. *If B and C are cofibrants, and A is any simplicial sets with maps $B \rightarrow A$, $C \rightarrow A$ then the pullback $B \times_A C$ is cofibrant.*

Proof. Sketch: we check it first for the product $B \times C$, and then observe that $B \times_A C \subset B \times C$ and a subobject of a cofibrant object is cofibrant. \square

Proposition 2.2. *If A, B, C are cofibrant simplicial sets, $A \hookrightarrow C$ is a cofibrations and $f: B \rightarrow C$ is any morphism, then the natural projection map $A \times_C B \rightarrow B$ is a cofibration.*

Proof. To be added. \square

Lemma 2.3. *For $0 \leq k \leq n$, the horn inclusion $h_n^k: \Lambda_k^n \rightarrow \Delta^n$ is a retract of the pushout product*

$$h_1^k \hat{\times} i^n: (\Delta^1 \times \partial\Delta^n) \cup (\{k\} \times \Delta^n) \rightarrow \Delta^1 \times \Delta^n$$

Proposition 2.4. *A map is a Kan fibration if and only if it has the right lifting property with respect to the pushout products $i \hat{\times} \delta^k$.*

3. Dependent products

The first aim of this section is to show that for every fibration $p: B \rightarrow A$ with A cofibrant, the pushforward functor $\Pi_p: \mathcal{E}_{/B} \rightarrow \mathcal{E}_{/A}$ preserves fibrations. Our second aim is to build on this result to define sufficient structure to interpret Martin-Löf's rules for Π -types.

For our first aim, let $p: B \rightarrow A$ be a fibration with cofibrant codomain and observe that $\Pi_p: \mathcal{E}_{/B} \rightarrow \mathcal{E}_{/A}$ preserves fibrations if and only if its left adjoint

$$p^*: \mathcal{E}_{/A} \rightarrow \mathcal{E}_{/B}$$

preserves trivial cofibrations. Furthermore,

Lemma 3.1.

- (i) $\mathcal{J} \subset \text{Cof} \cap \mathcal{S}$.
- (ii) $\text{Cof} \cap \mathcal{S} \subseteq \text{TrivCof}$.

Lemma 3.2. *For $k \in \{0, 1\}$, the pullback of a strong k -oriented homotopy equivalence with cofibrant codomain along a fibration is a strong k -oriented homotopy equivalence.*

Theorem 3.3. *The semi-model structure for Kan complexes on \mathbf{SSet} has the restricted Frobenius condition.*

Proof. Since the semi-model structure in which we are working is cofibrantly generated, it is sufficient [TO CHECK] that p^* sends generating trivial cofibrations to trivial cofibrations. So, let $p: B \rightarrow \Delta[n]$ be a fibration, $i: \Lambda^k[n] \rightarrow \Delta[n]$ be a horn inclusion, and define $j =_{\text{def}} p^*(i)$, given by the pullback diagram

$$\begin{array}{ccc} \bullet & \xrightarrow{\quad} & \Lambda^k[n] \\ j \downarrow & \lrcorner & \downarrow i \\ B & \xrightarrow[p]{} & \Delta[n] \end{array}$$

We need to show that j is a trivial cofibration. First, since i is a trivial cofibration, it is in particular a cofibration and therefore j is again a cofibration by Proposition 2.1. Secondly, since $i \in \mathcal{J}$, by part (i) of Lemma 3.1, it is a cofibration and a strong homotopy equivalence. Since its codomain is cofibrant, j is a strong homotopy equivalence by Lemma 3.2. But now j is both a cofibration and a strong homotopy equivalence and hence it is a trivial cofibration, as required, by part (ii) of Lemma 3.2. \square

We now discuss our second aim, namely showing how we can build on Theorem 3.3 to define sufficient structure to interpret Π -types. This is not immediate since the result of applying Π_p to a fibration with cofibrant domain need not have cofibrant domain. Suppose that we have fibrations $q: B \rightarrow A$ and $p: A \rightarrow \Gamma$, with all objects both fibrant and cofibrant. We begin by applying

$$\Pi_p: \mathbf{SSet}_{/A} \rightarrow \mathbf{SSet}_{/\Gamma}$$

to $q: B \rightarrow A$, so as to obtain $\Pi_p(q): \Pi_A(B) \rightarrow \Gamma$, which is a fibration by Theorem 3.3. The domain of this map is fibrant but not necessarily cofibrant, so we consider its cofibrant replacement

$$\tilde{\Pi}_A(B) =_{\text{def}} \mathbb{L}(\Pi_A(B)),$$

which comes equipped with a trivial fibration $\varepsilon: \tilde{\Pi}_A(B) \rightarrow \Pi_A(B)$. Finally, we define the fibration $\tilde{\Pi}_p(q): \tilde{\Pi}_A(B) \rightarrow \Gamma$ as the composite

$$\begin{array}{ccc} \tilde{\Pi}_A(B) & \xrightarrow{\varepsilon} & \Pi_A(B) \\ & \searrow \tilde{\Pi}_p(q) & \swarrow \Pi_p(q) \\ & \Gamma & \end{array}$$

This will satisfy the β -rule but not the η -rule for Π -types [TO BE ADDED].

4. The universe

Recall that we work in a constructive set theory with two universes u_1 and u_2 and that we refer to elements of u_1 as small sets. We then define a simplicial set X to be *small*

Definition 4.1.

- (i) We say that a simplicial set X is *small* if X_n is a small set for every $[n] \in \Delta$.
- (ii) We say that a map $f: Y \rightarrow X$ in **SSet** is *small* if for every $x: \Delta[n] \rightarrow X$ the simplicial set Y_x fitting in the pullback square

$$\begin{array}{ccc} Y_x & \longrightarrow & Y \\ \downarrow & \lrcorner & \downarrow f \\ \Delta[n] & \xrightarrow{x} & X \end{array}$$

is small.

By the results in [29] for arbitrary presheaf categories, small maps in **SSet** admit a weak classifier, i.e. a small map $\rho: \bar{V} \rightarrow V$ such that for every small map $f: Y \rightarrow X$ there exists a pullback diagram of the form

$$\begin{array}{ccc} Y & \longrightarrow & \bar{V} \\ \downarrow f & \lrcorner & \downarrow \rho \\ X & \longrightarrow & V \end{array}$$

Letting $X = \Delta[n]$ in this diagram suggests to define V_n as the set of all small maps with codomain $\Delta[n]$. In this way, however, one does not obtain a presheaf since the transition functions will satisfy the functorial laws only up to isomorphism rather than equality. To remedy this, the n -simplices of V are defined instead to be the functors $F: (\Delta/[n])^{\text{op}} \rightarrow \mathbf{Set}$ such that the corresponding map of simplicial sets $\text{El}(F) \rightarrow \Delta[n]$ is small. MORE TO BE ADDED.

Following [15, 39], we consider the pullback

$$\begin{array}{ccc} \bar{U} & \longrightarrow & \bar{V} \\ \downarrow \pi & \lrcorner & \downarrow \rho \\ U & \longrightarrow & V \end{array}$$

where $U \subseteq V$ is defined by letting

$$U_n = \{F \in V_n \mid \text{El}(F) \rightarrow \Delta[n] \text{ is a small Kan fibration} \}$$

Proposition 4.2.

- (i) $\pi: \bar{U} \rightarrow U$ is a small Kan fibration.
- (ii) $\pi: \bar{U} \rightarrow U$ classifies small Kan fibrations, i.e. for every small Kan fibration $f: Y \rightarrow X$ there exists a pullback diagram of the form

$$\begin{array}{ccc} Y & \longrightarrow & \bar{U} \\ \downarrow f & \lrcorner & \downarrow \pi \\ X & \longrightarrow & U \end{array}$$

- (iii) The simplicial set \bar{U} is cofibrant.

Proof. We prove the three claims separately.

- (i) Should follow by locality.
- (ii) Should be immediate.
- (iii) See handwritten notes. Key step is the constructive version of the Eilenberg-Zilber lemma. \square

However, the simplicial set \mathbf{U} does not appear to be cofibrant and hence it does not seem possible to show that $\pi: \overline{\mathbf{U}} \rightarrow \mathbf{U}$ is a weak classifier for small Kan fibrations with cofibrant codomain. In order to remedy this, we consider the cofibrant replacement \mathbf{U}_c of \mathbf{U} , which comes equipped with a trivial fibration $p: \mathbf{U}_c \rightarrow \mathbf{U}$, and the pullback

$$\begin{array}{ccc} \overline{\mathbf{U}}_c & \longrightarrow & \overline{\mathbf{U}} \\ \pi_c \downarrow & \lrcorner & \downarrow \pi \\ \mathbf{U}_c & \xrightarrow{p} & \mathbf{U} \end{array}$$

We can now prove that $\pi_c: \overline{\mathbf{U}}_c \rightarrow \mathbf{U}_c$ has the desired properties.

Proposition 4.3.

- (i) $\pi_c: \overline{\mathbf{U}}_c \rightarrow \mathbf{U}_c$ is a small Kan fibration with fibrant codomain.
- (ii) The map $\pi_c: \overline{\mathbf{U}}_c \rightarrow \mathbf{U}_c$ classifies small Kan fibrations with cofibrant domain, i.e. for every small Kan fibration $f: Y \rightarrow X$ with X cofibrant there exists a pullback diagram of the form

$$\begin{array}{ccc} Y & \longrightarrow & \overline{\mathbf{U}}_c \\ f \downarrow & & \downarrow \pi_c \\ X & \longrightarrow & \mathbf{U}_c \end{array}$$

- (iii) The simplicial set $\overline{\mathbf{U}}_c$ is cofibrant.

Proof. Part (i) follows from part (i) of Proposition 4.2. For part (ii), let $f: Y \rightarrow X$ be a small Kan fibration with X cofibrant. Since f is a small Kan fibration, we know from Proposition 4.2 that there is a pullback diagram of the form

$$\begin{array}{ccc} Y & \longrightarrow & \overline{\mathbf{U}} \\ f \downarrow & \lrcorner & \downarrow \pi \\ X & \longrightarrow & \mathbf{U} \end{array}$$

Since X is cofibrant, we have the lifting diagram

$$\begin{array}{ccc} 0 & \longrightarrow & \mathbf{U}_c \\ \downarrow & \nearrow & \downarrow p \\ X & \longrightarrow & \mathbf{U} \end{array}$$

which shows that the map $X \rightarrow \mathbf{U}$ factors via \mathbf{U}_c . We then obtain the diagram

$$\begin{array}{ccccc} Y & \longrightarrow & \overline{\mathbf{U}}_c & \longrightarrow & \overline{\mathbf{U}} \\ f \downarrow & & \downarrow \pi_c & \lrcorner & \downarrow \pi \\ X & \longrightarrow & \mathbf{U}_c & \xrightarrow{p} & \mathbf{U} \end{array}$$

Here, the right-hand side square and the rectangle are pullbacks and therefore the left-hand side square is also a pullback, as required. Part (iii) follows from the fact that both \mathcal{U}_c and $\bar{\mathcal{U}}$ are cofibrant, the latter being part (iii) of Proposition 4.2. \square

Note that we have not shown yet that \mathcal{U}_c fibrant. This will be done in Section 6, as a consequence of the equivalence extension property for fibrations, which we establish in Section 5.

5. The equivalence extension property

- Here we follow Kapulkin and Lumsdaine.

6. Fibrancy and univalence of the universe

- Fibrancy should follow directly from equivalence extension property, without using ‘composition vs filling’ but rather retract property for horns (see notes).
- Once we have established fibrancy of U_c , then one can prove univalence by showing that $t: \mathbf{Weq}(U_c) \rightarrow U_c$ is a trivial fibration.
- Question: do we need to know that $\mathbf{Weq}(U_c)$ is a cofibrant object to get univalence?

7. Semantics

- This should be essentially straightforward, following Kapulkin and Lumsdaine, but we may need to modify the notion of a Π -structure to accommodate the cofibrant replacements that we take for Π .

References

- [1] S. Awodey. Notes on cubical models of type theory. Draft manuscript, March 2015.
- [2] S. Awodey and M. A. Warren. Homotopy theoretic models of identity types. *Mathematical Proceedings of the Cambridge Philosophical Society*, 146:45–55, 2009.
- [3] M. Batanin, D.-C. Cisinski, and M. Weber. Multitensor lifting and strictly unital higher category theory. *Theory and Applications of Categories*, 28:804–856, 2013.
- [4] J. Bénabou and J. Roubaud. Monades et descente. *C. R. Acad. Sc. Paris*, 270, Serie A:96–98, 1970.
- [5] B. van den Berg and R. Garner. Types are weak ω -groupoids. *Proceedings of the London Mathematical Society*, 102(3):370–394, 2010.
- [6] B. van den Berg and R. Garner. Topological and simplicial models of identity types. *Transactions of the ACM on Computational Logic*, 13(1):3–44, 2012.
- [7] B. Berger and I. Moerdijk. On an extension of the notion of Reedy category. *Mathematische Zeitschrift*, 269(3-4):977–1004, 2011.
- [8] J. Bergner and C. Rezk. Reedy categories and the Θ -construction. *Mathematische Zeitschrift*, 274(1):499–514, 2013.
- [9] M. Bezem, T. Coquand, and S. Huber. A model of type theory in cubical sets. In Ralph Matthes and Aleksey Schubert, editors, *19th International Conference on Types for Proofs and Programs (TYPES 2013)*, volume 26, pages 107–128. Schloss Dagstuhl — Leibniz-Zentrum für Informatik, 2014.
- [10] M. Bezem, T. Coquand, and E. Parmann. Non-constructivity in Kan simplicial sets. In Thorsten Altenkirch, editor, *13th International Conference on Typed Lambda Calculi and Applications (TLCA 2015)*, volume 38, pages 92–106. Schloss Dagstuhl — Leibniz-Zentrum für Informatik, 2015.
- [11] J. Bourke and R. Garner. Algebraic weak factorisation systems I: accessible AWFS. *Journal of Pure and Applied Algebra*, 220:108–147, 2016.
- [12] J. Bourke and R. Garner. Algebraic weak factorisation systems II: categories of weak maps. *Journal of Pure and Applied Algebra*, 220:148–174, 2016.
- [13] A. K. Bousfield. Constructions of factorisation systems in categories. *Journal of Pure and Applied Algebra*, 9:207–220, 1977.
- [14] D.-C. Cisinski. Les préfaisceaux comme modèles des types d’homotopie. *Astérisque*, 308:xxiv+392, 2006.
- [15] D.-C. Cisinski. Univalent universes for elegant models of homotopy types. arXiv:1406.0058, 2014.
- [16] M. M. Clementino, E. Giuli, and W. Tholen. Topology in a category: compactness. *Portugaliae Mathematica*, 53(4):397–433, 1996.
- [17] C. Cohen, T. Coquand, S. Huber, and A. Mörtberg. Cubical type theory: a constructive interpretation of the univalence axiom. arXiv:1611.02108. To be published in the post-proceedings of the 21st International Conference on Types for Proofs and Programs, TYPES 2015, 2016.
- [18] J. Emmenegger. A category-theoretic version of the identity type weak factorization system. arXiv:1412.0153, 2014.
- [19] P. Gabriel and M. Zisman. *Calculus of fractions and homotopy theory*, volume 35 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, 1967.
- [20] N. Gambino and R. Garner. The identity type weak factorisation system. *Theoretical Computer Science*, 409:94–109, 2008.
- [21] N. Gambino and J. Kock. Polynomial functors and polynomial monads. *Mathematical Proceedings of the Cambridge Philosophical Society*, 154(1):153–192, 2013.
- [22] R. Garner. A homotopy-theoretic universal property of Leinster’s operad for weak ω -categories. *Mathematical Proceedings of the Cambridge Philosophical Society*, 147:615–628, 2009.
- [23] R. Garner. Understanding the small object argument. *Applied Categorical Structures*, 17(3):247–285, 2009.
- [24] R. Garner. Homomorphisms of higher categories. *Advances in Mathematics*, 224(6):2269–2311, 2010.
- [25] R. Garner and S. Lack. On the axioms for adhesive and quasiadhesive categories. *Theory and Applications of Categories*, 27(3):27–46, 2012.
- [26] P. Goerss and J. F. Jardine. *Simplicial homotopy theory*. Birkhäuser, 1999.
- [27] M. Grandis and W. Tholen. Natural weak factorisation systems. *Archivum Mathematicum*, 42:397–408, 2006.
- [28] P. Hirschhorn. *Model categories and their localizations*. American Mathematical Society, 2003.

- [29] M. Hofmann and T. Streicher. Lifting grothendieck universes. Available from the second-named author's web page, 1997.
- [30] M. Hovey. *Model categories*. American Mathematical Society, 1999.
- [31] S. Huber. A model of type theory in cubical sets. Licentiate of philosophy thesis, University of Gothenburg, 2015.
- [32] J. M. E. Hyland. First steps in synthetic domain theory. In M.-C. Pedicchio A. Carboni and G. Rosolini, editors, *Category Theory*, volume 1488 of *Lecture Notes in Mathematics*, pages 280–301. Springer, 1991.
- [33] P. T. Johnstone. *Sketches of an elephant: a Topos theory compendium*. Oxford Logic Guides. Oxford University Press, New York, NY, 2002.
- [34] A. Joyal. The theory of quasi-categories and its applications. Quaderns 45, Centre de Recerca Matemàtica, 2008.
- [35] A. Joyal and M. Tierney. An introduction to simplicial homotopy theory. Available from <http://hopf.math.purdue.edu/Joyal-Tierney/JT-chap-01.pdf>, 1999.
- [36] A. Joyal and M. Tierney. Quasi-categories vs Segal spaces. In *Categories in algebra, geometry and mathematical physics*, volume 431 of *Contemp. Math.*, pages 277–326. American Mathematical Society, 2007.
- [37] A. Joyal and M. Tierney. Notes on simplicial homotopy theory. Lecture notes, available at <http://mat.uab.cat/~kock/crm/hocat/advanced-course/Quadern47.pdf>, 2008.
- [38] K. Kamps and T. Porter. *Abstract homotopy and simple homotopy theory*. World Scientific Publishing Co., 1997.
- [39] C. Kapulkin and P. LeFanu Lumsdaine. The simplicial model of Univalent Foundations (after Voevodsky). arXiv:1211.2851v4, 2016.
- [40] G. M. Kelly and R. Street. Review of the elements of 2-categories. In *Category Seminar*, volume 420 of *Lecture Notes in Mathematics*. Springer, 1974.
- [41] S. Mac Lane. *Categories for the working mathematician*. Springer, second edition, 1998.
- [42] F. W. Lawvere. Adjointness in foundations. *Dialectica*, 23:281–296, 1969.
- [43] F. W. Lawvere. Equality in hyperdoctrines and comprehension schema as an adjoint functor. In *Proceedings of the American Mathematical Society Symposium on Pure Mathematics XVII*, pages 1–14, 1970.
- [44] J. Lurie. *Higher topos theory*. Number 170 in Annals of Mathematics Studies. Princeton University Press, 2009.
- [45] I. Moerdijk and J. Nuiten. Minimal fibrations of dendroidal sets. arXiv:1509.01073, 2015.
- [46] B. Nordström, K. Petersson, and J. Smith. Martin-löf type theory. In S. Abramsky, D. M. Gabbay, and T. S. E. Maibaum, editors, *Handbook of Logic in Computer Science*, Oxford Logic Guides, chapter V, pages 1–37. Oxford University Press, 2001.
- [47] I. Orton and A. M. Pitts. Axioms for modelling cubical type theory in a topos. In *25th EACSL Annual Conference on Computer Science Logic (CSL 2016)*, volume 62 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 24:1–24:19, Dagstuhl, Germany, 2016. Schloss Dagstuhl–Leibniz-Zentrum für Informatik.
- [48] A. M. Pitts. Nominal presentation of cubical sets models of type theory. In H. Herbelin, P. Letouzey, and M. Sozeau, editors, *20th International Conference on Types for Proofs and Programs (TYPES 2014)*. Schloss Dagstuhl — Leibniz-Zentrum für Informatik, 2015.
- [49] D. G. Quillen. *Homotopical algebra*, volume 43 of *Lecture Notes in Mathematics*. Springer, 1967.
- [50] E. Riehl. Monoidal algebraic model structures. *Journal of Pure and Applied Algebra*, 217:1069–1104, 2013.
- [51] E. Riehl and D. Verity. The theory and practice of Reedy categories. *Theory and Applications of Categories*, 29(9):256–301, 2014.
- [52] E. Riehl. Algebraic model structures. *New York Journal of Mathematics*, 17:173–231, 2011.
- [53] E. Riehl. *Categorical homotopy theory*. Cambridge University Press, 2014.
- [54] J. Rosický and W. Tholen. Factorization, fibration and torsion. *Journal of Homotopy and Related Structures*, 2(295–314), 2007.
- [55] G. Rosolini. *Continuity and effectiveness in topoi*. PhD thesis, University of Oxford, 1986.
- [56] M. Shulman. Univalence for inverse diagrams and homotopy canonicity. *Mathematical Structures in Computer Science*, 25:1203–1277, 2015.
- [57] M. Shulman. The univalence axiom for elegant Reedy presheaves. *Homology, Homotopy and Applications*, To appear.
- [58] A. Swan. An algebraic weak factorisation system on 01-substitution sets: a constructive proof. arXiv:1409.1829, 2014.
- [59] V. Voevodsky. Univalent foundations project. http://www.math.ias.edu/vladimir/files/univalent_foundations_project.pdf, 2010.
- [60] V. Voevodsky. The equivalence axiom and univalent models of type theory. (talk at cmu on february 4, 2010). arXiv:1402.5556v2, 2014.

- [61] M. A. Warren. *Homotopy theoretic aspects of constructive type theory*. PhD thesis, Carnegie Mellon University, 2008.