

6.7900: Machine Learning

Lecture 6

Lecture start: Tues/Thurs 2:35pm

Who's speaking today? Prof. Tamara Broderick

Course website: gradml.mit.edu

Questions? Ask here or on piazza.com/mit/fall2024/67900/

Materials: Slides, video, etc linked from gradml.mit.edu after the lecture (but there is no livestream)

Last Time

- I. Challenges with MLE/ERM for linear regression
- II. Bayesian linear regression
 - A. Bayes with Gaussians
 - B. Posterior

Today

- I. Visualizations
- II. Uncertainty
- III. Ridge regression
- IV. More flexible/complex features

6.7900: Machine Learning

Lecture 6

Lecture start: Tues/Thurs 2:35pm

Who's speaking today? Prof. Tamara Broderick

Course website: gradml.mit.edu

Questions? Ask here or on piazza.com/mit/fall2024/67900/

Materials: Slides, video, etc linked from gradml.mit.edu after the lecture (but there is no livestream)

Last Time

- I. Challenges with MLE/ERM for linear regression
- II. Bayesian linear regression
 - A. Bayes with Gaussians
 - B. Posterior

Today

- I. Visualizations
- II. Uncertainty
- III. Ridge regression
- IV. More flexible/complex features

See board
and demos

Some points related to uncertainty

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.
 - Before I shuffle, uncertainty is aleatoric. After shuffling but before looking, uncertainty is epistemic.

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.
 - Before I shuffle, uncertainty is aleatoric. After shuffling but before looking, uncertainty is epistemic.

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

predictive vs
posterior

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.
 - Before I shuffle, uncertainty is aleatoric. After shuffling but before looking, uncertainty is epistemic.

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

predictive vs
posterior

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.
 - Before I shuffle, uncertainty is aleatoric. After shuffling but before looking, uncertainty is epistemic.
- The Bayesian posterior uncertainty is conditional on our model. Goes to zero if enough data with enough x coverage.

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

predictive vs
posterior

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.
 - Before I shuffle, uncertainty is aleatoric. After shuffling but before looking, uncertainty is epistemic.
- The Bayesian posterior uncertainty is conditional on our model. Goes to zero if enough data with enough x coverage.
 - “Uncertainty” as a term of art vs colloquial meaning

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

predictive vs
posterior

Some points related to uncertainty

- Rather than return an error in the face of collinearity, the Bayesian approach expresses uncertainty over possibilities
- It's common to see publications discuss:
 - **Aleatoric uncertainty**: “intrinsic randomness”
 - **Epistemic uncertainty**: “lack of knowledge” about something that we could know for sure in principle
- Example: I plan to shuffle a deck of cards and then ask whether the top card is an ace.
 - Before I shuffle, uncertainty is aleatoric. After shuffling but before looking, uncertainty is epistemic.
- The Bayesian posterior uncertainty is conditional on our model. Goes to zero if enough data with enough x coverage.
 - “Uncertainty” as a term of art vs colloquial meaning
 - Not just a Bayesian issue: (frequentist) standard errors go to 0 as data grows. So in real-life (“all models are wrong”), common p-values get arbitrarily small from enough data

[Aldous, “Is it practical and useful to distinguish aleatoric and epistemic uncertainty?”]

predictive vs
posterior

MAP and regularization

- What if I just want a single regression line and not a whole distribution?

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP.

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D})$$

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta|\mathcal{D}) = \arg \min_{\theta} -\log p(\theta|\mathcal{D})$$

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta|\mathcal{D}) = \arg \min_{\theta} -\log p(\theta|\mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used,
now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

$$= \arg \min_{\theta} \left\{ (X\theta - Y)^{\top} (X\theta - Y) + \frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta \right\}$$

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used,
now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

$$= \arg \min_{\theta} \left\{ (X\theta - Y)^{\top} (X\theta - Y) + \frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta \right\}$$

the usual MLE/ERM
objective

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used,
now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

$$= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^\top (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^\top \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$$

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used,
now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge
regression

$$= \arg \min_{\theta} \left\{ (X\theta - Y)^{\top} (X\theta - Y) + \frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta \right\}$$

the usual MLE/ERM
objective

(ℓ_2 /ridge) penalty/
regularizer

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^{\top} (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^\top (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^\top \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^{\top} (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
 - Letting $\lambda = \sigma^2 / \sigma_0^2$,

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^{\top} (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
- Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^{\top} X)^{-1} (X^{\top} Y)$

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^\top (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^\top \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
- Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^\top X)^{-1} (X^\top Y)$

What happens as λ gets very small?

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^\top (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^\top \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
- Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^\top X)^{-1} (X^\top Y)$

What happens as λ gets very small? Very large?

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^\top (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^\top \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
 - Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^\top X)^{-1} (X^\top Y)$
- Avoids the inversion issue of the MLE by choosing a particular option among options “equally good” in MLE

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^\top (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^\top \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
 - Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^\top X)^{-1} (X^\top Y)$
- Avoids the inversion issue of the MLE by choosing a particular option among options “equally good” in MLE check: always invertible

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^{\top} (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
 - Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^{\top} X)^{-1} (X^{\top} Y)$
- Avoids the inversion issue of the MLE by choosing a particular option among options “equally good” in MLE check: always invertible
 - No uncertainty (e.g. when regression is underdetermined)

MAP and regularization

- What if I just want a single regression line and not a whole distribution? Some options: posterior mean or MAP. Lecture 3:

$$\hat{\theta}_{\text{MAP}} = \arg \max_{\theta} p(\theta | \mathcal{D}) = \arg \min_{\theta} -\log p(\theta | \mathcal{D})$$

- consider the linear regression model we used, now with $\mu_0 = 0_D$ & $\Sigma_0 = \sigma_0^2 I_{D \times D}$

ridge regression $= \arg \min_{\theta} \left\{ \underbrace{(X\theta - Y)^{\top} (X\theta - Y)}_{\text{the usual MLE/ERM objective}} + \underbrace{\frac{\sigma^2}{\sigma_0^2} \theta^{\top} \theta}_{(\ell_2/\text{ridge}) \text{ penalty/regularizer}} \right\}$

[“ridge analysis” RW Hoerl 2020, Ridge Regression: A Historical Context; can think of “ridge” here due to collinearity in likelihood]

- Since the Gaussian posterior is symmetric, the MAP is equal to the mean here. So we already computed it in Lecture 5.
 - Letting $\lambda = \sigma^2 / \sigma_0^2$, $\hat{\theta}_{\text{MAP}} = (\lambda I_{D \times D} + X^{\top} X)^{-1} (X^{\top} Y)$
- Avoids the inversion issue of the MLE by choosing a particular option among options “equally good” in MLE check: always invertible
 - No uncertainty (e.g. when regression is underdetermined)
 - May offer better generalization than vanilla MLE

A note on features

A note on features

- Linear models can be very flexible given non-trivial features

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
- E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
- E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ recall demo

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
 - E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ recall demo
 - More generally, for $x \in \mathbb{R}^{D_x}$, $\phi(x)$ could collect all polynomials of some degree r or smaller

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
 - E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ [recall demo](#)
 - More generally, for $x \in \mathbb{R}^{D_x}$, $\phi(x)$ could collect all polynomials of some degree r or smaller
 - Lots of other options: e.g. Fourier basis, logistic basis, wavelets, splines

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
 - E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ [recall demo](#)
 - More generally, for $x \in \mathbb{R}^{D_x}$, $\phi(x)$ could collect all polynomials of some degree r or smaller
 - Lots of other options: e.g. Fourier basis, logistic basis, wavelets, splines
 - For any fixed $\phi(x)$, basically all the math we did goes through with $\phi(x)$ in place of x

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
 - E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ [recall demo](#)
 - More generally, for $x \in \mathbb{R}^{D_x}$, $\phi(x)$ could collect all polynomials of some degree r or smaller
 - Lots of other options: e.g. Fourier basis, logistic basis, wavelets, splines
 - For any fixed $\phi(x)$, basically all the math we did goes through with $\phi(x)$ in place of x
 - E.g. for $N \times D$ matrix Φ with n th row $\phi(x^{(n)})^\top$

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
 - E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ [recall demo](#)
 - More generally, for $x \in \mathbb{R}^{D_x}$, $\phi(x)$ could collect all polynomials of some degree r or smaller
 - Lots of other options: e.g. Fourier basis, logistic basis, wavelets, splines
 - For any fixed $\phi(x)$, basically all the math we did goes through with $\phi(x)$ in place of x
 - E.g. for $N \times D$ matrix Φ with n th row $\phi(x^{(n)})^\top$
 - Then OLS: $\hat{\theta} = (\Phi^\top \Phi)^{-1} \Phi^\top Y$

A note on features

- Linear models can be very flexible given non-trivial features
- We've been considering $h(x) = \theta_1 x_1 + \cdots + \theta_D x_D = \theta^\top x$
- But we could take $h(x) = \theta_1 \phi_1(x) + \cdots + \theta_D \phi_D(x) = \theta^\top \phi(x)$
 - Now x can be any dimension (and need not be real-valued), and D is the dimension of the features $\phi(x)$
 - E.g. for $x \in \mathbb{R}$, $\phi(x) = [1, x, x^2]^\top$ [recall demo](#)
 - More generally, for $x \in \mathbb{R}^{D_x}$, $\phi(x)$ could collect all polynomials of some degree r or smaller
 - Lots of other options: e.g. Fourier basis, logistic basis, wavelets, splines
 - For any fixed $\phi(x)$, basically all the math we did goes through with $\phi(x)$ in place of x
 - E.g. for $N \times D$ matrix Φ with n th row $\phi(x^{(n)})^\top$
 - Then OLS: $\hat{\theta} = (\Phi^\top \Phi)^{-1} \Phi^\top Y$
- Deep neural networks perform nonlinear feature extraction/learning