

A brief of physical processes in the interstellar medium & in star formation

TNU/SAGI Summer School in Observational Astronomy 2024

Buon Ma Thuot - August 2024

Ngân LÊ

Simons Astrophysics Group at ICISE (SAGI)

Institute For Interdisciplinary Research in Science and Education (IFIRSE)

International Centre for Interdisciplinary Science and Education (ICISE), Quy Nhon, Vietnam


09 August 2024

Outline

1. The interstellar medium (ISM) & its main constituents
2. Physical processes in interstellar gas
3. Physical processes in star formation: accretion and ejection processes

The Interstellar Medium

The Interstellar Medium



- Interstellar medium (ISM): everything in the space between stars.
- The ISM is comprised of gas, dust, cosmic rays, interstellar radiation (starlight and CMB), magnetic fields, dark matter, *interstellar objects, and spacecraft*.
- The ISM is filamentary, magnetized, and turbulent.

Composition of the ISM: phases of gas

Basics physical properties of these different phases

Component	Temperature (K)	Density (cm ⁻³)	Ionization fraction	
Warm neutral medium (WNM)	10^4	~ 0.5	~ 0.1	HI gas (atomic)
Cold neutral medium (CNM)	100	20–50	10^{-4}	
Hot ionized medium (HM)	10^6	0.01	1.0	fully ionized HII gas
Warm ionized medium (WIM)	10^4	~ 0.5	1.0	
Molecular gas	10–20	$\gtrsim 100$	$\ll 10^{-4}$	H ₂ , CO, ...

- Atomic gas in the ISM: H (~70% *), He (28% *), and heavier elements (C, N, O, Si, Fe, ... - for 2% *)
- Gas is found in a variety of chemical forms

* by total mass

Composition of the ISM: phases of gas

Table 1.3 Phases of Interstellar Gas


Phase	T (K)	n_H (cm^{-3})	Comments
Coronal gas (HIM) $f_V \approx 0.5?$ $\langle n_H \rangle f_V \approx 0.002 \text{ cm}^{-3}$ ($f_V \equiv$ volume filling factor)	$\gtrsim 10^{5.5}$	~ 0.004	Shock-heated Collisionally ionized Either expanding or in pressure equilibrium Cooling by: ◊ Adiabatic expansion ◊ X ray emission Observed by: • UV and x ray emission • Radio synchrotron emission
H II gas $f_V \approx 0.1$ $\langle n_H \rangle f_V \approx 0.02 \text{ cm}^{-3}$	10^4	$0.3 - 10^4$	Heating by photoelectrons from H, He Photoionized Either expanding or in pressure equilibrium Cooling by: ◊ Optical line emission ◊ Free-free emission ◊ Fine-structure line emission Observed by: • Optical line emission • Thermal radio continuum
Warm HI (WNM) $f_V \approx 0.4$ $n_H f_V \approx 0.2 \text{ cm}^{-3}$	~ 5000	0.6	Heating by photoelectrons from dust Ionization by starlight, cosmic rays Pressure equilibrium Cooling by: ◊ Optical line emission ◊ Fine structure line emission Observed by: • HI 21 cm emission, absorption • Optical, UV absorption lines

Table 1.3 Phases of Interstellar Gas

Phase	T (K)	n_H (cm^{-3})	Comments
Cool HI (CNM) $f_V \approx 0.01$ $n_H f_V \approx 0.3 \text{ cm}^{-3}$	~ 100	30	Heating by photoelectrons from dust Ionization by starlight, cosmic rays Cooling by: ◊ Fine structure line emission Observed by: • HI 21-cm emission, absorption • Optical, UV absorption lines
Diffuse H ₂ $f_V \approx 0.001$ $n_H f_V \approx 0.1 \text{ cm}^{-3}$	$\sim 50 \text{ K}$	~ 100	Heating by photoelectrons from dust Ionization by starlight, cosmic rays Cooling by: ◊ Fine structure line emission Observed by: • HI 21-cm emission, absorption • CO 2.6-mm emission • optical, UV absorption lines
Dense H ₂ $f_V \approx 10^{-4}$ $\langle n_H \rangle f_V \approx 0.2 \text{ cm}^{-3}$	$10 - 50$	$10^3 - 10^6$	Heating by photoelectrons from dust Ionization and heating by cosmic rays Self-gravitating: $p > p(\text{ambient ISM})$ Cooling by: ◊ CO line emission ◊ CI fine structure line emission Observed by: • CO 2.6-mm emission • dust FIR emission
Cool stellar outflows	$50 - 10^3$	$1 - 10^6$	Observed by: • Optical, UV absorption lines • Dust IR emission • HI, CO, OH radio emission

Adapted from lecture by Thiem Hoang
(KASI & UST)


Composition of the ISM: dust (grains)



Herranen et al. 2019

- a mixture of primarily **silicate** and **carbonaceous** compounds, including polycyclic aromatic hydrocarbons (**PAHs**)
- have **aspherical** shapes & sub-micron sizes (~ $0.1 \mu\text{m}$), but can significantly grow in dense gas of protoplanetary disks
- absorb or scatter starlight (energy) in the UV, optical, or near-IR range
- re-emit energy in longer wavelength, from far-IR to sub-mm → thermal dust emission

Main sources of interstellar radiation field (ISRF)




- **Thermal emission from dust grains** heated by starlight (Far-IR to IR wavelength)
- **Starlight** from ionizing photons emitted by stellar photospheres (HI region, Photodissociation region - PDR)
- **X-ray emission** from hot plasma (10^5 - 10^8 K)
- **Emission from ionized plasma** (free-free, free-bound, and bound-bound transitions, 10^4 K)
- Cosmic microwave background (**CMB**) radiation (~ 2.73 K)
- **Galactic synchrotron radiation** from relativistic electrons

Klessen & Glover (2016)

Q1: What which wavelengths the energy density of the ISRF is mainly in?

Main sources of interstellar radiation field (ISRF)



- **Thermal emission from dust grains** heated by starlight (Far-IR to IR wavelength)
- **Starlight** from ionizing photons emitted by stellar photospheres (HI region, Photodissociation region - PDR)
- **X-ray emission** from hot plasma (10^5 - 10^8 K)
- **Emission from ionized plasma** (free-free, free-bound, and bound-bound transitions, 10^4 K)
- Cosmic microwave background (**CMB**) radiation (~ 2.73 K)
- **Galactic synchrotron radiation** from relativistic electrons

Klessen & Glover (2016)

Most of the energy density of the ISRF is in IR (**thermal dust and CMB emission**), optical and UV (**starlight**) range → key role in regulating ISM's properties

Physical processes in interstellar gas

Collisional Processes

- Collisions: fundamental to the physics of the ISM
- Collisional rate: a rate at which a particle can collide with other particle of different masses
- Collisional rate coefficient ?

$$\langle \sigma_{AB} \nu \rangle = \left(\frac{8}{\pi \nu k_B T} \right)^{1/2} \langle \sigma \rangle, \text{ with } \langle \sigma \rangle = \int_0^{\infty} \frac{\sigma_{AB} E}{k_B T} e^{-E/k_B T} \frac{dE}{dT}$$

?

Collisional Processes

- Assume a collision between 2 particles A and B, with masses of m_A and m_B , and density of n_A and n_B ; B is at rest and A is moving with velocity of v toward B
- reduced mass: $\mu = m_A m_B / (m_A + m_B)$, reduced energy: $E = 0.5 \mu v^2$
- Cross-section of collision: σ_{AB} as a function of v ,
- Number flux of A: $J_A = n_A v$
- Collision rate of A with B: $J_A \sigma_{AB} = n_A v \sigma_{AB}(v)$
- Number of collisions per second per volume: $J_A \sigma_{AB} n_B = n_A n_B v \sigma_{AB}(v)$
- Integration over velo. distribution of A: $I = \int_0^\infty n_A n_B v \sigma_{AB}(v) f(v) dv$
- Collision rate per unit volume $I \equiv n_A n_B \langle \sigma v \rangle = n_A n_B k_{AB}$ (event/cm³/s)
- Collisional rate coefficient: $k_{AB} = \langle \sigma v \rangle = \int_0^\infty v \sigma_v f(v) dv$ (cm⁻³s)
- Assume a thermal gas at temp. T , then the relative velocity will have Maxwellian distribution:

$$f(v) dv = \left(\frac{\mu}{2\pi k_B T} \right)^{3/2} 4\pi v^2 e^{-\mu v^2 / 2k_B T} dv$$

- Using $f(v) dv = f(E) dE$, one gets: $k_{AB} = \langle \sigma v \rangle = \left(\frac{8k_B T}{\pi \mu} \right)^{1/2} \int_0^\infty \sigma_{AB}(E) \frac{E}{k_B T} e^{-E/k_B T} \frac{dE}{dT}$

Collisional Processes

- Collisional rate coefficient: $\langle \sigma_{AB} \nu \rangle = \left(\frac{8}{\pi \nu k_B T} \right)^{1/2} \int_0^\infty \frac{\sigma_{AB} E}{k_B T} e^{-E/k_B T} \frac{dE}{dT}$

- If B is much heavier than A: $m_A \ll m_B, \mu \approx m_A$

- For e.g., electron collides with atoms:

$$\langle \sigma_{AB} \nu \rangle = \left(\frac{8}{\pi \nu k_B T} \right)^{1/2} \langle \sigma \rangle, \text{ with } \langle \sigma \rangle = \int_0^\infty \frac{\sigma_{AB} E}{k_B T} e^{-E/k_B T} \frac{dE}{dT}$$

Physical processes in interstellar gas:

- Four basic types of collisional interactions:
 - between charged particles: (ions-ions, ions-electrons, and electrons-electrons)
—> Coulomb field: $\sim r^{-1}$ —> use the “impact approximation”
 - between ions and neutral particles (atoms, molecules)
—> induced-dipole potential: $\sim r^{-4}$ —> use exact results for scattering
 - between electrons and neutrals —> use experimental data
 - between neutrals —> effective “hard-sphere” radius

Physical processes in interstellar gas:

- Collisional Processes – interaction between gas and gas
- Radiative Processes – interaction between gas and radiation

Physical processes in interstellar gas:

Radiative Transfer Equation:

$$\boxed{\frac{dI_\nu}{ds} = -\alpha_\nu I_\nu + j_\nu}$$

α_ν : absorption (attenuation) coefficient, j_ν : emission coefficient or emissivity

Emission coefficient: $j_\nu = \frac{1}{4\pi} n_u A_{ul} h\nu \phi(\nu)$

Absorption coefficient: $\alpha_\nu = n_l \sigma_{lu} - n_u \sigma_{ul} = -\frac{h\nu}{4\pi} (B_{lu} n_l - B_{ul} n_u) \phi(\nu)$

Source function:


$$S_\nu = \frac{j_\nu}{\alpha_\nu} = \frac{A_{lu} n_u}{(B_{lu} n_l - B_{ul} n_u)} = \frac{2h\nu^3}{c^2} \left(\frac{g_u n_l}{g_l n_u} - 1 \right)^{-1},$$



Solution: $I_\nu = I_\nu(0) e^{-\tau_\nu} + \int_0^{\tau_\nu} e^{-(\tau_\nu - \tau')} S_\nu d\tau'$


Physical processes in star formation

Molecular clouds – the stellar nurseries



- Stars like our Sun form in cold and dense parts of the molecular cloud, which consists of filaments, clumps, and dense cores

Evolutionary sequence of a low-mass star



- Gravitational fragment forms dense core

- Material accretes onto the forming star
- Collimated jets

- Wide-angle outflows dissipate the envelope
- Mass of the star increases, and an embedded disk is formed

- No envelope left, evolution within the gas-rich protoplanetary disk


- Planet formation inside the dust-rich disk, lack of accretion

André 2002

- Mass accretion rate: a key quantity to study evolution of pre-main sequence stars
- Jets and outflows: signature of star formation in earliest stages

Near-IR spectroscopy of YSOs

Lê+, in prep.




- SpeX spectra of YSO candidates cover the range of 0.8–2.5 μm
- sources show the typical signatures of YSOs (H Br γ , H Pa β , and/or CO bandhead)
- calculated from empirical relation with line luminosity of H Br γ and/or H Pa β

$$\log(L_{\text{acc}}) = -0.7 + 0.9 [\log(L_{\text{Br}\gamma}) + 4] \quad (\text{Calvet+2004})$$

$$\log(L_{\text{acc}}) = 1.8 + 1.03 \log(L_{\text{Pa}\beta}) \quad (\text{Muzerolle+1998})$$


Accretion luminosities



- calculated using empirical relation with line luminosity of H Br γ and/or H Pa β


Gaskinematics: low- J CO lines

Lê + in prep., APEX PI 230 (OGHReS, König +2021)



- Integrated line intensities of CO 2-1 show relatively low-density gas associated with the filament in the CMa- ℓ 224 region
- SPIRE 250 μm on 0.1 pc scales shows the region observed with FIFI-LS

Maps of CO 2–1 isotopologues



- ¹²CO 2-1 is extended perpendicular to the far-IR and C¹⁸O line emission; traces either the filament or some outflows