

Towards porting PLN to MeTTa

Nil Geisweiller, Hedra Yusuf

AGI-22 Workshop



SingularityNET



PLN Recall

$P, Q, \dots : Atom^n \rightarrow \{\top, \perp\}$ (possibly fuzzy)

Atomese	MeTTa	Math
$(P \text{ tv})$	$(\equiv P \text{ tv})$	$\Pr(\mathcal{S}(P)) \approx \text{tv.mean}$
$(\text{Not } \text{tv } P)$	$(\equiv (\neg P) \text{ tv})$	$\Pr(\overline{\mathcal{S}(P)}) \approx \text{tv.mean}$
$(\text{Or } \text{tv } P \text{ } Q)$	$(\equiv (\vee P \text{ } Q) \text{ tv})$	$\Pr(\mathcal{S}(P) \cup \mathcal{S}(Q)) \approx \text{tv.mean}$
$(\text{And } \text{tv } P \text{ } Q)$	$(\equiv (\wedge P \text{ } Q) \text{ tv})$	$\Pr(\mathcal{S}(P) \cap \mathcal{S}(Q)) \approx \text{tv.mean}$
$(\text{Implication } \text{tv } P \text{ } Q)$	$(\equiv (P \rightarrow Q) \text{ tv})$	$\Pr(\mathcal{S}(Q) \mathcal{S}(P)) \approx \text{tv.mean}$
$(\text{Evaluation } \text{tv } P \text{ (List } X_1 \dots X_n))$	$(\equiv (P X_1 \dots X_n) \text{ tv})$	$\Pr(P(X_1, \dots, X_n) = \top) \approx \text{tv.mean}$

PLN rules: Deduction

$$\frac{P \rightarrow Q \stackrel{m}{=} tv_{PQ} \quad Q \rightarrow R \stackrel{m}{=} tv_{QR} \quad P \stackrel{m}{=} tv_P \quad Q \stackrel{m}{=} tv_Q \quad R \stackrel{m}{=} tv_R}{P \rightarrow R \stackrel{m}{=} tv} \text{ (DED)}$$

$$tv.mean = tv_{PQ}.mean \times tv_{QR}.mean + \frac{(1 - tv_{PQ}.mean) \times (tv_R.mean - tv_Q.mean \times tv_{QR}.mean)}{1 - tv_Q.mean}$$

PLN rules: Implication Direct Introduction

$$\frac{(P \ a_1) \stackrel{m}{=} tv_{Pa_1} \quad (Q \ a_1) \stackrel{m}{=} tv_{Qa_1} \quad \dots \quad (P \ a_n) \stackrel{m}{=} tv_{Pa_n} \quad (Q \ a_n) \stackrel{m}{=} tv_{Qa_n}}{P \rightarrow Q \stackrel{m}{=} tv} \text{ (IDI)}$$

$$tv.mean = \frac{\sum_x tv_{Px}.mean \times tv_{Qx}.mean}{\sum_x tv_{Px}.mean}$$

Evidential Truth Value

```
;; TruthValue type and constructors
(: TruthValue Type)
(: Bl (-> Bool TruthValue))
(: Pr (-> Number TruthValue))
(: PrCnt (-> Number Number TruthValue))

;; TruthValue methods
(: mode (-> TruthValue Number))
(: mean (-> TruthValue Number))
(: pos_count (-> TruthValue Number))
(: neg_count (-> TruthValue Number))

;; EvidentialTruthValue type and constructors
(: EvidentialTruthValue Type)
(: ETV (-> (Set $a) TruthValue EvidentialTruthValue))
```

Atomese	MeTTa	Math
(stv s (count->confidence c))	(PrCnt s c)	<s c>
-	(ETV E tv)	(E, tv)

PLN rules: Implication Direct Introduction (Recursive Decomposition)

- Base

$$\frac{}{P \rightarrow Q \stackrel{m}{=} (\emptyset, \langle 1 \ 0 \rangle)} \text{ (IDI Axiom)}$$

- Recursion

$$\frac{(P \ a) \stackrel{m}{=} tv_{Pa} \quad (Q \ a) \stackrel{m}{=} tv_{Qa} \quad P \rightarrow Q \stackrel{m}{=} (E, tv_{PQ}) \quad a \notin E}{P \rightarrow Q \stackrel{m}{=} (\{a\} \cup E, tv)} \text{ (IDI Induction)}$$

$$tv.count = tv_{PQ}.count + tv_{Pa}.mean$$

$$tv.mean = \frac{tv_{PQ}.pos_count + tv_{Pa}.mean \times tv_{Qa}.mean}{tv.count}$$

Temporal Deduction \mapsto Deduction

$$\frac{
 \frac{P \rightsquigarrow^{T_1} Q}{P \rightarrow \tilde{Q}^{T_1}} \text{ (PI2I)} \quad
 \frac{
 \frac{Q \rightsquigarrow^{T_2} R}{Q \rightarrow \tilde{R}^{T_2}} \text{ (PI2I)} \quad
 \frac{\tilde{Q}^{T_1} \rightarrow \tilde{R}^{T_1+T_2}}{\tilde{Q}^{T_1} \rightarrow \tilde{R}^{T_1+T_2}} \text{ (TS)}
 }{P \rightarrow \tilde{R}^{T_1+T_2}} \text{ (I2PI)}
 }{
 \frac{
 \frac{Q}{\tilde{Q}^{T_1}} \text{ (TS)} \quad
 \frac{R}{\tilde{R}^{T_1+T_2}} \text{ (TS)}
 }{P \rightsquigarrow^{T_1+T_2} R} \text{ (DED)}
 }$$

- DED: Deduction
- TS: Temporal Shift
- PI2I: PredictiveImplication to Implication
- I2PI: Implication to PredictiveImplication

Temporal Procedural Deduction \mapsto Deduction

$$\begin{array}{c}
 \frac{P \wedge \hat{A} \rightsquigarrow^{T_1} Q}{P \wedge \hat{A} \rightarrow \tilde{Q}^{T_1}} \text{ (PI2I)} \quad \frac{\hat{B}}{\tilde{B}^{T_1}} \text{ (TS)} \quad \frac{Q \wedge \hat{B} \rightsquigarrow^{T_2} R}{Q \wedge \hat{B} \rightarrow \tilde{R}^{T_2}} \text{ (PI2I)} \\
 \frac{P \wedge \hat{A} \wedge \tilde{B}^{T_1} \rightarrow \tilde{Q}^{T_1} \wedge \tilde{B}^{T_1}}{P \wedge \hat{A} \wedge \tilde{B}^{T_1} \rightarrow \tilde{Q}^{T_1} \wedge \tilde{B}^{T_1}} \text{ (CI)} \quad \frac{\tilde{Q}^{T_1} \wedge \tilde{B}^{T_1} \rightarrow \tilde{R}^{T_1+T_2}}{\tilde{Q}^{T_1} \wedge \tilde{B}^{T_1} \rightarrow \tilde{R}^{T_1+T_2}} \text{ (TS)} \quad P \wedge \hat{A} \wedge \tilde{B}^{T_1} \quad \frac{Q \wedge \hat{B}}{\tilde{Q}^{T_1} \wedge \tilde{B}^{T_1}} \text{ (TS)} \quad \frac{R}{\tilde{R}^{T_1+T_2}} \text{ (TS)} \\
 \hline
 \frac{P \wedge \hat{A} \wedge \tilde{B}^{T_1} \rightarrow \tilde{R}^{T_1+T_2}}{((P \wedge \hat{A}) \wedge^{T_1} \hat{B}) \rightsquigarrow^{T_1+T_2} R} \text{ (I2PI)} \quad \frac{}{\tilde{R}^{T_1+T_2}} \text{ (DED)}
 \end{array}$$

- CI: Conjunction Introduction
- TS: Temporal Shift
- DED: Deduction
- PI2I: PredictiveImplication to Implication
- I2PI: Implication to PredictiveImplication

Conclusion

Demo Time