

# Partial Operator Induction with Beta Distribution

---

Nil Geisweiller

AGI-18

Prague



SingularityNET



NOVAMENTE

Problem:

Combining Models from Different Contexts

Theory:

Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

Problem:

Combining Models from Different Contexts

Theory:

Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

Problem:

Combining Models from Different Contexts

Theory:


Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

## Problem: Models from different contexts


How to combine models obtained from different contexts?



Large Contexts  $\rightarrow$  Underfit


Small Contexts  $\rightarrow$  Overfit

## Problem: Preserve Uncertainty



# Problem: Preserve Uncertainty

## Exploration vs Exploitation (Thompson Sampling)



# Problem: ImplicationLink

ImplicationLink <TV>

R

S

≡

Second Order

$P(S|R)$


Beta Distribution in disguise




# Solution

Bayesian Model Averaging / Solomonoff Operator Induction,  
modified to:

1. Support partial models



2. Produce a probability distribution estimate, rather than probability estimate.



3. Specialize for Beta distributions

Problem:

Combining Models from Different Contexts

Theory:

Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

Problem:

Combining Models from Different Contexts

Theory:

Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

# Solomonoff Operator Induction

## Bayesian Model Averaging + Universal Distribution

Probability Estimate:

$$\hat{P}(A_{n+1}|Q_{n+1}) = \sum_j a_0^j \prod_{i=1}^{n+1} O^j(A_i|Q_i)$$

where:

- $Q_i = i^{th}$  question
- $A_i = i^{th}$  answer
- $O^j = j^{th}$  operator
- $a_0^j =$  prior of  $j^{th}$  operator

## Specialization of Solomonoff Operator Induction

OpenCog implication link

```
ImplicationLink <TV>  
  R  
  S
```

≡

Class of parameterized operators

$$O_p^j(A_i|Q_i) = \text{if } R^j(Q_i) \text{ then } \begin{cases} p, & \text{if } A_i = A_{n+1} \\ 1 - p, & \text{otherwise} \end{cases}$$

# Beta Distribution

Probability Density Function:

$$pdf_{\alpha,\beta}(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)}$$


Beta Function:

$$B_x(\alpha, \beta) = \int_0^x p^{\alpha-1}(1-p)^{\beta-1} dp$$

$$B(\alpha, \beta) = B_1(\alpha, \beta)$$

Conjugate Prior:

$$pdf_{m+\alpha, n-m+\beta}(x) \propto x^m(1-x)^{n-m} pdf_{\alpha,\beta}(x)$$



$$O_p^j(A_i|Q_i) = \text{if } R^j(Q_i) \text{ then } \begin{cases} p, & \text{if } A_i = A_{n+1} \\ 1 - p, & \text{otherwise} \end{cases}$$

---

Data

$$O_{p,C}^j(A_i|Q_i) = \begin{cases} \text{if } R^j(Q_i) \text{ then } \begin{cases} p, & \text{if } A_i = A_{n+1} \\ 1 - p, & \text{otherwise} \end{cases} \\ \text{else } C(A_i|Q_i) \end{cases}$$

---

Data




# Second Order Solomonoff Operator Induction

Probability Estimate:

$$\hat{P}(A_{n+1}|Q_{n+1}) = \sum_j a_0^j \prod_{i=1}^{n+1} O^j(A_i|Q_i)$$

Probability Distribution Estimate:

$$\hat{cdf}_{(A_{n+1}|Q_{n+1})}(x) = \sum_{O^j(A_{n+1}|Q_{n+1}) \leq x} a_0^j \prod_{i=1}^n O^j(A_i|Q_i)$$



# Combing Solomonoff Operator Induction and Beta Distributions

$$\hat{cdf}_{(A_{n+1}|Q_{n+1})}(x) \propto \sum_j a_0^j r^j B_x(m^j + \alpha, n^j - m^j + \beta) B(m^j + \alpha, n^j - m^j + \beta)$$

where

- $n^j$  = number of observations explained by  $j^{th}$  model
- $m^j$  = number of true observations explained by  $j^{th}$  model
- $r^j$  = likelihood of the unexplained data

$r^j = ???$

# Combing Solomonoff Operator Induction and Beta Distributions

$$\hat{cdf}_{(A_{n+1}|Q_{n+1})}(x) \propto \sum_j a_0^j r^j B_x(m^j + \alpha, n^j - m^j + \beta) B(m^j + \alpha, n^j - m^j + \beta)$$

where

- $n^j$  = number of observations explained by  $j^{th}$  model
- $m^j$  = number of true observations explained by  $j^{th}$  model
- $r^j$  = likelihood of the unexplained data

$$r^j = ??? \approx 2^{-v(1-c)}$$

- $v = n - n^j$  = number of unexplained observations
- $c$  = compressability parameter
  - $c = 1 \rightarrow$  explains remaining data
  - $c = 0 \rightarrow$  can't explain remaining data

Problem:

Combining Models from Different Contexts

Theory:

Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

Problem:

Combining Models from Different Contexts

Theory:

Solomonoff Operator Induction and Beta Distribution

Practice:

Inference Control Meta-Learning

# Inference Control Meta-learning

Learn how to reason efficiently

# Inference Control Meta-learning

Learn how to reason efficiently

Methodology:

1. Solve sequence of problems (via reasoning)

# Inference Control Meta-learning

Learn how to reason efficiently

Methodology:

1. Solve sequence of problems (via reasoning)
2. Store inference traces



Learn how to reason efficiently

Methodology:

1. Solve sequence of problems (via reasoning)
2. Store inference traces
3. Mine traces to discover patterns

# Inference Control Meta-learning

Learn how to reason efficiently

Methodology:

1. Solve sequence of problems (via reasoning)
2. Store inference traces
3. Mine traces to discover patterns
4. Build control rules

Implication <TV>

And

<inference-pattern>

<rule>

<good-inference>

# Inference Control Meta-learning

Learn how to reason efficiently

Methodology:

1. Solve sequence of problems (via reasoning)
2. Store inference traces
3. Mine traces to discover patterns
4. Build control rules

Implication <TV>

And

<inference-pattern>

<rule>

<good-inference>

5. Combine control rules to guide future reasoning

# Combine Control Rules


Implication <TV1>

And

<inference-pattern-1>

deduction-rule

<good-inference>



$$c = 1$$


Implication <TV2>

And


<inference-pattern-2>

deduction-rule

<good-inference>



=



# Combine Control Rules


Implication <TV1>

And

<inference-pattern-1>

deduction-rule

<good-inference>



$$c = 0.5$$


Implication <TV2>

And


<inference-pattern-2>

deduction-rule

<good-inference>



=



# Combine Control Rules


Implication <TV1>

And

<inference-pattern-1>

deduction-rule

<good-inference>



$$c = 0.1$$


Implication <TV2>

And


<inference-pattern-2>

deduction-rule

<good-inference>



=



## Contribution:

- Second Order Solomonoff Operator Induction
- Specialized for Beta Distribution
- Attempt to Deal with Partial Models

## Contribution:

- Second Order Solomonoff Operator Induction
- Specialized for Beta Distribution
- Attempt to Deal with Partial Models

## Future Work:

- Improve Likelihood of Unexplained Data
- More Experiments (Inference Control Meta-learning)



## Contribution:

- Second Order Solomonoff Operator Induction
- Specialized for Beta Distribution
- Attempt to Deal with Partial Models

## Future Work:

- Improve Likelihood of Unexplained Data
- More Experiments (Inference Control Meta-learning)

Thank you!