Breast Cancer

6

- A new system with:

“a remarkable ability to distinguish benign from malignant human breast histopathology”

- Sensitivity and specificity over 85%



PLOS | ONE

RESEARCH ARTICLE

Pigeons (*Columba livia*) as Trainable Observers of Pathology and Radiology Breast Cancer Images

Richard M. Levenson[1]*, Elizabeth A. Krupinski[2], Victor M. Navarro[2], Edward A. Wasserman[3]*

1 Department of Pathology and Laboratory Medicine, University of California Davis Medical Center, Sacramento, California, United States of America, **2** Department of Psychological and Brain Sciences, The University of Iowa, Iowa City, Iowa, United States of America, **3** Department of Radiology & Imaging Sciences, College of Medicine, Emory University, Atlanta, Georgia, United States of America

* leverson@ucdavis.edu (RML); ed-wasserman@uiowa.edu (EAW)

Abstract

Breast Cancer

- A new system with:

  "*a remarkable ability to distinguish benign from malignant human breast histopathology*"
- Sensitivity and specificity over 85%



PLOS ONE

RESEARCH ARTICLE

Pigeons (*Columba livia*) as Trainable Observers of Pathology and Radiology Breast Cancer Images

Richard M. Levenson[1]*, Elizabeth A. Krupinski[2], Victor M. Navarro[2], Edward A. Wasserman[3]*

1 Department of Pathology and Laboratory Medicine, University of California Davis Medical Center, Sacramento, California, United States of America, 2 Department of Psychological and Brain Sciences, The University of Iowa, Iowa City, Iowa, United States of America, 3 Department of Radiology & Imaging Sciences, College of Medicine, Emory University, Atlanta, Georgia, United States of America

* levenson@ucdavis.edu (RML); ed-wasserman@uiowa.edu (EAW)

Abstract

Facial Recognition