

Calculus Workshop

Integration Problem Set

Nathan Grimes

2023-09-19

Answer the following questions to the best of your ability. Feel free to work with anyone in the cohort, though I would encourage attempting on your own first to make sure you fully understand the concepts.

Logarithms

1) Solve for x

$$\ln(x + 2) = 12$$

$$e^{3x+1} = 16$$

$$6e^{4x} = 41e^{2x}$$


2) Find the derivative:

$$G(a) = \frac{2 \ln(4a - 2)}{e^{3a}}$$

Riemann Sum

- 1) Approximate the area under the curve $f(x) = 3x^2 - 6x + 10$ on the interval $[6, 12]$ with 6 uniform rectangles.

Does the position of the rectangle make a difference? Describe in words, show mathematically, or draw on the graph how evaluating the rectangle in different ways might lead to slightly different approximations.



Integrals

1)

The marginal benefit of abatement (e.g. reducing) for carbon is given by:

$$MB = 31 - 2Q$$

However there is also a marginal cost for carbon abatement given by:

$$MC = 6 + 3Q$$

Find the total net benefit of carbon abatement to society at equilibrium. (Hint: To get equilibrium and the bounds of the integral, first set marginal benefit equal to marginal cost and solve for Q^* . Then your integral bounds should be from 0 to Q^*)

2)

Take the Integrals

A) $y = \frac{3}{x^2}, y(0) = 5$

B) $g(t) = 3t^5 - 2t^3 + 16t - 7$

C) $\int_2^4 \frac{1}{2}x \quad (1)$

3)

A model for the rate of change in ozone concentrations over time between 1962-1984 is given by $\frac{dC}{dt} = 2t + 20$. Where C is the ozone concentration (ppm) and t is the elapsed time in years. Given that in 1964 the ozone concentration was 30 ppm, what was the ozone concentration in 1982?