

Canalization and plasticity in psychopathology

R.L. Carhart-Harris^{a,b,*}, S. Chandaria^{b,c,d}, D.E. Erritzoe^{b,e}, A. Gazzaley^a, M. Girn^f,
H. Kettner^{a,b}, P.A.M. Mediano^{g,h}, D.J. Nutt^b, F.E. Rosas^{b,c,i,j}, L. Roseman^{b,e},
C. Timmermann^{b,e}, B. Weiss^{b,e}, R.J. Zeifman^{b,k}, K.J. Friston^l

^a Psychedelics Division – Neuroscape, Department of Neurology, University of California, San Francisco, USA

^b Centre for Psychedelic Research, Imperial College London, UK

^c Centre for Eudaimonia and Human Flourishing, Linacre College, University of Oxford, UK

^d Institute of Philosophy, School of Advanced Study, University of London, UK

^e CNWL-Imperial Psychopharmacology and Psychedelic Research Clinic (CIPPRS), UK

^f Department of Neurology and Neurosurgery, Montreal Neurological Institute, McGill University, Montreal, QC, Canada

^g Department of Computing, Imperial College London, London, UK

^h Department of Psychology, University of Cambridge, UK

ⁱ Department of Informatics, University of Sussex, UK

^j Centre for Complexity Science, Imperial College London, UK

^k NYU Langone Center for Psychedelic Medicine, NYU Grossman School of Medicine, USA

^l Wellcome Centre for Human Neuroimaging, University College London, UK

ARTICLE INFO

Handling Editor: Bruno Frenguelli

ABSTRACT

This theoretical article revives a classical bridging construct, *canalization*, to describe a new model of a general factor of psychopathology. To achieve this, we have distinguished between two types of plasticity, an early one that we call ‘TEMP’ for ‘Temperature or Entropy Mediated Plasticity’, and another, we call ‘canalization’, which is close to Hebbian plasticity. These two forms of plasticity can be most easily distinguished by their relationship to ‘precision’ or inverse variance; TEMP relates to increased model variance or decreased precision, whereas the opposite is true for canalization. TEMP also subsumes increased learning rate, (Ising) temperature and entropy. Dictionary definitions of ‘plasticity’ describe it as the property of being easily shaped or molded; TEMP is the better match for this. Importantly, we propose that ‘pathological’ phenotypes develop via mechanisms of canalization or increased model precision, as a defensive response to adversity and associated distress or dysphoria. Our model states that canalization entrenches in psychopathology, narrowing the phenotypic state-space as the agent develops expertise in their pathology. We suggest that TEMP – combined with gently guiding psychological support – can counter canalization. We address questions of whether and when canalization is adaptive versus maladaptive, furnish our model with references to basic and human neuroscience, and offer concrete experiments and measures to test its main hypotheses and implications.

This article is part of the Special Issue on "National Institutes of Health Psilocybin Research Speaker Series".

1. Introduction

The construct of ‘canalization’ was introduced in biology in 1942 by British evolutionary scientist, Conrad Hal Waddington, in the context of phenotypic variation in development, where it refers to phenotypic stabilization (Waddington, 1959); see also (Peterson, 2011; Posteraro and Canalization, 2022). Canalization can be used in reference to the development of psychopathology, where it bears relevance to environmental or perturbational sensitivity (Hartman and Belsky, 2016; Belsky

and Pluess, 2009), being in this sense, the converse of ‘phenotypic plasticity’ (Dewitt et al., 1998; Belsky and Pluess, 2013), i.e., increased phenotypic plasticity relates to increased sensitivity, but canalization relates to the opposite of this.

A generalized, standard dictionary definition of ‘plasticity’ describes it as *the property of being easily shaped or molded*. One definition of neuroplasticity describes it as the brain’s ‘change capacity’: “[neuroplasticity is] the functional and structural reorganization capacity of the brain” (Assaf et al., 2019). If we are true to either, it is apparent that

* Corresponding author. Psychedelics Division – Neuroscape, Department of Neurology, University of California, San Francisco, USA.

E-mail address: robin.carhart-harris@ucsf.edu (R.L. Carhart-Harris).

<https://doi.org/10.1016/j.neuropharm.2022.109398>

Received 12 October 2022; Received in revised form 1 December 2022; Accepted 20 December 2022

Available online 27 December 2022

0028-3908/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

canalization is not a good fit. This is because canalization describes how features of the mind, brain, or behavior become less able to change in a non-specific way.

As readers will discover, this distinction is key to the present paper and requires a reconsideration of some popular assumptions regarding plasticity, including a challenge to a traditional, often implicit convention (e.g., in basic neuroscience) of equating plasticity with neuroplasticity – and Hebbian plasticity in particular. Hebbian plasticity can be defined as the principle that neurons or synapses that consistently functionally interact, change their properties in order to facilitate future interactions. To quote psychologist, Donald Hebb, “any two cells or systems of cells that are repeatedly active at the same time will tend to become ‘associated,’ so that activity in one facilitates activity in the other” (Hebb, 1949). This specifically describes the positive i.e., ‘potentiation’ aspect of what has long been called ‘Hebbian plasticity’, but we recognize that terms can also be inverted to describe ‘long-term depression,’ a related mechanism also serving associative plasticity.

1.1. Precision, plasticity, and learning

In a population, for a given ‘phenotype’ (i.e., a physical, behavioural, or cognitive feature) relevant to psychopathology, we can compute a mean value and a variance around the mean. Such variance can be called ‘inverse precision’. An increase in the volatility of a particular phenotype – such as an internalizing style of cognition and behavior (Kotov et al., 2022), over time, or in response to perturbation, would manifest as an increase in phenotypic plasticity and decrease in its ‘precision’. In other words, when precision is reduced, the individual is sensitized and able to explore broader regimes of phenotypic state space.

Couched in terms of learning, an increase in phenotypic volatility or plasticity manifests as an increase in *learning-rate* (Iigaya et al., 2018). Trait uniformity or consistency around a mean, despite environmental pressures, would reflect phenotypic canalization, i.e., expressed with high precision, the phenotype is more consistent and resilient. On the role of learning and canalization in evolutionary development - see (Hinton and Nowlan, 1987), and on canalization in habit formation, see (Dickinson, 1985). See also the notion of ‘learning traps’ and consider how it relates to canalization (Gopnik, 2020).

1.2. A one factor approach to psychopathology

“There may be a factor that accounts for meaningful variance across major forms of psychopathology and ... research could benefit from probing its origins.” (Caspi et al., 2014)

Caspi et al. *The p Factor: One General Psychopathology Factor in the Structure of Psychiatric Disorders?*

The aim of the present article is to attempt a new psychological and neurobiological model of a single factor or dimension of mental health or psychopathology (Caspi et al., 2014), inspired by the apparent dialectic between canalization and plasticity. In doing so, we are explicitly treating this ‘p factor’ as substantive (Levin-Aspenson et al., 2021).

Our general theory states that: cognitive and behavioral phenotypes that are regarded as psychopathological, are canalized features of mind, brain, or behavior that have come to dominate an individual’s psychological state space. We propose that the canalized features develop as responses to adversity, distress, and dysphoria, and endure despite, rather than because of, evidence. We also propose that their depth of expression or entrenchment determines, to a large extent, the severity of the psychopathology, including its degree of treatment-resistance and susceptibility to relapse.

1.3. Nosological challenges in psychiatry

For reasons of expediency, measurements in psychiatry have, in the main, been cross-sectional. That is, measurements are rarely

longitudinal and neither do they track an individual person’s dynamics across extended periods of time e.g., hour-to-hour or day-to-day over periods of months or years. Instead, most major outcome measures, such as patient or clinician rating scales of symptom severity, offer a retrospective and subjective ‘snapshot’ of the strength of a feature psychopathology in relation to a one-to-four-week period. Accordingly, the ensuing impression is temporally and contextually isolated. It captures the *product* of a process but not how it came to be. This may explain why psychiatry has struggled to explain etiology in terms of a patient’s lived trajectory, and as a result, find an effective nosology and accompanying treatments.

1.4. Psychopathologies are acquired but are they adaptive or maladaptive?

We argue that the brain, mind and behavior sometimes develop in putatively ‘maladaptive’ ways, but still to minimize a variational ‘free-energy’ (i.e., a quantitative, statistical score of surprise or uncertainty) – which we take to be a universal imperative of all living systems (Friston, 2013) (and see Box 1 regarding ‘free-energy’). Thus, psychopathologies remain cognitive and behavioral strategies for minimizing uncertainty; they are not exceptions to this ‘free-energy principle’ (Friston, 2010). In the sense that they match a universal imperative, we recognize that the styles of thinking and behaving associated with psychopathology may not be intrinsically abnormal or maladaptive.

Thus, we accept that certain cases of psychopathology could be regarded as adaptive (e.g., from the perspective of the individual), given a fuller understanding and appreciation of the adverse contextual conditions out of which the pathology has arisen (Conway et al., 2018; Cleck and Blendy, 2008; Dias-Ferreira et al., 2009; Felitti et al., 2019; Edmiston et al., 2011) – see also (Olthof et al., 2020) and (Sroufe, 1997). Moreover, apparently maladaptive phenotypes may initially have been adaptive for a given context - but critically, have become too entrenched, influential, and insensitive to new or changing conditions, i.e., they become, through canalization, insufficiently adaptive.

In a sense, individuals that develop a psychopathology at any given time are merely obeying a natural imperative to reduce *negatively valenced feeling states*. The colloquial terms ‘suffering’ and ‘pain’ are relevant here, as are the more academic constructs of psychological distress and dysphoria. We argue that, while unpleasant, these states are not intrinsically pathological; rather, they are fundamental and universal elements of the human experience.

“The First Noble Truth is suffering.” (Nhāt, 1999)

Thich Nhat Hanh. *The Heart of the Buddha’s Teaching*.

This universality of states of distress or dysphoria, does not, however, extend to psychopathological *phenotypes*; we argue that these are acquired. Thus, according to our model, pathological phenotypes come secondary to basic feeling states. More specifically, we propose that they are defensive reactions to unpleasant states – featuring e.g., distress or dysphoria. What classifies a psychological or neurobiological phenotype as pathological, is its degree of entrenchment, influence, and resistance to revision e.g., in response to new contexts or evidence.

1.5. Canalization and Hebbian plasticity

Challenging an absolutism regarding the value of Hebbian plasticity for healthy brain development, we propose that pathological cognitive and behavioral strategies for minimizing free-energy can be the product of classical Hebbian learning process that have gone awry. See (Song et al., 2000) and (Sumner et al., 2020a) for a modern account of Hebbian neuroplasticity. Developmental learning - like inter-generational learning (see (Boyd et al., 2011) and (Gopnik, 2020)) - can be generally regarded as a slow process involving the adaptation of the parameters and structure of a person’s (generative) world models underwriting

Box 1**Complexity science, free-energy, psychopathology, and psychedelics**

This article has taken inspiration from complexity science, dynamical systems theory, and the free-energy principle, but it is not its remit to explain these here. Rather, we refer the interested reader to previous explanations of these frameworks and how they have been applied to psychopathology (Olthof et al., 2020) and psychedelic action (Hipólito et al., 2022; Carhart-Harris and Friston, 2019).

The notion of canalization began with the philosophers Henri-Louis Bergson and Alfred North Whitehead (Peterson, 2011; Posteraro and Canalization, 2022) but was given greater visibility by the evolutionary biologist, Conrad Hal Waddington, who applied it to phenotypic development (Waddington, 1959). In what became known as the ‘Waddington Landscape’ (Fig. 1), Waddington invites us to imagine a ball rolling down a gradient featuring various valleys of differing depth. Within this metaphorical image, the steepness and depth of the valley or ‘canal’ walls, is meant to reflect the strength or depth of canalization, encoding phenotypic precision.

Translating this image into a more modern, energy landscape representation, the valleys or canals represent dynamical attractors, whose gravitational pull is also encoded by the steepness of their walls and overall depth. Within a free-energy scheme, the landscape represents a gradient descent, and the steepness and depth of the valleys relates to their precision-weighting, i.e., steep and deep valleys encode precise models. Translating to psychology, we can imagine a valley as representing a cognitive or behavioral phenotype, feature, or ‘style’, and its depth and steepness is intended to encode its strength of expression, robustness, influence, and resilience to influence and change.

Importantly, these state-space representations are over-simplifications. Moreover, although they depict dynamical phenomena, they remain static images. As is true of living phenomena more generally, the actual reality is dynamic, however, and individuals will be able to switch between free-energy minimizing strategies in a context-dependent fashion, as is true of the cycling nature of psychopathology. Dynamical methods will be required to better capture the emergence of different free-energy minimizing processes, in order to better understand their properties and signatures. Fortunately, interesting candidates are already being developed and explored (Eichenbaum et al., 2021), including in the context of psychedelic action (Lord et al, 2019).

perceptual inference and planning as inference, the essence of sentient behavior (Friston, 2022). However, we argue that such developmental mechanisms can go awry in psychopathology, as we shall explain.

Neurobiologically, experience-dependent learning requires synaptic plasticity mediated by experience or activity-dependent changes in neuronal connections strengths (Hebb, 1949). Under the free-energy principle, classical Hebbian plasticity can be cast as Bayesian belief-updating about the parameters of a lived world that correspond to its contingencies and its causal structure. The ensuing parameter updates depend on the precision (inverse variance or volatility) ascribed to sensory transients. The higher the learning rate, the greater potential for the modulation of precision. In a complex, unpredictable world, it may serve an organism’s adaptability if learning rate is sufficiently high such that ‘mind, brain and behavior’ can be appropriately plastic and adaptable (Potts et al., 2020). Indeed, it has been argued before that adaptability is our specie’s defining trait – ascribable to our disproportionately expanded transmodal cortex (Fjell et al., 2015; Changeux et al., 2021; Sneve et al., 2019; Buckner and Krienen, 2013; Luppi et al., 2022).

Cast in Bayesian terms, the pathology we have chosen to highlight in this paper pertains to when the precision (or confidence) of prior beliefs (a prediction or model) becomes inappropriately high, leading to a failure of adaptability and the perpetuation of cognitive or behavioural entrenchment. From a purely theoretical and technical perspective, inference and learning can be thought of as a gradient descent (a mathematical optimization algorithm for finding a local minimum i.e., the nearest lowest value, of a differential function) on a variational free-energy landscape. This landscape representation implies that, for every belief state, there is an accompanying free-energy that scores the confidence or certainty of that belief state. As new experiential evidence is garnered, inference and learning typically move belief states in a direction that reduces free-energy (or uncertainty). This developmental direction is encoded by experience-dependent plasticity and learning – and in most cases, it underlies healthy – or even ‘wise’ development (Moran et al., 2014). However, in certain cases and contexts, the process can ‘overshoot’ – creating extreme phenotypes that are (too) resistant to change.

Perhaps the most important component of our model is ‘precision’, which corresponds to the curvature of the free-energy landscape. This is an appealing conceptualization that speaks directly to canalization. In this view, canalization is like a river, carving its own riverbed, where the

depth of the eroded valley corresponds to precision and curvature. This means that to be over-canalized in one’s sentient behavior is to hold overly-precise beliefs in a specific state-space.

As stated above, canalization is conceptually close to Hebbian learning and plasticity; we appreciate it may be unusual for some to see Hebbian mechanisms equated with pathology, but this is not the first time it has been done (Torregrossa et al., 2011; Ungless et al., 2001; Chistiakova et al., 2014; Luscher and Malenka, 2011; Luscher, 2013; Pascoli et al., 2018). Like normal Hebbian learning, canalization can be regarded as an experience-dependent process that reinforces specific associations. However, our model states that such classical learning mechanisms are recruited in a problematic way in psychopathology. We propose that while the initial learning could be seen as an adaptive response to adversity, distress or dysphoria, the product is maladaptive if it becomes too dominating of a person’s future cognitive and behavioural state-space.

1.6. Non-Hebbian (neuro)plasticity

Basic neuroscience has identified and defined types of neuroplasticity other than classical Hebbian-plasticity; examples include: synaptic scaling and homeostatic plasticity (Turrigiano and Nelson, 2004), metaplasticity (Jedlicka et al., 2022; Abraham, 2008), heterosynaptic plasticity (Chistiakova et al., 2014), and critical-period plasticity (Lepow et al., 2021). All of the above could, and have been, related to the therapeutic action of novel treatments for psychopathology e.g., (Lepow et al., 2021; Kavalali and Monteggia, 2022; Fattore et al., 2018; Nardou et al., 2019), and they are relevant to the mechanisms that we are concerned with here.

However, with the possible exception of heterosynaptic (Olson, 2018) and critical period-like plasticity, mere increases in these types of plasticity may not be sufficient for a therapeutic action and outcome. For example, certain pro-plasticity drugs have been shown to augment stimulus-induced Hebbian plasticity (Sumner et al., 2020b) – which is an example of a metaplastic action i.e., increasing the ease or rate of Hebbian plasticity. Synaptic scaling is generally thought to preserve synapse-specific weighting (Kavalali and Monteggia, 2022) – and thus, the products of Hebbian learning. Reactivated critical period plasticity (or increased learning rate) combined with adverse or trauma-related conditions, could drive aberrant learning mechanisms enroute to

iatrogenic outcomes (Brouwer and Carhart-Harris, 2021). Indeed, given that acute stress has been found to increase neuroplasticity markers in the cortex and aid learning (Brivio et al., 2020) (see also (McEwen, 2019) and (Epel, 2020)), one can conceive how an initial increase in a nonspecific type of plasticity (such as heterosynaptic or critical period plasticity) could be hijacked by Hebbian mechanisms to canalize cognition or behavior in a particular way. See (Brouwer and Carhart-Harris, 2021; Timmermann et al., 2022) for a relevant discussion.

1.7. TEMP and neural annealing

At this point, we are compelled to add an essential qualifier to a potentially fallacious assumption that increases in (an undefined) plasticity are intrinsically health-promoting, salutogenic or therapeutic (see also (Dewitt et al., 1998) and (Brouwer and Carhart-Harris, 2021)). The computational construct we wish to invoke here is that of ‘simulated annealing’ (Kirkpatrick et al., 1983), see also (Gómez-Emilsson, 2021). Our model states that an effective intervention for psychopathology should, through an acute action analogous to an increase in system temperature or entropy – as per the Ising model (Suzuki et al., 2007; Ruffini et al., 2022) (see also (Singleton et al., 2022a)), trigger a downstream sub or post-acute effect that is analogous to a rebalancing or recalibration of synaptic weights, i.e., a counteraction to canalization.

To aid the construction of this model and its subsequent testing, we have felt it necessary to introduce a new construct that we call ‘Temperature of Entropy Mediated Plasticity’ or ‘TEMP’. To be more concrete, we propose that increases in the complexity or entropy of on-going neural ensemble activity indexes a key early phase of TEMP (Schartner et al., 2017; Carhart-Harris, 2018; Mediano et al., 2022) (see also (Ruffini et al., 2022)). Moreover, we have also created a novel measure to index the psychological counterparts of TEMP – i.e., in this case, decreased confidence in prior assumptions or beliefs (Zeifman et al., 2022). Indeed, this measure could be used to index the degree of precision (confidence) of strongly held beliefs relevant to a person’s particular pathological presentation. Critically, this measure will also allow us to quantify intervention-induced changes in belief precision or




Fig. 1. The ‘Waddington Landscape’ – first introduced in 1957 (Waddington, 2014). In our scheme, the depth and steepness of valleys encodes the precision of beliefs. The implied gradient descent of the ball symbolizes the implied trajectory toward canalization. This image can guide our conceptualizing of development on multiple scales and in multiple domains, speaking to the value of canalization as a powerful bridging construct e.g., for translating between psychology and biology. In a similar way, computational (Corlett and Fletcher, 2014) and complexity science approaches (Hipólito et al., 2022) offer appealing methods for translating between disciplines within psychiatry, including psychology and biology.




Fig. 2. A simple neuroimaging protocol for assessing the hypothesized action of interventions capable of triggering measurable plasticity processes. EEG is a powerful modality due to its affordability, tolerability, temporal resolution, usability and translatability across populations and species. See (Schartner et al., 2017; Zeifman et al., 2022) and (Mediano et al., 2022) for some strong candidate markers of early, upstream plastic effects – which we refer to as ‘TEMP’. EEG could be replaced by other indices of population level activity such as local-field potentials, in non-human animal experiments. MRI approaches are well-suited to pre versus post intervention scanning, where they can be used to detect both functional and anatomical brain changes e.g., (Lyons et al., 2022).

confidence. The schematic shown in Fig. 2 offers a general (neuroimaging-centric) experimental approach for assessing the effect and consequences of an increase in TEMP.

We are intrigued to speculate that heterosynaptic plasticity (Chistiakova et al., 2014) – seemingly the type of plasticity induced by the synaptogenic effect of psychedelics (Ly et al., 2018; Shao et al., 2021) – occurs downstream of an initial increase in the entropy of on-going (e.g., cortical) neuronal ensembles. Indeed, if this link was to be empirically established, it would contribute to validating the construct of ‘Temperature or Entropy Mediated Plasticity’ or ‘TEMP’.

Returning to the model of annealing, transient increase in TEMP can be conceived of as enabling neuronal representations of belief states to escape entrenched minima or ‘canals’ that constitute the free-energy landscape such that they can explore other free-energy minima or phenotypic possibilities. Note, precision is often associated with an ‘inverse temperature’ parameter in this setting, see Box 1 – and (Schneider and Sagan, 2005). Importantly, in the annealing model, increased temperature will have an acute and later phase carryover effect, equivalent to rebalancing the landscape by reducing its average skew or curvature, i.e., its precision. In this way, increased temperature would alter not just the energy or exploration rate within a state space, but also the shape of the space itself, i.e., after the system has ‘cooled’. Indeed, this two-phase process of heating and cooling to change a system’s structure is the essence of annealing.

Thus, according to our model, a transient increase in TEMP within a system should help rebalance its global state-space (Singleton et al., 2022b; Daws et al., 2022). Here, we invoke the analogies of ‘wiping the slate clean’, ‘starting afresh’ or ‘rebirth’. The idea is that by ‘heating’ the system, its corresponding free-energy landscape, underwriting neuronal dynamics and accompanying belief precision, will be effectively flattened (e.g., see (Singleton et al., 2022b; Girm et al., 2022)). Moreover, if we translate this to psychopathology, the existence of a skew, deformation, or bias in the system state-space – i.e., reflecting excessive precision on a specific domain within it, will imply that it is sensitive to the flattening effect of an increase in TEMP (e.g., see (Zeifman et al., 2022)).

This process may depend on how the acute entropic action relates to increased post-synaptic gain in deep-layer pyramidal neurons translating into an enhanced sensitivity to bottom-up signaling or prediction error (Carhart-Harris and Friston, 2019) – i.e., the essence of belief-updating and a major component of psychological ‘insight’ (Peill

et al., 2022) and gains in learning (Friston, 2010).

Consistent with conventional wisdom for psychedelic psychotherapy (Richards, 2015), allowing such a process to play out in a supportive (e.g., ‘gently guiding’ (Timmermann et al., 2022; Murphy et al., 2021)) environment, both acutely and post acutely, may be most conducive to positive therapeutic outcomes (Murphy et al., 2021; Carhart-Harris et al., 2018a; Melanson et al., 2022). Post-acutely, Socratic questioning (Braun et al., 2015) and inquiry (Hook et al., 2021) may aid patient realizations by challenging canalized beliefs.

1.8. Is canalization intrinsically ‘bad’, and plasticity, intrinsically ‘good’?

The answer here is, of course, “no”. In the same way that Hebbian plasticity is essential for healthy learning and development, so is canalization and consolidation of “well-trodden paths” – see also (Moran et al., 2014). Thus, we must address *why* and *when* canalization can become maladaptive. To provide a meaningful answer to this, information is needed on: 1) the nature of the phenotype that is canalized; 2) the extent to which it is canalized; 3) the context in which processes of canalization first occurred; and 4) whether that current context has changed.

1.9. Some specific examples of pathological canalization

Let us consider the example of negatively biased cognition, a common aspect of the ‘internalizing’ phenotypic style (Kotov et al., 2022), and a hallmark of major depressive disorder (Beck, 1971) – one of the most prevalent psychiatric disorders (Sansone and Sansone, 2010). Here, our formulation states that negatively biased cognition is a canalized phenotype, i.e., it is a cognitive style that a person can develop after a psychological ‘injury’ (or non-distinct injuries), and that can reinforce over time, with (implicit) practice, such that it is progressively easier to enter.

Figuratively speaking, for the ‘internalizer’, belief-updating has ‘fallen into a rut’ (Holtzheimer and Mayberg, 2011); this person’s thinking and/or behavior has become entrenched in overly precise beliefs; they ‘go there’, likely because, given their experiences and circumstances (Maté, 2010), it once made sense – or still makes sense (at least implicitly) – for them to do so. However, if deeply ingrained or allowed to reinforce over time, this canalized style will dominate the individual’s phenotypic state-space and be difficult to escape or undo – and when it is left, it will be easy to return to (Holtzheimer and Mayberg, 2011). Indeed, technically, the regimes of belief space that become canalized constitute an ‘attracting’ set, i.e., they attract or ‘pull-in’ (confirming) evidence.

Sufficient contextual information would be required to understand whether and why canalization of a given phenotype once provided, or continues to provide, an adaptive function – even if only implicitly (e.g., see (Fisher et al., 2017)). The prevalence of internalizing traits and their apparently high loading onto ‘p’ (Kim and Eaton, 2015) could be interpreted as implying that ‘dark rooming’ (Friston et al., 2012a) is a particularly common strategy for minimizing free-energy in response to adversity and distress. In a sense, the individual may (implicitly) ‘dark room’ because it is intolerable ‘out there’, and an alternative to harmful (e.g., aggressive) acts, such as suicide.

This internalizing type of canalization is particularly pernicious because reshaping the phenotypic landscape requires an updating of prior beliefs with new sensory evidence. However, avoidant phenotypic styles will preclude exposure to new experiences that could guide the process of narrative revision. Such a picture is familiar one for cognitive behavioral therapy that aims (e.g., through cognitive restructuring and exposures) to challenge prior beliefs by providing contrary evidence to certain prior convictions: e.g., “if I do this or that — or forget to do this or that — some awful thing will happen”.

1.10. Generalizing the model to other examples of pathological canalization

One could easily translate the above formulation into another major spectrum of psychopathology – the externalizing dimension (Kotov et al., 2022). Various styles of cognitive and behavioral addictions, such as cravings and compulsions, can be conceived of as involving Hebbian learning ‘gone wrong’ (Ungless et al., 2001; Ostroumov and Dani, 2018). In a purely psychological sense, the external objects of ‘relief’ or ‘escape’ in addiction, come to entice and enslave thought and behavior (Maté, 2010).

“When I was using, I had tunnel vision.” (Maté, 2010)

Gabor Maté. *In the Realm of Hungry Ghosts*.

As with ‘depressive dark rooming’, the ‘addictive strategy’ could be seen as adaptive, if it works to effectively minimize free-energy in a psychosocial milieu that engenders intolerable suffering (Maté, 2010). This free-energy minimization might be most effective within a brief time horizon, e.g., by inducing a transient emotional escape (Fernandez-Rodriguez et al., 2018), but such relief may still represent a logical option for a person not otherwise able to tolerate their experiences (Maté, 2010). Again, the pathological style could be seen as adaptive for the individual – at least in the short-term, yet dissonant with, and potentially threatening to, community norms and values. In other words, the psychopathology may be Bayes-optimal for a person with a particular history; however, their emergent prior beliefs may nevertheless dissonate with new contexts or diverge from community norms and values in a way that is harmful for the individual or the community (Wakefield, 1992) – yet they ‘cannot help it’.

Several other cognitive and behavioral strategies for minimizing free-energy across psychiatric diagnostic categories can be identified (Ainley et al., 2012, 2016; Limanowski et al., 2017; Parr et al., 2018; Limanowski, 2022; Fotopoulou and Tsakiris, 2017; Duquette, 2017). For example, in body-image and eating disorders, image becomes canalized in a distorted way (Riva and Gaudio, 2012) and there is excessive cognitive and behavioral control – typified by caloric control in anorexia. Obsessions and compulsions in obsessive compulsive disorder (OCD) are characteristically repetitive and canalizing. Context-specific triggers in anxiety and stress or trauma-related disorders, including post-traumatic disorder (PTSD), also speak to runaway Hebbian mechanisms (Chistiakova et al., 2014). Again, it’s easy to appreciate how certain cognitive and behavioural manifestations may seem appropriate for extreme environments such as military combat, but clash with the norms and values of civilian or post-military life.

Although rarer by prevalence (Sansone and Sansone, 2010), in addition to the internalizing and externalizing dimensions, the spectrum of ‘thought disorder’, which includes psychotic disorders, is not excepted from our formulation. Entrenched delusional complexes fit the principle of free-energy minimization (Lyndon and Corlett, 2020), again via a strategy that could be felt (by the individual) to be functional (if not essential) for them – but nevertheless remains intensely dissonant with, and potentially threatening to, community norms and values (Leader, 2011). A somatoform category, incorporating functional neurological syndromes such as fibromyalgia and chronic fatigue syndrome, as well as hypochondriasis, is also captured by our formulation. This category encompasses cases in which organic causes of physical symptoms are difficult to identify and isolate, yet come to consume thought and behavior, and can, like so many ‘disorders’, become inculcated into self-schemas (Edwards et al., 2012).

Some more challenging cases for our canalization model of psychopathology include symptoms of a disinhibition – such as in attention deficit hyperactivity disorder (ADHD) and psychotic disorders. Bipolar disorder is a particularly interesting case, as it straddles internalizing, thought disorder and externalizing dimensions (Kotov et al., 2022).

Returning to an earlier point regarding the cross-sectional nature of

psychiatric diagnosis, the symptoms associated with diagnostic categories may be more accurately regarded as dynamic, phenotypic products of (arguably easier to measure) state-specific phenomena, which clinicians usually do not have sufficient information about due to an absence of individual patient-level longitudinal information – but see (Fisher et al., 2017) and (Fortuna et al., 2022; Bos et al., 2022; Yerushalmi et al., 2021; Jones et al., 2021). A category like bipolar disorder features highly distinct states that are typically cycling, thus more state-sensitive measures are required to capture its mechanisms (Fortuna et al., 2022; Bos et al., 2022; Yerushalmi et al., 2021; Jones et al., 2021).

Regardless, we adopt the position that all of the relevant cognitive and behavioral presentation within a categorical palette or dimensional spectra of psychopathology are consistent with the free-energy minimizing imperative of all living systems (Ramstead et al., 2018). For example, in bipolar disorder, the agent oscillates between a depressive ‘dark room’, and a frenzied high-energy search; all done for same ‘purpose’ of minimizing distress or uncertainty. Indeed, notions of ‘manic defense’ (Baruch, 1997) can extend to other transient states of psychological instability – such as intense emotionality in certain personality disorders, where the individual seeks to suspend distress or uncertainty e.g., via projective or splitting defense mechanisms aimed at simplifying a psychosocial milieu that is experienced as overwhelming, ambiguous, or threatening – e.g., see (Herzog et al., 2022) for a relevant discussion.

It is tempting to speculate that high arousal ‘hot states’ interact with Hebbian mechanisms to canalize defensive responses, making them easier to find in the future; a process akin to kindling (Young and Dulcis, 2015). Such a process would be consistent with a metaplastic development of pathological habits, whereby endogenous processes involving elevated stress (Brouwer and Carhart-Harris, 2021; Brivio et al., 2020) and increase system TEMP are hijacked by Hebbian mechanisms. In such cases as acute psychoses in psychotic disorders, the ‘hot states’ may paradoxically represent endogenous efforts to escape local minima ‘carved’ by canalization – as has been discussed here (Brouwer and Carhart-Harris, 2021); however, because extreme stress and distress typically accompany extreme adversity, it is more likely that the surrounding context is traumatogenic than therapeutic.

Thus, to summarize, according to our model, symptoms of psychopathology are canalized phenotypes, i.e., over-potentialized styles of thinking or behaving that develop as defensive responses to unpleasant states featuring distress or dysphoria. They come to dominate an individual’s global psychological state space; they are compelling, difficult to give-up, easy to fall into and resistant to revision.

1.11. When is canalization not pathological?

What of the devoted expert, religious person, political activist, obsessional academic, artist or sportsperson? Do these individuals not exhibit canalized thought and/or behavior? Indeed, it would not be difficult to pathologize some cases, but in-and-of themselves, they do not necessarily meet criteria for psychopathology.

According to our model, an accurate determination of pathology requires information on the context in which the phenotype first arose, the contemporaneous context in which it exists now, and whether it could be deemed to be harmful to the individual or the larger community in which the individual resides. For example, within a micro-community of unhoused individuals with drug addictions, the individual’s addictive behaviors will not be abnormal, yet they will nevertheless diverge from the norms and values of the wider community and be regarded as generally harmful, both to the individual and the broader community (Wakefield, 1992).

In the case of a resolute expert, their canalized skillset may be valued – if not admired – by the wider culture. Ironically, their expertise, and perceived need to maintain it, may nevertheless draw them away from other precious micro-communities, such as their family. A ‘monomaniacal’ dedication could easily spillover into pathology if the

individual cannot resist its pull in order to be available for ‘other things’. Rigid perfectionism, ruminative deliberation and workaholism are all relevant in this regard, which, while naturally dimensional, become pathological by way of degree and context appropriateness – as well as via the potential recruitment of ‘comorbid’ ancillary symptoms that are more classically recognized as pathological.

Thus, the question of whether or not a canalized phenotype is pathological, relates to the degree to which it endures or enslaves the individual, constricting and dominating their global psychological state space, while clashing with new situations and evidence as well as wider societal norms and values. For example, a devoted religious person or meditator may retain hobbies, interests and inter-personal relationships that are extraneous to their life of practice, whereas an addicted person may become consumed by their object(s) of addiction, such that their capacity to be engaged and rewarded by anything else, is severely depleted.

1.12. Psychedelic therapy for psychopathology

“Factors that make treatments effective can give insight into the cause of a condition.” (Andrews and Thomson, 2009)

Andrews & Thomson. The bright side of being blue: depression as an adaptation for analyzing complex problems.

One approach for assessing the usefulness of our model is to ask: *how can we most effectively treat a presentation defined by problematic canalization?*

Here, we make special reference to psychedelic therapy. There is growing evidence from controlled (Carhart-Harris et al., 2021), naturalistic observational (Nygart et al., 2022) and population studies (Hendricks et al., 2015) of robust and diverse improvements in mental health outcomes after psychedelic use. By ‘psychedelics’ we are referring to direct serotonin 2A receptor (5-HT_{2A}) agonists such as psilocybin/psilocin and related drugs. If we focus on gold-standard criteria for quality evidence, i.e., from randomized controlled trials, there is consistent evidence of rapid and sustained improvements in depression, anxiety and addiction disorder symptoms, typically after just one or two administrations of a psychedelic alongside psychological support – e.g., (Carhart-Harris et al., 2021).

Buoyed by the promising nature of these findings and related industry investment, psychedelic therapy trials are now underway in obsessive compulsive disorder, various addiction disorders, eating and image disorders, functional neurological syndromes, various anxiety disorders and several other conditions. After promising phase 2B results of psilocybin therapy for treatment-resistant depression (Goodwin et al., 2022), phase-3 licensing trials are set to read out and challenge for regulatory approval in coming years. The development of related interventions – e.g., MDMA assisted psychotherapy (Mitchell et al., 2021), and ketamine (Alnefeesi et al., 2022), is even more advanced.

Does this purposeful activity reflect an optimism overshooting evidence? It seems appropriate to acknowledge that media hype surrounding psychedelic medicine has created high expectations (Yaden et al., 2022). Nevertheless, we propose that a compelling rationale underlies the psychedelic therapy paradigm – including the notion that the intervention can be transdiagnostically effective (Kocarova et al., 2021).

Here, as elsewhere (Carhart-Harris and Friston, 2019), we argue that the core therapeutic action of psychedelic therapy is the relaxation of inappropriately precise beliefs underpinning pathological phenotypes, thereby facilitating (re)learning mechanisms that can generate a healthy recalibration of an individual’s psychological state-space. However, we extend this model by further invoking the process of simulated annealing (Kirkpatrick et al., 1983), whereby a direct entropic action on spontaneous cortical activity is followed by downstream synaptic reweighting, effectively de-weighting relevant phenotypes – speculatively, by increasing heterosynaptic plasticity (Chistiakova et al., 2014). Some candidate neuroimaging metrics may also be sensitive to the

effects of this process (Ly et al., 2018; Shao et al., 2021; Moda-Sava et al., 2019; Jones et al., 2009).

1.13. Psychedelics and plasticity

There is converging evidence that serotonergic psychedelics, historically referred to as ‘classic’ psychedelics, elicit their characteristic behavioral effects via agonism of the 5-HT_{2A}R, potentially via biased signaling (Kim et al., 2020; Kaplan et al., 2022). It has been implied that the promotion of (downstream) anatomical markers of plasticity — such as the growth of new synaptic spines — may be necessary and sufficient for achieving therapeutic outcomes via psychedelic therapy (Vargas et al., 2021). Psychedelics, and 5-HT_{2A}R agonism more specifically, have been found to promote the expression of plasticity genes (Desouza et al., 2021), brain derived neurotrophic factor (BDNF) (Vaidya et al., 1997) as well as spine formation in vitro (Ly et al., 2018; Jones et al., 2009) and in vivo (Shao et al., 2021), with especially robust and reliable effects seen in the cortex (Vaidya et al., 1997; Mason et al., 2020) — see (Calder and Hasler, 2022) for a review.

When viewed through a lens of simulated annealing, other neuronal phenomena such as *increased*: asynchronous glutamate release (Aghajanian and Marek, 1999), expression of neural activity markers (Gresch et al., 2002), complexity of spontaneous oscillatory activity (Schartner et al., 2017), entropy of connectivity motifs (Tagliazucchi et al., 2014), high-frequency harmonics (Atasoy et al., 2017), near (Toker et al., 2022), or supercritical dynamics (Ruffini et al., 2022) sensitivity to perturbation (Jobst et al., 2021), and global integration (Tagliazucchi et al., 2016) — can all be linked to either an *early*, upstream temperature or entropy increase (Schartner et al., 2017) or a *later*, downstream synaptic reweighting effect — e.g., as implied by preclinical data (Shao et al., 2021; Daws et al., 2022; Moda-Sava et al., 2019; Raval et al., 2021).

Future research could examine the link between these distinct phases of a hypothesized annealing process. Relevantly, a recent multimodal healthy human imaging study found increased brain entropy under the effects of high-dose psilocybin and decreased prefrontal cortex-to-subcortex white matter diffusivity one-month later (Lyons et al., 2022). The increase in brain entropy was predictive of long-term improvements in psychological well-being.

The classic serotonergic psychedelic, LSD, has been shown to accelerate learning rate in rabbits via a 5-HT_{2A}R-dependent action (Harvey, 2003), and an increase in learning rate was recently observed in humans under LSD (Kanen et al., 2022). See also (Buchborn et al., 2014; De Gregorio et al., 2022). These findings are complemented by other human studies highlighting *acute increases* in: sensitivity to music (Kaelen et al., 2015) and hypnotic suggestion (Carhart-Harris et al., 2015) under serotonergic psychedelics; plus *sub-acute increases* in cognitive flexibility (Murphy-Beiner and Soar, 2020; Doss et al., 2021); and *long-term changes* in: personality structure (Weiss et al., 2021a, 2021b; Erritzoe et al., 2019; Carhart-Harris et al., 2016), political perspective (Nour et al., 2017; Lyons and Carhart-Harris, 2018), and philosophical (Timmermann et al., 2021) and religious belief (Griffiths et al., 2019); as well as *long-term improvements* in nature relatedness or connectedness (Nour et al., 2017; Lyons and Carhart-Harris, 2018; Forstmann and Sagioglou, 2017), psychological flexibility (Agin-Liebes et al., 2022; Davis et al., 2020; Close et al., 2020), and generic and specific mental health outcomes (Nygart et al., 2022; Mans et al., 2021; Carhart-Harris et al., 2018b).

In many cases, these psychological changes could be viewed as consistent with a rebalancing of a skewed psychological state space — i. e., a ‘resetting’ (Carhart-Harris et al., 2017) type effect. However, this rule is not absolute, and concerns have been raised over the possibility of exploiting early plasticity mechanisms to instill or reinforce certain beliefs (Brouwer and Carhart-Harris, 2021). The latter scenario would be an example of a drug-induced metaplastic action, where increased TEMP is hijacked to serve canalization rather than recalibration. We

believe that such deliberate shaping of phenotypic presentations, as opposed to a rebalancing of global dynamics, is more the exception than the rule, however, but nevertheless remains a risk with pro-plasticity interventions (Safron, 2020) — see also (Timmermann et al., 2022).

1.14. REBUS model

The Relaxed Beliefs Under pSychedelics (REBUS) model (Carhart-Harris and Friston, 2019) is inspired by the free-energy principle (Friston, 2013) — and the ‘the entropic brain’ hypothesis (Carhart-Harris, 2018; Carhart-Harris et al., 2014). The main tenet of REBUS is that psychedelics decrease the precision-weighting (i.e., inverse variance or ‘confidence’) of internal predictive models (i.e., ‘beliefs’ or ‘priors’) encoded in brain activity. More specifically, REBUS proposes that precision-weighting on priors is decreased *under* psychedelics; effectively flattening the free-energy landscape.

Indeed, REBUS maintains that relaxed model precision is the basic *acute* action of psychedelics, which, within the free-energy framework, is implied by their entropic action — and *vice versa*. Supportive evidence for REBUS was outlined in the foundational paper (Carhart-Harris and Friston, 2019) and more evidence has accrued since (Girn et al., 2022; Alamia et al., 2020); including a recent finding of decreased self-rated ‘confidence’ in negative self-beliefs after an inaugural high-dose psychedelic experience; an effect explained and predicted by increased entropy in EEG recordings of spontaneous scalp potentials (Zeifman et al., 2022) — see also Fig. 2.

The REBUS model relates well to the canalization model of psychopathology (or ‘CANAL’, for short), presented here. The former is specific to the action of psychedelics, whereas the latter has evolved from this specific application to appeal beyond it, i.e., to the notion of a single dimension of ‘psychopathology’ (Caspi et al., 2014) and how this might be understood and treated. The two models are consistent, in the sense that REBUS offers a mechanistic explanation for how psychedelic therapy can ‘treat’ canalized phenotypes, i.e., the model states that via a drug-induced entropic action, psychedelics plasticize reinforced patterns of cognition and behavior.

The combination of REBUS and CANAL can help to explain the putative transdiagnostic efficacy of psychedelic therapy (Kocarova et al., 2021), and it also inspires speculations about psychiatric symptoms that may be *especially* amenable to psychedelic therapy; i.e., those that most compellingly fit the model of pathology by canalization. The examples considered above include reinforced or entrenched cognitive and behavioral styles or biases. More specifically, ruminative, obsessive, controlling, compulsive cognitive and behavioral styles — such as are exhibited in anxiety and mood disorders, OCD, addiction disorders, and eating and image disorders, somatic syndromes, and presentations of delusion. As we search for an underlying mechanism for the principal component to psychopathology (Caspi et al., 2014), it should not be lost on the reader that the list just provided is relevant to the most prevalent psychiatric disorders.

Despite meshing well with psychedelic therapy, our model may, however, also invite reflections on cases where psychedelic-use would be risky. For example, we speculate that cases exhibiting high cognitive and behavioral *instability* — or implying special risk for psychological *decompensation* (Kluft, 1996) — may be contraindicated for psychedelic use, if not psychedelic therapy. The reason being that the entropic or TEMP increasing action of psychedelics may precipitate or augment a psychological instability that cannot be sufficiently well tolerated or recovered from, particularly when there are constraints on the quantity and quality of care available for guiding healthy re-learning. Granted, destabilization in the psychotherapeutic process (Olthof et al., 2020) has been found to predict sudden therapeutic gains (Shalom and Aderka, 2020) — but this principle may be precarious in severe psychopathologies, where there is a threat of decompensation.

A speculative list of especially challenging or risky presentations for psychedelic therapy includes: the at-risk mental state, early or acute

psychosis, schizophrenia, certain personality disorders and bipolar 1 disorder. Our unpublished research also suggests that young age may be a risk factor for certain potential side effects that have been linked to psychedelic-use (Zhou et al., 2022). Future research is required to test hypotheses that, e.g., personality and bipolar disorders are contraindications for psychedelics, or whether an adapted version of psychedelic therapy, e.g., using a more customized and prolonged treatment protocol — with more extensive psychological support and sophisticated dosing — could yield a positive cost-to-benefit ratio.

1.15. Testing CANAL

The CANAL model is speculative and must perform well under testing to establish its value. Despite notable initiatives to discover brain biomarkers of specific psychiatric symptoms (i.e., Research Domain Criteria or R-DoC), results have been largely disappointing (Weinberger et al., 2015). There could be several reasons for this, some of which may relate to cross-sectional sampling, questionable nosology and the “ubiquitous non-specificity” problem within psychiatry (Caspi and Moffitt, 2018).

Briefly, we can list some more specific potential explanations for failures in the discovery of biomarkers of psychopathology: high heterogeneity of symptoms within a given (e.g., sum scored) diagnostic category (Fried and Nesse, 2015; Fried et al., 2022), poor psychological measurement of the focal feature of psychopathology (Fried et al., 2022; Fried, 2017), high comorbidity between psychiatric disorders (Newman et al., 1998), unimpressive diagnostic reliability (Chmielewski et al., 2015), high inter-subject variability in brain function (Mueller et al., 2013), high diurnal intra-subject variability in psychological presentation (Shiffman et al., 2008), an inability to capture pathological states due to an absence of regular and longitudinal monitoring (Shiffman et al., 2008), as well as an absence of symptom provocation paradigms (D'Souza et al., 1999) or other emotionally engaging stimuli capable of ‘drawing out’ relevant phenotypes (Finn and Bandettini, 2021).

To propose one general psychopathological factor (‘p’) was to do something bold and disruptive (Caspi et al., 2014). The same is true of proposing a unified psychological and neurobiological explanatory model of ‘p’. Our model rests, to a large extent, on the legitimacy of ‘p’ — and thus, we refer the interested reader to the original text that introduced the notion (Caspi et al., 2014) — as well as other appraisals (Levin-Aspenson et al., 2021; Kelley et al., 2019; Lahey et al., 2012; van Bork et al., 2017) and supportive evidence (Selzam et al., 2018; Brainstorm et al., 2018; Goodkind et al., 2015; Sha et al., 2019; Elliott et al., 2018; Barch, 2017; Menon, 2011; Patalay et al., 2015). The CANAL model is intentionally parsimonious. There are natural strengths and weaknesses to parsimonious models (Ruffini and Lopez-Sola, 2022). Put simply, a useful model should explain a lot with few parameters (Ruffini and Lopez-Sola, 2022), and if it can do this, it will be a powerful data assimilation tool. It will not however, explain everything, but that is not its purpose.

To test CANAL, we must move out of the abstract and into the concrete. In what follows, we hypothesize some specific features of both the *process* and *products* of canalization, and we consider experimental manipulations that could up or down-regulate the relevant features. If entrenched canalization is a useful model of psychopathology, then intentionally up-regulating it could be regarded as unethical. Symptom provocation paradigms are relevant in this regard, but naturalistic sampling — e.g., in longitudinal studies — can track disease progression without intervening on it (Bos et al., 2022).

Many animal models of psychopathology do, however, aim to induce ‘pathological’ presentations. The most classic method for this is the chronic stress paradigm (Duman and Duman, 2015). Drug reinforcement learning is another classic method, in this case for modelling addiction processes (Robbins and Everitt, 2002). With drug-reinforcement, we recognize an important role for dopaminergic modulation in potentiating reward learning and thus, canalization

(Ungless et al., 2001; FitzGerald et al., 2015; Dobi et al., 2011), and the potential for serotonergic neuromodulation to counteract this (Walsh et al., 2018; Li et al., 2021; Boureau and Dayan, 2011) — see also (Bogenschutz et al., 2022; Johnson et al., 2014). We regard these pre-clinical paradigms as valid and useful models of the canalization processes that underlie the development of human psychopathologies. Aberrant learning mechanisms are readily apparent in these models (Ruan and Yao, 2017; Seligman et al., 1968). We are, however, also mindful of the limitations of rodent to human translational research in psychiatry, and have highlighted the usefulness of reverse translation and healthy human to clinical population translation (Carhart-Harris, 2022).

1.16. Synaptic atrophy and growth

Moving to more specific neuronal encoding, evidence is compelling that chronic stress causes dendritic atrophy in the prefrontal cortex (Duman and Duman, 2015) and hippocampus (Duman and Duman, 2015; Qiao et al., 2020) — including in a fashion that biases behavior towards habit formation (Dias-Ferreira et al., 2009) (see also (Piazza and Le Moal, 1996)). Stress is a common risk factor for psychopathology (Kessler, 1997). It is relevant that the severity of general psychopathology has been found to correlate with reduced cortical thickness in a large sample of middle-aged individuals (Romer et al., 2021). Moreover, generalized psychopathology was found to relate to accelerated brain ageing, as well as impaired cognitive and sensory functioning (Wertz et al., 2021), and in two independent studies, estimates of past adversity correlated with reduced gray matter volume in various brain regions (Edmiston et al., 2011; Ansell et al., 2012).

The classic serotonergic psychedelic, psilocybin, has been found to promote rapid and enduring increases in cortical dendritic spinogenesis *in vivo* (Shao et al., 2021), replicating prior findings of increased cortical synaptogenesis with a variety of 5-HT_{2A}R agonist psychedelics (Ly et al., 2018; Jones et al., 2009). This effect appears to be consistent with increased heterosynaptic plasticity (Chistiakova et al., 2014).

1.17. The complex case of brain network modularity

Brain network modularity is a measure of the extent to which the brain's inter-regional functional connectivity structure can be decomposed into distinct modules, wherein each module consists of regions that are more connected to each other than to other regions or modules. Recent human functional neuroimaging work on the therapeutic action of psychedelics found evidence of decreased brain network modularity after psilocybin therapy for depression across two independent samples. Decreases in brain modularity (synonymous with ‘increased global brain integrity’ or decreased segregation) correlated with improvements in symptom severity in both datasets (Daws et al., 2022).

Elevated brain network modularity or segregation has been seen in depression (Li et al., 2022; Ye et al., 2015), bipolar disorder (Chang et al., 2022) and cocaine addiction (Liang et al., 2015) — suggesting greater whole-brain functional segregation in these populations. In a similar fashion to high baseline symptom severity in antidepressant trials (Henkel et al., 2011), high baseline modularity has been found to predict response to treatment for schizophrenia (Doucet et al., 2020), OCD (Reggente et al., 2018) and cognitive training (Gallen and D'Esposito, 2019). Moreover, as with psilocybin therapy (Daws et al., 2022), brain modularity has been found to decrease after electroconvulsive therapy (ECT) for depression and schizophrenia (Xu et al., 2020; Huang et al., 2018). ECT is another rapid and robustly acting intervention, which, like 5-HT_{2A}R agonist psychedelics (Vaidya et al., 1997), has been found to increase cortical BDNF mRNA expression (Nibuya et al., 1995).

However, extant findings have not unequivocally linked high modularity to psychopathology. For example, see (Sinha et al., 2019) for a contradictory finding of increased brain network modularity after ECT,

as well as another study suggesting that age-related increases in modularity may be delayed in young people with psychopathology (Kaufmann et al., 2017). See also a particularly difficult to reconcile finding of low brain network modularity in a large sample of patients with depression, bipolar disorder or schizophrenia (Ma et al., 2020).

Neurodevelopmentally, brain modularity increases (i.e., modules become more differentiated from each other) throughout development into adulthood (Baum et al., 2017; Cui et al., 2020), but after reaching adulthood, brain modularity decreases (Chan et al., 2014) – perhaps suggesting a parallel relationship between developments in brain modularity and goal-directed cognition or cognitive aptitude. Indeed, in the case of early brain development or recovery from stroke, observed changes in global network modularity may be dominated by the (re) emergence of patterns of canalization e.g., related to the differentiation of high-level association networks, and healthy brain and cognitive functioning (Baum et al., 2017; Cui et al., 2020; Siegel et al., 2018). Delayed large-scale network differentiation may reflect immature brain network development (Kaufmann et al., 2017; Vanes and Dolan, 2021), but this does not imply that problematic canalization is absent in these cases or a special risk. Indeed, it is interesting to reflect whether immature brain development may relate to elevated environmental sensitivity or susceptibility (Greven et al., 2019) – a hypothetical primer for canalization.

It is important to highlight that the typical method for computing brain network modularity estimates the relative amount of functional segregation in the brain as a whole and computes modularity for a set of so-called ‘canonical resting-state networks’ or RSNs (Newman, 2006). As such, a similar modularity score may be underpinned by distinct patterns of inter-regional or inter-modular connectivity, which would, in turn, have different neuropsychological consequences. Moreover, it is important to state that there is no compelling reason to believe that the canonical RSNs (e.g., identified through independent components analyses on fMRI data) should be especially relevant to entrenched canalization in psychopathology. We speculate that stress, reward, or frontal-subcortical circuitry or networks (Pascoli et al., 2018; Pizzagalli and Roberts, 2022) may be more relevant to psychopathology – and yet are often not focused on in standard network analyses. See (Beam et al., 2021) for just one alternative method for defining brain networks.

Thus, a better justified identification and examination of candidate brain circuitry for assessing markers of canalization in psychopathology could help advance discovery, not just in non-human animal research – where progress has been made (Dias-Ferreira et al., 2009) – but critically, in human neuroimaging (though see (Pizzagalli and Roberts, 2022)). Once compelling candidate systems or circuitry have been identified, it will be easier to use human functional neuroimaging (and fMRI in particular) to test hypotheses inspired by CANAL, including its predictions regarding treatment targets and mechanisms.

1.18. Synaptic replenishment

The picture is arguably clearer and more compelling at the synaptic level. For example, the psychedelic-related dissociative anesthetic, ketamine, was recently found to recover dendritic spines atrophied by chronic stress in an animal model of psychopathology (Moda-Sava et al., 2019). Moreover, recovery of the atrophied spines was selective and related to behaviors reflective of improved functioning (Moda-Sava et al., 2019). See also (Holmes et al., 2022) for relevant findings in humans, using a synaptic density positron emission tomography radio-tracer measured prior to and 1-day after a single dose of ketamine.

Given these and related synaptic ‘replenishing’ findings with classic serotonergic psychedelics (Shao et al., 2021; De Gregorio et al., 2022) – combined with the compelling literature on synaptic atrophy with chronic stress (Duman and Duman, 2015), reduced gray matter in humans reporting histories of adversity (Edmiston et al., 2011; Ansell et al., 2012), and reduced cortical thickness in humans related to psychopathology (Romer et al., 2021) – we can see the emergence of a

model of the transdiagnostic therapeutic action of psychedelic therapy that relates to the counteraction of over-potentiated canalization. We speculate that the canalization could occur via (at least) three mechanisms: 1) an atrophy-related reduction in abundant synaptic connections, selectively sparing well-reinforced synapses and associated circuits, 2) top-down inhibition or neglect of specific circuitry (e.g., linked to traumatic memories), or 3) strong potentiation of selective synapses and circuits (Bliss, 1990). All scenarios should reduce freedom within the global system by biasing certain circuits or sub-states.

Finally, we speculate that psychedelic therapy can work to counteract over-potentiated canalization via drug-induced increases in TEMP, replenishing abundant connections i.e., heterosynaptic plasticity (Moda-Sava et al., 2019), to free dynamics within the global system. It matches clinical reports and observations (Carhart-Harris et al., 2018c; Watts et al., 2017) to assume that this effect will be best mediated via gently guiding psychotherapy (Timmermann et al., 2022).

Saying more about the brain mechanisms, we speculate that through a 5-HT_{2A}R agonism-induced modulation of synaptic efficacy and population-level activity – that can be recorded electrophysiologically across species (Schartner et al., 2017; Pascovich et al., 2022) – psychedelics trigger an early phase of plasticity – detectable via EEG as increased signal entropy (Schartner et al., 2017) – that translates downstream into a heterosynaptic plasticity and corresponding rebalancing of synaptic weights. This hypothesized sequence could be detectable at multiple scales and phases e.g., via human (Daws et al., 2022) and non-human animals (Shao et al., 2021; Moda-Sava et al., 2019) imaging techniques.

More work is required to test a hypothesized link between cortical atrophy and canalization (though see (Dias-Ferreira et al., 2009)) as well as the hypotheses that TEMP relates not just to psychological plasticity (Zeifman et al., 2022) but downstream anatomical neuroplasticity, with heterosynaptic plasticity being a compelling candidate in this regard (Shao et al., 2021).

1.19. Stress and canalization

Conceptually, we speculate that there are at least two ways in which stress may contribute to canalization in brain circuits and synaptic connections, one would entail chronic stress-induced cortical atrophy (Dias-Ferreira et al., 2009; Duman and Duman, 2015) and another would entail a Hebbian ‘hijacking’ of stress-induced increases in TEMP (Brivio et al., 2020), see also (Parr et al., 2020; Friston et al., 2012b, 2021) – and (Dias-Ferreira et al., 2009; Moda-Sava et al., 2019). Thus, canalization could narrow the psychological state-space either via removing or inhibiting association non-specific synapses (Moda-Sava et al., 2019) or strengthening association specific ones (Dias-Ferreira et al., 2009). Preclinical examples of Hebbian plasticity serving stimulant addiction can be seen here (Ungless et al., 2001; Dobi et al., 2011) and a separate example of stress-induced circuit reorganization biasing behavior toward habit – can be seen here (Dias-Ferreira et al., 2009).

1.20. Positive psychology and environmental enrichment

While we have focused on the pathology end of the mental health spectrum, a speculative marker of psychological health and/or positively valenced mood states may be self-organized criticality (SOC) (Friston et al., 2012b, 2021) – across scales – within and between brain and body (Carhart-Harris et al., 2014). Stated simply, we speculate that freer dynamics and information transfer across systems and scales, generally correspond to healthier and happier states of being (Carhart-Harris et al., 2014).

Separately, we are intrigued by evidence that environmental enrichment aids neuroplasticity (Baroncelli et al., 2010; Lee and Soya, 2017; Cancedda et al., 2004; Jha et al., 2016; Grech et al., 2018; Huang et al., 1999; Kazlauskas et al., 2011) and that individuals with above-average phenotypic plasticity or environmental-sensitivity

flourish in enriched environments – i.e., so-called ‘vantage sensitivity’ (Jolicœur-Martineau et al., 2020).

1.21. Limitations

Due to empirical gaps between the inter-species and inter-modality findings, we acknowledge that our model is extrapolative and speculative. However, we have been explicit about candidate biomarkers of key measures (Schartner et al., 2017; Mediano et al., 2022) and processes (e.g. (Zeifman et al., 2022)), and can envisage the types of experiments needed to test and potentially validate our hypotheses and constructs. We can be even more explicit: we predict that psychedelic-induced acute increases in spontaneous brain ‘temperature’ (Ruffini et al., 2022) or ‘entropy’ (Carhart-Harris, 2018; Carhart-Harris et al., 2014)—as scored through Lempel-Ziv complexity (Schartner et al., 2017) or a recently improved version of it - known as ‘Complexity of State-Space Entropy Rate’ or ‘CSER’ (Mediano et al., 2022) applied to electrophysiological data recorded from the neocortex of mammalian species - including humans — should predict increases in downstream markers of neuroplasticity, such as heterosynaptic plasticity in rodents (Shao et al., 2021), plus other anatomical and molecular markers (Raval et al., 2021) across species.

Separately, we hope to see others test whether decreased brain network modularity -as recently observed post psychedelic therapy for depression (Daws et al., 2022)—replicates across trials, including those in other psychiatric populations e.g., (Spriggs et al., 2021). We also recognize that behavioural measures are required to operationalize canalization, but refer the interested reader to one relevant measure in development (Zeifman et al., 2022). Cognitive flexibility and associative learning paradigms may be relevant and useful also, given their translatability across species (Harvey, 2003; Murphy-Beiner and Soar, 2020; Doss et al., 2021; Lu et al., 2021; Kanen et al., 2021).

1.22. Near neighbors

Finally, we believe that our model of psychopathology as canalization occurring in response to adversity-related distress or dysphoria, is novel, but we also recognize that we may have inadvertently overlooked prior models that share similarities with it. In this regard, we recognize the constructs of ‘capture’ (Kessler, 2016) and ‘branching’ (Sroufe, 1997), and see also (Klaus, 2002). We also recognize literature on genetic contributions to differential susceptibility or sensitivity (Jolicœur-Martineau et al., 2020; Belsky et al., 2019). Speculatively, there may be a polygenic contribution to inter-subject variability in TEMP and associated phenotypic plasticity (Greven et al., 2019; Pluess, 2015) – which could prime for canalization, in a context-dependent fashion (Brouwer and Carhart-Harris, 2021) -see also (Hopwood et al., 2022; Musci et al., 2019). Future research could assess individual differences in susceptibility to canalization, where individual differences in propensity for TEMP may be an early but insufficient component of this. In this regard, the question of *why* early increases in TEMP become transformed into canalization in some people (e.g., rather than system freedom or flexibility), seems important. The relevance of individual differences in tolerance of uncertainty (Einstein, 2014), psychological flexibility (Hayes, 2019) or experiential acceptance, may be relevant to examine in this regard. Thankfully, learnable behaviors for countering canalization have been proposed (Hayes, 2019).

2. Conclusion

The present work has introduced a new and intentionally parsimonious model of mental illness, based on the phenomenon of canalization. The model proposes that entrenched canalization, acquired in response to adversity, distress or dysphoria, is a principal component of psychopathology. The model takes inspiration from the (re)emergence of a putative transdiagnostic treatment for psychopathology; namely,

psychedelic therapy. We propose that psychedelics promote an early form of plasticity (TEMP), which, when combined with gently guiding psychological support, can serve to counter entrenched canalization.

“Return to simplicity ... [Do] not carve up” (Tzu, 1993)

Lao-Tzu. *Tao Te Ching*. Translated by Addis and Lombardo.

Funding

The time and effort of the authors on this article was not supported by grants or any commercial interests. Funding open-access publication of this article for the Neuropharmacology Special Issue was through a gift from the Steven & Alexandra Cohen Foundation to the Foundation for the National Institutes of Health.

Declaration of competing interest

RCH, DE, AG, DJN report being a scientific advisor to a number of new companies and not-for-profits that are seeking to develop psychedelic therapy and bring it to market. RCH is grateful to Meg Spriggs for comments on an earlier draft of this manuscript, Courtney Gallen for guidance on referencing brain network modularity findings, Rayyan Zafar for references on dopamine, stimulants and Hebbian mechanisms, Lindsay Cameron for references on environmental enrichment and neuroplasticity, and Max Wolff for the connection to the ideas of Klaus Grawe.

Data availability

No data was used for the research described in the article.

References

- Abraham, W.C., 2008. Metaplasticity: tuning synapses and networks for plasticity. *Nat. Rev. Neurosci.* 9 (5), 387.
- Aghajanian, G.K., Marek, G.J., 1999. Serotonin, via 5-HT_{2A} receptors, increases EPSCs in layer V pyramidal cells of prefrontal cortex by an asynchronous mode of glutamate release. *Brain Res.* 825 (1–2), 161–171.
- Agin-Liebes, G., et al., 2022. Prospective examination of the therapeutic role of psychological flexibility and cognitive reappraisal in the ceremonial use of ayahuasca. *J. Psychopharmacol.* 36 (3), 295–308.
- Ainley, V., et al., 2012. Looking into myself: changes in interoceptive sensitivity during mirror self-observation. *Psychophysiology* 49 (11), 1672–1676.
- Ainley, V., et al., 2016. ‘Bodily precision’: a predictive coding account of individual differences in interoceptive accuracy. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 371 (1708).
- Alamia, A., et al., 2020. DMT alters cortical travelling waves. *Elife* 9.
- Alnefeesi, Y., et al., 2022. Real-world effectiveness of ketamine in treatment-resistant depression: a systematic review & meta-analysis. *J. Psychiatr. Res.* 151, 693–709.
- Andrews, P.W., Thomson Jr., J.A., 2009. The bright side of being blue: depression as an adaptation for analyzing complex problems. *Psychol. Rev.* 116 (3), 620–654.
- Ansell, E.B., et al., 2012. Cumulative adversity and smaller gray matter volume in medial prefrontal, anterior cingulate, and insula regions. *Biol. Psychiatry* 72 (1), 57–64.
- Assaf, Y., Johansen-Berg, H., Thiebaut de Schotten, M., 2019. The role of diffusion MRI in neuroscience. *NMR Biomed.* 32 (4), e3762.
- Atasoy, S., et al., 2017. Connectome-harmonic decomposition of human brain activity reveals dynamical repertoire re-organization under LSD. *Sci. Rep.* 7 (1), 17661.
- Barch, D.M., 2017. The neural correlates of transdiagnostic dimensions of psychopathology. *Am. J. Psychiatry* 174 (7), 613–615.
- Baroncelli, L., et al., 2010. Nurturing brain plasticity: impact of environmental enrichment. *Cell Death Differ.* 17 (7), 1092–1103.
- Baruch, G., 1997. The manic defence in analysis: the creation of a false narrative. *Int. J. Psychoanal.* 78 (Pt 3), 549–559.
- Baum, G.L., et al., 2017. Modular segregation of structural brain networks supports the development of executive function in youth. *Curr. Biol.* 27 (11), 1561–1572 e8.
- Beam, E., et al., 2021. A data-driven framework for mapping domains of human neurobiology. *Nat. Neurosci.* 24 (12), 1733–1744.
- Beck, A.T., 1971. Cognition, affect, and psychopathology. *Arch. Gen. Psychiatry* 24 (6), 495–500.
- Belsky, J., Pluess, M., 2009. Beyond diathesis stress: differential susceptibility to environmental influences. *Psychol. Bull.* 135 (6), 885–908.
- Belsky, J., Pluess, M., 2013. Beyond risk, resilience, and dysregulation: phenotypic plasticity and human development. *Dev. Psychopathol.* 25 (4 Pt 2), 1243–1261.
- Belsky, J., et al., 2019. Polygenic differential susceptibility to prenatal adversity. *Dev. Psychopathol.* 31 (2), 439–441.

- Bliss, T.V., 1990. Long-term potentiation. *Science* 249 (4972), 973.
- Bogenschutz, M.P., et al., 2022. Percentage of heavy drinking days following psilocybin-assisted psychotherapy vs placebo in the treatment of adult patients with alcohol use disorder: a randomized clinical trial. *JAMA Psychiatr.*
- Bos, F.M., et al., 2022. Anticipating manic and depressive transitions in patients with bipolar disorder using early warning signals. *Int. J. Bipolar. Disord.* 10 (1), 12.
- Boureau, Y.L., Dayan, P., 2011. Opponency revisited: competition and cooperation between dopamine and serotonin. *Neuropsychopharmacology* 36 (1), 74–97.
- Boyd, R., Richerson, P.J., Henrich, J., 2011. The cultural niche: why social learning is essential for human adaptation. *Proc. Natl. Acad. Sci. U. S. A.* 108 (Suppl. 2), 10918–10925.
- Brainstorm, C., et al., 2018. Analysis of shared heritability in common disorders of the brain. *Science* 360 (6395).
- Braun, J.D., et al., 2015. Therapist use of Socratic questioning predicts session-to-session symptom change in cognitive therapy for depression. *Behav. Res. Ther.* 70, 32–37.
- Brivio, P., et al., 2020. Acute stress induces cognitive improvement in the novel object recognition task by transiently modulating bdnf in the prefrontal cortex of male rats. *Cell. Mol. Neurobiol.* 40 (6), 1037–1047.
- Brouwer, A., Carhart-Harris, R.L., 2021. Pivotal mental states. *J. Psychopharmacol.* 35 (4), 319–352.
- Buchhorn, T., et al., 2014. Repeated lysergic acid diethylamide in an animal model of depression: normalisation of learning behaviour and hippocampal serotonin 5-HT₂ signalling. *J. Psychopharmacol.* 28 (6), 545–552.
- Buckner, R.L., Krienen, F.M., 2013. The evolution of distributed association networks in the human brain. *Trends Cognit. Sci.* 17 (12), 648–665.
- Calder, A.E., Hasler, G., 2022. Towards an understanding of psychedelic-induced neuroplasticity. *Neuropsychopharmacology*.
- Cancedda, L., et al., 2004. Acceleration of visual system development by environmental enrichment. *J. Neurosci.* 24 (20), 4840–4848.
- Carhart-Harris, R.L., 2018. The entropic brain - revisited. *Neuropharmacology* 142, 167–178.
- Carhart-Harris, R., 2022. Translational challenges in psychedelic medicine. *N. Engl. J. Med.*
- Carhart-Harris, R.L., Friston, K.J., 2019. REBUS and the anarchic brain: Toward a unified Model of the brain Action of psychedelics. *Pharmacol. Rev.* 71 (3), 316–344.
- Carhart-Harris, R.L., et al., 2014. The entropic brain: a theory of conscious states informed by neuroimaging research with psychedelic drugs. *Front. Hum. Neurosci.* 8, 20.
- Carhart-Harris, R.L., et al., 2015. LSD enhances suggestibility in healthy volunteers. *Psychopharmacology (Berl)* 232 (4), 785–794.
- Carhart-Harris, R.L., et al., 2016. The paradoxical psychological effects of lysergic acid diethylamide (LSD). *Psychol. Med.* 46 (7), 1379–1390.
- Carhart-Harris, R.L., et al., 2017. Psilocybin for treatment-resistant depression: fMRI-measured brain mechanisms. *Sci. Rep.* 7 (1), 13187.
- Carhart-Harris, R.L., et al., 2018a. Psychedelics and the essential importance of context. *J. Psychopharmacol.* 32 (7), 725–731.
- Carhart-Harris, R.L., et al., 2018b. Psilocybin with psychological support for treatment-resistant depression: six-month follow-up. *Psychopharmacology (Berl)* 235 (2), 399–408.
- Carhart-Harris, R.L., et al., 2018c. Psychedelics and connectedness. *Psychopharmacology (Berl)* 235 (2), 547–550.
- Carhart-Harris, R., et al., 2021. Trial of psilocybin versus escitalopram for depression. *N. Engl. J. Med.* 384 (15), 1402–1411.
- Caspi, A., Moffitt, T.E., 2018. All for one and one for all: mental disorders in one dimension. *Am. J. Psychiatr.* 175 (9), 831–844.
- Caspi, A., et al., 2014. The p factor: one general psychopathology Factor in the Structure of psychiatric disorders? *Clin. Psychol. Sci.* 2 (2), 119–137.
- Chan, M.Y., et al., 2014. Decreased segregation of brain systems across the healthy adult lifespan. *Proc. Natl. Acad. Sci. U. S. A.* 111 (46), E4997–5006.
- Chang, Z., et al., 2022. Segregation, integration and balance in resting-state brain functional networks associated with bipolar disorder symptoms. *Hum. Brain Mapp.*
- Changeux, J.P., Goulas, A., Hilgetag, C.C., 2021. A connectomic Hypothesis for the Homination of the brain. *Cerebr. Cortex* 31 (5), 2425–2449.
- Chistiakova, M., et al., 2014. Heterosynaptic plasticity: multiple mechanisms and multiple roles. *Neuroscientist* 20 (5), 483–498.
- Chmielewski, M., et al., 2015. Method matters: understanding diagnostic reliability in DSM-IV and DSM-5. *J. Abnorm. Psychol.* 124 (3), 764–769.
- Cleck, J.N., Blendy, J.A., 2008. Making a bad thing worse: adverse effects of stress on drug addiction. *J. Clin. Invest.* 118 (2), 454–461.
- Close, J.B., et al., 2020. Psychedelics and psychological flexibility – results of a prospective web-survey using the Acceptance and Action Questionnaire II. *J. Contextual Behav. Sci.* 16, 37–44.
- Conway, C.C., et al., 2018. Transdiagnostic pathways from early social stress to psychopathology: a 20-year prospective study. *JCPP (J. Child Psychol. Psychiatry)* 59 (8), 855–862.
- Corlett, P.R., Fletcher, P.C., 2014. Computational psychiatry: a Rosetta Stone linking the brain to mental illness. *Lancet Psychiatr.* 1 (5), 399–402.
- Cui, Z., et al., 2020. Individual variation in functional topography of association networks in youth. *Neuron* 106 (2), 340–353 e8.
- D'Souza, D.C., et al., 1999. Symptom provocation studies in psychiatric disorders: scientific value, risks, and future. *Biol. Psychiatr.* 46 (8), 1060–1080.
- Davis, A.K., Barrett, F.S., Griffiths, R.R., 2020. Psychological flexibility mediates the relations between acute psychedelic effects and subjective decreases in depression and anxiety. *J. Contextual Behav. Sci.* 15, 39–45.
- Daws, R.E., et al., 2022. Increased global integration in the brain after psilocybin therapy for depression. *Nat. Med.* 28 (4), 844–851.
- De Gregorio, D., et al., 2022. Repeated lysergic acid diethylamide (LSD) reverses stress-induced anxiety-like behavior, cortical synaptogenesis deficits and serotonergic neurotransmission decline. *Neuropsychopharmacology* 47 (6), 1188–1198.
- Desouza, L.A., et al., 2021. The hallucinogenic Serotonin2A receptor agonist, 2,5-Dimethoxy-4-Iodoamphetamine, Promotes cAMP response element binding protein-dependent gene Expression of specific plasticity-associated Genes in the rodent neocortex. *Front. Mol. Neurosci.* 14, 790213.
- Dewitt, T.J., Sih, A., Wilson, D.S., 1998. Costs and limits of phenotypic plasticity. *Trends Ecol. Evol.* 13 (2), 77–81.
- Dias-Ferreira, E., et al., 2009. Chronic stress causes frontostriatal reorganization and affects decision-making. *Science* 325 (5940), 621–625.
- Dickinson, A., 1985. Actions and habits: the development of behavioural autonomy. *Philosophical Transactions of the Royal Society B* 308, 67–78.
- Dobi, A., et al., 2011. Cocaine-induced plasticity in the nucleus accumbens is cell specific and develops without prolonged withdrawal. *J. Neurosci.* 31 (5), 1895–1904.
- Doss, M.K., et al., 2021. Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder. *Transl. Psychiatry* 11 (1), 574.
- Doucet, G.E., et al., 2020. Baseline brain structural and functional predictors of clinical outcome in the early course of schizophrenia. *Mol. Psychiatr.* 25 (4), 863–872.
- Duman, C.H., Duman, R.S., 2015. Spine synapse remodeling in the pathophysiology and treatment of depression. *Neurosci. Lett.* 601, 20–29.
- Duquette, P., 2017. Increasing our insular world view: interoception and psychopathology for psychotherapists. *Front. Neurosci.* 11 (135), 135.
- Edmiston, E.E., et al., 2011. Corticostriatal-limbic gray matter morphology in adolescents with self-reported exposure to childhood maltreatment. *Arch. Pediatr. Adolesc. Med.* 165 (12), 1069–1077.
- Edwards, M.J., et al., 2012. A Bayesian account of 'hysteria'. *Brain* 135 (Pt 11), 3495–3512.
- Eichenbaum, A., et al., 2021. Differential contributions of static and time-varying functional connectivity to human behavior. *Netw. Neurosci.* 5 (1), 145–165.
- Einstein, D.A., 2014. Extension of the transdiagnostic model to focus on intolerance of uncertainty: a review of the literature and implications for treatment. *Clin Psychol (New York)* 21 (3), 280–300.
- Elliott, M.L., et al., 2018. A connectome-wide functional Signature of transdiagnostic Risk for mental illness. *Biol. Psychiatr.* 84 (6), 452–459.
- Epel, E.S., 2020. The geroscience agenda: Toxic stress, hormetic stress, and the rate of aging. *Ageing Res. Rev.* 63, 101167.
- Erritzoe, D., et al., 2019. Recreational use of psychedelics is associated with elevated personality trait openness: exploration of associations with brain serotonin markers. *J. Psychopharmacol.* 33 (9), 1068–1075.
- Fattore, L., et al., 2018. Psychedelics and reconsolidation of traumatic and appetitive maladaptive memories: focus on cannabinoids and ketamine. *Psychopharmacology (Berl)* 235 (2), 433–445.
- Felitti, V.J., et al., 2019. Reprint of: Relationship of childhood Abuse and household Dysfunction to Many of the leading Causes of Death in adults: the adverse childhood experiences (ACE) study. *Am. J. Prev. Med.* 56 (6), 774–786.
- Fernandez-Rodriguez, C., et al., 2018. Activation vs. *Experiential Avoidance as a Transdiagnostic Condition of Emotional Distress: An Empirical Study*. *Front Psychol* 9, 1618.
- Finn, E.S., Bandettini, P.A., 2021. Movie-watching outperforms rest for functional connectivity-based prediction of behavior. *Neuroimage* 235, 117963.
- Fisher, A.J., et al., 2017. Exploring the idiographic dynamics of mood and anxiety via network analysis. *J. Abnorm. Psychol.* 126 (8), 1044–1056.
- FitzGerald, T.H., Dolan, R.J., Friston, K., 2015. Dopamine, reward learning, and active inference. *Front. Comput. Neurosci.* 9, 136.
- Fjell, A.M., et al., 2015. High-expanding cortical regions in human development and evolution are related to higher intellectual abilities. *Cerebr. Cortex* 25 (1), 26–34.
- Forstmann, M., Sagioglou, C., 2017. Lifetime experience with (classic) psychedelics predicts pro-environmental behavior through an increase in nature relatedness. *J. Psychopharmacol.* 31 (8), 975–988.
- Fortuna, K.L., et al., 2022. Feasibility, acceptability, and potential Utility of peer-supported ecological momentary assessment among People with serious mental illness: a pilot study. *Psychiatr. Q.*
- Fotopoulou, A., Tsakiris, M., 2017. Mentalizing homeostasis: The social origins of interoceptive inference. *Neuro-psychoanalysis* 19 (1), 3–28.
- Fried, E.I., 2017. The 52 symptoms of major depression: Lack of content overlap among seven common depression scales. *J. Affect. Disord.* 208, 191–197.
- Fried, E.I., Nesse, R.M., 2015. Depression sum-scores don't add up: why analyzing specific depression symptoms is essential. *BMC Med.* 13, 72.
- Fried, E., Flake, J.K., Robinaugh, D.J., 2022. Revisiting the theoretical and methodological foundations of depression measurement. *Nat. Rev. Psychol.*
- Friston, K., 2010. The free-energy principle: a unified brain theory? *Nat. Rev. Neurosci.* 11 (2), 127–138.
- Friston, K., 2013. Active inference and free energy. *Behav. Brain Sci.* 36 (3), 212–213.
- Friston, K., 2022. Computational psychiatry: from synapses to sentence. *Mol. Psychiatr.*
- Friston, K., Thornton, C., Clark, A., 2012a. Free-energy minimization and the dark-room problem. *Front. Psychol.* 3, 130.
- Friston, K., Breakpear, M., Deco, G., 2012b. Perception and self-organized instability. *Front. Comput. Neurosci.* 6, 44.
- Friston, K.J., et al., 2021. Parcels and particles: Markov blankets in the brain. *Netw. Neurosci.* 5 (1), 211–251.
- Gallen, C.L., D'Esposito, M., 2019. Brain modularity: a Biomarker of intervention-related plasticity. *Trends Cognit. Sci.* 23 (4), 293–304.
- Girn, M., et al., 2022. Serotonergic psychedelic drugs LSD and psilocybin reduce the hierarchical differentiation of unimodal and transmodal cortex. *Neuroimage* 256, 119220.

- Gómez-Emilsson, A., 2021. Healing trauma with neural annealing. *QRI*.
- Goodkind, M., et al., 2015. Identification of a common neurobiological substrate for mental illness. *JAMA Psychiatr.* 72 (4), 305–315.
- Goodwin, G.M., et al., 2022. Single-dose Psilocybin for a treatment-resistant Episode of major depression. *N. Engl. J. Med.* 387 (18), 1637–1648.
- Gopnik, A., 2020. Childhood as a solution to explore-exploit tensions. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 375 (1803), 20190502.
- Grech, A.M., et al., 2018. Sex-dependent Effects of environmental Enrichment on spatial Memory and brain-derived neurotrophic factor (BDNF) Signaling in a developmental "Two-Hit" mouse model combining BDNF Haploinsufficiency and chronic glucocorticoid stimulation. *Front. Behav. Neurosci.* 12, 227.
- Gresch, P.J., Strickland, L.V., Sanders-Bush, E., 2002. Lysergic acid diethylamide-induced Fos expression in rat brain: role of serotonin-2A receptors. *Neuroscience* 114 (3), 707–713.
- Greven, C.U., et al., 2019. Sensory Processing Sensitivity in the context of Environmental Sensitivity: a critical review and development of research agenda. *Neurosci. Biobehav. Rev.* 98, 287–305.
- Griffiths, R.R., et al., 2019. Survey of subjective "God encounter experiences": comparisons among naturally occurring experiences and those occasioned by the classic psychedelics psilocybin, LSD, ayahuasca, or DMT. *PLoS One* 14 (4), e0214377.
- Hartman, S., Belsky, J., 2016. An evolutionary perspective on family studies: differential susceptibility to environmental influences. *Fam. Process* 55 (4), 700–712.
- Harvey, J.A., 2003. Role of the serotonin 5-HT_{2A} receptor in learning. *Learn. Mem.* 10 (5), 355–362.
- Hayes, S.C., 2019. *A Liberated Mind: How to Pivot toward what Matters*. Avery, New York xvi, 428 pages.
- Hebb, D.O., 1949. *The Organization of Behavior; a Neuropsychological Theory*. A Wiley Book in Clinical Psychology. Wiley, New York, xix, 335 pp.
- Hendricks, P.S., et al., 2015. Classic psychedelic use is associated with reduced psychological distress and suicidality in the United States adult population. *J. Psychopharmacol.* 29 (3), 280–288.
- Henkel, V., et al., 2011. Relationship between baseline severity of depression and antidepressant treatment outcome. *Pharmacopsychiatry* 44 (1), 27–32.
- Herzog, P., Kube, T., Fassbinder, E., 2022. How childhood maltreatment alters perception and cognition - the predictive processing account of borderline personality disorder. *Psychol. Med.* 1–18.
- Hinton, G.E., Nowlan, S.J., 1987. How learning can guide evolution. *Complex Syst.* 1. Hipólito, I., Mago, J., Rosas, F., Carhart-Harris, R.L., 2022. Pattern breaking: a complex systems approach to psychedelic medicine. *PsyArXiv*.
- Holmes, S.E., et al., 2022. Imaging the effect of ketamine on synaptic density (SV2A) in the living brain. *Mol. Psychiatr.* 27 (4), 2273–2281.
- Holtzheimer, P.E., Mayberg, H.S., 2011. Stuck in a rut: rethinking depression and its treatment. *Trends Neurosci.* 34 (1), 1–9.
- Hook, J.N., et al., 2021. Inquiry-based stress reduction: a systematic review of the empirical literature. *J. Clin. Psychol.* 77 (6), 1280–1295.
- Hopwood, C.J., Wright, A.G.C., Bleidorn, W., 2022. Person–environment transactions differentiate personality and psychopathology. *Nat. Rev. Psychol.* 1 (1), 55–63.
- Huang, Z.J., et al., 1999. BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual cortex. *Cell* 98 (6), 739–755.
- Huang, H., et al., 2018. Increased resting-state global functional connectivity density of default mode network in schizophrenia subjects treated with electroconvulsive therapy. *Schizophr. Res.* 197, 192–199.
- Ilgaya, K., et al., 2018. An effect of serotonergic stimulation on learning rates for rewards apparent after long intertrial intervals. *Nat. Commun.* 9 (1), 2477.
- Jedlicka, P., et al., 2022. Contributions by metaplasticity to solving the catastrophic forgetting problem. *Trends Neurosci.* 45 (9), 656–666.
- Jha, S., et al., 2016. Antidepressant and BDNF effects of enriched environment treatment across ages in mice lacking BDNF expression through promoter IV. *Transl. Psychiatry* 6 (9), e896.
- Jobst, B.M., et al., 2021. Increased sensitivity to strong perturbations in a whole-brain model of LSD. *Neuroimage* 230, 117809.
- Johnson, M.W., et al., 2014. Pilot study of the 5-HT_{2A} agonist psilocybin in the treatment of tobacco addiction. *J. Psychopharmacol.* 28 (11), 983–992.
- Jolicœur-Martineau, A., et al., 2020. Distinguishing differential susceptibility, diathesis-stress, and vantage sensitivity: Beyond the single gene and environment model. *Dev. Psychopathol.* 32 (1), 73–83.
- Jones, K.A., et al., 2009. Rapid modulation of spine morphology by the 5-HT_{2A} serotonin receptor through kalirin-7 signaling. *Proc. Natl. Acad. Sci. U. S. A.* 106 (46), 19575–19580.
- Jones, S.E., et al., 2021. Daily Ecological Momentary Assessments of happy and sad moods in people with schizophrenia and bipolar disorders: what do participants who are never sad think about their activities and abilities? *Schizophr Res Cogn* 26, 100202.
- Kaelin, M., et al., 2015. LSD enhances the emotional response to music. *Psychopharmacology (Berl)* 232 (19), 3607–3614.
- Kanen, J.W., et al., 2021. Effect of lysergic acid diethylamide (LSD) on reinforcement learning in humans. *bioRxiv*, 2020.12.04.412189.
- Kanen, J.W., et al., 2022. Effect of lysergic acid diethylamide (LSD) on reinforcement learning in humans. *Psychol. Med.* 1–12.
- Kaplan, A.L., et al., 2022. Bespoke library docking for 5-HT_{2A} receptor agonists with antidepressant activity. *Nature*.
- Kaufmann, T., et al., 2017. Delayed stabilization and individualization in connectome development are related to psychiatric disorders. *Nat. Neurosci.* 20 (4), 513–515.
- Kavalali, E.T., Monteggia, L.M., 2022. Rapid homeostatic plasticity and neuropsychiatric therapeutics. *Neuropsychopharmacology*.
- Kazlauskas, V., et al., 2011. Enriched environment effects on behavior, memory and BDNF in low and high exploratory mice. *Physiol. Behav.* 102 (5), 475–480.
- Kelley, T.M., et al., 2019. A new "inside-out" perspective on general factor p. *Eur. Psychiatr.* 61, 85–87.
- Kessler, R.C., 1997. The effects of stressful life events on depression. *Annu. Rev. Psychol.* 48, 191–214.
- Kessler, D.A., 2016. *Capture: Unraveling the Mystery of Mental Suffering*, first ed. Harper Wave, an imprint of HarperCollinsPublishers, New York, NY. ix, 406 pages.
- Kim, H., Eaton, N.R., 2015. The hierarchical structure of common mental disorders: connecting multiple levels of comorbidity, bifactor models, and predictive validity. *J. Abnorm. Psychol.* 124 (4), 1064–1078.
- Kim, K., et al., 2020. Structure of a hallucinogen-activated Gq-coupled 5-HT_{2A} serotonin receptor. *Cell* 182 (6), 1574–1588 e19.
- Kirkpatrick, S., Gelatt Jr., C.D., Vecchi, M.P., 1983. Optimization by simulated annealing. *Science* 220 (4598), 671–680.
- Klaus, G., 2002. *Psychological Therapy*. Hogrefe & Huber Publishers, Seattle, WA, p. 639 xvi.
- Kluft, R.P., 1996. Treating the traumatic memories of patients with dissociative identity disorder. *Am. J. Psychiatr.* 153 (7 Suppl. 1), 103–110.
- Kocanova, R., Horacek, J., Carhart-Harris, R., 2021. Does psychedelic therapy have a transdiagnostic action and prophylactic potential? *Front. Psychiatr.* 12, 661233.
- Kotov, R., et al., 2022. The hierarchical taxonomy of psychopathology (HiTOP) in psychiatric practice and research. *Psychol. Med.* 52 (9), 1666–1678.
- Lahey, B.B., et al., 2012. Is there a general factor of prevalent psychopathology during adulthood? *J. Abnorm. Psychol.* 121 (4), 971–977.
- Leader, D., 2011. *What Is Madness? Hamish Hamilton*, London, p. 359.
- Lee, M., Soya, H., 2017. Effects of acute voluntary loaded wheel running on BDNF expression in the rat hippocampus. *J. Exerc. Nutr. Biochem.* 21 (4), 52–57.
- Lepow, L., Morishita, H., Yehuda, R., 2021. Critical Period plasticity as a framework for psychedelic-assisted psychotherapy. *Front. Neurosci.* 15, 710004.
- Levin-Aspenson, H.F., et al., 2021. What is the general factor of psychopathology? Consistency of the p factor across samples. *Assessment* 28 (4), 1035–1049.
- Li, Y., et al., 2021. Synaptic mechanism underlying serotonin modulation of transition to cocaine addiction. *Science* 373 (6560), 1252–1256.
- Li, N., et al., 2022. Functional brain abnormalities in major depressive disorder using a multiscale community detection approach. *Neuroscience* 501, 1–10.
- Liang, X., et al., 2015. Interactions between the salience and default-mode networks are disrupted in cocaine addiction. *J. Neurosci.* 35 (21), 8081–8090.
- Limanowski, J., 2022. Precision control for a flexible body representation. *Neurosci. Biobehav. Rev.* 134, 104401.
- Limanowski, J., 2017. (Dis-)Attending to the body. In: Metzinger, T.K., Wiese, W. (Eds.), *Philosophy and Predictive Processing*. MIND Group, Frankfurt am Main.
- Lord, L.D., et al., 2019. Dynamical exploration of the repertoire of brain networks at rest is modulated by psilocybin. *Neuroimage* 199, 127–142.
- Lu, J., et al., 2021. An analog of psychedelics restores functional neural circuits disrupted by unpredictable stress. *Mol. Psychiatr.* 26 (11), 6237–6252.
- Luppi, A.I., Girm, M., Rosas, F.E., Timmermann, C., Roseman, L., Erritzoe, D., Nutt, D.J., Stamatakis, E.A., Spreng, R.N., Xing, L., Huttner, W.B., Carhart-Harris, R.L., 2022. A key role for serotonin 2A receptor signaling in the expansion and functioning of the human transmodal association cortex. *Neuron*.
- Luscher, C., 2013. Drug-evoked synaptic plasticity causing addictive behavior. *J. Neurosci.* 33 (45), 17641–17646.
- Luscher, C., Malenka, R.C., 2011. Drug-evoked synaptic plasticity in addiction: from molecular changes to circuit remodeling. *Neuron* 69 (4), 650–663.
- Ly, C., et al., 2018. Psychedelics promote structural and functional neural plasticity. *Cell Rep.* 23 (11), 3170–3182.
- Lyndon, S., Corlett, P.R., 2020. Hallucinations in posttraumatic stress disorder: insights from predictive coding. *J. Abnorm. Psychol.* 129 (6), 534–543.
- Lyons, T., Carhart-Harris, R.L., 2018. Increased nature relatedness and decreased authoritarian political views after psilocybin for treatment-resistant depression. *J. Psychopharmacol.* 32 (7), 811–819.
- Lyons, T., S.M., Kerkelä, L., Rosas, F.E., Timmermann, C., Roseman, L., Douglas, H., Girm, M., Mediano, P.A.M., Sharif, F., Espasiano, L., Erritzoe, D., Nutt, D.J., Carhart-Harris, R.L., 2022. Enduring Brain Changes after Psilocybin.
- Ma, Q., et al., 2020. Transdiagnostic dysfunctions in brain modules across patients with schizophrenia, bipolar disorder, and major depressive disorder: a connectome-based study. *Schizophr. Bull.* 46 (3), 699–712.
- Mans, K., et al., 2021. Sustained, multifaceted Improvements in mental well-being following psychedelic Experiences in a prospective opportunity sample. *Front. Psychiatr.* 12, 647909.
- Mason, N.L., et al., 2020. Me, myself, bye: regional alterations in glutamate and the experience of ego dissolution with psilocybin. *Neuropsychopharmacology* 45 (12), 2003–2011.
- Maté, G., 2010. *In the Realm of Hungry Ghosts: Close Encounters with Addiction*. Calif., Berkeley. North Atlantic Books. xviii, 495 pp.
- McEwen, B.S., 2019. The good side of "stress". *Stress* 22 (5), 524–525.
- Mediano, P.A.M., R. F., Luppi, A.I., Noreika, V., Carhart-Harris, R.L., Barnett, L., Bor, D., 2022. Spectrally and Temporally Resolved Estimation of Neural Signal Diversity.
- Melanson, B., et al., 2022. Conditioned anti-immobility by ketamine: a comparison to escitalopram and bupropion. *Exp. Clin. Psychopharmacol.*
- Menon, V., 2011. Large-scale brain networks and psychopathology: a unifying triple network model. *Trends Cognit. Sci.* 15 (10), 483–506.
- Mitchell, J.M., et al., 2021. MDMA-assisted therapy for severe PTSD: a randomized, double-blind, placebo-controlled phase 3 study. *Nat. Med.* 27 (6), 1025–1033.
- Moda-Sava, R.N., et al., 2019. Sustained rescue of prefrontal circuit dysfunction by antidepressant-induced spine formation. *Science* 364 (6436).

- Moran, R.J., et al., 2014. The brain ages optimally to model its environment: evidence from sensory learning over the adult lifespan. *PLoS Comput. Biol.* 10 (1), e1003422.
- Mueller, S., et al., 2013. Individual variability in functional connectivity architecture of the human brain. *Neuron* 77 (3), 586–595.
- Murphy, R., et al., 2021. Therapeutic alliance and rapport modulate responses to psilocybin assisted therapy for depression. *Front. Pharmacol.* 12, 788155.
- Murphy-Beiner, A., Soar, K., 2020. Ayahuasca's 'afterglow': improved mindfulness and cognitive flexibility in ayahuasca drinkers. *Psychopharmacology (Berl)* 237 (4), 1161–1169.
- Musci, R.J., Augustinavicius, J.L., Volk, H., 2019. Gene-environment Interactions in psychiatry: recent Evidence and clinical implications. *Curr. Psychiatr. Rep.* 21 (9), 81.
- Nardou, R., et al., 2019. Oxytocin-dependent reopening of a social reward learning critical period with MDMA. *Nature* 569 (7754), 116–120.
- Newman, M.E., 2006. Modularity and community structure in networks. *Proc. Natl. Acad. Sci. U. S. A.* 103 (23), 8577–8582.
- Newman, D.L., et al., 1998. Comorbid mental disorders: implications for treatment and sample selection. *J. Abnorm. Psychol.* 107 (2), 305–311.
- Nhat, H.N., 1999. The heart of the Buddha's teaching: transforming suffering into peace, joy & liberation: the four noble truths, the noble eightfold path, and other basic Buddhist teachings. In: 1st Broadway Books Trade Pbk. Broadway Books, New York, p. 292.
- Nibuya, M., Morinobu, S., Duman, R.S., 1995. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J. Neurosci.* 15 (11), 7539–7547.
- Nour, M.M., Evans, L., Carhart-Harris, R.L., 2017. Psychedelics, Personality and political perspectives. *J. Psychoact. Drugs* 49 (3), 182–191.
- Nygart, V.A., et al., 2022. Antidepressant effects of a psychedelic experience in a large prospective naturalistic sample. *J. Psychopharmacol.* 36 (8), 932–942.
- Olson, D.E., 2018. Psychoplastogens: a promising Class of plasticity-promoting neurotherapeutics. *J. Exp. Neurosci.* 12, 1179069518800508.
- Olthof, M., Hasselman, M., Maatman, F.O., Bosman, A.M.T., Lichtwarck-Aschoff, A., 2020. Complexity theory of psychopathology. *PsyArXiv*.
- Olthof, M., et al., 2020. Destabilization in self-ratings of the psychotherapeutic process is associated with better treatment outcome in patients with mood disorders. *Psychother. Res.* 30 (4), 520–531.
- Ostrovskiy, A., Dani, J.A., 2018. Convergent neuronal plasticity and metaplasticity mechanisms of stress, nicotine, and alcohol. *Annu. Rev. Pharmacol. Toxicol.* 58, 547–566.
- Parr, T., et al., 2018. Precision and false perceptual inference. *Front. Integr. Neurosci.* 12, 39.
- Parr, T., Sajid, N., Friston, K.J., 2020. Modules or mean-fields? *Entropy* 22 (5), 552.
- Pascoli, V., et al., 2018. Stochastic synaptic plasticity underlying compulsion in a model of addiction. *Nature* 564 (7736), 366–371.
- Pascovich, C., et al., 2022. Ketamine and sleep modulate neural complexity dynamics in cats. *Eur. J. Neurosci.* 55 (6), 1584–1600.
- Patalay, P., et al., 2015. A general psychopathology factor in early adolescence. *Br. J. Psychiatry* 207 (1), 15–22.
- Peill, J.M., et al., 2022. Validation of the Psychological Insight Scale: a new scale to assess psychological insight following a psychedelic experience. *J. Psychopharmacol.* 36 (1), 31–45.
- Peterson, E.L., 2011. The excluded philosophy of evo-devo? Revisiting C.H. Waddington's failed attempt to embed Alfred North Whitehead's "organicism" in evolutionary biology. *Hist. Philos. Life Sci.* 33 (3), 301–320.
- Piazza, P.V., Le Moal, M.L., 1996. Pathophysiological basis of vulnerability to drug abuse: role of an interaction between stress, glucocorticoids, and dopaminergic neurons. *Annu. Rev. Pharmacol. Toxicol.* 36, 359–378.
- Pizzagalli, D.A., Roberts, A.C., 2022. Prefrontal cortex and depression. *Neuropsychopharmacology* 47 (1), 225–246.
- Pluess, M., 2015. Individual differences in environmental sensitivity. *Child Development Perspectives* 9 (3), 138–143.
- Posteraro, T., Canalization, 2022. Creative evolution: images of life from Bergson to Whitehead and beyond. *Bergsonian*.
- Potts, R., et al., 2020. Increased ecological resource variability during a critical transition in hominin evolution. *Sci. Adv.* 6 (43).
- Qiao, Y., et al., 2020. Effect of combined chronic predictable and unpredictable stress on depression-like symptoms in mice. *Ann. Transl. Med.* 8 (15), 942.
- Ramstead, M.J.D., Badcock, P.B., Friston, K.J., 2018. Answering Schrodinger's question: a free-energy formulation. *Phys. Life Rev.* 24, 1–16.
- Raval, N.R., et al., 2021. A single Dose of psilocybin increases synaptic Density and decreases 5-HT_{2A} receptor Density in the pig brain. *Int. J. Mol. Sci.* 22 (2).
- Reggente, N., et al., 2018. Multivariate resting-state functional connectivity predicts response to cognitive behavioral therapy in obsessive-compulsive disorder. *Proc. Natl. Acad. Sci. U. S. A.* 115 (9), 2222–2227.
- Richards, W.A., 2015. Sacred Knowledge: Psychedelics and Religious Experiences. Columbia University Press, New York xxx, 244 pages.
- Riva, G., Gaudio, S., 2012. Allostatic lock in anorexia nervosa: new evidences from neuroimaging studies. *Med. Hypotheses* 79 (1), 113–117.
- Robbins, T.W., Everitt, B.J., 2002. Limbic-striatal memory systems and drug addiction. *Neurobiol. Learn. Mem.* 78 (3), 625–636.
- Romer, A.L., et al., 2021. Pervasively thinner neocortex as a transdiagnostic feature of general psychopathology. *Am. J. Psychiatr.* 178 (2), 174–182.
- Ruan, H., Yao, W.D., 2017. Cocaine promotes coincidence detection and lowers induction threshold during hebbian associative synaptic potentiation in prefrontal cortex. *J. Neurosci.* 37 (4), 986–997.
- Ruffini, G., Lopez-Sola, E., 2022. AIT foundations of structured experience. *PsyArXiv*.
- Ruffini, G., et al., 2022. LSD-induced increase of Ising temperature and algorithmic complexity of brain dynamics. *bioRxiv*, 2022.08.27.505518.
- Safron, A., 2020. On the Varieties of Conscious Experiences: Altered Beliefs under Psychedelics (ALBUS) *PsyArXiv*.
- Sansone, R.A., Sansone, L.A., 2010. Psychiatric disorders: a global look at facts and figures. *Psychiatry (Edmont)* 7 (12), 16–19.
- Schartner, M.M., et al., 2017. Increased spontaneous MEG signal diversity for psychoactive doses of ketamine, LSD and psilocybin. *Sci. Rep.* 7, 46421.
- Schneider, E.D., Sagan, D., 2005. Into the Cool: Energy Flow, Thermodynamics, and Life. University of Chicago Press, Chicago, p. 362 xviii.
- Seligman, M.E., Maier, S.F., Geer, J.H., 1968. Alleviation of learned helplessness in the dog. *J. Abnorm. Psychol.* 73 (3), 256–262.
- Selzam, S., et al., 2018. A polygenic p factor for major psychiatric disorders. *Transl. Psychiatry* 8 (1), 205.
- Sha, Z., et al., 2019. Common dysfunction of large-scale neurocognitive networks across psychiatric disorders. *Biol. Psychiatr.* 85 (5), 379–388.
- Shalom, J.G., Aderka, I.M., 2020. A meta-analysis of sudden gains in psychotherapy: Outcome and moderators. *Clin. Psychol. Rev.* 76, 101827.
- Shao, L.X., et al., 2021. Psilocybin induces rapid and persistent growth of dendritic spines in frontal cortex in vivo. *Neuron* 109 (16), 2535–2544 e4.
- Shiffman, S., Stone, A.A., Hufford, M.R., 2008. Ecological momentary assessment. *Annu. Rev. Clin. Psychol.* 4, 1–32.
- Siegel, J.S., et al., 2018. Re-emergence of modular brain networks in stroke recovery. *Cortex* 101, 44–59.
- Singleton, S.P., et al., 2022a. LSD and psilocybin flatten the brain's energy landscape: insights from receptor-informed network control theory. *bioRxiv*, 2021.05.14.444193.
- Singleton, S.P., et al., 2022b. Receptor-informed network control theory links LSD and psilocybin to a flattening of the brain's control energy landscape. *Nat. Commun.* 13 (1), 5812.
- Sinha, P., et al., 2019. Network neurobiology of electroconvulsive therapy in patients with depression. *Psychiatry Res. Neuroimaging* 287, 31–40.
- Sneve, M.H., et al., 2019. High-Expanding Regions in primate cortical brain evolution support supramodal cognitive flexibility. *Cerebr. Cortex* 29 (9), 3891–3901.
- Song, S., Miller, K.D., Abbott, L.F., 2000. Competitive Hebbian learning through spike-timing-dependent synaptic plasticity. *Nat. Neurosci.* 3 (9), 919–926.
- Spriggs, M.J., et al., 2021. Study protocol for "psilocybin as a treatment for anorexia nervosa: a pilot study". *Front. Psychiatr.* 12, 735523.
- Sroufe, L.A., 1997. Psychopathology as an outcome of development. *Dev. Psychopathol.* 9 (2), 251–268.
- Sumner, R.L., et al., 2020a. The role of Hebbian learning in human perception: a methodological and theoretical review of the human Visual Long-Term Potentiation paradigm. *Neurosci. Biobehav. Rev.* 115, 220–237.
- Sumner, R.L., et al., 2020b. Ketamine enhances visual sensory evoked potential long-term potentiation in patients with major depressive disorder. *Biol. Psychiatr. Cogn. Neurosci. Neuroimaging* 5 (1), 45–55.
- Suzuki, S., Nishimori, H., Suzuki, M., 2007. Quantum annealing of the random-field Ising model by transverse ferromagnetic interactions. *Phys. Rev. E - Stat. Nonlinear Soft Matter Phys.* 75 (5 Pt 1), 051112.
- Tagliazucchi, E., et al., 2014. Enhanced repertoire of brain dynamical states during the psychedelic experience. *Hum. Brain Mapp.* 35 (11), 5442–5456.
- Tagliazucchi, E., et al., 2016. Increased global functional connectivity correlates with LSD-induced ego dissolution. *Curr. Biol.* 26 (8), 1043–1050.
- Timmermann, C., et al., 2021. Psychedelics alter metaphysical beliefs. *Sci. Rep.* 11 (1), 22166.
- Timmermann, C., Watts, R., Dupuis, D., 2022. Towards psychedelic apprenticeship: Developing a gentle touch for the mediation and validation of psychedelic-induced insights and revelations. *Transcult. Psychiatr.* 13634615221082796.
- Toker, D., et al., 2022. Consciousness is supported by near-critical slow cortical electro-dynamics. *Proc. Natl. Acad. Sci. U. S. A.* 119 (7).
- Torregrossa, M.M., Corlett, P.R., Taylor, J.R., 2011. Aberrant learning and memory in addiction. *Neurobiol. Learn. Mem.* 96 (4), 609–623.
- Turrigiano, G.G., Nelson, S.B., 2004. Homeostatic plasticity in the developing nervous system. *Nat. Rev. Neurosci.* 5 (2), 97–107.
- Tzu, L., 1993. Tao Te Ching.
- Ungless, M.A., et al., 2001. Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. *Nature* 411 (6837), 583–587.
- Vaidya, V.A., et al., 1997. 5-HT_{2A} receptor-mediated regulation of brain-derived neurotrophic factor mRNA in the hippocampus and the neocortex. *J. Neurosci.* 17 (8), 2785–2795.
- van Bork, R., et al., 2017. What is the p-factor of psychopathology? Some risks of general factor modeling. *Theor. Psychol.* 27 (6), 759–773.
- Vanes, L.D., Dolan, R.J., 2021. Transdiagnostic neuroimaging markers of psychiatric risk: a narrative review. *Neuroimage Clin.* 30, 102634.
- Vargas, M.V., et al., 2021. Psychedelics and other psychoplastogens for treating mental illness. *Front. Psychiatr.* 12, 727117.
- Waddington, C.H., 1959. Canalization of development and genetic assimilation of acquired characters. *Nature* 183 (4676), 1654–1655.
- Waddington, C.H., 2014. The Strategy of the Genes. Routledge.
- Wakefield, J.C., 1992. The concept of mental disorder. On the boundary between biological facts and social values. *Am. Psychol.* 47 (3), 373–388.
- Walsh, J.J., et al., 2018. 5-HT release in nucleus accumbens rescues social deficits in mouse autism model. *Nature* 560 (7720), 589–594.
- Watts, R., et al., 2017. Patients' accounts of increased "connectedness" and "acceptance" after psilocybin for treatment-resistant depression. *J. Humanist. Psychol.* 57 (5), 520–564.

- Weinberger, D.R., Glick, I.D., Klein, D.F., 2015. Whither research domain criteria (RDoC)? the good, the bad, and the ugly. *JAMA Psychiatr.* 72 (12), 1161–1162.
- Weiss, B., et al., 2021a. Examining psychedelic-induced Changes in social Functioning and Connectedness in a naturalistic online sample Using the five-factor Model of personality. *Front. Psychol.* 12, 749788.
- Weiss, B., et al., 2021b. Examining changes in personality following shamanic ceremonial use of ayahuasca. *Sci. Rep.* 11 (1), 6653.
- Wertz, J., et al., 2021. Association of history of psychopathology with accelerated aging at midlife. *JAMA Psychiatr.* 78 (5), 530–539.
- Xu, J., et al., 2020. Electroconvulsive therapy modulates functional interactions between submodules of the emotion regulation network in major depressive disorder. *Transl. Psychiatry* 10 (1), 271.
- Yaden, D.B., Potash, J.B., Griffiths, R.R., 2022. Preparing for the bursting of the psychedelic hype bubble. *JAMA Psychiatr.*
- Ye, M., et al., 2015. Changes of functional brain networks in major depressive disorder: a graph theoretical analysis of resting-state fMRI. *PLoS One* 10 (9), e0133775.
- Yerushalmi, M., et al., 2021. Ecological momentary assessment of bipolar disorder symptoms and partner affect: longitudinal pilot study. *JMIR Form Res* 5 (9), e30472.
- Young, J.W., Dulcis, D., 2015. Investigating the mechanism(s) underlying switching between states in bipolar disorder. *Eur. J. Pharmacol.* 759, 151–162.
- Zeifman, R., Spriggs, M.J., Kettner, H., Lyons, T., Rosas, F., Mediano, P., et al., 2022. From Relaxed Beliefs under Psychedelics (REBUS) to Revised Beliefs after Psychedelics (REBAS): Preliminary Development of the RELaxed Beliefs Questionnaire (REB-Q). *PsyArXiv*.
- Zhou, K., D, W.D., Carhart-Harris, R.L., Kettner, H., 2022. Long Term Effects On Cognition And Perception Following Psychedelic Use In a Naturalistic Setting.