

A straight line is the shortest distance between two points

Nghia Doan

March 4, 2023

In this article, we explore some basic properties of broken lines.

Fact (Triangle Inequality). For any three points A, B, C ,


$$AB + BC \geq AC,$$

The equality holds if and only if A, B , and C are collinear.

Fact (Broken Line Inequality). For any points A_1, A_2, \dots, A_n , $A_1A_2 + A_2A_3 + \dots + A_{n-1}A_n \geq A_1A_n$. The equality holds if and only if A_1, A_2, \dots, A_{n-1} , and A_n are collinear.


Lemma (Heron's Problem)

Two points A and B lie on one side of a straight line l . C is a point on l . The sum $CA + CB$ is minimal if and only if $C = BC' \cap l$, where B' is the reflection of B over l .




Example (Cross-section of a cube)

Lilian cuts a cube with side length 1. She got a with a hexagon cross-section as shown below. What is the minimal value of the hexagon perimeter $AB + BC + CD + DE + EF + FA$?




Solution. The diagram below is obtained by unfolding the cube into a net. The hexagon perimeter forms a broken line $ABCDEFA'$.



This is always larger or equal the distance AA' , which is same as three times the diagonal of the unit square. Hence the perimeter is always at least $3\sqrt{2}$. \square

Example (Diagonal of a hexagon)


$ABCDEF$ is a convex hexagon, where $\angle A \geq 90^\circ$ and $\angle D \geq 90^\circ$. Prove that $BC + CE + EF + FB \geq 2AD$.



Proof. Let M, N , and P be the midpoints of BF, BE , and CE , respectively. Since any broken line is longer or equal the distance between two endpoints, so $AD \leq AM + MN + NP + PD$. MN is the median segment in $\triangle BEF$, thus $FE = 2MN$. Similarly $BC = 2NP$. In $\triangle ABF$, $\angle A \geq 90^\circ$, thus $BF \geq 2AM$. Similarly $CE \geq 2DP$. Therefore $BC + CE + EF + FB \geq 2(AM + MN + NP + PD) = 2AD$. \square

Example (Romanian Math Olympiad)

Let $ABCD$ be a convex quadrilateral. It is known that the circles with diameter AB and CD are externally tangent, and so are the circles with diameters AD and BC . Prove that $ABCD$ is a rhombus.



Proof. We first prove a claim.

Claim — Let M and N be the midpoint of AB and CD , respectively, then $AD + BC \geq 2MN$.

Proof. Let E be the midpoint of AC . It is easy to see that

$$MN \leq ME + EN = \frac{BC}{2} + \frac{AD}{2} = \frac{AD + BC}{2}$$

The equality can happen if and only if MN intersect AC at the midpoint of AC , so $MN \parallel AD \parallel BC$. ■

By the claim $AD + BC \geq 2MN$, similarly $AB + CD \geq 2PQ$, thus

$$AB + BC + CD + DA \geq 2(MN + PQ) \quad (*)$$

Now, let P and Q be the midpoints of BC and AD , respectively. Since the circles of diameters AB and CD are externally tangent so $AB + CD = 2MN$, similarly $AD + BC = 2PQ$. Thus

$$AB + BC + CD + DA = 2(MN + PQ) \quad (**)$$

(**) implies the existence of equality in (*), so $MN \parallel AD \parallel BC$ and $PQ \parallel AB \parallel CD$. Thus $ABCD$ is a parallelogram, and $MN = AD = BC$. Similarly $AB = CD$. Since $AB + CD = 2MN$ (see above), therefore

$$AD = BC = MN = AB = CD.$$

Hence, $ABCD$ is a rhombus. □