

ON THE CONTINUITY OF PARTIAL ACTIONS OF HAUSDORFF GROUPS ON METRIC SPACES

J. GÓMEZ, H. PINEDO, AND C. UZCÁTEGUI

ABSTRACT. We provide a sufficient condition for a topological partial action of a Hausdorff group on a metric space is continuous, provide that it is separately continuous.

1. INTRODUCTION

The notion of partial action of a group is a weakening of the classical concept of group action and was introduced in [6] and [11], and was then developed in [1] and [10], in which the authors provided examples in different guises. Since then partial actions have been an important tool in C^* -algebras and dynamical systems, and in the developing of new cohomological theories [4], [5] and [12]. Every partial action of a group G on a set \mathbb{X} can be obtained, roughly speaking, as a restriction of a global action (see [1] and [10]) on some bigger set \mathbb{X}_G , called the enveloping space of \mathbb{X} . Nevertheless, in the category of topological spaces, when G acts partially on a space \mathbb{X} , the superspace \mathbb{X}_G does not necessarily inherit its topological properties; for instance the globalization of a partial action of a group on a Hausdorff space is not in general Hausdorff (see e.g. [1, Example 1.4], [1, Proposition 1.2]).

On the other hand, actions of Polish groups have important connections with many areas of mathematics (see [2], [8] and the references therein). Recall that a Polish space is a topological space which is separable and completely metrizable, and a Polish group is a topological group whose topology is Polish. Partial actions of Polish groups have been recently considered in the works [9, 13, 14].

It is known that an action of a Polish group G on a metric space \mathbb{X} is continuous provided that it is separately continuous (see [8, Theorem 3.1.4]). In this note, under a mild restriction, we generalize [8, Theorem 3.1.4] in two directions, first we only assume that G is Hausdorff and Baire, and second we only assume that G acts partially on \mathbb{X} .

2. THE NOTIONS

Let G be a topological group with identity element 1 and \mathbb{X} a topological space. A partially defined map $G \times \mathbb{X} \rightarrow \mathbb{X}$ is a map whose domain is a subset of $G \times \mathbb{X}$. Let $m: G \times \mathbb{X} \rightarrow \mathbb{X}$, $(g, x) \mapsto m(g, x) = g \cdot x \in \mathbb{X}$ be a partial map, that is m is a map whose domain is contained in $G \times \mathbb{X}$. We write $\exists g \cdot x$ to mean that (g, x) is in the domain of m . Then, following [1, 10] m defines a (set theoretic) *partial action* of G on \mathbb{X} , if for all $g, h \in G$ and $x \in \mathbb{X}$ the following assertions hold:

- (PA1) $\exists g \cdot x$ implies $\exists g^{-1} \cdot (g \cdot x)$ and $g^{-1} \cdot (g \cdot x) = x$,
- (PA2) $\exists g \cdot (h \cdot x)$ implies $\exists (gh) \cdot x$ and $g \cdot (h \cdot x) = (gh) \cdot x$,
- (PA3) $\exists 1 \cdot x$, and $1 \cdot x = x$.

Date: February 26, 2018.

1991 Mathematics Subject Classification. 54H15, 54E50, 54E35.

Key words and phrases. Partial action, separately continuity, Hausdorff groups.

The second and third authors thank La Vicerrectoría de Investigación y Extensión de la Universidad Industrial de Santander for the financial support for this work, which is a part of the VIE project # 5761.

Let $G * \mathbb{X} = \{(g, x) \in G \times \mathbb{X} \mid \exists g \cdot x\}$ be the domain of m , set $\mathbb{X}_{g^{-1}} = \{x \in \mathbb{X} \mid \exists g \cdot x\}$, and $m_g: \mathbb{X}_{g^{-1}} \ni x \mapsto g \cdot x \in \mathbb{X}_g$. Then a partial action $m: G * \mathbb{X} \rightarrow \mathbb{X}$ induces a family of bijections $\{m_g: \mathbb{X}_{g^{-1}} \rightarrow \mathbb{X}_g\}_{g \in G}$, and we denote $m = \{m_g: \mathbb{X}_{g^{-1}} \rightarrow \mathbb{X}_g\}_{g \in G}$.

Proposition 2.1. [15, Lemma 1.2] *A partial action m of G on \mathbb{X} is a family $m = \{m_g: \mathbb{X}_{g^{-1}} \rightarrow \mathbb{X}_g\}_{g \in G}$, where $\mathbb{X}_g \subseteq \mathbb{X}$, $m_g: \mathbb{X}_{g^{-1}} \rightarrow \mathbb{X}_g$ is bijective, for all $g \in G$, and such that:*

- (i) $\mathbb{X}_1 = \mathbb{X}$ and $m_1 = \text{Id}_{\mathbb{X}}$;
- (ii) $m_g(\mathbb{X}_{g^{-1}} \cap \mathbb{X}_h) = \mathbb{X}_g \cap \mathbb{X}_{gh}$;
- (iii) $m_g m_h: \mathbb{X}_{h^{-1}} \cap \mathbb{X}_{h^{-1}g^{-1}} \rightarrow \mathbb{X}_g \cap \mathbb{X}_{gh}$, and $m_g m_h = m_{gh}$ in $\mathbb{X}_{h^{-1}} \cap \mathbb{X}_{h^{-1}g^{-1}}$;

for all $g, h \in G$.

Notice that conditions (ii) and (iii) in Lemma 2.1 say that m_{gh} is an extension of $m_g m_h$ for all $g, h \in G$. We consider $G \times \mathbb{X}$ with the product topology and the subset $G * \mathbb{X}$ of $G \times \mathbb{X}$ inherits the subspace topology.

Definition 2.2. *A topological partial action of the group G on the topological space \mathbb{X} is a partial action $m = \{m_g: \mathbb{X}_{g^{-1}} \rightarrow \mathbb{X}_g\}_{g \in G}$ on the underlying set \mathbb{X} , such that each \mathbb{X}_g is open in \mathbb{X} , and each m_g is a homeomorphism, for any $g \in G$. If $m: G * \mathbb{X} \rightarrow \mathbb{X}$ is continuous, we say that the partial action is continuous.*

Example 2.3. Induced partial action: *Let G be a topological group, and \mathbb{Y} a topological space and let $u: G \times \mathbb{Y} \rightarrow \mathbb{Y}$ be a continuous action of G on \mathbb{Y} and $\mathbb{X} \subseteq \mathbb{Y}$ be an open set. For $g \in G$, set $\mathbb{X}_g = \mathbb{X} \cap u_g(\mathbb{X})$ and let $m_g = u_g \upharpoonright \mathbb{X}_{g^{-1}}$ (the restriction of u_g to $\mathbb{X}_{g^{-1}}$). Then $m: G * \mathbb{X} \ni (g, x) \mapsto m_g(x) \in \mathbb{X}$ is a topological partial action of G on \mathbb{X} .*

The interested reader may consult others examples of topological partial actions in [1, Example 1.2, Remark 1.1, Example 1.3, Example 1.4] and [10, p. 108].

3. TOPOLOGICAL PARTIAL ACTIONS OF HAUSDORFF GROUPS ON METRIC SPACES

Let $m: G * \mathbb{X} \rightarrow \mathbb{X}$ be a topological partial action. For $x \in \mathbb{X}$, we set $G^x = \{g \in G \mid (g, x) \in G * \mathbb{X}\}$, then by (PA3) $1 \in G^x$, we also set $m^x: G^x \ni g \rightarrow m(g, x) \in \mathbb{X}$, m is called *separately continuous* if the maps m^x are continuous, for all $x \in \mathbb{X}$.¹ It is known that group actions of Polish groups on metric spaces are continuous, if and only if, they are separately continuous. We provide a mild condition on G^x , $x \in \mathbb{X}$, for which separately continuous partial actions of Hausdorff-Baire groups are continuous, the proof we present is inspired by the one given in [8, Theorem 3.1.4], for classical group actions.

Theorem 3.1. *Let G be a Hausdorff group, (\mathbb{X}, d) a metric space, and m a topological partial action of G on \mathbb{X} . Suppose that G is Baire G^x is open in G , for any $x \in \mathbb{X}$. Then m is continuous, if and only if, it is separately continuous.*

Proof. It is clear that continuous partial actions are separately continuous. For the converse, suppose that m is separately continuous and let $(g_0, x_0) \in G * \mathbb{X}$, we check that m is continuous at (g_0, x_0) . Let $l, n \in \mathbb{N}$ and set

$$F_{n,l} = \{g \in G^{x_0} \mid \forall x \in \mathbb{X}_{g^{-1}} (d(x, x_0) < 2^{-n} \Rightarrow d(m(g, x), m(g, x_0)) \leq 2^{-l})\}.$$

We shall check that $F_{n,l}$ is a closed subset of G^{x_0} . Indeed, let $g \in G^{x_0}$ and $\{g_i\}_{i \in I}$ a net with $g_i \rightarrow g$ such that $g_k \in F_{n,l}$, for all $k \in \mathbb{N}$. Then

$$(\forall k \in \mathbb{N}) \left(\forall x \in \mathbb{X}_{g_i^{-1}} \right) \left(d(x, x_0) < \frac{1}{2^n} \Rightarrow d(m(g_i, x), m(g_i, x_0)) \leq \frac{1}{2^l} \right).$$

¹Notice that for any $g \in G$ the map m_g is continuous by definition of topological partial action.

Let $x \in \mathbb{X}_{g^{-1}}$, that is $g \in G^x$, since G^x is open one may assume that $\{g_i\}_{i \in I} \subseteq G^x$, and $x \in \mathbb{X}_{g^{-1}} \cap \mathbb{X}_{g_i^{-1}}$, for all $i \in I$. Thus, by the continuity of m^x and m^{x_0} , we have that $m(g_i, x) \rightarrow m(g, x)$ and $m(g_i, x_0) \rightarrow m(g, x_0)$, which gives

$$(\forall x \in \mathbb{X}_{g^{-1}}) \left(d(x, x_0) < \frac{1}{2^n} \Rightarrow d(m(g, x), m(g, x_0)) \leq \frac{1}{2^l} \right),$$

and we conclude that $g \in F_{n,l}$. Now, we check that

$$(3.1) \quad G^{x_0} = \bigcap_l \bigcup_n F_{n,l}.$$

It is clear that $G^{x_0} \supseteq \bigcap_l \bigcup_n F_{n,l}$. Conversely, take $g \in G^{x_0}$ and $l \in \mathbb{N}$. Since m_g is continuous at x_0 , for $\varepsilon = \frac{1}{2^l}$, there exists $\delta > 0$ such that

$$(\forall x \in \mathbb{X}_{g^{-1}}) \left(d(x, x_0) < \delta \Rightarrow d(m(g, x), m(g, x_0)) \leq \frac{1}{2^l} \right),$$

Let $n \in \mathbb{N}$ with $\frac{1}{2^n} < \delta$, then $g \in F_{n,l}$. Thus $G^{x_0} \subseteq \bigcap_l \bigcup_n F_{n,l}$, and we have (3.1).

Since $F_{l,n}$ is closed in G^{x_0} , for all n, l ; the set $D = \bigcup_l \bigcup_n (F_{n,l} \setminus \text{int}(F_{n,l}))$ is meager. But G^{x_0} is a non empty open set, since G is Baire, there is $g_1 \in G^{x_0} \setminus D$. We shall check that m is continuous at (g_1, x_0) . Indeed, let $\{(h_\alpha, y_\alpha)\}_{\alpha \in \Lambda} \subseteq G * \mathbb{X}$ be a net converging to (g_1, x_0) . Take $\varepsilon > 0$ and $l \in \mathbb{N}$ such that $\frac{1}{2^{l-1}} < \varepsilon$. By (3.1), there exists $n \in \mathbb{N}$ such that $g_1 \in F_{n,l}$. Since $g_1 \notin D$ then $g_1 \in \text{int}(F_{n,l})$. The fact that $h_\alpha \rightarrow g_1$, implies that there is $\alpha_1 \in \Lambda$ such that $h_\alpha \in \text{int}(F_{n,l})$, for all $\alpha \geq \alpha_1$. Also, since $y_\alpha \rightarrow x_0$, there exists $\alpha_2 \in \mathbb{N}$ such that $d(y_\alpha, x_0) < \frac{1}{2^n}$, for all $\alpha \geq \alpha_2$. Additionally, by the continuity of m^{x_0} , we have that $m(h_\alpha, x_0) \rightarrow m(g_1, x_0)$. Thus, there exists $\alpha_3 \in \Lambda$ such that $d(m(h_\alpha, x_0), m(g_1, x_0)) < \frac{1}{2^l}$, for all $\alpha \geq \alpha_3$.

Let $\alpha \geq \max\{\alpha_1, \alpha_2, \alpha_3\}$. Then $h_\alpha \in F_{n,l}$ and $d(y_\alpha, x_0) < \frac{1}{2^n}$, hence

$$d(m(h_\alpha, y_\alpha), m(g_1, x_0)) \leq d(m(h_\alpha, y_\alpha), m(h_\alpha, x_0)) + d(m(h_\alpha, x_0), m(g_1, x_0)) < \frac{1}{2^{l-1}} < \varepsilon.$$

Thus, m is continuous at (g_1, x_0) .

Now, since $x_0 \in \mathbb{X}_{g_1^{-1}} \cap \mathbb{X}_{g_0^{-1}}$, we have, by Proposition 2.1(ii), that $g_1 \cdot x_0 \in \mathbb{X}_{g_1} \cap \mathbb{X}_{g_1 g_0^{-1}}$, then $(g_0 g_1^{-1}) \cdot (g_1 \cdot x_0)$ is defined and $(g_0 g_1^{-1}) \cdot (g_1 \cdot x_0) = g_0 \cdot x_0$, by (PA2). That is

$$(3.2) \quad m(g_0, x_0) = m(g_0 g_1^{-1}, m(g_1, x_0)).$$

Finally take a net $\{(h_j, y_j)\}_{j \in J}$ in $G * \mathbb{X}$ converging to (g_0, x_0) . Then $g_1 g_0^{-1} h_j \rightarrow g_1 \in G^{x_0}$, and since G^{x_0} is open we assume that the net $\{g_1 g_0^{-1} h_j\}_{j \in J}$ is contained in G^{x_0} , thus $x_0 \in \mathbb{X}_{(g_1 g_0^{-1} h_j)^{-1}}$, for all $j \in J$. But $y_j \rightarrow x_0$, then there is $j_0 \in J$ such that $y_j \in \mathbb{X}_{(g_1 g_0^{-1} h_j)^{-1}}$, for all $j \geq j_0$. Now $y_j \in \mathbb{X}_{(g_1 g_0^{-1} h_j)^{-1}} \cap \mathbb{X}_{h_j^{-1}}$, then $h_j \cdot y_j \in \mathbb{X}_{(g_1 g_0^{-1})^{-1}}$ (by Proposition 2.1(ii)). By (PA2) we get

$$(g_1 g_0^{-1} h_j) \cdot y_j = (g_1 g_0^{-1}) \cdot (h_j \cdot y_j) \in \mathbb{X}_{g_1 g_0^{-1}}.$$

Thus, by (PA1),

$$h_j \cdot y_j = (g_0 g_1^{-1}) \cdot [(g_1 g_0^{-1} h_j) \cdot y_j].$$

Notice that $(g_1 g_0^{-1} h_j, y_j) \rightarrow (g_1, x_0)$. Then by continuity of $m_{g_0 g_1^{-1}}$ and the continuity of m at (g_1, x_0) one gets

$$m(h_j, y_j) = (g_0 g_1^{-1}) \cdot [(g_1 g_0^{-1} h_j) \cdot y_j] \rightarrow (g_0 g_1^{-1}) \cdot (g_1 \cdot x_0) = m(g_0 g_1^{-1}, m(g_1, x_0)),$$

and by (3.2) we obtain $m(h_j, y_j) \rightarrow m(g_0, x_0)$. \square

Corollary 3.2. *Let G be a topological Hausdorff group, and $a: G \times \mathbb{X} \rightarrow \mathbb{X}$ an action of G on a metric space \mathbb{X} . If G is Baire, then a is continuous, if and only if, a is separately continuous.*

Corollary 3.3. *Let G be a countable discrete group and m a topological partial action of G on a metric space \mathbb{X} . Then m is continuous, if and only if, m is separately continuous.*

Example 3.4. Möbius transformations [3, p. 175] *The group $G = \text{GL}(2, \mathbb{R})$ is Polish and acts partially on \mathbb{R} by setting*

$$g \cdot x = \frac{ax + b}{cx + d}, \quad \text{where } g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G.$$

Notice that for all $g \in G$ the set $\mathbb{R}_g = \{x \in \mathbb{R} \mid cx + d \neq 0\}$ is open, and $m = \{m_g: \mathbb{R}_{g^{-1}} \ni x \rightarrow g \cdot x \in \mathbb{R}_g\}_{g \in G}$ is a topological partial action. For $x \in \mathbb{R}$ let $t_x = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$, then for $y \in \mathbb{R}$ one has that $t_{y-x} \cdot x = y$. Since

$$G^0 = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G : d \neq 0 \right\}$$

is open, and

$$G^x = G^{t_x \cdot 0} = G^0 t_x^{-1} = G^0 \begin{pmatrix} 1 & -x \\ 0 & 1 \end{pmatrix},$$

then G^x is open. Finally, since

$$m^x: G^x \ni \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ax + b}{cx + d} \in \mathbb{R},$$

is continuous, then m is continuous.

Acknowledgements

The authors thank the referee for his/her suggestions and comments that helped to improve the quality of the paper. In particular, for suggesting the current version of Theorem 3.1, that was initially established only for Polish groups.

REFERENCES

- [1] F. Abadie, Enveloping actions and Takai duality for partial actions. *Journal of Func. Anal.* (2003) 197: 14-67.
- [2] H. Becker, A. Kechris, The Descriptive Set Theory of Polish Group Actions London Math. Soc. Lect. Notes, 1996.
- [3] K. Choi, Y. Lim, Transitive partial actions of groups, *Period. Math. Hung.* **56**, (2008), No. 2, 169-181.
- [4] M. Dokuchaev, B. Novikov, H. Pinedo, The partial Schur Multiplier of a group, *J. Algebra*, **392** (2013), 199-225.
- [5] M. Dokuchaev, M. Khrypchenko, Partial cohomology of groups, *J. Algebra*, **427** (2015), 142-182.
- [6] R. Exel, Partial actions of groups and actions of inverse semigroups, *Proc. Am. Math. Soc.* **126** (1998), (12), 3481-3494.
- [7] R. Exel, T. Giordano, D. Gonçalves, Enveloping algebras of partial actions as groupoid C^* -algebras. *J. Operator Theory* (2011) 65: 197-210.
- [8] Su Gao, Invariant Descriptive Set Theory, *Chapmann & Hall* (2009).
- [9] J. Gómez, H. Pinedo and C. Uzcátegui, The open mapping principle for partial actions of Polish groups. *Preprint*.
- [10] J. Kellendonk and M. V. Lawson, Partial actions of groups. *Internat. J. Algebra Comput.* (2004) 14: 87-114.
- [11] K. Mc Clanahan, K -theory for partial crossed products by discrete groups, *J. Funct. Anal.* (1995) 130: 77-117.

- [12] H. Pinedo, Partial projective representations and the partial Schur multiplier: a survey. *Bol. Mat.* (2015) (22) 2 167–175.
- [13] H. Pinedo and C. Uzcátegui, Polish globalization of polish group partial actions. *Math. Log. Quart.* To appear.
- [14] H. Pinedo and C. Uzcátegui, Borel globalization of partial actions of polish groups. *arxiv:1702.02611v1* .
- [15] J.C. Quigg, I. Raeburn, Characterizations of crossed products by partial actions. *J.Operator Theory* (1997) 37: 311-340.

ESCUELA DE MATEMÁTICAS, UNIVERSIDAD INDUSTRIAL DE SANTANDER, CRA. 27 CALLE 9 UIS EDIFICIO 45,
BUCARAMANGA, COLOMBIA

E-mail address: `jorge.gomez14@correo.uis.edu.co`

ESCUELA DE MATEMÁTICAS, UNIVERSIDAD INDUSTRIAL DE SANTANDER, CRA. 27 CALLE 9 UIS EDIFICIO 45,
BUCARAMANGA, COLOMBIA

E-mail address: `hpinedot@uis.edu.co`

ESCUELA DE MATEMÁTICAS, UNIVERSIDAD INDUSTRIAL DE SANTANDER, CRA. 27 CALLE 9 UIS EDIFICIO 45,
BUCARAMANGA, COLOMBIA

E-mail address: `cuzcatea@uis.edu.co`