

PROJECTIVE COCYCLES OVER $SL(2, \mathbb{R})$ ACTIONS: MEASURES INVARIANT UNDER THE UPPER TRIANGULAR GROUP.

CHRISTIAN BONATTI, ALEX ESKIN, AND AMIE WILKINSON

ABSTRACT. We consider the action of $SL(2, \mathbb{R})$ on a vector bundle \mathbf{H} preserving an ergodic probability measure ν on the base X . Under an irreducibility assumption on this action, we prove that if $\hat{\nu}$ is any lift of ν to a probability measure on the projectivized bundle $\mathbb{P}(\mathbf{H})$ that is invariant under the upper triangular subgroup, then $\hat{\nu}$ is supported in the projectivization $\mathbb{P}(\mathbf{E}_1)$ of the top Lyapunov subspace of the positive diagonal semigroup. We derive two applications. First, the Lyapunov exponents for the Kontsevich-Zorich cocycle depend continuously on affine measures, answering a question in [MMY]. Second, if $\mathbb{P}(\mathbf{V})$ is an irreducible, flat projective bundle over a compact hyperbolic surface Σ , with hyperbolic foliation \mathcal{F} tangent to the flat connection, then the foliated horocycle flow on $T^1\mathcal{F}$ is uniquely ergodic if the top Lyapunov exponent of the foliated geodesic flow is simple. This generalizes results in [BG] to arbitrary dimension.

1. INTRODUCTION

Let G denote the group $SL(2, \mathbb{R})$, and for $t \in \mathbb{R}$, let

$$g^t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}, \quad u_+^t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, \quad \text{and} \quad u_-^t = \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}.$$

Let $A = \{g^t : t \in \mathbb{R}\}$, $U_+ = \{u_+^t : t \in \mathbb{R}\}$, and $U_- = \{u_-^t : t \in \mathbb{R}\}$. The group G is generated by A, U_+ and U_- . We denote by $P = AU_+$ the group of upper triangular matrices, and we recall that the group P is solvable and hence *amenable*: any action of P on a compact metric space admits an invariant probability measure (see, e.g. [EW, Section 8.4]). Recall that $PSL(2, \mathbb{R})$ admits a natural identification with the unit tangent bundle $T^1\mathbb{H}^2$ where \mathbb{H}^2 is the upper half plane endowed with the hyperbolic metric. Through this identification, the left action of $SL(2, \mathbb{R})$ on itself induces an identification of $A = \{g^t\}$ with the geodesic flow, and U_+ (resp. U_-) with the unstable (resp. stable) horocycle flow of \mathbb{H}^2 .

In this paper, we consider the following situation. Suppose G acts on a separable metric space X , preserving an ergodic Borel probability measure ν . By the Mautner phenomenon, ν is ergodic with respect to the action of the diagonal group A . Let $\mathbf{H} \rightarrow X$ be a continuous vector bundle over X with fiber a finite dimensional vector space H , and write $\mathbf{H}(x)$ for the fiber of \mathbf{H} over $x \in X$. Suppose that G acts on \mathbf{H} by linear automorphisms on the fibers and the given action on the base. We denote the action of $g \in G$ on \mathbf{H} by g_* , so for $\mathbf{v} \in \mathbf{H}(x)$, we have $g_*\mathbf{v} \in \mathbf{H}(gx)$.

Assume that \mathbf{H} is equipped with a Finsler structure (that is, a continuous choice of norm $\{\|\cdot\|_x : x \in X\}$ on the fibers of \mathbf{H}), and that with respect to this Finsler, the action of G on \mathbf{H} satisfies the following integrability condition:

$$(1) \quad \int_X \sup_{t \in [-1,1]} (\log \|g_*^t\|_x) d\nu(x) < +\infty,$$

where, for $g \in G$, $\|g_*\|_x$ denotes the operator norm of the linear action of g_* on the fiber over x with respect to the Finsler structure:

$$\|g_*\|_x := \sup_{\{\mathbf{v} \in H : \|\mathbf{v}\|_x = 1\}} \|g_*\mathbf{v}\|_{gx}.$$

Since the A -action on X is ergodic with respect to ν , the Oseledets multiplicative ergodic theorem implies that there exists a g^t -equivariant splitting

$$(2) \quad \mathbf{H}(x) = \bigoplus_{j=1}^m \mathbf{E}_j(x),$$

defined for ν -almost every $x \in X$, and real numbers $\lambda_1 > \dots > \lambda_m$ (called the *Lyapunov exponents* of the A -action) such that for ν -a.e. $x \in X$ and all $\mathbf{v} \in \mathbf{E}_j(x)$,

$$\lim_{|t| \rightarrow \infty} \frac{1}{t} \log \frac{\|g_*^t \mathbf{v}\|_{g^t x}}{\|\mathbf{v}\|_x} = \lambda_j.$$

Definition 1.1 (ν -measurable invariant subbundle). A ν -measurable invariant subbundle for the G action on \mathbf{H} is a measurable linear subbundle \mathbf{W} of \mathbf{H} with the property that $g_*(\mathbf{W}(x)) = \mathbf{W}(gx)$, for ν -a.e. $x \in X$ and every $g \in G$.

Definition 1.2 (irreducible). We say that the G action on \mathbf{H} is *irreducible with respect to the G -invariant measure ν on X* if it does not admit ν -measurable invariant subbundles: that is, if \mathbf{W} is a ν -measurable invariant subbundle, then either $\mathbf{W}(x) = \{0\}$, ν -a.e. or $\mathbf{W}(x) = \mathbf{H}(x)$, ν -a.e.

We note that the bundles \mathbf{E}_j in the splitting (2) are not invariant in the sense of Definition 1.1, since they are (in general) equivariant only under the subgroup A and not under all of G .

Let $\mathbb{P}(H)$ be the projective space of H (i.e., the space of lines in H), and let $\mathbb{P}(\mathbf{H})$ be the projective bundle associated to \mathbf{H} with fiber $\mathbb{P}(H)$. Then G also acts on $\mathbb{P}(\mathbf{H})$ via the induced projective action on the fibers. The space $\mathbb{P}(\mathbf{H})$ may not support a G -invariant measure, but since P is amenable and $\mathbb{P}(H)$ is compact, it will always support a P -invariant measure. In particular, for any P -invariant measure μ on X , there will be a P -invariant measure $\hat{\mu}$ on $\mathbb{P}(\mathbf{H})$ that projects to μ under the natural map $\mathbb{P}(\mathbf{H}) \rightarrow X$. For such a μ , denote by $\mathcal{M}_P^1(\mu)$ the (nonempty) set of all P -invariant Borel probability measures on $\mathbb{P}(\mathbf{H})$ projecting to μ on X .

We can now state our main theorem.

Theorem 1.3. *Suppose that the G -action on $\pi: \mathbf{H} \rightarrow X$ is irreducible with respect to the G -invariant (and therefore P -invariant) measure ν on X , and let $\hat{\nu} \in \mathcal{M}_P^1(\nu)$. Disintegrating $\hat{\nu}$ along the fibers of $\mathbb{P}(\mathbf{H})$, write*

$$d\hat{\nu}([\mathbf{v}]) = d\eta_{\pi(\mathbf{v})}([\mathbf{v}]) d\nu(\pi(\mathbf{v})),$$

where $[\mathbf{v}] \in \mathbb{P}(\mathbf{H})$ denotes the line determined by $\{0\} \neq \mathbf{v} \in \mathbf{H}$. Then for ν -a.e. $x \in X$, the measure η_x on $\mathbb{P}(\mathbf{H})(x)$ is supported on $\mathbb{P}(\mathbf{E}_1(x))$, where as in (2), $\mathbf{E}_1(x)$ is the Lyapunov subspace corresponding to the top Lyapunov exponent of the A -action.

In particular, if \mathbf{E}_1 is one-dimensional, then $\#\mathcal{M}_P^1(\nu) = 1$.

The same conclusions of Theorem 1.3 hold when $G = SL(2, \mathbb{R})$ is replaced by any rank 1 semisimple Lie group, and P denotes the Borel (i.e. minimal parabolic) subgroup of G .

P -invariant measures are a natural object of study, as they are closely related to stationary measures of a G -action. Suppose that G acts on a space Ω , and let m be a Borel probability measure on G . Recall that a probability measure ρ on Ω is m -stationary if $m * \rho = \rho$, where for $A \subset \Omega$ measurable, we define

$$m * \rho(A) = \int_G \rho(gA) dm(g).$$

A compactly supported Borel probability measure m on G is *admissible* if the following two conditions hold: first, there exists a $k \geq 1$ such that the k -fold convolution m^{*k} is absolutely continuous with respect to Haar measure; second, the smallest closed subgroup containing $\text{supp}(m)$ is G itself. Furstenberg proved that there is a 1–1 correspondence between P -invariant measures on Ω and m -stationary measures for admissible m . In fact any m -stationary measure on Ω is of the form $\lambda * \hat{\nu}$ where λ is the unique m -stationary measure on the Furstenberg boundary of G (which is in our case the circle G/P) and $\hat{\nu}$ is a P -invariant measure on Ω .

Remark. In light of the discussion above, for any admissible measure m on G , Theorem 1.3 also gives a classification of the m -stationary measures on \mathbf{H} projecting to ν .

In the context where $X = SL(2, \mathbb{R})/\Gamma$, with Γ cocompact (or of finite covolume), \mathbf{H} a flat H -bundle over X , and $H = \mathbb{R}^2$, \mathbb{C}^2 or \mathbb{C}^3 , Theorem 1.3 was proved by Bonatti and Gomez-Mont [BG], where irreducibility is replaced with the equivalent hypothesis that $\rho(\Gamma)$ is Zariski dense, where ρ is the monodromy representation.

Some constructions used in the proof of Theorem 1.3 are also used in [EM] in their classification of $SL(2, \mathbb{R})$ invariant probability measures on moduli spaces.

2. APPLICATIONS AND THE IRREDUCIBILITY CRITERION

The irreducibility hypothesis in Theorem 1.3 is not innocuous. Checking for the non-existence of invariant *measurable* subbundles is in general an impossible task, but there are two restricted contexts where it is feasible, on which we focus here:

- Suppose that $X = G/\Gamma$, for some discrete subgroup $\Gamma \subset G$, and \mathbf{H} is a flat bundle over $X = G/\Gamma$ with monodromy representation $\rho: \Gamma \rightarrow GL(H)$. Then G acts transitively on X , the P -invariant measures on X are all algebraic by [Moz] (which uses Ratner's Theorem), and irreducibility is then equivalent to the condition that there are no invariant algebraic subbundles of \mathbf{H} . In the case where ν is Haar measure, irreducibility of the associated G -action reduces to the condition that ρ is an irreducible representation. In Subsection 2.1 we derive some consequences of Theorem 1.3 in this context.
- If the bundle \mathbf{H} admits a Hodge structure (not necessarily G -invariant) then checking irreducibility can sometimes be reduced to showing that there are no invariant subbundles that are compatible with the Hodge structure, a much simpler task (since such subbundles must be real-analytic). In particular, the condition that G acts transitively on the base in the previous setting can be relaxed. This has been established rigorously by Simion Filip for the Kontsevich-Zorich action, and we use Theorem 1.3 in Subsection 2.2 to deduce further results in that context.

The mantra here is that for such Hodge bundles whose base supports a G -invariant measure, any measurable G -invariant subbundle must come from algebraic geometry. For the Kontsevich-Zorich action, this has been established in [EFW].

We now describe the applications in more detail.

2.1. Linear representations of G -lattices.

The first application concerns the dynamics of the horocycle flow of a foliation with hyperbolic leaves. If a manifold M has a 2-dimensional foliation \mathcal{F} whose leaves carry continuously varying hyperbolic structures, then there is a natural G -action on the unit tangent bundle $T^1\mathcal{F} \subset T^1M$ to the leaves, induced by the identification of $T^1\mathbb{H}^2$ with $PSL(2, \mathbb{R})$ mentioned in the introduction. This gives rise to the foliated geodesic and horocyclic flows on $T^1\mathcal{F}$.

The dynamical properties of these flows have geometric consequences for the foliations, including properties of harmonic measures (A probability measure on M is *harmonic* with respect to the foliation \mathcal{F} if it is invariant under leafwise heat flow: see [Ga]). In particular, a probability measure on M is harmonic along the leaves of \mathcal{F} if and only if it is the projection of a P -invariant measure on $T^1\mathcal{F}$ ([Mar, BaMa]). We consider here the special case of foliations induced by representations of surface groups into linear groups. First, we discuss the suspension construction.

2.1.1. The suspension construction and a criterion for simplicity.

Let $\Gamma < SL(2, \mathbb{R})$ be a lattice, and let $\rho: \Gamma \rightarrow GL(H)$ be a representation, where $H = \mathbb{R}^k$ or \mathbb{C}^k , $k \geq 2$. Then Γ acts on $SL(2, \mathbb{R}) \times H$ by the diagonal action

$$\gamma: (g, v) \mapsto (g\gamma^{-1}, \rho(\gamma)v).$$

The group $G = SL(2, \mathbb{R})$ also acts on $SL(2, \mathbb{R}) \times H$, by left multiplication in the first factor, and trivially in the second factor. The Γ and G actions commute.

Define the *suspension* of ρ :

$$\mathbf{H}_\rho = SL(2, \mathbb{R}) \times H/\Gamma;$$

it is a H -bundle over $SL(2, \mathbb{R})/\Gamma$ admitting the quotient (left) G -action. The orbits of this G action foliate \mathbf{H}_ρ ; this foliation is the quotient of the horizontal foliation of $SL(2, \mathbb{R}) \times H$ by Γ .

The suspension construction is quite general: if Γ is the covering group of a normal cover \hat{M} of a manifold M , and $\rho: \Gamma \rightarrow \text{Aut}(X)$ is a homomorphism into the automorphism group of an object X , then there is an associated fiber bundle $X_\rho \rightarrow M$ with fiber X obtained by taking the quotient of $\hat{M} \times X$ by the diagonal action of Γ , acting in the first factor by deck translations and in the second factor by ρ . If a group G acts Γ -equivariantly on \hat{M} , then it acts Γ -equivariantly on $\hat{M} \times X$, trivially in the second component. This induces a G -action on X_ρ . This suspension construction is also used to construct the Kontsevich-Zorich cocycle in the next section.

Returning to lattices in $SL(2, \mathbb{R})$, the following result gives a concrete criterion for establishing when the Lyapunov exponents of the A -action on \mathbf{H}_ρ with respect to Haar measure on $SL(2, \mathbb{R})/\Gamma$ are all distinct, a property often referred to as *simple Lyapunov spectrum*.

Theorem 2.1. [BF, Corollary 5.5] *Let $\Gamma < SL(2, \mathbb{R})$ be a lattice and $\rho: \Gamma \rightarrow SL(H)$ be a representation. If $\rho(\Gamma)$ is Zariski dense in $SL(H)$, then the Lyapunov spectrum for the A -action on \mathbf{H}_ρ , with respect to Haar measure on G/Γ , is simple.*

Theorem 2.1 is proved in the context of Kontsevich-Zorich cocycles in [EMat, Theorem 1], by combining the main result of [GM] and [GR] with [Fur67, Theorem 3], (whose proof is given in [Fu71]). It is straightforward to adapt the proof in [EMat] to give a proof of Theorem 2.1. The formulation in [BF] is more general than stated here and fits into a general theory (building on a long tradition initiated by Furstenberg) connecting Lyapunov spectrum with the theory of boundary actions.

2.1.2. Foliated geodesic and horocyclic flows. The suspension construction gives a natural way to construct a foliated bundle with hyperbolic leaves and compact leaf space. Namely, given a representation $\rho: \Gamma \rightarrow GL(H)$ of a lattice $\Gamma < SL(2, \mathbb{R})$, one constructs the projective bundle $\mathbb{P}(\check{\mathbf{H}}_\rho)$ over the hyperbolic surface \mathbb{H}/Γ by suspending the induced projective action of Γ on $\mathbb{P}(H)$, with Γ acting by isometries on the first factor of $\check{\mathbf{H}} = \mathbb{H} \times \mathbb{P}(H)$.

The bundle $\mathbb{P}(\check{\mathbf{H}}_\rho)$ does not carry a $SL(2, \mathbb{R})$ action, but it does carry a natural *foliation* (equivalently, a flat connection). In particular, the (trivial) horizontal foliation of $\check{\mathbf{H}}$ by hyperbolic planes descends under the Γ -quotient to a foliation \mathcal{F}_ρ of $\mathbb{P}(\check{\mathbf{H}}_\rho)$ by hyperbolic surfaces. The leaf space of this foliation is $\mathbb{P}(H)$, and the monodromy representation for \mathcal{F}_ρ is ρ . Since the leaves of \mathcal{F}_ρ carry a Riemannian structure, the

unit tangent bundle $T^1\mathcal{F}_\rho$ to the foliation \mathcal{F}_ρ carries a natural flow, the foliated geodesic flow. The construction of the foliated geodesic flow of a representation can be carried out quite generally, whenever $\Gamma = \pi_1(M)$, with M a Riemannian manifold (see, e.g. [BF, Example 5.2]).

Now recalling that the unit tangent bundle to \mathbb{H}/Γ is $SL(2, \mathbb{R})/\Gamma$, we note that in the present setting $T^1\mathcal{F}_\rho$ is naturally identified with the bundle $\mathbb{P}(\mathbf{H}_\rho)$, obtained by again suspending the action of Γ on $\mathbb{P}(H)$, this time with Γ acting by right multiplication on the first factor of $\mathbf{H} = SL(2, \mathbb{R}) \times \mathbb{P}(H)$. There is a natural projection $\mathbb{P}(\mathbf{H}_\rho)$ to $\mathbb{P}(\check{\mathbf{H}}_\rho)$ induced by the projection $SL(2, \mathbb{R})/\Gamma \rightarrow \mathbb{H}/\Gamma$. The bundle $\mathbb{P}(\mathbf{H}_\rho)$ admits an $SL(2, \mathbb{R})$ action; the action of A on $\mathbb{P}(\mathbf{H}_\rho)$ is the foliated geodesic flow, and the action of U_+ is the foliated (positive) horocycle flow.

Note that the bundle $\mathbb{P}(\mathbf{H}_\rho)$ is the projectivization of the bundle $\mathbf{H}_\rho \rightarrow SL(2, \mathbb{R})/\Gamma$, which is the suspension of the ρ -action on H defined above. This bundle also admits a $SL(2, \mathbb{R})$ action. Bonatti and Gomez-Mont proved the following result, which connects the dynamics of the foliated horocycle flow U_+ on $\mathbb{P}(\mathbf{H}_\rho)$ with the Lyapunov exponents of the A -action on \mathbf{H}_ρ .

Theorem 2.2 ([BG]). *Let $\rho: \Gamma \rightarrow GL(3, \mathbb{C})$ satisfy the integrability condition (1), which holds automatically if G/Γ is compact. If there is no $\rho(\Gamma)$ -invariant measure on \mathbb{CP}^2 , and the largest Lyapunov exponent for the A -action on \mathbf{H}_ρ has multiplicity 1, then there is a unique probability measure invariant under the foliated horocycle flow on $\mathbb{P}(\mathbf{H}_\rho)$ and projecting to Lebesgue/Haar measure on $SL(2, \mathbb{R})/\Gamma$. In particular, if $SL(2, \mathbb{R})/\Gamma$ is compact, then the foliated horocycle flow is uniquely ergodic.*

Our result generalizes the Bonatti–Gomez-Mont theorem to linear representations of Γ in arbitrary dimension:

Theorem 2.3. *Let $H = \mathbb{C}^k$ or \mathbb{R}^k , for $k \geq 2$, and let $\rho: \Gamma \rightarrow GL(H)$ satisfy the integrability condition (1), which holds automatically if G/Γ is compact. If the representation ρ is irreducible (i.e., if $\rho(\Gamma)$ has no proper invariant subspace of H), and the largest Lyapunov exponent for the A -action on \mathbf{H}_ρ has multiplicity 1, then there is a unique probability measure invariant under the horocycle flow on $\mathbb{P}(\mathbf{H})$ and projecting to Lebesgue/Haar measure on $SL(2, \mathbb{R})/\Gamma$. In particular, if $SL(2, \mathbb{R})/\Gamma$ is compact, then the foliated horocycle flow is uniquely ergodic.*

Proof. We first observe that the condition that ρ is an irreducible representation implies that the G -action on \mathbf{H}_ρ is irreducible. Suppose that there were a proper, nontrivial G -invariant ν -measurable subbundle of \mathbf{H} . Since the base $G/\Gamma = T^1\Sigma$ consists of a unique G -orbit, this bundle must be continuous (indeed analytic). This bundle projects to a continuous subbundle in $\mathbf{H}_{\Sigma, \rho}$ invariant under the monodromy of \mathcal{F}_ρ . Fixing a point $x \in X$ and considering the monodromy representation of $\pi_1(x, \Sigma)$ on the fiber of $\mathbf{H}_{\Sigma, \rho}$ over x , this monodromy-invariant subbundle of $\mathbf{H}_{\Sigma, \rho}$ restricts to a $\rho(\Gamma)$ -invariant subspace of H , which gives a contradiction.

Since \mathbf{E}^1 is one-dimensional, the fibers of $\mathbb{P}(\mathbf{E}^1)$ consist almost everywhere of a single point, and so $\mathbb{P}(\mathbf{E}^1)$ is the image of a measurable section $\sigma: SL(2, \mathbb{R})/\Gamma \rightarrow \mathbb{P}(\mathbf{H})$. Let $\mu^+ := \sigma_*\nu$ be the pushforward of Lebesgue/Haar measure ν on $SL(2, \mathbb{R})/\Gamma$ to $\mathbb{P}(\mathbf{H})$ under σ , which is invariant under the P action and ergodic under both A and U_+ actions.

Suppose that μ is another measure on $\mathbb{P}(\mathbf{E}^1)$ invariant under U_+ and projecting to ν . If μ is absolutely continuous with respect to μ^+ , then ergodicity of μ^+ with respect to U_+ implies that $\mu = \mu_+$. We may assume that μ is not absolutely continuous with respect to μ_+ . Taking the singular part of μ , saturating by the U_+ action, and renormalizing to be a probability, we obtain a measure μ_1 that is U_+ invariant and singular with respect to μ^+ . The projection of this measure to G/Γ is absolutely continuous with respect to ν and hence is equal to ν . The disintegration of μ_1 along fibers assigns measure 0 to $\mathbb{P}(\mathbf{E}^1)$.

Now push forward μ_1 along the action of g^{-t} and average, obtaining a limit measure ν that is P -invariant and projects to ν . The details of this construction are worked out in [BG, Sections 4 and 5].

Since we took averages over negative time, Oseledets' theorem implies that ν is not supported on $\mathbb{P}(\mathbf{E}^1)$. But this is a contradiction, by Theorem 1.3, since the G -action on H is irreducible. \square

In light of Theorem 2.1, we obtain a simple criterion for unique ergodicity of the foliated horocycle flow when Γ is cocompact.

Corollary 2.4. *Let $H = \mathbb{C}^k$ or \mathbb{R}^k , for $k \geq 2$, and let $\rho: \Gamma \rightarrow GL(H)$ satisfy the integrability condition (1). If the image $\rho(\Gamma)$ is Zariski dense in $GL(H)$, then there is a unique probability measure invariant under the horocycle flow on $\mathbb{P}(\mathbf{H})$ and projecting to Lebesgue/Haar measure on $SL(2, \mathbb{R})/\Gamma$.*

In particular, if $SL(2, \mathbb{R})/\Gamma$ is compact and $\rho: \Gamma \rightarrow GL(H)$ is any representation with Zariski dense image, then the foliated horocycle flow is uniquely ergodic.

In light of the results in [BaMa], we also obtain the following immediate corollary to Theorem 1.3, which we state for simplicity in the cocompact setting.

Corollary 2.5. *Let $\Gamma < SL(2, \mathbb{R})$ be a cocompact lattice, let $H = \mathbb{C}^k$ or \mathbb{R}^k , for $k \geq 2$, and let $\rho: \Gamma \rightarrow GL(H)$ be a representation such that $\rho(\Gamma)$ is Zariski dense.*

Consider the compact manifold $M = \mathbb{P}(\check{\mathbf{H}}_\rho) \rightarrow \mathbb{H}/\Gamma$, and let \mathcal{F}_ρ be the horizontal foliation by hyperbolic surfaces. Then there is a unique probability measure on M that is harmonic with respect to \mathcal{F}_ρ .

Matsumoto [M] recently showed that the same results do not hold for general foliations by hyperbolic surfaces: on a (3-dimensional) solvmanifold, he constructs a foliation with hyperbolic leaves whose foliated horocycle flow is not uniquely ergodic and admitting more than one harmonic measure.

2.2. The Kontsevich-Zorich cocycle.

In our second application of Theorem 1.3, we deduce the continuity of Lyapunov exponents of affine invariant measures.

Flat surfaces and strata. Suppose $g \geq 1$, and let $\beta = (\beta_1, \dots, \beta_m)$ be a partition of $2g - 2$. Let the Teichmüller space $\Omega(\beta)$ be the space of pairs (M, ω) where M is a Riemann surface of genus g and ω is a holomorphic 1-form on M (i.e. an Abelian differential) whose zeroes are labelled from 1 to m , and such that the zero labeled i has multiplicity β_i . In $\Omega(\beta)$, (M, ω) and (M', ω') are identified if there exists a holomorphic diffeomorphism $f : M \rightarrow M'$ isotopic to the identity, pulling back ω' to ω and preserving the labeling of the zeros. The space $\Omega(\beta)$ is smooth and contractible.

We also consider the moduli space $\mathcal{H}(\beta)$ whose definition is the same as that of $\Omega(\beta)$ except that in $\mathcal{H}(\beta)$, (M, ω) and (M', ω') are identified if there exists a holomorphic diffeomorphism $f : M \rightarrow M'$ (not necessarily isotopic to the identity) pulling back ω' to ω and preserving the labeling of the zeroes. We will refer to $\mathcal{H}(\beta)$ as a *stratum of Abelian differentials*.

Let S be a closed topological surface of genus g , and let Σ be a collection of m pairwise distinct points in S . The group $\text{Diff}^+(S, \Sigma)$ of diffeomorphisms of S fixing the points in Σ acts on $\Omega(\beta)$, with stabilizer containing the subgroup $\text{Diff}_0(S, \Sigma)$ of null-isotopic diffeomorphisms. Considering the quotient action, one obtains a properly discontinuous action of the mapping class group $\text{Mod}(S, \Sigma) := \text{Diff}^+(S, \Sigma)/\text{Diff}_0(S, \Sigma)$ on $\Omega(\beta)$, with quotient $\mathcal{H}(\beta)$. Thus, the space $\mathcal{H}(\beta)$ is an orbifold with universal cover $\Omega(\beta)$. For any β , there is a k -fold finite cover $\mathcal{H}(\beta)_k$ of $\mathcal{H}(\beta)$ in which these orbifold points disappear: $\mathcal{H}(\beta)_k$ is a smooth manifold covered by $\Omega(\beta)$.

For each $(M, \omega) \in \mathcal{H}(\beta)$, the form ω defines a canonical flat metric on M with conical singularities at the zeros of ω . Thus we refer to points of $\mathcal{H}(\beta)$ as *flat surfaces* or *translation surfaces*. For an introduction to this subject, see the survey [Zo2].

Affine measures and manifolds. There is a well-studied action of $G = SL(2, \mathbb{R})$ on $\Omega(\beta)$; in this action, $g \in G$ acts linearly on translation structures via post-composition in charts. This action preserves the subset $\Omega_1(\beta) \subset \Omega(\beta)$ of marked surfaces of (flat) area 1. This G -action commutes with the action of $\text{Mod}(S, \Sigma)$ and so descends to an action on $\mathcal{H}(\beta)$ preserving the space $\mathcal{H}_1(\beta)$ of flat surfaces of area 1.

An *affine invariant manifold* is a closed subset of $\mathcal{H}_1(\beta)$ that is invariant under this G action and which in *period coordinates* (see [Zo2, Chapter 3]) looks like an affine subspace. Each affine invariant manifold \mathcal{N} is the support of an ergodic $SL(2, \mathbb{R})$ -invariant probability measure $\nu_{\mathcal{N}}$. Locally, in period coordinates, this measure is (up to normalization) the restriction of Lebesgue measure to the subspace \mathcal{N} , see [EM] for the precise definitions. It is proved in [EMM] that the closure of any $SL(2, \mathbb{R})$ orbit is an affine invariant manifold; this is analogous to one of Ratner's theorems in the theory of unipotent flows. (In genus 2, this result was previously proved by McMullen [McM]).

To state our result, we will need the following:

Theorem 2.6 ([[EMM](#), Theorem 2.3]). *Let \mathcal{N}_n be a sequence of affine manifolds in $\mathcal{H}(\beta)$, and suppose $\nu_{\mathcal{N}_n} \rightarrow \nu$. Then ν is a probability measure. Furthermore, ν is the affine measure $\nu_{\mathcal{N}}$, where \mathcal{N} is the smallest submanifold with the following property: there exists some $n_0 \in \mathbb{N}$ such that $\mathcal{N}_n \subset \mathcal{N}$ for all $n > n_0$.*

In particular, the space of ergodic P -invariant probability measures on $\mathcal{H}_1(\beta)$ is compact in the weak-star topology.

Since $\mathcal{H}_1(\beta)$ is not compact, the weak star topology is defined using compactly supported functions. In particular, $\nu_{\mathcal{N}_n} \rightarrow \nu$ if and only if for any compactly supported continuous function ϕ on $\mathcal{H}_1(\beta)$, we have that $\int \phi d\nu_{\mathcal{N}_n} \rightarrow \int \phi d\nu$.

Remark 2.7. In the setting of unipotent flows, Theorem [2.6](#) is due to Mozes and Shah [[MS](#)].

The Kontsevich-Zorich cocycle. The *Hodge bundle* \mathbf{H} over $\mathcal{H}(\beta)$ is constructed in a similar fashion to the suspension construction in Subsection [2.1](#). We start with the bundle $\hat{\mathbf{H}}$ over $\Omega(\beta)$ whose fiber over (M, ω) is the cohomology group $H^1(M, \mathbb{R})$ (viewed as a $2g$ -dimensional real vector space). As $\Omega(\beta)$ is contractible, this bundle is trivial, diffeomorphic to $\Omega(\beta) \times \mathbb{R}^{2g}$. We then consider the natural (right) action of the group $\Gamma = \text{Mod}(M, \Sigma)$ on $\hat{\mathbf{H}}$, where $\gamma \in \Gamma$ acts by pullback in the fibers, and we set $\mathbf{H} := \hat{\mathbf{H}}/\Gamma$, which is a bundle over $\mathcal{H}(\beta)$. The Hodge bundle fails to be a vector bundle over the orbifold points of $\mathcal{H}(\beta)$, but passing to a finite branched cover, the corresponding Hodge bundle over $\mathcal{H}(\beta)_k$ is a smooth vector bundle, and the G -action lifts to this bundle. As the results below concern affine measures, which behave well under finite branched covers, *we will henceforth assume that \mathbf{H} is a vector bundle over $\mathcal{H}(\beta)$.*

There is a natural (left) G -action on $\hat{\mathbf{H}}$ that is the standard G action on $\Omega(\beta)$ and trivial on the fibers. This descends as in Subsection [2.1](#) to a G action on \mathbf{H} . This G action is often described by measurably trivializing \mathbf{H} and describing elements of this action using elements of the symplectic group $Sp(2g, \mathbb{Z})$. To be precise, if we choose a fundamental domain in $\Omega(\beta)$ for the action of the mapping class group Γ , then we have the cocycle $\tilde{\alpha} : SL(2, \mathbb{R}) \times \mathcal{H}_1(\beta) \rightarrow \Gamma$ where for x in the fundamental domain, $\tilde{\alpha}(g, x)$ is the element of Γ needed to return the point gx to the fundamental domain. Then, we define the Kontsevich-Zorich cocycle $\alpha(g, x)$ by

$$\alpha(g, x) = \rho(\tilde{\alpha}(g, x)),$$

where $\rho : \Gamma \rightarrow Sp(2g, \mathbb{Z})$ is the homomorphism given by the action on cohomology. The Kontsevich-Zorich cocycle can be interpreted as the monodromy of the Gauss-Manin connection restricted to the orbit of $SL(2, \mathbb{R})$, see e.g. [[Zo2](#), page 64].

Henceforth we will conflate action and cocycle and simply refer to this G -action on \mathbf{H} as the Kontsevich-Zorich cocycle. We further restrict (i.e. pull back) the bundle \mathbf{H} to the space $\mathcal{H}_1(\beta)$ of flat surfaces of area 1. As the G -action on $\mathcal{H}(\beta)$ preserves

$\mathcal{H}_1(\beta)$, we thus have a restricted G action on the bundle $\mathbf{H} \rightarrow \mathcal{H}_1(\beta)$. The discussion that follows concerns this action (cocycle) on this bundle.

There is a way to choose a Finsler structure on the Hodge bundle \mathbf{H} over $\mathcal{H}_1(\beta)$ so that the Kontsevich-Zorich cocycle satisfies the integrability condition (1) with respect to *any* affine measure ν on any such \mathcal{F} . Moreover this integrability is uniform: there exists C depending only on the genus such that for any $x \in \mathcal{H}_1(\beta)$, any $t \in [-1, 1]$ and any $\mathbf{v} \in \mathbf{H}(x)$,

$$C^{-1} \|\mathbf{v}\|_x \leq \|g_*^t \mathbf{v}\|_{g^t x} \leq C \|\mathbf{v}\|_x.$$

(see [AGY, Subsection 2.2.2] or [EMM, Section 7.2]). In particular,

$$(3) \quad \int_Z \sup_{t \in [-1, 1]} (\log \|g_*^t\|_x) d\nu(x) < \epsilon.$$

The Lyapunov exponents of the Kontsevich-Zorich cocycle are defined with respect to this Finsler structure; they exist and are constant almost everywhere with respect to any ergodic affine measure. These exponents appear in other areas of dynamics, for example in the study of deviations of ergodic averages of interval exchange transformations [Zo] [Fo1], [Bu] and in the study of the diffusion rate of the periodic wind-tree model [DHL].

The Lyapunov spectrum of the Kontsevich-Zorich cocycle has been studied extensively. Building on numerical experiments by Zorich, it was conjectured by Kontsevich in [K] that for the Masur-Veech measures, the Lyapunov spectrum is simple: this Kontsevich-Zorich Conjecture has been proved partially by Forni in [Fo1] and then fully by Avila and Viana [AV]. For other affine measures ν the situation is more complicated (and in particular the spectrum need not be simple). Also the sum of the Lyapunov exponents (and in some cases the Lyapunov exponents themselves) can be computed explicitly. Recent papers in which the Lyapunov spectrum of the Kontsevich-Zorich cocycle plays a major role include [Au1, Au2, Au3, AEZ, Ba1, Ba2, BoMö, CM1, CM2, DM, DZ, DD, EKMZ, EKZ, EKZ2, EMat, Fi1, Fi2, Fi3, FFM, Fo2, F3, FM, FMZ1, FMZ2, FMZ3, KZ1, GH1, GH2, Mat, MY, MYZ, MMY, MW, Mö, P1, P2, P3, T, Yu, YZ, W].

The following theorem answers a question asked in [MMY].

Theorem 2.8. *Let \mathcal{N}_n be a sequence of affine manifolds, and suppose the affine measures $\nu_{\mathcal{N}_n}$ converge to the (affine) measure ν (as in Theorem 2.6). Then the Lyapunov exponents of $\nu_{\mathcal{N}_m}$ (with respect to g^t) converge to the Lyapunov exponents of ν .*

The proof of Theorem 2.8 depends on Theorem 1.3 and the following theorem of S. Filip:

Theorem 2.9 ([Fil]). *Let α denote (some exterior power of) the Kontsevich-Zorich cocycle restricted to an affine invariant submanifold \mathcal{N} . Let $\nu_{\mathcal{N}}$ be the affine measure whose support is \mathcal{N} , and suppose α has a $\nu_{\mathcal{N}}$ -measurable almost-invariant subbundle*

\mathbf{W} (see Definition 1.1). Then \mathbf{W} agrees ν_N -almost everywhere with a continuous one.

In fact it is proved in [Fi1] that the dependence of any such $\mathbf{W}(x)$ on x is polynomial in the period coordinates.

Simplicity of Lyapunov Spectrum of Teichmüller curves. We recall that a *Teichmüller curve* is a closed $SL(2, \mathbb{R})$ orbit on $\mathcal{H}_1(\beta)$. Teichmüller curves (which are submanifolds of real dimension 3) are the smallest possible affine manifolds; any other type of affine manifold has dimension greater than 3.

By the Kontsevich-Zorich classification [KZ2], the stratum $\mathcal{H}(4)$ (in genus 3) has two connected components $\mathcal{H}(4)^{hyp}$ and $\mathcal{H}(4)^{odd}$. The connected component $\mathcal{H}(4)^{hyp}$ consists entirely of hyperelliptic surfaces.

As a consequence of some recent results, we obtain the following:

Theorem 2.10. *All but finitely many Teichmüller curves in $\mathcal{H}(4)$ have simple Lyapunov spectrum.*

Theorem 2.10 was shown in [MMY] to follow from a conjecture of Delecroix and Lelièvre (stated in [MMY]). Our proof below is unconditional; however it is much less explicit and is completely ineffective. It also depends on the very recent results of Filip [Fi1], Nguyen-Wright [NW], and [ANW].

Proof of Theorem 2.10. Suppose there exist infinitely many Teichmüller curves $\mathcal{N}_n \subset \mathcal{H}_1(4)$ with multiplicities in the Lyapunov spectrum. By Theorem 2.6, the \mathcal{N}_n have to converge to an affine manifold \mathcal{N} (in the sense that the affine measures $\nu_{\mathcal{N}_n}$ will converge to the affine measure $\nu_{\mathcal{N}}$). Furthermore, \mathcal{N} cannot be a Teichmüller curve (since by Theorem 2.6, \mathcal{N} must contain all the \mathcal{N}_n for n sufficiently large, and thus $\dim \mathcal{N} > 3$). By the main theorems of [NW] and [ANW] the only proper affine submanifolds of $\mathcal{H}_1(4)$ that are not Teichmüller curves are the two connected components $\mathcal{H}_1(4)^{hyp}$ and $\mathcal{H}_1(4)^{odd}$ and the Prym locus \mathcal{P} of $\mathcal{H}_1(4)^{odd}$. Thus \mathcal{N} must be either $\mathcal{H}_1(4)^{hyp}$ or $\mathcal{H}_1(4)^{odd}$ or \mathcal{P} . The Lyapunov spectrum of both $\mathcal{H}_1(4)^{hyp}$ and $\mathcal{H}_1(4)^{odd}$ is simple by [AV], and the Lyapunov spectrum of \mathcal{P} is simple by [MMY, §6.6]. This contradicts Theorem 2.8. \square

Remark 2.11. We expect that the applicability of the proof of Theorem 2.10 to increase as our knowledge of the classification of affine invariant manifolds increases.

Remark 2.12. It remains a challenge to find an approach to the Delecroix-Lelièvre conjecture itself which would leverage some equidistribution results.

3. CONSTRUCTION OF AN INVARIANT SUBSPACE

We write $\mathcal{B} := \{u_+^s : s \in [-1, 1]\}$ for the “unit ball” in U_+ , and \mathcal{B}_t for $g^t \mathcal{B} g^{-t} = \{u_+^s : s \in [-e^{2t}, e^{2t}]\}$. We also write

$$\mathcal{B}[x] = \{ux : u \in \mathcal{B}\} \subset X, \quad \mathcal{B}_t[x] = \{ux : u \in \mathcal{B}_t\} \subset X.$$

Similarly, we use the notation

$$U_+[x] = \{ux : u \in U_+\} \subset X.$$

Recall that ν denotes a G -invariant measure on X that is ergodic for the action of $A = \{g^t\}$. In what follows, when we write “almost all x ” we mean “all x outside a set of ν -measure 0.”

3.1. The forward and backward flags. We consider the forward and backward flags for the action of g^t on fibers, which define the Oseledets decomposition; let

$$\{0\} = \mathbf{E}_{\geq n+1}(x) \subset \mathbf{E}_{\geq n}(x) \subset \cdots \subset \mathbf{E}_{\geq 1}(x) = \mathbf{H}(x)$$

be the forward flag, and let

$$\{0\} = \mathbf{E}_{\leq 0}(x) \subset \mathbf{E}_{\leq 1}(x) \subset \cdots \subset \mathbf{E}_{\leq n}(x) = \mathbf{H}(x)$$

be the backward flag. This means that for almost all x and for $\mathbf{v} \in \mathbf{E}_{\geq i}(x)$ such that $\mathbf{v} \notin \mathbf{E}_{\geq i+1}(x)$,

$$(4) \quad \lim_{t \rightarrow +\infty} \frac{1}{t} \log \frac{\|g_*^t \mathbf{v}\|}{\|\mathbf{v}\|} = \lambda_i,$$

and for $\mathbf{v} \in \mathbf{E}_{\leq i}(x)$ such that $\mathbf{v} \notin \mathbf{E}_{\leq i-1}(x)$,

$$(5) \quad \lim_{t \rightarrow -\infty} \frac{1}{t} \log \frac{\|g_*^t \mathbf{v}\|}{\|\mathbf{v}\|} = \lambda_i.$$

By e.g. [GM, Lemma 1.5], for a.e. x and every i , we have

$$(6) \quad \mathbf{H}(x) = \mathbf{E}_{\leq i}(x) \oplus \mathbf{E}_{\geq i+1}(x).$$

The bundles $\mathbf{E}_1, \dots, \mathbf{E}_n$ in the Oseledets decomposition are then defined by the almost everywhere transverse intersections $\mathbf{E}_i := \mathbf{E}_{\leq i} \cap \mathbf{E}_{\geq i}$. Note that $\mathbf{E}_{\leq 1} = \mathbf{E}_1$, $\mathbf{E}_{\geq n} = \mathbf{E}_n$ and in general, $\mathbf{E}_{\leq j} = \mathbf{E}_1 \oplus \mathbf{E}_2 \oplus \cdots \oplus \mathbf{E}_j$ and $\mathbf{E}_{\geq j} = \mathbf{E}_j \oplus \mathbf{E}_{j+1} \oplus \cdots \oplus \mathbf{E}_n$ almost everywhere.

The bundles $\mathbf{E}_{\geq i}$ and $\mathbf{E}_{\leq i}$ are evidently A -equivariant; for almost all $x \in X$ and every $t \in \mathbb{R}$, we have

$$(7) \quad g_*^t(\mathbf{E}_{\leq j}(x)) = \mathbf{E}_{\leq j}(g^t x), \quad g_*^t(\mathbf{E}_{\geq j}(x)) = \mathbf{E}_{\geq j}(g^t x).$$

Remark 3.1. The bundles have some additional invariance properties; it follows from (4) and (5) that for almost all $x \in X$, $u \in U_+$ and $\bar{u} \in U_-$, we have

$$(8) \quad u_*(\mathbf{E}_{\leq j}(x)) = \mathbf{E}_{\leq j}(ux), \quad \bar{u}_*(\mathbf{E}_{\geq j}(x)) = \mathbf{E}_{\geq j}(\bar{u}x).$$

Proof. Equation (4) means that the bundle $\mathbf{E}_{\geq j}(x)$ is characterized by the fact that the forward rate of expansion of vectors in $\mathbf{E}_{\geq j}(x)$ under g_*^t is no more than λ_j . For any $s, t \in \mathbb{R}$ the vectors $\mathbf{v} \in H_x$, for some $x \in X$, and $\mathbf{w} = (u_-^s)_*(\mathbf{v})$ satisfy

$$g_*^t(\mathbf{w}) = (u_-^{\exp(-t) \cdot s})_*(g_*^t(\mathbf{v})).$$

Since the measure ν is G -invariant, almost every point x is recurrent for g^t . Thus the sequence of linear maps $(u_-^{\exp(-t) \cdot s})_*$ tend to the identity for large $t > 0$ for which $g_*^t x$ is close to x . As a consequence, if the rates of expansion of the vectors \mathbf{v} and \mathbf{w} are defined, they are the same. This proves the invariance of the bundle $\mathbf{E}_{\geq j}(x)$ under U_- . \square

Remark 3.2. If for some $j \geq 2$, the bundle $\mathbf{E}_{\geq j}$ were also invariant under almost every $u \in U_+$, then because G is generated by A, U_+ and U_- , the bundle would be invariant under G , violating the irreducibility hypothesis on the action of G .

As remarked before, the bundles \mathbf{E}_i and therefore $\mathbf{E}_{\geq i}$ and $\mathbf{E}_{\leq i}$ are measurable. This implies that, for any Borel probability measure for which these bundles are almost everywhere defined, they are continuous when restriction to a set whose measure is arbitrarily close to 1. This implies the following statement.

Lemma 3.3. *Let $\mathbf{E} \subset \mathbf{H}$ be a measurable subbundle of \mathbf{H} . For any compact subset $K \subset X$ and any probability measure θ on K with respect to which the bundle is defined θ -almost everywhere, there is a set $Y \subset X$ of full θ -measure such that any $y \in Y$ admits a subset $G_y \subset Y$ with the following properties:*

- $\theta(G_y \cap U) > 0$, for any neighborhood U of y ;
- the bundle \mathbf{E} is continuous on G_y .

The map from \mathbb{R} to U_+ defined by $t \mapsto u_t^+$ defines a measure on U_+ by the pushforward of normalized Lebesgue measure on \mathbb{R} . This measure coincides with the Haar measure on U_+ . We refer to this measure as Lebesgue measure on U_+ and denote it by $|\cdot|$. The previous lemma, U_+ -invariance of ν and Fubini's theorem imply:

Corollary 3.4. *In our setting there is a subset $Y \subset X$ of full ν -measure such that for all $y \in Y$:*

- the bundles $\mathbf{E}_{\geq j}(uy)$ are well-defined for Lebesgue-almost every $u \in \mathcal{B}$;
- there exists a measurable set $\mathcal{G}_y \subset \mathcal{B}$ such that $|\mathcal{G}_y \cap U| > 0$, for any neighborhood U of the identity in \mathcal{B} , and such that $u \mapsto \mathbf{E}_{\geq j}(uy)$ is continuous on \mathcal{G}_y .

3.2. The forward flag and the unstable horocycle flow: defining the inert flag. Our strategy for proving Theorem 1.3 is to show that if the measure $\hat{\nu}$ is not supported in \mathbf{E}_1 , then there is a nontrivial subbundle of some $\mathbf{E}_{\geq j}$ that is G -invariant, obtaining a contradiction.

To this end, we define the *inert flag*

$$(9) \quad \{0\} = \mathbf{F}_{\geq n+1}(x) \subseteq \mathbf{F}_{\geq n}(x) \subseteq \mathbf{F}_{\geq n-1}(x) \subseteq \dots \mathbf{F}_{\geq 2}(x) \subseteq \mathbf{F}_{\geq 1}(x) = \mathbf{H}(x).$$

where $\mathbf{F}_{\geq j}(x)$, for $j = 1, \dots, m$ are defined by

$$(10) \quad \mathbf{F}_{\geq j}(x) = \{\mathbf{v} \in \mathbf{H}(x) : \text{for almost all } u \in \mathcal{B}, u_* \mathbf{v} \in \mathbf{E}_{\geq j}(ux)\}.$$

In other words, $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$ if and only if, for almost all $u \in \mathcal{B}$ we have

$$(11) \quad \limsup_{t \rightarrow \infty} \frac{1}{t} \log \|g_*^t u_* \mathbf{v}\| \leq \lambda_j.$$

From the definition it follows that the $\mathbf{F}_{\geq j}(x)$ are vector subspaces of $\mathbf{H}(x)$ that decrease with j , and thus they indeed form a flag.

Lemma 3.5. *For almost every $x \in X$, the following hold:*

- (a) $\mathbf{F}_{\geq j}(x) \subset \mathbf{E}_{\geq j}(x)$;
- (b) $\mathbf{F}_{\geq j}(x) = \{\mathbf{v} \in \mathbf{H}(x) : \text{for almost all } u \in U_+, u_* \mathbf{v} \in \mathbf{E}_{\geq j}(ux)\}$;
- (c) $\mathbf{F}_{\geq j}(x)$ is A -equivariant; i.e., $g_*^t(\mathbf{F}_{\geq j}(x)) = \mathbf{F}_{\geq j}(g^t x)$, for all $t \in \mathbb{R}$; and
- (d) for almost all $u \in U_+$, we have $u_*(\mathbf{F}_{\geq j}(x)) = \mathbf{F}_{\geq j}(ux)$.




FIGURE 1. Invariance properties of $\mathbf{E}_{\leq j-1}$, $\mathbf{E}_{\geq j}$, and $\mathbf{F}_{\geq j}$ under almost every $u \in U_+$.

Proof.

(a) Consider $x \in X$ such that (6) holds both for x and for almost every point in $\mathcal{B}[x]$ and such that x belongs to the set Y described in Corollary 3.4. Since (6) holds almost everywhere, Fubini's theorem implies that the set of such x has full measure in X . Pick such an x , and let $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$. Then we can write $\mathbf{v} = \mathbf{v}_{\geq j} + \mathbf{v}_{\leq j-1}$, where $\mathbf{v}_{\geq j} \in \mathbf{E}_{\geq j}(x)$, and $\mathbf{v}_{\leq j-1} \in \mathbf{E}_{\leq j-1}(x)$. We have $u_* \mathbf{v} = u_* \mathbf{v}_{\geq j} + u_* \mathbf{v}_{\leq j-1}$.

If $\mathbf{v}_{\leq j-1} \neq 0$, then (8) implies that for almost every $u \in U_+$, we have $0 \neq u_* \mathbf{v}_{\leq j-1} \in \mathbf{E}_{\leq j-1}(ux)$.

By our choice of x , we have $\mathbf{E}_{\leq j-1}(ux) \cap \mathbf{E}_{\geq j}(ux) = \emptyset$ for almost every $u \in U_+$. Now suppose $u \in \mathcal{G}_x$, where $\mathcal{G}_x \subset \mathcal{B}$ given by Corollary 3.4. We decompose $u_* \mathbf{v}_{\geq j}$ as the sum of a vector in $\mathbf{E}_{\geq j}(ux)$ and a vector in $\mathbf{E}_{\leq j-1}(ux)$; since both of these spaces vary continuously in $u \in \mathcal{G}_x$, if $u \in \mathcal{G}_x$ is sufficiently close to the identity in U_+ , then the component of $u_* \mathbf{v}_{\geq j}$ in $\mathbf{E}_{\leq j-1}(ux)$ is arbitrarily small, and in particular smaller than $u_* \mathbf{v}_{\leq j-1}$.

This shows that $u_*\mathbf{v} \notin \mathbf{E}_{\geq j}(ux)$ for $u \in \mathcal{G}_x$ sufficiently close to the identity. But Corollary 3.4 implies that the set of such u has positive Lebesgue measure in \mathcal{B} ; this contradicts the assumption that $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$. Therefore $\mathbf{v}_{\leq j-1} = 0$, and $\mathbf{v} \in \mathbf{E}_{\geq j}(x)$, proving (a).

(c) Note that for $t \geq 0$, $g^t\mathcal{B}[x] = \mathcal{B}_t[g^tx] \supset \mathcal{B}[g^tx]$. It follows that

$$(12) \quad g_*^t(\mathbf{F}_{\geq j}(x)) \subset \mathbf{F}_{\geq j}(g^tx).$$

In particular, the dimension of $\mathbf{F}_{\geq j}(x)$ is a measurable function, with values in $\{0, \dots, \dim(H)\}$, and which is nondecreasing along g^t -orbits. Thus the sets $\{x \in X : \dim(\mathbf{F}_{\geq j}(x)) \geq i\}$ are positively invariant by g^t , $t \geq 0$. As the measure ν is assumed to be A -ergodic, it follows that there exists $i \in \{0, \dots, \dim(H)\}$ such that, for ν -almost every x , the dimension of $\mathbf{F}_{\geq j}(x)$ is equal to i . For such a ν -generic x , the inclusion in (12) becomes an equality:

$$g_*^t(\mathbf{F}_{\geq j}(x)) = \mathbf{F}_{\geq j}(g^tx),$$

proving item (c).

(b) Consider now an x such that (c) holds, a vector $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$, and a large $t > 0$. Now $g_*^{-t}\mathbf{v}$ belongs to $\mathbf{F}_{\geq j}(g^{-t}x)$. Since $g^t\mathcal{B}[g^{-t}x] = \mathcal{B}_t[x]$, we deduce that for every $t > 0$,

$$\mathbf{F}_{\geq j}(x) = \{\mathbf{v} \in \mathbf{H}(x) : \text{for almost all } u \in \mathcal{B}_t(x), u_*\mathbf{v} \in \mathbf{E}_{\geq j}(ux)\}.$$

As $t > 0$ was arbitrary, this establishes (b).

(d) Consider x satisfying (a) and (b), and let $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$. Then for almost every $u \in U_+$, we have $u_*\mathbf{v} \in \mathbf{E}_{\geq j}(ux)$. Fix such a u , and suppose that $u_*\mathbf{v} \notin \mathbf{F}_{\geq j}(ux)$. This implies there exists a positive measure set of $u' \in U_+$ such that $u'u_*\mathbf{v} \notin \mathbf{E}_{\geq j}(u'ux)$. But this violates the assumption that $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$. Thus for almost all $u \in U_+$, we have $u_*(\mathbf{F}_{\geq j}(x)) \subset \mathbf{F}_{\geq j}(ux)$. The reverse inclusion is proved by applying u^{-1} . This gives (d). \square

3.3. The inert flag is G -invariant.

Lemma 3.6. *For almost all $\bar{u} \in U_-$ and almost all $x \in X$, $\bar{u}_*(\mathbf{F}_{\geq j}(x)) = \mathbf{F}_{\geq j}(\bar{u}x)$.*

Proof. It is enough to show that (outside a set of measure 0), for all $\mathbf{v} \in \mathbf{F}_{\geq j}(x)$, we have $\bar{u}_*\mathbf{v} \in \mathbf{F}_{\geq j}(\bar{u}x)$. Hence, in view of the definition (10) of $\mathbf{F}_{\geq j}$, it is enough to show that for almost all $u \in U_+$ and $\bar{u} \in U_-$,

$$(13) \quad (u\bar{u})_*\mathbf{v} \in \mathbf{E}_{\geq j}(u\bar{u}x).$$

We may write $u\bar{u} = \bar{u}'u'a$ where $\bar{u}' \in U_-$, $u' \in U_+$, $a \in A$. Let

$$\mathbf{w} = (u'a)_*\mathbf{v}.$$

Then, by Lemma 3.5 (c) and (d), $\mathbf{w} \in \mathbf{F}_{\geq j}(u'ax)$. By Lemma 3.5 (a), $\mathbf{F}_{\geq j}(u'ax) \subset \mathbf{E}_{\geq j}(u'ax)$. Therefore

$$(u\bar{u})_*\mathbf{v} = (\bar{u}'u'a)_*\mathbf{v} = \bar{u}'_*\mathbf{w} \in \bar{u}'_*(\mathbf{F}_{\geq j}(u'ax)) \subset \bar{u}'_*(\mathbf{E}_{\geq j}(u'ax)).$$

But, by (8),

$$\bar{u}'_*(\mathbf{E}_{\geq j}(u'ax)) = \mathbf{E}_{\geq j}(\bar{u}'u'ax) = \mathbf{E}_{\geq j}(u\bar{u}x).$$

Thus, (13) holds. \square

In view of Lemma 3.5 (c), (d) and Lemma 3.6, we have proved the following:

Proposition 3.7. *The subspaces $\mathbf{F}_{\geq j}(x)$ are equivariant under the action of $SL(2, \mathbb{R})$.*

3.4. Relaxing the definition of the components of the inert flag.

Lemma 3.8. *There exists a subset $\Omega \subset X$ with $\nu(\Omega) = 1$ with the following property. For $j \in \{0, \dots, n\}$, $x \in \Omega$, and $\mathbf{v} \in \mathbf{H}(x)$, let*

$$(14) \quad Q_j(\mathbf{v}) = \{u \in \mathcal{B} : u_*\mathbf{v} \in \mathbf{E}_{\geq j}(ux)\}.$$

Then either $|Q_j(\mathbf{v})| = 0$, or $|Q_j(\mathbf{v})| = |\mathcal{B}|$ (and thus $\mathbf{v} \in \mathbf{F}_j(x)$).

Proof. Fix $j \in \{0, \dots, n\}$. For a subspace $\mathbf{V} \subset \mathbf{H}(x)$, let

$$Q_j(\mathbf{V}) = \{u \in \mathcal{B} : u_*\mathbf{V} \subset \mathbf{E}_{\geq j}(ux)\}.$$

Let d be the maximal number such that there exists $Y \subset X$ with $\nu(Y) > 0$ such that for $x \in Y$ there exists a subspace $\mathbf{V} \subset \mathbf{H}(x)$ of dimension d with $|Q_j(\mathbf{V})| > 0$. For a fixed $x \in Y$, let $\mathcal{W}(x)$ denote the set of subspaces \mathbf{V} of dimension d for which $|Q_j(\mathbf{V})| > 0$. Then, by the maximality of d , if \mathbf{V} and \mathbf{V}' are distinct elements of $\mathcal{W}(x)$ then $Q_j(\mathbf{V}) \cap Q_j(\mathbf{V}') = \emptyset$. Fix a measurable collection of subspaces $\mathbf{V}_x \in \mathcal{W}(x)$, for $x \in Y$, such that $|Q_j(\mathbf{V}_x)|$ is maximal (among elements of $\mathcal{W}(x)$).

For $x \in Y$, $u \in \mathcal{B}$ and $t > 0$, consider the set

$$D(u, x) := \{z \in \mathcal{B}_{-t} : zu \in Q_j(\mathbf{V}_x)\}.$$

Let $\epsilon > 0$ be arbitrary, and suppose $x \in Y$. The Vitali covering lemma implies that there exists $t_0 > 0$ and a subset $Q(\mathbf{V}_x)^* \subset Q(\mathbf{V}_x) \subset \mathcal{B}$ such that

$$(15) \quad |Q(\mathbf{V}_x) \setminus Q(\mathbf{V}_x)^*| < \epsilon |Q(\mathbf{V}_x)|,$$

and for all $u \in Q(\mathbf{V}_x)^*$ and all $t > t_0$, we have

$$(16) \quad |D(u, x)| \geq (1 - \epsilon) |\mathcal{B}_{-t}|.$$

(In other words, $Q(\mathbf{V}_x)^*$ are “points of $(1 - \epsilon)$ density” for $Q(\mathbf{V}_x)$, relative to the collection of (small) balls $\{\mathcal{B}_{-t}u : u \in Q(\mathbf{V}_x), t > t_0\}$.) Let

$$Y^* = \{ux : x \in Y, u \in Q(\mathbf{V}_x)^*\}.$$

Then, since $\nu(Y) > 0$, the U_+ -invariance of ν and (15) imply that $\nu(Y^*) > 0$. Let $\Omega = \{x \in X : g^{-t}x \in Y^* \text{ infinitely often}\}$. Poincaré recurrence implies that $\nu(\Omega) = 1$. Suppose $x \in \Omega$. We can choose $t > t_0$ such that $g^{-t}x \in Y^*$. Note that

$$(17) \quad \mathcal{B}[x] = g^t \mathcal{B}_{-t} [g^{-t}x].$$

Let $x' = g^{-t}x \in Y^*$, and let $\mathbf{V}_{t,x} = (g^t)_* \mathbf{V}_{x'}$. Then in view of (16) and (17), we have

$$(18) \quad |Q(\mathbf{V}_{t,x})| \geq (1 - \epsilon) |\mathcal{B}|.$$

By the maximality of d (and assuming $\epsilon < 1/2$), $\mathbf{V}_{t,x}$ does not depend on t . Hence, for every $x \in \Omega$, there exists $\mathbf{V} \subset \mathbf{H}(x)$ such that $\dim \mathbf{V} = d$ and $|Q(\mathbf{V})| \geq (1 - \epsilon)|\mathcal{B}|$. Since $\epsilon > 0$ is arbitrary, for each $x \in \Omega$, there exists $\mathbf{V} \subset \mathbf{H}(x)$ with $\dim \mathbf{V} = d$, and $|Q(\mathbf{V})| = |\mathcal{B}|$. Now the maximality of d implies that if $\mathbf{v} \notin \mathbf{V}$ then $|Q(\mathbf{v})| = 0$. \square

4. PROOF OF THEOREM 1.3

Now suppose that there is more than one Lyapunov exponent on \mathbf{H} (so that $\lambda_2 < \lambda_1$). Then for almost all x , $\mathbf{F}_{\geq 2}(x)$ is a proper, nontrivial subspace of $\mathbf{F}_{\geq 1}(x) = \mathbf{H}(x)$.

Suppose also that the cocycle is irreducible with respect to the measure ν (so there are no non-trivial proper equivariant ν -measurable subbundles). Then it follows from Proposition 3.7 that for all $j \geq 2$ and ν -almost all x ,

$$(19) \quad \mathbf{F}_{\geq j}(x) = \{0\}.$$

Let $\hat{\nu} \in \mathcal{H}_P(\nu)$ be any P -invariant measure on the total space of the projectivized bundle $\mathbb{P}^1(\mathbf{H})$ that projects to ν . We may disintegrate $\hat{\nu}$ as follows:

$$d\hat{\nu}([\mathbf{v}]) = d\nu(\pi(\mathbf{v})) d\eta_{\pi(\mathbf{v})}([\mathbf{v}]).$$

Lemma 3.8 and (19) imply that for almost every $x \in X$ and every $\mathbf{v} \in \mathbf{H}(x) \setminus \{0\}$:

$$(20) \quad |\{u \in \mathcal{B} : u_*\mathbf{v} \in \mathbf{E}_{\geq 2}(ux)\}| = 0.$$

In other words, for almost every x and every nonzero $\mathbf{v} \in \mathbf{H}(x)$, we have $u_*\mathbf{v} \notin \mathbf{E}_{\geq 2}(ux)$, for almost every $u \in \mathcal{B}$.

The U_+ -invariance of $\hat{\nu}$ then implies the following claim.

Claim 4.1. *For ν -almost all x , the disintegration η_x of $\hat{\nu}$ is supported in the set $\mathbb{P}(\mathbf{H}(x) \setminus \mathbf{E}_{\geq 2}(x))$.*

Proof. First notice that almost all P -invariant measures in the P -ergodic disintegration of $\hat{\nu}$ project on ν . Therefore it suffices to prove the claim under the extra assumption that $\hat{\nu}$ is ergodic for P .

We argue by contradiction, assuming that there is a measurable set Z contained in $\mathbb{P}(\mathbf{E}_{\geq 2})$ and a positive ν -measure set $Y \subset X$ such that $\eta_y(Z(y)) > 0$ for $y \in Y$, where $Z(y) = Z \cap \mathbb{P}(\mathbf{H}(y))$. In other words $\hat{\nu}(Z) > 0$. As the bundle $\mathbf{E}_{\geq 2}$ and the disintegrations η_y are equivariant under the action of $\{g^t\}$, we may assume that Y and Z are invariant under the action of g^t .

Since P is amenable, and $\hat{\nu}$ is ergodic with respect to P , the Mean Ergodic Theorem for amenable actions (see [EW, Theorem 8.13]) implies that for $\hat{\nu}$ -almost every point $[\mathbf{v}]$ in $\mathbb{P}(\mathbf{H})$ there is a set $\mathcal{C}_{[\mathbf{v}]} \subset P$ of positive measure with respect to Haar measure on P , such that for $h \in \mathcal{C}_{[\mathbf{v}]}$, we have $h_*([\mathbf{v}]) \in Z$, and therefore $h_*([\mathbf{v}]) \in \mathbb{P}(\mathbf{E}_{\geq 2}(hx))$, where $x \in X$ is the base point of $[\mathbf{v}]$.

As Z is invariant under the action of g^t , the set $\mathcal{C}_{[\mathbf{v}]}$ is invariant under the action of g^t on the group P ; it follows that $\mathcal{C}_{[\mathbf{v}]}$ intersects U_+ in a set of positive Lebesgue measure. According to Lemma 3.8, this implies that $[\mathbf{v}] \in \mathbf{F}_{\geq 2}(x)$. In particular,

one gets that $\mathbf{F}_{\geq 2}(x)$ is not trivial for ν -almost every point $x \in X$, contradicting the assumption of ν -irreducibility. \square

Using the A -invariance of $\hat{\nu}$ and Oseledets' theorem, we obtain the next claim, which gives the main conclusion of Theorem 1.3:

Claim 4.2. η_x must in fact be supported in $\mathbb{P}(\mathbf{E}_1(x))$ for η -a.e. x .

Proof. To any point $[\mathbf{v}]$ of $\mathbb{P}\mathbf{H}(x) \setminus \mathbb{P}(\mathbf{E}_{\geq 2}(x))$ at a point x generic for ν for which the Lyapunov spaces are well defined, we associate a real number $\alpha([\mathbf{v}])$ as follows: there is a vector $\mathbf{v} \in [\mathbf{v}]$ so that $\mathbf{v} = \mathbf{v}_1 + \alpha([\mathbf{v}])\mathbf{v}_{\geq 2}$ where $\mathbf{v}_1 \in \mathbf{E}_1(x)$ and $\mathbf{v}_{\geq 2} \in \mathbf{E}_{\geq 2}(x)$ are unit vectors.

As g^t preserves the Lyapunov spaces, the set where α is defined is invariant under the action of g^t on $\mathbb{P}\mathbf{H}$. Furthermore, by definition of the Lyapunov spaces one gets

$$\lim_{t \rightarrow +\infty} \alpha((g^t)_*([\mathbf{v}])) = 0,$$

for any $[\mathbf{v}] \in \mathbb{P}\mathbf{H}(x) \setminus \mathbb{P}(\mathbf{E}_{\geq 2}(x))$, where x is generic for ν .

The previous claim said that α is well defined $\hat{\nu}$ -almost everywhere. We want to prove that α vanishes $\hat{\nu}$ -almost everywhere.

Otherwise, there is a compact interval $[a, b] \subset (0, +\infty)$ so that $\hat{\nu}(Z_{a,b}) > 0$ where $Z_{a,b} = \{(x, [\mathbf{v}]) : \alpha([\mathbf{v}]) \in [a, b]\}$. Thus $Z_{a,b}$ is a set of positive measure but every point in $Z_{a,b}$ has only finitely many positive return times for g^t in $Z_{a,b}$, contradicting the Poincaré recurrence theorem. This contradiction concludes the proof. \square

If, moreover, the top Lyapunov exponent λ_1 is simple, it follows that for almost all x , η_x must be the Dirac measure supported on $\mathbb{P}(\mathbf{E}_1(x))$. Hence, in this case, there is only one invariant measure. \square

5. PROOF OF THEOREM 2.8

We begin with the general observation (due to Furstenberg) that Lyapunov exponents of a cocycle on \mathbf{H} can be computed explicitly as integrals over $\mathbb{P}(\mathbf{H})$.

Let $\pi: \mathbf{H} \rightarrow X$ be a bundle with an $A = \{g^t\}$ action, and let $\|\cdot\|_x$ be a Finsler on \mathbf{H} . Let ν be an ergodic A -invariant probability measure on X satisfying the integrability condition (1). Define $\sigma: A \times \mathbb{P}(\mathbf{H}) \rightarrow \mathbb{R}$ by

$$(21) \quad \sigma(g^t, [\mathbf{v}]) = \log \frac{\|g_*^t(\mathbf{v})\|_{g(\pi(\mathbf{v}))}}{\|\mathbf{v}\|_{\pi(\mathbf{v})}},$$

where \mathbf{v} is any nonzero vector representing the projective class $[\mathbf{v}] := \mathbb{F}\mathbf{v} \in \mathbb{P}(\mathbf{H})$ (where \mathbb{F} is the base field). Note that σ is a real-valued cocycle over the action of A on $\mathbb{P}(\mathbf{H})$. By (1), this cocycle is integrable with respect to *any* probability measure $\hat{\nu}$ on $\mathbb{P}(\mathbf{H})$ projecting to ν .

Denote by $\lambda_1(\nu)$ the top Lyapunov exponent for g^t with respect to ν , as in (2). Let

$$Z_1(\nu) = \{\mathbf{v} \in \mathbb{P}^1(\mathbf{H}) : \mathbf{v} \in \mathbb{P}(\mathbf{E}_1(\pi(\mathbf{v})))\},$$

where \mathbf{E}_1 is as in (2).

We recall the following elementary result:

Lemma 5.1. *Let $\hat{\nu}$ be any g^t -invariant measure on $\mathbb{P}(\mathbf{H})$ that projects to ν and is supported in $Z_1(\nu)$. Then*

$$\lambda_1(\nu) = \int_{\mathbb{P}^1(\mathbf{H})} \sigma(g^1, [\mathbf{v}]) d\hat{\nu}([\mathbf{v}]).$$

(Here g^1 means g^t for $t = 1$).

Proof. Suppose $t \in \mathbb{N}$. Since σ is a cocycle,

$$\sigma(g^t, [\mathbf{v}]) = \sum_{n=1}^t \sigma(g^1, [g_*^{t-n} \mathbf{v}]).$$

Since $\hat{\nu}$ is g^t -invariant, integrating both sides over $\mathbb{P}^1(\mathbf{H})$ with respect to $\hat{\nu}$ and dividing both sides by t , we get

$$\begin{aligned} (22) \quad \frac{1}{t} \int_{\mathbb{P}^1(\mathbf{H})} \sigma(g^t, [\mathbf{v}]) d\hat{\nu}([\mathbf{v}]) &= \frac{1}{t} \sum_{n=1}^t \int_{\mathbb{P}^1(\mathbf{H})} \sigma(g^1, [\mathbf{v}]) d\hat{\nu}([g_*^{t-n} \mathbf{v}]) = \\ &= \int_{\mathbb{P}^1(\mathbf{H})} \sigma(g^1, [\mathbf{v}]) d\hat{\nu}([\mathbf{v}]). \end{aligned}$$

However, in view of the assumption that $\hat{\nu}$ is supported in Z_1 , it follows from ergodicity of ν and the multiplicative ergodic theorem that the left-hand side of (22) tends to $\lambda_1(\nu)$ as $t \rightarrow \infty$. \square

Proof of Theorem 2.8.

Let $\mathcal{N}_n, \mathcal{N}, \nu_{\mathcal{N}_n}, \nu_{\mathcal{N}}$ be as in Theorem 2.8. Fix the Finsler structure $\|\cdot\|_x$ on the Hodge bundle over $\mathcal{H}_1(\beta)$ so that the Kontsevich-Zorich cocycle satisfies the uniform integrability condition in (3).

By [EM, Theorem A.6] (which is essentially due to Forni [Fo1]), the Kontsevich-Zorich cocycle restricted to the affine manifold \mathcal{N} is semisimple. This means that after passing to a finite cover, we have the direct sum decomposition

$$(23) \quad H^1(M, \mathbb{R}) = \bigoplus_{i=1}^m W_i(x)$$

where the W_i are G -equivariant, and the restriction of the cocycle to each W_i is strongly irreducible. The map $x \mapsto W_i(x)$ is *a priori* only measurable, but Theorem 2.9 implies that it is continuous (and in fact real analytic).

Note that by Theorem 2.6, we have $\mathcal{N}_n \subset \mathcal{N}$ for sufficiently large n . Then, for sufficiently large n , the decomposition (23) also holds for each \mathcal{N}_n .

It is clearly enough to prove the theorem for the restriction of the cocycle to each W_i . We thus let $\mathbf{H}(x) = W_i(x)$ (for some fixed i), and we may now assume that

the restriction $\alpha_{\mathbf{H}} : G \times \mathbf{H} \rightarrow \mathbf{H}$ of the Kontsevich-Zorich cocycle to \mathbf{H} is strongly irreducible on \mathcal{N} .

Let $\lambda_1(\mathcal{N}_n)$ and $\lambda_1(\mathcal{N})$ denote the top Lyapunov exponents of $\alpha_{\mathbf{H}}$ with respect to the affine measures $\nu_{\mathcal{N}_n}$ and $\nu_{\mathcal{N}}$.

Note that for $\nu_{\mathcal{N}_n}$ -a.e. $x \in X$ the set $\mathbb{P}(\mathbf{E}_1(x)) \subset \mathbb{P}(\mathbf{H})(x)$ is closed, and the set $Z_1(\nu_{\mathcal{N}_n})$ is P -invariant. Then by the amenability of P , for each n there exists a P -invariant measure $\hat{\nu}_n$ on $\mathbb{P}^1(\mathbf{H})$ such that $\hat{\nu}_n$ projects to $\nu_{\mathcal{N}_n}$ and is supported in $Z_1(\nu_{\mathcal{N}_n})$. By Lemma 5.1,

$$(24) \quad \lambda_1(\mathcal{N}_n) = \int_{\mathbb{P}^1(\mathbf{H})} \sigma(g^1, [\mathbf{v}]) d\hat{\nu}_n([\mathbf{v}]),$$

where $\sigma : A \times \mathbb{P}(\mathbf{H}) \rightarrow \mathbb{R}$ is the cocycle defined by (21).

Let $\hat{\nu}$ be any weak-star limit of the measures $\hat{\nu}_n$. Then $\hat{\nu}$ is a P -invariant measure that projects to $\nu_{\mathcal{N}}$. By Theorem 1.3, (each ergodic component of) $\hat{\nu}$ is supported in $Z_1(\nu_{\mathcal{N}})$.

Therefore, by Lemma 5.1,

$$(25) \quad \lambda_1(\mathcal{N}) = \int_{\mathbb{P}^1(\mathbf{H})} \sigma(g^1, [\mathbf{v}]) d\hat{\nu}([\mathbf{v}]).$$

Although $X = \mathcal{H}_1(\beta)$ is not compact, it follows from [EMas] that for every $\delta > 0$ there exists a compact set $K_\delta \subset \mathcal{H}_1(\beta)$ such that for every G -invariant measure ν , we have $\nu(K_\delta) > 1 - \delta$. Given $\epsilon > 0$, let $\delta > 0$ be given by the uniform integrability assumption (3), and fix the compact set K_δ .

Consider the restriction $\mathbb{P}_{K_\delta}(\mathbf{H})$ of $\mathbb{P}(\mathbf{H})$ to this compact set. The weak convergence $\hat{\nu}_n \rightarrow \hat{\nu}$ implies that the integral of the continuous, compactly supported function $\mathbf{1}_{\mathbb{P}_{K_\delta}(\mathbf{H})} \cdot \sigma$ with respect to $\hat{\nu}_n$ converges to the integral with respect to $\hat{\nu}$, and the integral of $\mathbf{1}_{\mathbb{P}(\mathbf{H}) \setminus \mathbb{P}_{K_\delta}(\mathbf{H})} \cdot \sigma$ with respect to all $\hat{\nu}_n$ and $\hat{\nu}$ is less than ϵ , by the uniform integrability assumption (3). Since $\epsilon > 0$ is arbitrary, we conclude that the integrals in (25) converge as $n \rightarrow \infty$ to the integral in (24), and so $\lambda_1(\mathcal{N}_n) \rightarrow \lambda_1(\mathcal{N})$.

To show convergence of other Lyapunov exponents, it suffices to repeat the argument for the cocycle acting on exterior powers of $H^1(M, \mathbb{R})$. We note that the cocycle remains semisimple in this setting and the analogue of (23) still holds (see [Fil]). \square

REFERENCES

- [AEZ] J. Athreya, A. Eskin, Alex, A. Zorich. Right-angled billiards and volumes of moduli spaces of quadratic differentials on \mathbb{CP}^1 . With an appendix by Jon Chaika. *Ann. Sci. Éc. Norm. Supér.* (4) **49** (2016), no. 6, 1311–1386. [10](#)
- [Au1] D. Auricino. Teichmüller Discs with Completely Degenerate Kontsevich-Zorich Spectrum. *Comment. Math. Helv.* **90** (2015), no. 3, 573–643. [10](#)
- [Au2] D. Auricino. Affine Invariant Submanifolds with Completely Degenerate Kontsevich-Zorich Spectrum. [arXiv:1302.0913](#) (2013). To appear in *Ergodic Theory and Dynamical Systems*. [10](#)

- [Au3] D. Auricino. Affine manifolds and zero Lyapunov exponents in genus 3. *Geom. Funct. Anal.* **25** (2015), no. 5, 1333–1370. [10](#)
- [ANW] D. Auricino, D-M. Nguyen, A. Wright Classification of higher rank orbit closures in $H^{odd}(4)$. *J. Eur. Math. Soc. (JEMS)* **18** (2016), no. 8, 1855–1872. [11](#)
- [AGY] A. Avila, S. Gouëzel, J-C. Yoccoz. Exponential mixing for the Teichmüller flow. *Publ. Math. Inst. Hautes Études Sci.* **104** (2006), 143–211. [10](#)
- [AV] A. Avila, M. Viana, Simplicity of Lyapunov spectra: proof of the Zorich–Kontsevich conjecture, *Acta Math.*, **198**, No. 1, (2007), 1–56. [10](#), [11](#)
- [BF] U. Bader, A. Furman, Boundaries, rigidity of representations, and Lyapunov exponents, *Proceedings of ICM 2014, Invited Lectures*, (2014), 71 – 96. [5](#), [6](#)
- [Ba1] M. Bainbridge, Euler Characteristics of Teichmüller Curves in Genus Two, *Geom. Topol.*, **11** (2007), 1887–2073. [10](#)
- [Ba2] M. Bainbridge, Billiards in L-shaped tables with barriers, *GAF*, **20** (2010), no. 2, 299–356; **20** (2010), no. 5, 1306. [10](#)
- [BaMa] Y. Bakhtin, M. Martínez, A characterization of harmonic measures on laminations by hyperbolic Riemann surfaces. *Ann. Inst. Henri Poincaré Probab. Stat.* **44** (2008), 1078–1089. [4](#), [7](#)
- [BG] C. Bonatti and X. Gómez-Mont, Sur le comportement statistique des feuilles de certains feuilletages holomorphes. *Essays on geometry and related topics*, Vol. 1, 2, 15–41, Monogr. Enseign. Math., 38, Enseignement Math., Geneva, 2001. [1](#), [3](#), [6](#), [7](#)
- [BoMö] I. Bouw, M. Möller, Teichmüller curves, triangle groups, and Lyapunov exponents, *Annals of Math.*, (2) **172** (2010), no. 1, 139–185. [10](#)
- [Bu] A. Bufetov. Limit theorems for translation flows. *Ann. of Math.* (2) **179** (2014), no. 2, 431–499. [10](#)
- [CE] J. Chaika, A. Eskin, Every flat surface is Birkhoff and Osceledets generic in almost every direction. *J. Mod. Dyn.* **9** (2015), 1–23.
- [CM1] D. Chen, M. Möller, Non-varying sums of Lyapunov exponents of Abelian differentials in low genus. *Geom. Topol.* **16** (2012), 2427–2479. [10](#)
- [CM2] D. Chen, M. Möller. Quadratic differentials in low genus: exceptional and non-varying. *Ann. Sci. Éc. Norm. Supér. (4)* **47** (2014), no. 2, 309–369. [10](#)
- [DD] J. Daniel, B. Deroin. Lyapunov exponents of the Brownian motion on a Kähler manifold. [arXiv:1702.02551 \[math.DS\]](#) (2017). [10](#)
- [DHL] V. Delecroix, P. Hubert, S. Lelièvre. Diffusion for the periodic wind-tree model. *Ann. Sci. Éc. Norm. Supér. (4)* **47** (2014), no. 6, 1085–1110. [10](#)
- [DM] V. Delecroix, C. Matheus Santos. Un contre-exemple de la réciproque du critère de Forni pour la positivité des exposants de Lyapunov du cocycle de Kontsevich-Zorich. (French) [[A counterexample to the reciprocal to Forni’s criterion for the positivity of Lyapunov exponents of the Kontsevich-Zorich cocycle]] *Math. Res. Lett.* **22** (2015), no. 6, 1667–1678. [10](#)
- [DZ] Cries and whispers in wind-tree forests. [arxiv:1502.06405 \[math.DS\]](#) (2015). [10](#)
- [EW] M. Einsiedler, T. Ward, Ergodic theory with a view towards number theory. Graduate Texts in Mathematics, **259**. Springer-Verlag London, Ltd., London, 2011. [1](#), [17](#)
- [EKMZ] A. Eskin, M. Kontsevich, M. Moeller, A. Zorich. Lower bounds for Lyapunov exponents of flat bundles on curves. [arXiv:1609.01170 \[math.GT\]](#) (2016). To appear in *Geometry and Topology*. [10](#)
- [EKZ] A. Eskin, M. Kontsevich, A. Zorich. Lyapunov spectrum of square-tiled cyclic covers, *Journal of Modern Dynamics*, **5:2** (2011), 319–353. [10](#)

- [EKZ2] A. Eskin, M. Kontsevich, A. Zorich. Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow. *Publ. Math. Inst. Hautes Études Sci.* **120** (2014), 207–333. [10](#)
- [EMas] A. Eskin, H. Masur, Asymptotic formulas on flat surfaces. *Ergodic Theory and Dynamical Systems*, **21** (2) (2001), 443–478. [20](#)
- [EMat] A. Eskin, C. Matheus Santos. A coding-free simplicity criterion for the Lyapunov exponents of Teichmüller curves. *Geom. Dedicata* **179** (2015), 45–67. [5](#), [10](#)
- [EM] A. Eskin, M. Mirzakhani. “Invariant and stationary measures for the $SL(2, \mathbb{R})$ action on moduli space.” [arXiv:1302.3320 \[math.DS\]](#) (2013). [3](#), [8](#), [19](#)
- [EMM] A. Eskin, M. Mirzakhani, A. Mohammadi. “Isolation, equidistribution and orbit closures for the $SL(2\mathbb{R})$ action on moduli space.” *Ann. of Math.* (2) **182** (2015), no. 2, 673–721. [8](#), [9](#), [10](#)
- [EFW] A. Eskin, S. Filip, A. Wright. “The algebraic hull of the Kontsevich-Zorich cocycle.” [arXiv:1702.02074 \[math.DS\]](#) (2017). [4](#)
- [Fi1] S. Filip. Semisimplicity and rigidity of the Kontsevich-Zorich cocycle. *Invent. Math.* **205** (2016), no. 3, 617–670. [10](#), [11](#), [20](#)
- [Fi2] S. Filip. Splitting mixed Hodge structures over affine invariant manifolds. *Ann. of Math.* (2) **183** (2016), no. 2, 681–713. [10](#)
- [Fi3] S. Filip. “Zero Lyapunov exponents and monodromy of the Kontsevich-Zorich cocycle.” *Duke Math. J.* **166** (2017), no. 4, 657–706. [10](#)
- [Fi4] S. Filip. Families of K3 surfaces and Lyapunov exponents. [arXiv:1412.1779 \[math.DS\]](#) (2014).
- [FFM] Quaternionic covers and monodromy of the Kontsevich-Zorich cocycle in orthogonal groups. [arXiv:1502.07202 \[math.DS\]](#) (2015). [10](#)
- [Fo1] G. Forni, Deviation of ergodic averages for area-preserving flows on surfaces of higher genus, *Annals of Math.*, **155**, no. 1, 1–103 (2002) [10](#), [19](#)
- [Fo2] G. Forni, On the Lyapunov exponents of the Kontsevich-Zorich cocycle. Handbook of dynamical systems. Volume 1B. Editors: B. Hasselblatt and A. Katok. Elsevier, Amsterdam, 549–580, 2006. [10](#)
- [F3] G. Forni, A geometric criterion for the non-uniform hyperbolicity of the Kontsevich–Zorich cocycle, *Journal of Modern Dynamics*, **5:2** (2011), 355–395. [10](#)
- [FM] G. Forni, C. Matheus Santos. “An example of a Teichmüller disk in genus 4 with degenerate Kontsevich-Zorich spectrum.” [arXiv:0810.0023 \[math.DS\]](#) (2008). [10](#)
- [FMZ1] G. Forni, C. Matheus, A. Zorich, Square-tiled cyclic covers, *Journal of Modern Dynamics*, **5:2** (2011), 285–318. [10](#)
- [FMZ2] G. Forni, C. Matheus, A. Zorich, Lyapunov spectra of covariantly constant subbundles of the Hodge bundle. *Ergodic Theory Dynam. Systems* **34** (2014), no. 2, 353–408. [10](#)
- [FMZ3] G. Forni, C. Matheus, A. Zorich, Zero Lyapunov exponents of the Hodge bundle. *Comment. Math. Helv.* **89** (2014), no. 2, 489–535. [10](#)
- [Fur67] H. Furstenberg. Poisson boundaries and envelopes of discrete groups. *Bull. Amer. Math. Soc.* **73** (1967), 350–356. [5](#)
- [Fu71] H. Furstenberg, Random walks and discrete subgroups of Lie groups, *Advances in Probability and Related Topics*, Vol. 1 (1971), 1–63, Dekker, New York. [5](#)
- [Ga] L. Garnett. Foliations, the ergodic theorem and Brownian motion. *J. Funct. Anal.* **51** (1983) 285–311. [4](#)
- [GH1] J. Grivaux, P. Hubert, Exposants de Lyapunov du flot de Teichmüller (d’après Eskin–Kontsevich–Zorich). (French) [[The Lyapunov exponents of Teichmüller flow (following

- Eskin-Kontsevich-Zorich)] *Astérisque* No. 361, (2014), Exp. No. 1060, vii, 43–75. ISBN: 978-285629-785-8 Séminaire Nicolas Bourbaki, Octobre 2012. [10](#)
- [GH2] J. Grivaux, P. Hubert, *Loci in strata of meromorphic differentials with fully degenerate Lyapunov spectrum*, *J. Mod. Dyn.* **8** (2014), no. 1, 61–73. [10](#)
- [GM] I. Goldsheid and G. Margulis, Lyapunov exponents of a product of random matrices, *Uspekhi Mat. Nauk*, **44**, No. 5 (1989), 13–60 (Russian); English translation in *Russian Math. Surveys*, **44**, No. 5 (1989), 11–71. [5](#), [12](#)
- [GR] Y. Guivarc’h and A. Raugi, *Products of random matrices: convergence theorems*, in Random Matrices and their Applications (Brunswick, ME, 1984), Contemp. Math., 50, pp. 31–54. Amer. Math. Soc., Providence, RI, 1986. [5](#)
- [K] M. Kontsevich, Lyapunov exponents and Hodge theory. “The mathematical beauty of physics” (Saclay, 1996), (in Honor of C. Itzykson) 318–332, Adv. Ser. Math. Phys., 24, World Sci. Publishing, River Edge, NJ, 1997. [10](#)
- [KZ1] M. Kontsevich and A. Zorich, Lyapunov exponents and Hodge theory, Preprint IHES M/97/13, pp. 1–16; [arXiv:hep-th/9701164](#). [10](#)
- [KZ2] M. Kontsevich and A. Zorich, Connected components of the moduli spaces of Abelian differentials, *Inventiones Math.*, **153:3**, (2003), 631–678. [11](#)
- [Mar] M. Martínez, Measures on hyperbolic surface laminations. *Ergodic Theory Dynam. Systems*, **26** (2006), 847–867. [4](#)
- [Mat] C. Matheus Santos, Appendix to the paper of G. Forni, A geometric criterion for the nonuniform hyperbolicity of the Kontsevich–Zorich cocycle, *Journal of Modern Dynamics*, **4** (2010), no. 3, 453–486. [10](#)
- [MW] C. Matheus and G. Weitze-Schmithüsen. Some examples of isotropic $SL(2, \mathbb{R})$ -invariant subbundles of the Hodge bundle. *Int. Math. Res. Not. IMRN* (2015), no. 18, 8657–8679. [10](#)
- [MY] C. Matheus Santos, J.-C. Yoccoz, The action of the affine diffeomorphisms on the relative homology group of certain exceptionally symmetric origamis. *Journal of Modern Dynamics*, **5** (2011), no. 2, 386–395. [10](#)
- [MYZ] C. Matheus Santos, J.-C. Yoccoz, D. Zmiaikou, Homology of origamis with symmetries, *Ann. Inst. Fourier (Grenoble)*, **64** (2014), no. 3, 1131–1176. [10](#)
- [MMY] C. Matheus Santos, M. Möller, J.-C. Yoccoz. “A criterion for the simplicity of the Lyapunov spectrum of square-tiled surfaces.” *Invent. Math.* **202** (2015), no. 1, 333–425. [1](#), [10](#), [11](#)
- [M] S. Matsumoto, Weak form of equidistribution theorem for harmonic measures of foliations by hyperbolic surfaces. *Proc. Amer. Math. Soc.*, **144** (2016), no. 3, 1289–1297. [7](#)
- [McM] C. McMullen, Dynamics of SL over moduli space in genus two, *Ann. of Math.*, **165** (2007), no. 2, 397–456. [8](#)
- [Mö] M. Möller, Shimura- and Teichmüller curves, *Journal of Modern Dynamics*, **5:1** (2011), 1–32. [10](#)
- [Moz] S. Mozes. Epimorphic subgroups and invariant measures. *Ergodic Theory Dynam. Systems* **15** (1995), no. 6, 1207–1210. [4](#)
- [MS] S. Mozes, N. Shah, *On the space of ergodic invariant measures of unipotent flows*, *Ergodic Theory Dynam. Systems* **15** (1995), no. 1, 149–159. [9](#)
- [NW] D.-M. Nguyen, A. Wright. Non-Veech surfaces in $\mathcal{H}(4)^{hyp}$ are generic. *Geom. Funct. Anal.* **24** (2014), no. 4, 1316–1335. [11](#)
- [P1] A. Pardo, Counting problem on wind-tree models. [arXiv:1604.05654](#) [math.DS] (2016). [10](#)

- [P2] A. Pardo, Quantitative error term in the counting problem on Veech wind-tree models. [arXiv:1704.07428 \[math.DS\]](#) (2017). [10](#)
- [P3] A. Pardo, A non-varying phenomenon with an application to the wind-tree model. [arXiv:1704.07682 \[math.DS\]](#) (2017). [10](#)
- [T] R. Trevino, On the Non-Uniform Hyperbolicity of the Kontsevich-Zorich Cocycle for Quadratic Differentials. *Geom. Dedicata* **163** (2013), 311–338. [10](#)
- [W] A. Wright. Schwarz triangle mappings and Teichmüller curves: abelian square-tiled surfaces. *J. Mod. Dyn.* **6** (2012), no. 3, 405–426. [10](#)
- [Yu] F. Yu. Eigenvalues of curvature, Lyapunov exponents and Harder-Narasimhan filtrations. [arXiv:1408.1630 \[math.AG\]](#) (2014). [10](#)
- [YZ] F. Yu and K. Zuo. Weierstrass filtration on Teichmüller curves and Lyapunov exponents. *J. Mod. Dyn.* **7** (2013), no. 2, 209–237. [10](#)
- [Zo] A. Zorich, Deviation for interval exchange transformations, *Ergodic Theory and Dynamical Systems*, **17** (1997), 1477–1499. [10](#)
- [Zo2] A. Zorich, Flat surfaces. In collection “Frontiers in Number Theory, Physics and Geometry. Vol. 1: On random matrices, zeta functions and dynamical systems”; Ecole de physique des Houches, France, March 9–21 2003, P. Cartier; B. Julia; P. Moussa; P. Vanhove (Editors), Springer-Verlag, Berlin, (2006), 439–586. [8](#), [9](#)