# ON THE PROPERTIES OF NEW FAMILIES OF GENERALIZED FIBONACCI NUMBERS

#### GAMALIEL CERDA-MORALES

ABSTRACT. In this paper, new families of generalized Fibonacci and Lucas numbers are introduced. In addition, we present the recurrence relations and the generating functions of the new families for k=2.

#### 1. Introduction

There are a lot of integer sequences such as Fibonacci, Pell, Jacobsthal, Balancing, Lucas, Pell-Lucas, Jacobsthal-Lucas, Lucas-balancing, etc. For example, Horadam numbers are used as a generalized of Fibonacci and Lucas sequences by scientists for basic theories and their applications. For interest application of these numbers in science and nature, one can see [1, 6, 8, 10, 11]. For instance, some special cases of the Horadam numbers could be derived directly using a new matrix representation, see [3, 4].

In 1965, Horadam studied some properties of sequences of the type,  $\{W_n\}_{n\geq 0}$  or  $\{W_n(a,b;p,q)\}$ , where a,b are nonnegative integers and p,q are arbitrary integers, see [6]. Such sequences are defined by the recurrence relations of second order

$$(1.1) W_n = pW_{n-1} - qW_{n-2}, \ n \ge 2,$$

with  $W_0 = a$  and  $W_1 = b$ . We are interested in the following two special cases of  $\{W_n\}$ :  $\{U_n\}$  is defined by  $U_0 = 0$ ,  $U_1 = 1$ , and  $\{V_n\}$  is defined by  $V_0 = 2$ ,  $V_1 = p$ . It is well known that  $\{U_n\}$  and  $\{V_n\}$  can be expressed in the form

(1.2) 
$$U_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \ V_n = \alpha^n + \beta^n,$$

where  $\alpha = \frac{p+\sqrt{\Delta}}{2}$ ,  $\beta = \frac{p-\sqrt{\Delta}}{2}$  and the discriminant is  $\Delta = p^2 - 4q$ . Especially, if p = -q = 1, p = -2q = 2, 2p = -q = 2 and p = 6q = 6, then  $\{U_n\}_{n \geq 0}$  is the usual Fibonacci, Pell, Jacobsthal and Balancing sequence, respectively.

The aim of this work is to study some properties of two new sequences that generalize the  $\{U_n\}_{n\geq 0}$  and  $\{V_n\}_{n\geq 0}$  numbers. In this work we will follow closely the work of El-Mikkawy and Sogabe (see [5]) where the authors give a new family that generalizes the Fibonacci numbers and establish relations with the ordinary Fibonacci numbers.

So, in the next section we start giving the new definition of generalized Fibonacci and Lucas numbers, and we exhibit some elements of them. We also present relations of these sequences with ordinary  $U_n$  and  $V_n$ . We deduce some properties of

1

<sup>2000</sup> Mathematics Subject Classification. Primary 11B37, 11B50, 11B83 Secondary 05A15. Key words and phrases. Horadam sequences, generalized Fibonacci numbers, generating matrix, generating function, recurrence relations.

these new families, but using different methods. Furthermore, we study a particular case, that is two sequences of the new defined families for k=2. For these sequences we present some recurrence relations and generating functions.

#### 2. Main Results

We define the new generalized Fibonacci and Lucas numbers and obtain some results related to these numbers by using [5].

**Definition 2.1.** Let n and  $k \neq 0$  be natural numbers. There exist unique numbers m and r such that n = mk + r  $(0 \le r < k)$ . Using these parameters, we define new generalized Fibonacci numbers  $U_n^{(k)}$  and generalized Lucas numbers  $V_n^{(k)}$  by

(2.1) 
$$U_n^{(k)} = \frac{1}{(\alpha - \beta)^k} (\alpha^{m+1} - \beta^{m+1})^r (\alpha^m - \beta^m)^{k-r}$$

and

(2.2) 
$$V_n^{(k)} = (\alpha^{m+1} + \beta^{m+1})^r (\alpha^m + \beta^m)^{k-r}$$

where  $\alpha = \frac{p+\sqrt{\Delta}}{2}$  and  $\beta = \frac{p-\sqrt{\Delta}}{2}$ , respectively.

The first few numbers of the new generalized Fibonacci and generalized Lucas numbers as follows

$$(2.3) \qquad \{U_n^{(1)}\}_{n=0}^3 = \{0,1,p,p^2-q\}, \ \{V_n^{(1)}\}_{n=0}^3 = \{2,p,p^2-2q,p^3-3pq\},$$

$$\{U_n^{(2)}\}_{n=0}^3 = \{0,0,1,p\}, \ \{V_n^{(2)}\}_{n=0}^3 = \{4,2p,p^2,p^3-2pq\},$$

$$\{U_n^{(3)}\}_{n=0}^3 = \{0, 0, 0, 1\}, \ \{V_n^{(3)}\}_{n=0}^3 = \{8, 4p, 2p^2, 8\}$$

for k = 1, 2, 3.

Given n a nonnegative integer and k a natural number, then

$$U_{mk+r}^{(k)} = U_m^{k-r} U_{m+1}^r$$
 and  $V_{mk+r}^{(k)} = V_m^{k-r} V_{m+1}^r$ ,

where m and r are nonnegative integers such that n = mk + r  $(0 \le r < k)$ . Note that  $U_n^{(1)}$  is the generalized Fibonacci numbers  $U_n$  and  $V_n^{(1)}$  is the generalized Lucas numbers  $V_n$ .

In the following, we give some properties of the new generalized Fibonacci and generalized Lucas numbers.

**Theorem 2.1.** Let  $k, m \in \{1, 2, 3, ...\}$  be fixed numbers. The new generalized Fibonacci numbers and generalized Lucas numbers satisfy

(i) 
$$\sum_{t=0}^{k-1} {k-1 \choose t} (-p)^{-t} U_{mk+t}^{(k)} = \left(\frac{q}{p}\right)^{k-1} U_m U_{(m-1)(k-1)}^{(k-1)},$$

(ii) 
$$\sum_{t=0}^{k-1} {k-1 \choose t} (\frac{-p}{q})^t U_{mk+t}^{(k)} = (-q)^{1-k} U_m U_{(m+2)(k-1)}^{(k-1)}$$
 and

$$\begin{array}{l} \text{(i)} \ \, \sum_{t=0}^{k-1} {k-1 \choose t} (-p)^{-t} U_{mk+t}^{(k)} = \left(\frac{q}{p}\right)^{k-1} U_m U_{(m-1)(k-1)}^{(k-1)}, \\ \text{(ii)} \ \, \sum_{t=0}^{k-1} {k-1 \choose t} (\frac{-p}{q})^t U_{mk+t}^{(k)} = (-q)^{1-k} U_m U_{(m+2)(k-1)}^{(k-1)} \ \, \text{and} \\ \text{(iii)} \ \, \sum_{t=0}^{k-1} p^{-t} U_{mk+t}^{(k)} = \left(-\frac{p}{q}\right) p^{-k} \left(\frac{U_m}{U_{m-1}}\right) \left(U_{(m+1)k}^{(k)} - p^k U_{mk}^{(k)}\right). \end{array}$$

*Proof.* Note of the definition 1,  $U_n^{(k)} = U_m^{k-r} U_{m+1}^r$ , where n = mk + r and  $0 \le r < k$ . Then, (i).

$$\begin{split} \sum_{t=0}^{k-1} \binom{k-1}{t} (-p)^{-t} U_{mk+t}^{(k)} &= (-p)^{1-k} U_m \sum_{t=0}^{k-1} \binom{k-1}{t} (-p)^{k-1-t} U_m^{k-1-t} U_{m+1}^t \\ &= (-p)^{1-k} U_m \sum_{t=0}^{k-1} \binom{k-1}{t} (-p U_m)^{k-1-t} U_{m+1}^t. \end{split}$$

Using the binomial theorem,  $\sum_{t=0}^{k-1} {k-1 \choose t} (-pU_m)^{k-1-t} U_{m+1}^t = (-pU_m + U_{m+1})^{k-1}$ . Then,

$$\begin{split} \sum_{t=0}^{k-1} \binom{k-1}{t} (-p)^{-t} U_{mk+t}^{(k)} &= (-p)^{1-k} U_m (-p U_m + U_{m+1})^{k-1} \\ &= (-p)^{1-k} U_m (-q U_{m-1})^{k-1} \\ &= \left(\frac{q}{p}\right)^{k-1} U_m U_{(m-1)(k-1)}^{(k-1)}, \end{split}$$

is obtained and this completes the proof.

(ii).

$$\begin{split} \sum_{t=0}^{k-1} \binom{k-1}{t} (\frac{-p}{q})^t U_{mk+t}^{(k)} &= U_m \sum_{t=0}^{k-1} \binom{k-1}{t} (-q)^{-t} p^t U_m^{k-1-t} U_{m+1}^t \\ &= (-q)^{1-k} U_m \sum_{t=0}^{k-1} \binom{k-1}{t} (-qU_m)^{k-1-t} (pU_{m+1})^t. \end{split}$$

Using the binomial theorem,  $\sum_{t=0}^{k-1} {k-1 \choose t} (-qU_m)^{k-1-t} (pU_{m+1})^t = (pU_{m+1} - qU_m)^{k-1}$ . Then,

$$\sum_{t=0}^{k-1} {k-1 \choose t} (\frac{-p}{q})^t U_{mk+t}^{(k)} = (-q)^{1-k} U_m (pU_{m+1} - qU_m)^{k-1}$$
$$= (-q)^{1-k} U_m U_{m+2}^{k-1}$$
$$= (-q)^{1-k} U_m U_{(m+2)(k-1)}^{(k-1)},$$

is obtained

(iii). 
$$\sum_{t=0}^{k-1} p^{-t} U_{mk+t}^{(k)} = \sum_{t=0}^{k-1} p^{-t} U_m^{k-t} U_{m+1}^t = U_m^k \sum_{t=0}^{k-1} \left( \frac{U_{m+1}}{pU_m} \right)^t$$
, but

$$\sum_{t=0}^{k-1} \left( \frac{U_{m+1}}{pU_m} \right)^t = \frac{\left( \frac{U_{m+1}}{pU_m} \right)^k - 1}{\frac{U_{m+1}}{pU_m} - 1} = (pU_m)^{1-k} \left( \frac{U_{m+1}^k - pU_m}{U_{m+1} - pU_m} \right).$$

Then.

$$\begin{split} \sum_{t=0}^{k-1} p^{-t} U_{mk+t}^{(k)} &= U_m^k (p U_m)^{1-k} \left( \frac{U_{m+1}^k - p^k U_m^k}{U_{m+1} - p U_m} \right) \\ &= p^{1-k} U_m \left( \frac{U_{m+1}^k - p^k U_m^k}{-q U_{m-1}} \right) \\ &= \left( -\frac{p}{q} \right) p^{-k} \left( \frac{U_m}{U_{m-1}} \right) \left( U_{(m+1)k}^{(k)} - p^k U_{mk}^{(k)} \right). \end{split}$$

## 3. A particular case $U_n^{(2)}$ and $V_n^{(2)}$

In [3], the author use a matrix method for generating the generalized Fibonacci numbers by defining the W(p,q)-matrix and proved that

(3.1) 
$$W(p,q)^n = \begin{bmatrix} p & -q \\ 1 & 0 \end{bmatrix}^n = \begin{bmatrix} U_{n+1} & -qU_n \\ U_n & -qU_{n-1} \end{bmatrix},$$

for any natural number n.

Thus, for any  $n, l \ge 0$  and  $n + l \ge 2$ , we have

$$\left[ \begin{array}{cc} U_{n+l} & -qU_{n+l-1} \\ U_{n+l-1} & -qU_{n+l-2} \end{array} \right] = W(p,q)^{n+l-2} \left[ \begin{array}{cc} p & -q \\ 1 & 0 \end{array} \right].$$

If we compute the determinant of both sides of the previous equality we obtain

$$(3.2) -qU_{n+l}U_{n+l-2} + qU_{n+l-1}^2 = q \det(W(p,q)^{n+l-2}).$$

which is equivalent to

$$U_{n+l-1}^2 - U_{n+l}U_{n+l-2} = q^{n+l-2}.$$

Since, by Definition 2.1 (where m = n + l - 1, k = 2 and r = 0)

$$U_{2(n+l-1)}^{(2)} = U_{n+l-1}^2.$$

We have proved the following result:

**Theorem 3.1.** We have the following relation for the new family of generalized Fibonacci numbers  $\{U_n^{(2)}\}$ 

(3.3) 
$$U_{2(n+l-1)}^{(2)} = q^{n+l-2} + U_{n+l}U_{n+l-2},$$

where  $n + l \ge 2$ .

We study the particular case of the sequences  $\{U_n^{(2)}\}_{n\geq 0}$  and  $\{V_n^{(2)}\}_{n\geq 0}$  defined by 2.1 and 2.2, respectively. First, we present a recurrence relation for these sequences.

**Theorem 3.2.** The sequences  $\{U_n^{(2)}\}$  and  $\{V_n^{(2)}\}$  satisfy, respectively, the following recurrence relations:

$$U_n^{(2)} = pU_{n-1}^{(2)} - pqU_{n-3}^{(2)} + q^2U_{n-4}^{(2)}, \ n \ge 4,$$

and

$$(3.5) V_n^{(2)} = pV_{n-1}^{(2)} - pqV_{n-3}^{(2)} + q^2V_{n-4}^{(2)}, \ n \ge 4,$$

$$\begin{split} U_{2m}^{(2)} &= U_m^2 \\ &= U_m \left( p U_{m-1} - q U_{m-2} \right) \\ &= p U_m U_{m-1} - q \left( p U_{m-1} - q U_{m-2} \right) U_{m-2} \\ &= p U_{2m-1}^{(2)} - p q U_{2m-3}^{(2)} + q^2 U_{2m-4}^{(2)}. \end{split}$$

If we take n=2m we have the following equation  $U_n^{(2)}=pU_{n-1}^{(2)}-pqU_{n-3}^{(2)}+q^2U_{n-4}^{(2)}$ . If n is odd integer, then we have

$$\begin{split} U_{2m+1}^{(2)} &= U_m U_{m+1} \\ &= U_m \left( p U_m - q U_{m-1} \right) \\ &= p U_m U_m - q \left( p U_{m-1} - q U_{m-2} \right) U_{m-1} \\ &= p U_{2m}^{(2)} - p q U_{2m-2}^{(2)} + q^2 U_{2m-3}^{(2)}. \end{split}$$

In a similar way we can prove the result for  $\{V_n^{(2)}\}_{n\geq 0}$ . This theorem is completed.

We also note that if we consider separately the even and the odd terms of the above defined sequences we can obtain shorter recurrence relations. In fact, for n = 2m, for any natural number m, by Theorem 3.1 (with n = m and l = 1) we have

$$\begin{split} U_{2m}^{(2)} &= U_{m+1}U_{m-1} + q^{m-1} \\ &= U_{m-1} \left( pU_m - qU_{m-1} \right) + q^{m-1} \\ &= pU_{m-1}U_m - qU_{m-1}^2 + q^{m-1} \\ &= pU_{2m-1}^{(2)} - qU_{2m-2}^{(2)} + q^{m-1}. \end{split}$$

Similarly, for the odd case n = 2m + 1, we have

$$U_{2m+1}^{(2)} = U_m U_{m+1}$$

$$= pU_m U_m - qU_m U_{m-1}$$

$$= pU_{2m}^{(2)} - qU_{2m-1}^{(2)}.$$

Therefore we can conclude the following. A shorter recurrence relation for the sequence  $\{U_n^{(2)}\}$  is given by

$$\begin{cases} U_{2m}^{(2)} = pU_{2m-1}^{(2)} - qU_{2m-2}^{(2)} + q^{m-1} \\ U_{2m+1}^{(2)} = pU_{2m}^{(2)} - qU_{2m-1}^{(2)} \end{cases}$$

for the even and the odd terms.

Next we find generating functions for these sequences.

**Theorem 3.3.** The generating function  $g^{(2)}(x)$  for  $\{U_n^{(2)}\}$  is given by

(3.6) 
$$g^{(2)}(x) = \frac{x^2}{1 - px + pqx^3 - q^2x^4}.$$

Г

*Proof.* To the sum of this power series,  $g^{(2)}(x) = \sum_{n=0}^{\infty} U_n^{(2)} x^n$ , we call generating function of the new generalized Fibonacci sequence of numbers  $\{U_n\}_{n\geq 0}$ . Then,

$$\begin{split} &(1-px+pqx^3-q^2x^4)g^{(2)}(x)\\ &=(U_0^{(2)}+U_1^{(2)}x+U_2^{(2)}x^2+U_3^{(2)}x^3+\cdots)\\ &-(pU_0^{(2)}x+pU_1^{(2)}x^2+pU_2^{(2)}x^3+pU_3^{(2)}x^4+\cdots)\\ &+(pqU_0^{(2)}x^3+pqU_1^{(2)}x^4+pqU_2^{(2)}x^5+pqU_3^{(2)}x^6+\cdots)\\ &-(q^2U_0^{(2)}x^4+q^2U_1^{(2)}x^5+q^2U_2^{(2)}x^6+q^2U_3^{(2)}x^7+\cdots)\\ &=U_0^{(2)}+(U_1^{(2)}-pU_0^{(2)})x+(U_2^{(2)}-pU_1^{(2)})x^2+(U_3^{(2)}-pU_2^{(2)}+pqU_0^{(2)})x^3. \end{split}$$

Hence, taking into account the initial conditions of the sequence  $\{U_n^{(2)}\}_{n\geq 0}$  and by Theorem 3.2, this is equivalent to

$$(1 - px + pqx^3 - q^2x^4)g^{(2)}(x) = x^2$$

and therefore

$$g^{(2)}(x) = \frac{x^2}{1 - px + pqx^3 - q^2x^4},$$

which completes the proof.

**Theorem 3.4.** The generating function  $h^{(2)}(x)$  for  $\{V_n^{(2)}\}$  is given by

(3.7) 
$$h^{(2)}(x) = \frac{4 - 2px - p^2x^2 + 2pqx^3}{1 - px + pqx^3 - q^2x^4}$$

*Proof.* The proof can be obtained in a similar way to the proof of above theorem.

### References

- [1] A. Atalay, K. Uslu and H. Gokkaya, New families of s-fibonacci and s-lucas numbers, Selcuk Journal of Applied Mathematics, 14(2) (2013), 34–42.
- [2] P. Catarino et al., New families of Jacobsthal and Jacobsthal-Lucas numbers, Algebra and Discrete Mathematics, 20(1) (2015), 40–54.
- [3] G. Cerda-Morales, Matrix methods in Horadam sequences, Bol. Mat. 19 (2) (2012), 97–106.
- [4] G. Cerda-Morales, On generalized Fibonacci and Lucas numbers by matrix methods, Hacettepe Journal of Mathematics and Statistics, 42(2) (2013), 173–179.
- [5] M. El-Mikkawy and T. Sogabe, A new family of k-Fibonacci numbers, Applied Mathematics and Computation 215 (2010), 4456–4461.
- [6] A.F. Horadam, Basic properties of a certain generalized sequence of numbers, Fibonacci Quart. 3 (1965), 161–176.
- [7] A.F. Horadam, Jacobsthal representation numbers, Fibonacci Quart. Vol. 34(1) (1996), 40–53.
- [8] D. Kalman and R. Mena, The Fibonacci numbers-exposed, Math. Mag., 76 (2003), 81-167.
- [9] F. Koken and D. Bozkurt, On the Jacobsthal numbers by matrix methods, Int. J. Contemp. Math. Sci. 3(13) (2008), 605-614.
- [10] E. Karaduman, On determinants of matrices with general Fibonacci numbers entries, Appl. Math. Comp. 167 (2005), 670–676.
- [11] N. Yilmaz, Y. Yazlik and N. Taskara, On the k-Generalized Fibonacci Numbers, Selcuk Journal of Applied Mathematics, 13(1) (2012), 83–88.

Centro Superior de Docencia en Ciencias Básicas, Escuela de Pedagogía en Matemáticas, Universidad Austral de Chile, Puerto Montt, Chile.

E-mail address: gamaliel.cerda@uach.cl