

1

$$\mu = [-3, 1, 4]^T, \quad \Sigma = \begin{bmatrix} 1 & -2 & 0 \\ -2 & 5 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$a) \text{Cor}(X_1, X_2) = \frac{-2}{\sqrt{1 \cdot 5}} \neq 0$$

$\rightarrow X_1, X_2$ are not linearly independent

$$b) \text{Cor}(X_2, X_3) = \frac{0}{\sqrt{5 \cdot 2}} = 0$$

$\rightarrow X_2, X_3$ are linearly dependent ①

$$c) Y = [X_1, X_2]^T$$

$$\text{Cor}(X_1, X_3) = \frac{0}{\sqrt{1 \cdot 2}} = 0 \rightarrow X_1, X_3 \text{ are linearly independent } ②$$

①, ② $\rightarrow Y, X_3$ are linearly independent

$$d) Y = \frac{X_1 + X_2}{2} \sim N(\mu_Y = -1, \sigma_Y^2 = \frac{3}{2})$$

$$\text{Cov}(Y, X_3) = \frac{1}{2} \text{Cov}(X_1, X_3) + \frac{1}{2} \text{Cov}(X_2, X_3) = 0$$

$$\rightarrow \text{Cor}(Y, X_3) = 0$$

$\rightarrow Y, X_3$ linearly independent

$$e) \text{Cov}(Y, X_2) = \text{Cov}(X_2, X_2) - \frac{5}{2} \text{Cov}(X_1, X_2) - \text{Cov}(X_1, X_3) = 5 - \frac{5}{2}(-2) = 10 \neq 0$$

$\rightarrow Y, X_2$ not linearly independent

2 Mahalanobis distance

$$d_M(y, \mu) = \sqrt{(y - \mu)^T \Sigma^{-1} (y - \mu)}$$

(a)

$$d_E(y, \mu_1) = \sqrt{1+4+4} = \boxed{3}$$

$$d_E(y, \mu_2) = \sqrt{4+4+2.25} \approx \boxed{3.20}$$

(b)

$$\Sigma = \begin{bmatrix} 9.0 & 8.1 & -3.6 \\ 8.1 & 9.0 & -4.8 \\ -3.6 & -4.8 & 4.0 \end{bmatrix} \rightarrow \Sigma^{-1} = \begin{bmatrix} 0.741 & -0.864 & -0.370 \\ -0.864 & 1.317 & 0.802 \\ -0.370 & 0.802 & 0.880 \end{bmatrix}$$

$$d_M(y, \mu_1) = \sqrt{(y - \mu)^T \Sigma^{-1} (y - \mu)} = \boxed{1.132}$$

$$d_M(y, \mu_2) = \boxed{0.703}$$

(c) $\bar{X} \sim \text{MVN}(\mu, \frac{1}{n}\Sigma)$

$$(\bar{X} - \mu)^T \left(\frac{1}{n}\Sigma\right)^{-1} (\bar{X} - \mu) = (\bar{X} - \mu)^T \left(\frac{1}{n}\Sigma\right)^{-1/2} \left(\frac{1}{n}\Sigma\right)^{1/2} (\bar{X} - \mu)$$

$$= \underbrace{\left[\left(\frac{1}{n}\Sigma\right)^{-1/2} (\bar{X} - \mu)\right]^T}_{\sim \text{MVN}(\vec{0}, I_n)} \underbrace{\left[\left(\frac{1}{n}\Sigma\right)^{1/2} (\bar{X} - \mu)\right]}_{\sim \text{MVN}(\vec{0}, I_n)}$$

$$\rightarrow (\bar{X} - \mu)^T \left(\frac{1}{n}\Sigma\right)^{-1} (\bar{X} - \mu) \sim \chi_n^2$$

\parallel
 $n \times$ Mahalanobis distance of \bar{X} from μ

(d) μ_2 is more plausible because it has smaller Mahalanobis distance from the sample mean.

3

$$(a) E(Y_1) = \frac{1}{5} \sum_{i=1}^5 E(X_i) = \boxed{\mu}$$

$$\begin{aligned} \Sigma_{Y_1} &= \text{Cov}\left(\frac{1}{5} \sum_{i=1}^5 X_i, \frac{1}{5} \sum_{j=1}^5 X_j\right) \\ &= \frac{1}{25} \sum_{i=1}^5 \sum_{j=1}^5 \text{Cov}(X_i, X_j) \end{aligned}$$

$$X_1, X_2, \dots, X_5 \text{ iid} \Rightarrow \text{Cov}(X_i, X_j) = 0 \quad \forall i \neq j$$

$$\Rightarrow \Sigma_{Y_1} = \boxed{\frac{1}{25} \left[\sum_{i=1}^5 \text{Cov}(X_i, X_i) \right]}$$

$$(b) E(Y_2) = \sum_{i=1}^5 (i-1) E(X_i) = \boxed{\mu}$$

$$\Sigma_{Y_2} = \boxed{\sum_{i=1}^5 \text{Cov}(X_i, X_i)}$$

4

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \\ X_4 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 4 & 4 \\ 5 & 7 \\ 4 & 7 \end{bmatrix}$$

$$\text{Likelihood: } L(\mu, \sigma^2 | X) = \prod_{i=1}^n f(X_i) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x_i - \mu)^2}{2\sigma^2}}$$

$$\begin{aligned} \text{Log-likelihood: } LL(\mu, \sigma^2 | X) &= \sum_{i=1}^n \left[-\frac{1}{2} \log(2\pi) - \frac{1}{2} \log(\sigma^2) - \frac{(x_i - \mu)^2}{2\sigma^2} \right] \\ &= -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \end{aligned}$$

$$\hat{\mu}_{MLE} = \arg\min \sum_{i=1}^n (x_i - \mu)^2 = \arg\min \sum_{i=1}^n x_i^2 - 2n\bar{x}\mu + n\mu^2 = \arg\min f(\mu)$$

$$f'(\mu) = 2n\mu - 2n\bar{x}, \quad f'(\mu) = 0 \Leftrightarrow \boxed{\mu = \bar{x}}; \quad f''(\bar{x}) = 2n > 0 \rightarrow \text{minimum} \checkmark$$

$$\rightarrow \hat{\mu}_{MLE} = \bar{x} = \boxed{\begin{bmatrix} 4 \\ 6 \end{bmatrix}}$$

4(cont)

$$\text{Similarly, } \hat{\Sigma}_{MLE} = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})(X_i - \bar{X})^T$$

$$\rightarrow \hat{\Sigma}_{MLE} = \frac{1}{4} \left\{ \begin{bmatrix} -1 \\ 0 \end{bmatrix} \begin{bmatrix} -1 & 0 \end{bmatrix} + \begin{bmatrix} 0 \\ -2 \end{bmatrix} \begin{bmatrix} 0 & -2 \end{bmatrix} + \begin{bmatrix} 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} \right\}$$

$$= \frac{1}{4} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 4 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

$$\hat{\Sigma}_{MLE} = \begin{bmatrix} 0.5 & 0.25 \\ 0.25 & 1.5 \end{bmatrix}$$

Ngoc Ha

HW 2 - ST 557

Problem 5

$$r_Q = \frac{\sum_{i=1}^n (x_{(j)} - \bar{x})(q_{(j)} - \bar{q})}{\sqrt{\sum_{i=1}^n (x_{(j)} - \bar{x})^2} \sqrt{\sum_{i=1}^n (q_{(j)} - \bar{q})^2}}; \text{ where } q_{(j)} = \phi^{-1} \left(\frac{j - \frac{1}{2}}{n} \right)$$

Prepare functions

```
In [1]: inv_cdf <- function(i,n){  
  return(qnorm((i-0.5)/n))  
}
```

```
In [2]: sampDiffVec <- function(sample){  
  return(as.matrix(sort(sample)-mean(sample)))  
}
```

```
In [3]: theoDiffVec <- function(n){  
  quantiles = rep(0,n)  
  for (i in c(1:n)){  
    quantiles[i] = inv_cdf(i,n)  
  }  
  return(as.matrix(quantiles-mean(quantiles)))  
}
```

```
In [4]: r_Q <- function(sampDiffVec, theoDiffVec){  
  return((t(sampDiffVec)%*%theoDiffVec)/(norm(sampDiffVec, type='2')*norm(th  
eoDiffVec, type='2')))  
}
```

(5a)

```
In [5]: rQVec_a <- rep(0,10000)  
for (i in c(1:10000)){  
  samp <- runif(10)  
  rQVec_a[i] <- r_Q(sampDiffVec(samp),theoDiffVec(length(samp)))  
}
```

```
In [6]: cat("Part a's rejection rate:", length(rQVec_a[rQVec_a<0.9198])/length(rQVec_a
))
```

Part a's rejection rate: 0.056

(5b)

```
In [11]: rQVec_b <- rep(0,10000)
for (i in c(1:10000)){
  samp <- rchisq(n=5,df=5)
  rQVec_b[i] <- r_Q(sampDiffVec(samp),theoDiffVec(length(samp)))
}
```

```
In [12]: cat("Part b's rejection rate:", length(rQVec_b[rQVec_b<0.8788])/length(rQVec_b
))
```

Part b's rejection rate: 0.0847

(5c)

```
In [9]: rQVec_c <- rep(0,10000)
for (i in c(1:10000)){
  samp <- rchisq(n=20,df=2)
  rQVec_c[i] <- r_Q(sampDiffVec(samp),theoDiffVec(length(samp)))
}
```

```
In [10]: cat("Part c's rejection rate:", length(rQVec_c[rQVec_c<0.9508])/length(rQVec_c
))
```

Part c's rejection rate: 0.8067