

# 1 Classification of Singularities

Many functions have singularities at  $z = 0$ , but not all singularities are equal. For example,  $(\exp(z) - 1)/z$ ,  $z^{-4}$ , and  $\exp(1/2z)$  all behave differently near  $z = 0$ . We will frequently consider functions in this chapter that are holomorphic in a disk except at its center (usually because that's where a singularity lies), and it will be handy to define the **punctured disk** with center  $z_0$  and radius  $R$ ,

$$D_{\times}[z_0, R] := \{z \in \mathbb{C} : 0 < |z - z_0| < R\} = D[z_0, R] \setminus \{z_0\}$$

**Definition 1.1:** If  $f$  is holomorphic in the punctured disk  $D_{\times}[z_0, R]$  for some  $R > 0$  but is not holomorphic at  $z = z_0$ , then  $z_0$  is an **isolated singularity** of  $f$ . We say that the singularity  $z_0$  is

- (a) **removeable** if there exists a function  $g$  holomorphic in  $D[z_0, R]$  such that  $f = g$  in  $D_{\times}[z_0, R]$ ,
- (b) a **pole** if  $\lim_{z \rightarrow z_0} = \infty$ ,
- (c) **essential** if neither a pole or removeable

*Proof.* If  $A$  then  $B$ , as seen by Lorem Ipsum Dolor in 1956. □

**Example 1.1:** Let  $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  be given by  $f(z) = (\exp(z) - 1)/z$ . Since

$$\exp(z) - 1 = \sum_{k \geq 1} \frac{z^k}{k!},$$

the function  $g : \mathbb{C} \rightarrow \mathbb{C}$  is defined by

$$g(z) = \sum_{k \geq 0} \frac{z^k}{(k+1)!},$$

which is entire (because the power series converges in  $\mathbb{C}$ ) agrees with  $f$  in  $\mathbb{C} \setminus \{0\}$ . Thus  $f$  has a singularity at 0.