

# Automatic Monte Carlo Methods for Bayesian Inference

Noah Grudowski

Applied Mathematics, Illinois Institute of Technology

## Bayesian Statistics


- Model parameters are random
- Prior** distribution,  $\pi$ , reflects beliefs about the parameters
- Sampling yields a likelihood function,  $L$
- Posterior** density: product of the prior and the likelihood
- Estimates involve integrals, e.g.

$$\hat{\beta}_j = \frac{\int_{\mathbb{R}^{d+1}} b_j L(\mathbf{b}) \pi(\mathbf{b}) d\mathbf{b}}{\int_{\mathbb{R}^{d+1}} L(\mathbf{b}) \pi(\mathbf{b}) d\mathbf{b}} =: \frac{\mu_j}{\mu} \text{ for } j = 0, 1, 2, \dots, d$$

## Quasi-Monte Carlo Cubature

- Approximates integrals:  

$$\int_{[0,1]^{d+1}} f(\mathbf{x}) d\mathbf{x} \approx \frac{1}{n} \sum_{i=1}^n f(\mathbf{x}_i)$$
 where  $\{\mathbf{x}_i\}_{i=1}^\infty$  is a low-discrepancy sequence
- A **low-discrepancy sequence** is a **correlated** sequence whose empirical distribution matches the target distribution better than an IID sequence



## Model Problem

- Bayesian inference is applied to logistic regression.
- $t_i \sim \text{Ber} \left( \frac{\exp(\beta_0 + \sum_{j=1}^d \beta_j s_{ij})}{1 + \exp(\beta_0 + \sum_{j=1}^d \beta_j s_{ij})} \right)$ , for  $i = 1, 2, \dots, M$
- Standard norm prior for  $\beta$

$$\pi(\mathbf{b}) = \frac{\exp(-\frac{1}{2}\mathbf{b}^T \mathbf{b})}{\sqrt{(2\pi)^{d+1}}}$$

- Posterior mean estimates of  $\beta$

$$\hat{\beta}_j = \frac{\int_{\mathbb{R}^{d+1}} b_j L(\mathbf{b}) \pi(\mathbf{b}) d\mathbf{b}}{\int_{\mathbb{R}^{d+1}} L(\mathbf{b}) \pi(\mathbf{b}) d\mathbf{b}} =: \frac{\mu_j}{\mu}, \text{ for } j = 0, 1, 2, \dots, d$$

A special appreciation and thank you to the IIT College of Science for providing the funding behind this research project.

## Choice of Density for Integration

- $\mu_j/\mu$  cannot be calculated analytically
- Rewrite  $\mu$  as (similarly for  $\mu_j$ ):  

$$\mu = \int_{\mathbb{R}^{d+1}} L(\mathbf{b}) \pi(\mathbf{b}) d\mathbf{b} = \int_{\mathbb{R}^{d+1}} \frac{L(\mathbf{b}) \pi(\mathbf{b})}{\rho(\mathbf{b})} \rho(\mathbf{b}) d\mathbf{b} = \int_{[0,1]^{d+1}} f(\mathbf{x}) d\mathbf{x},$$
 where  $f(\mathbf{R}(\mathbf{b})) = \frac{L(\mathbf{b}) \pi(\mathbf{b})}{\rho(\mathbf{b})}$   
 and  $\rho(\mathbf{b}) = |\partial \mathbf{R} / \partial \mathbf{b}|$  for some suitable  $\mathbf{R} : \mathbb{R}^{d+1} \rightarrow [0, 1]^{d+1}$
- Now,  $\mu_j/\mu$  can be approximated by sampling from  $\rho$  and applying quasi-Monte Carlo cubature
- We made the following choices for  $\rho$ :  
 1)  $\rho = \pi = \text{prior}$   
 2)  $\rho = \rho_{\text{MLE}} = \text{Gaussian approximation to the likelihood}$   
 3)  $\rho \propto \pi \cdot \rho_{\text{MLE}}$


## New MATLAB Function

- Developed **function** `BayesianInference` that **automatically** estimates the unknown parameters  $\hat{\beta}$  using GAIL (Choi Et al, 2015)
- User inputs error tolerance,  $d$ ,  $\rho(\mathbf{b})$ , and more

## Uniform s-Data

```
M = 100; %number of s-Data values
s = (6+2)*rand(1,M) - 2; % s-Data uniform in [-2,6]
dim=2; absTol=0.0005; n=5;
densityChoice = [true true true]; %Use all 3 densities
BayesianInference(M,s,dim,absTol,densityChoice,n)
```


Density	Runtime (seconds)	Samples Taken
$\rho = \pi$	59.0479	1572864
$\rho = \rho_{\text{MLE}}$	1.728	40960
$\rho \propto \pi \cdot \rho_{\text{MLE}}$	0.3396	40960



## Normal s-Data

```
M = 100; % number of s-Data values
s = 2.*randn(1,M)+2; % s-Data normal with mean and sd 2
dim=2; absTol=0.0005; n=5;
densityChoice = [true true true]; % Use all 3 densities
BayesianInference(M,s,dim,absTol,densityChoice,n)
```

Density	Runtime (seconds)	Samples Taken
$\rho = \pi$	22.5585	655360
$\rho = \rho_{\text{MLE}}$	0.81697	20480
$\rho \propto \pi \cdot \rho_{\text{MLE}}$	1.1508	20480



## Conclusions

- Each choice of  $\rho$  was successful in approximating posterior mean estimates within the desired tolerance level
- The choice of  $\rho$  directly affects the time taken
- Further work should be done to improve the run time of `BayesianInference` and generalize the function to solve a greater variety of problems
- These results will be compared in detail to other methods used today, specifically Markov Chain Monte Carlo

## References

- Choi, S.C.T., Ding, Y., Hickernell, F.J., Jiang, L., Jimenez Rugama, I.I.A., Tong, X., Zhang, Y., Zhou, X. *GAIL: Guaranteed Automatic Integration Library (version 2.1)*, MATLAB software, 2015.
- Hickernell, F.J., Jimenez Rugama, I.I.A., Li, D. *Contemporary Computational Mathematics — A Celebration of the 80th Birthday of Ian Sloan*. Adaptive Quasi-Monte Carlo Methods for Cubature. pages 597-619, Springer-Verlag, 2018.