




$$GMm/R^2 = mV^2/R$$


$$M(< R) \propto V^2 R$$

$$v_c(R)=\sqrt{GM(< R)/R}$$


$$\scriptscriptstyle 1$$



Solid Body Rotation




$$V \propto R$$



Wheel-like rotation


Rotation curve for wheel-like rotation

Flat Rotation Curve






Vera Rubin measured that stars in galaxies are moving faster than they should be, inferring the existence of an unseen mass we call “dark matter”






Vera Rubin measuring galaxy rotation curves (~1970)



Resulting spectrum of light within aperture



Copyright © Addison Wesley.

Milky Way Rotation Curve (updates since book)






FIG. 2.— Rotation curve of the Milky Way for Cepheids assuming $R_0 = 8.09$ kpc and $\Theta_0 = 233.6 \text{ km s}^{-1}$ (model 2). Red data points represent high mass star forming regions (Reid et al. 2014). Grey data points are taken from Sofue et al. (2009) and were rescaled to new (R_0, Θ_0) using formula $V_{\text{new}} = V_{\text{old}} + \frac{R}{8.0} (\Theta_0 - 200)$. Solid and dashed lines show the best-fitting models (linear and universal, respectively).

Rotation curve and the dark matter






https://en.wikipedia.org/wiki/Oort_constants

Klypin+2002
Data needs to be Scaled to
account for new R0




Try for yourself:

<http://wittman.physics.ucdavis.edu/Animations/RotationCurve/GalacticRotation.html>







Uniform Sphere



Isothermal Sphere

$$\rho(r) = \frac{V_o^2}{4\pi G r^2}$$




$$\Phi(r) = 4\pi G \rho_o \ln(r) = V_o^2 \ln(r)$$





Plummer model

$$\rho(r) = \frac{M}{4/3\pi a^3} (1.0 + r^2/a^2)^{-5/2}$$

$$\Phi(r) = -GM/\sqrt{r^2 + a^2}$$






NFW


$$\rho \sim 1/r^3$$

$$\rho(r) = \frac{\rho_o}{r/r_s(1+r/r_s)^2}$$



$$\Phi_{\text{NFW}}(r) = -\sigma_N^2 \frac{\ln(1 + r/r_s)}{r/r_s}$$

$$\sigma_N = 4\pi G \rho_o r_s^2$$



complications

Cores vs. Cusps

McGaugh & dBlok (2002)




Fig. 1.— The rotation curve of the low surface brightness galaxy UGC 5750. Also shown are the best fitting NFW halo parameters ($c = 2.6$, $V_{200} = 123$ km s $^{-1}$; dashed line) for the limiting case of a zero mass (minimum) disk, and what the NFW halo should look like for a galaxy of this rotation velocity in the standard Λ CDM cosmology ($c = 10$, $V_{200} = 67$ km s $^{-1}$; solid line). The excess of the solid line over the data illustrates the cuspy halo problem. Though an NFW fit can be made (dashed line), it is a poor description of the data, and requires a very low concentration ($c = 2.6$ does not occur in any plausible cosmology). These problems become more severe as allowance is made for stars (BMR; BB).

More complications

