

## 6.5 Review

### Complement

Given an event  $E$ ,

$$Prob(E) + Prob(\bar{E}) = 1$$

Where  $\bar{E}$  is the complement of the event  $E$ .

**Example question** What is the probability that for a six-sided die rolled three times the same result comes up more than once?

- What is the sample space  $S$ ?  
 $\{1, 2, 3, 4, 5, 6\}$
- What is the event  $E$  (in English)?  
The set of outcomes that use the same # more than once.
- What is the complement of  $\bar{E}$  (in English)?  
The set of outcomes that are all different numbers.
- What *structure type* is  $\bar{E}$ ? What is  $n$  and  $r$ ?  
Permutation,  $n = 6$ ,  $r = 3$
- Calculate  $Prob(\bar{E})$   
 $Prob(\bar{E}) = n(\bar{E})/n(S) = \frac{P(6,3)}{6^3} = \frac{5}{9} = 0.\bar{5}$
- Calculate the probability for the Event  $Prob(E)$  using the proposition.  
 $1 - Prob(\bar{E}) = 1 - 0.55 \approx 0.44$

### Expected (average) value

For a given probability experiment, let  $X$  be a random variable whose possible values come from the set of numbers  $x_1, \dots, x_n$ . Then the **expected value of  $X$** , denoted by  $E[X]$ , is the sum

$$E[X] = (x_1) \cdot Prob(X = x_1) + \dots + (x_n) \cdot Prob(X = x_n)$$

## 6.6 Recursion revisited

### Average trials until getting first value

A common type of problem in this section is to find the average amount of trials run until you get some value for the first time... For example, rolling a die until you get a “1” for the first time.

Let’s say there’s some probability  $p$  of success, and  $X$  is the amount of trials (rolls, flips, etc.) until the first value is received, and  $E[X]$  is the average amount of trials that will run.

We start with this formula:

$$E[X] = p(1) + (1 - p)(1 + E[X])$$

The probability should be known, so by simplifying you can solve for  $E[X]$  to find the result.

**Example 1** Find the average number of tosses of a fair coin that it takes to get a result of heads for the first time.

The probability of getting a heads is  $p = (1/2)$ , so we can write this out as:

$$E[X] = \frac{1}{2}(1) + (1 - \frac{1}{2})(1 + E[X])$$

and simplify...

$$E[X] = \frac{1}{2} + \frac{1}{2} + \frac{1}{2}(E[X])$$

$$E[X] - \frac{1}{2}(E[X]) = 1$$

$$\frac{1}{2}(E[X]) = 1$$

$$E[X] = 2$$

**Question 1**

\_\_\_\_\_ / 2

What is the expected number of rolls of a six-sided die that is rolled until a 1 appears?

---

**Question 2**

\_\_\_\_\_ / 2

A pair of dice are thrown until at least one of the die comes up 1 for the first time. How many tosses, on average, are required?

We are rolling two die, which comes out to:

(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
(4,1)	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
(5,1)	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
(6,1)	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

- What is the sample size? \_\_\_\_\_
- How many of these rules have at least one 1? \_\_\_\_\_
- What is the probability of getting at least one? \_\_\_\_\_  
( $Prob(E) = n(E)/n(S)$ )
- Use the formula to find the expected value (average trials).