

# Homework 1

Quan Nguyen

January 2026

See <https://github.com/nguyencquan/Markov2026> for code used in simulations.

## 1 Power Law Sampling

Assume we have some distribution

$$\begin{cases} Cx^{-\gamma} & x \geq x_0, \\ 0, & x < x_0 \end{cases} \quad (1)$$

**a)** Where  $x_0 > 0$  and  $\gamma > 3$

A valid pdf must have a total probability of 1, which is calculated below:

$$1 = \int_{-\infty}^{\infty} Cx^{-\gamma} dx = C \int_{x_0}^{\infty} x^{-\gamma} dx = \frac{C}{x^{\gamma-1}(\gamma-1)} \Big|_{x_0}^{\infty} = \frac{C}{x_0^{\gamma-1}(\gamma-1)} \quad (2)$$

$$C = x_0^{\gamma-1}(\gamma-1) \quad (3)$$

**b)** First calculating the cumulative distribution we take a step from equation 2 except take the integral from  $x_0$  to  $x$  instead.

$$1 = \int_{-\infty}^{\infty} Cx^{-\gamma} dx = C \int_{x_0}^x x^{-\gamma} dx = \frac{C}{x^{\gamma-1}(\gamma-1)} \Big|_{x_0}^x = \frac{C}{x_0^{\gamma-1}(\gamma-1)} + \frac{C}{x^{\gamma-1}(\gamma-1)} \quad (4)$$

Substituting in C

$$F(x) = \begin{cases} 1 - \frac{x_0^{\gamma-1}}{x^{\gamma-1}}, & x \geq x_0 \\ 0, & x < x_0 \end{cases} \quad (5)$$

To calculate the inverse, note we can do root math since  $x_0 > 0$

$$p = 1 - \frac{x_0^{\gamma-1}}{x^{\gamma-1}} \quad (6)$$

$$\frac{x_0^{\gamma-1}}{x^{\gamma-1}} = 1 - p \quad (7)$$

$$\frac{x_0}{x} = (1 - p)^{\frac{1}{\gamma-1}} \quad (8)$$

$$x = \frac{x_0}{(1 - p)^{\frac{1}{\gamma-1}}} \quad (9)$$

**c)** Also we can note that we can use  $U(0, 1)$  without  $1 - p$  to speed it up since their distribution is exactly the same.

1. Sample from a normal distribution  $U(0, 1)$

2. substitute value into p for  $\frac{x_0}{p^{\frac{1}{\gamma-1}}}$

**d)**




Figure 1: Graph of pdf and random sampling of 100 points, given  $\gamma = 4$  and  $x_0 = 10$




Figure 2: Graph of pdf and random sampling of 1000 points, given  $\gamma = 4$  and  $x_0 = 10$




Figure 3: Graph of pdf and random sampling of 10000 points, given  $\gamma = 4$  and  $x_0 = 10$

## 2 Gamma Sampling

Suppose we have  $X \sim \text{Gamma}(2, 1)$  with pdf

$$f(x) = \begin{cases} xe^{-x}, & x > 0 \\ 0, & x \leq 0 \end{cases} \quad (10)$$

**a)** To calculate the CDF we take the integral of the pdf from 0 to x. Using integration by part with  $u = x$  and  $dv = e^{-x}$

$$\int_{-\infty}^x f(x)dx = \int_0^x xe^{-x}dx = -xe^{-x} + \int_0^x e^{-x}dx \quad (11)$$

$$= -e^{-x}(x+1)|_0^x \quad (12)$$

$$= 1 - e^{-x}(x+1) \quad (13)$$

Hence we have

$$F(x) = \begin{cases} 1 - e^{-x}(x+1), & x > 0 \\ 0, & x \leq 0 \end{cases} \quad (14)$$

**b** I will be using Newtons method to find the inverse of the CDF. The method will guess a root at one and runs 5 times then repeated until the root is positive. There is a sanity check where if the root stays negative an error occurs. Fortunately the code works. The solution to the CDF will be calculated by solving the root of  $F(x) - u = 0$ . The function took 5.1287 seconds to run.

**c** To solve for the optimal value for c, I will use the inequality:

$$C \frac{1}{2} e^{-\frac{x}{2}} \geq xe^{-x} \quad (15)$$

$$C \geq 2xe^{-\frac{x}{2}} = h(x) \quad (16)$$

$$(17)$$

To solve for the critical point of the equation, take the derivative and solve for the roots

$$h'(x) = 2e^{-\frac{x}{2}} - xe^{-\frac{x}{2}} \quad (18)$$

$$= e^{-\frac{x}{2}}(2-x) = 0 \quad (19)$$

$$(20)$$

Since the derivative is a strictly decreasing function and is positive when  $x < 2$ , the critical value is the point where  $h(x)$  is the absolute maximum yielding

$$C \geq h(2) = 4/e \quad (21)$$

Since efficiency is determined by  $1/C$  the smallest possible value of C to maximize efficiency is  $4/e$ .

Also note that the given distribution  $g(x)$  is aa pdf for an exponential distribution with  $\lambda = 1/2$  Using an acceptace rejection algorithm, it took 3.1873 seconds to complete.