

# Univariate times series

In practice

V. Lefieux

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Box and Jenkins methodology

1. stationarity (if needed),
2. order selection,
3. estimation,
4. diagnostic checking,
5. final order selection,
6. forecasting,
7. ex post analysis.

## Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

Methodology

**Stationarity**

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Methods

- ▶ trend-seasonality decomposition,
- ▶ differencing transformations,
- ▶ Box-Cox transformation.

Methodology

**Stationarity**

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Differencing transformations

Methodology

**Stationarity**

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

The two most common cases:

- ▶ Time series with a  $d$ -degree polynomial:  $\nabla^d$ .
- ▶ Time series with a  $s$ -period seasonality:  $\nabla_s^D$ .

# Box-Cox transformation

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

In order to remove a “variance problem”, the Box-Cox transformation is sometimes used:

$$\frac{X_t^\lambda - 1}{\lambda}$$

with  $\lambda \in \mathbb{R}$ .

Note that:

$$\frac{X_t^\lambda - 1}{\lambda} \xrightarrow{\lambda \rightarrow 0} \ln(X_t).$$



# Remark

One can find some stationarity tests that aren't comprehensive.

Methodology

**Stationarity**

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# In practice

- ▶ A very slowly decrease of the time series “ACF” suggests differencing at lag 1.
- ▶ A very slowly decrease of the time series “ACF” every  $s$  multiple lags suggests differencing at lag  $s$ .

Differencing is iteratively done, and generally :  $d \leq 2$  and  $D \leq 2$ .

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

Methodology

Stationarity

**Order selection**

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

We consider now that the transformed (or not) time series can potentially be fitted by a zero-mean ARMA model.

To select the model order, one can use:

- ▶ confidence intervals,
- ▶ corner method,
- ▶ information criterium,
- ▶ heuristic.

# Use of confidence intervals

Idea: empirically establish maximum values for  $p$  and  $q$ .

- For a  $AR(p)$  process:

$$\forall h > p : \sqrt{n}\hat{r}(h) \xrightarrow{\mathcal{L}} \mathcal{N}(0, 1).$$

One can define a confidence interval of level 95% and search the number of lags for which 95% of the  $\hat{r}(h)$  are in  $\left[-\frac{1.96}{\sqrt{n}}, \frac{1.96}{\sqrt{n}}\right]$ .

- For a  $MA(q)$  process:

$$\forall h > q : \sqrt{n}\hat{\rho}(h) \xrightarrow{\mathcal{L}} \mathcal{N}\left(0, 1 + 2 \sum_{k=1}^q \rho^2(k)\right).$$

One can define a confidence interval of level 95% and search the number of lags for which 95% of the  $\hat{\rho}(h)$  are in:

$$\left[-\frac{1.96}{\sqrt{n}} \left(1 + 2 \sum_{k=1}^q \hat{\rho}^2(k)\right)^{\frac{1}{2}}, \frac{1.96}{\sqrt{n}} \left(1 + 2 \sum_{k=1}^q \hat{\rho}^2(k)\right)^{\frac{1}{2}}\right].$$

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

# Information criterium

Aim : search a model for which a chosen information criterium is minimal.

Methodology

Stationarity

**Order selection**

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Heuristic

Methodology

Stationarity

**Order selection**

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

In order to draw hypothesis, find “significant” autocorrelations:

- ▶ For the AR part: partial autocorrelations.
- ▶ For the MA part: (simple) autocorrelations.

# Table of contents

Methodology

Stationarity

Order selection

**Estimation**

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

Methodology

Stationarity

Order selection

**Estimation**

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis



# Estimation

Use of the maximum likelihood estimation after a preliminary estimation.

Methodology

Stationarity

Order selection

**Estimation**

Diagnostic  
checking

Final order  
selection

Forecasting

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

**Diagnostic checking**

Final order selection

Forecasting

Ex post analysis

Methodology

Stationarity

Order selection

Estimation

**Diagnostic  
checking**

Final order  
selection

Forecasting

Ex post analysis

# Aim

- ▶ significance test of the parameters,
- ▶ whiteness and normality of the residuals.

Methodology

Stationarity

Order selection

Estimation

**Diagnostic  
checking**

Final order  
selection

Forecasting

Ex post analysis

# Significance test of the parameters

Methodology

Stationarity

Order selection

Estimation

**Diagnostic  
checking**

Final order  
selection

Forecasting

Ex post analysis

For example, we consider the following test for  $\varphi_p$ :

$$\begin{cases} H_0 : \text{the process order is ARMA}(p-1, q) \ (\varphi_p = 0) \\ H_1 : \text{the process order is ARMA}(p, q) \ (\varphi_p \neq 0) \end{cases}.$$

We use the Student statistic:

$$t = \frac{|\hat{\varphi}_p|}{\sqrt{\widehat{\text{Var}}(\hat{\varphi}_p)}}.$$

We reject  $H_0$  at the 5% level if  $|t| > 1.96$ .

# Whiteness and normality of the residuals

- ▶ Whiteness: LjungBox test.
- ▶ Normality: Shapiro-Wilk.

Note that if the normality of the residuals is rejected, it could be useful to add an ARCH or GARCH part. . .

Methodology

Stationarity

Order selection

Estimation

**Diagnostic  
checking**

Final order  
selection

Forecasting

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

**Final order selection**

Forecasting

Ex post analysis

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

**Final order  
selection**

Forecasting

Ex post analysis

# Final order selection

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

**Final order  
selection**

Forecasting

Ex post analysis

There are two main ways to finally select a model:

- ▶ Information criterium (e.g. AIC or BIC)).
- ▶ Predictive power.

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

**Forecasting**

Ex post analysis

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

**Forecasting**

Ex post analysis



# Forecasting

The forecasting and the forecasting interval are obtained using the autoregressive and moving average representations.

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

**Forecasting**

Ex post analysis

# Table of contents

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis

Methodology

Stationarity

Order selection

Estimation

Diagnostic  
checking

Final order  
selection

Forecasting

**Ex post analysis**

# Ex post analysis

Idea: truncate the time series and measure the forecasting error with an indicator such as the Root Mean Square Error (RMSE) or the Mean Average Percentage Error (MAPE):

$$\text{RMSE} = \sqrt{\frac{1}{T} \sum_{t=1}^T (x_t - \hat{x}_t)^2},$$

$$\text{MAPE} = \frac{1}{T} \sum_{t=1}^T \left| \frac{x_t - \hat{x}_t}{x_t} \right|.$$

Methodology

Stationarity

Order selection

Estimation

Diagnostic checking

Final order selection

Forecasting

Ex post analysis