

52nd—24th INTERNATIONAL-RUDOLF ORTVAY PROBLEM SOLVING CONTEST IN PHYSICS

Problem 14

Nguyen Thanh Long

April 30, 2022

Using the Ampere's law, we know that the magnetic field inside the soleloid (by the soleloid) is

$$B_0 = \frac{\mu_0 N I}{l}. \quad (1)$$

We can denote easily that in the homogeneous magnetic field, a tiny small iron sphere with relative permeability μ_r will have the magnetic dipole [1] :

$$m = V \frac{3(\mu_r - 1) B_0}{\mu_0 (\mu_r + 2)}. \quad (2)$$

The magnetic flux that the dipole effect on a circle wire distance z from the dipole is:

$$\frac{d\Phi(z)}{dN} = \int_0^r \frac{\mu_0}{4\pi} \frac{3z^2 - (z^2 + r^2)}{(z^2 + r^2)^{\frac{5}{2}}} \cdot 2\pi r dr = \frac{\mu_0 m r^2}{2(z^2 + r^2)^{\frac{3}{2}}}.$$

Where $dN = \frac{N}{l} dz$, so the total magnetic flux of the soleloid (caused by dipole) will be:

$$\Phi = \int_{-\infty}^{\infty} \frac{N}{l} \frac{\mu_0 m r^2}{2(z^2 + r^2)^{\frac{3}{2}}} dz = \frac{N \mu_0 m}{l}. \quad (3)$$

From (1), (2) and (3), we have:

$$\Phi = \frac{3\mu_0 (\mu_r - 1) V N^2}{(\mu_r + 2) l^2} I.$$

Therefore, the self-inductance change of the solenoid is;

$$\Delta L = \frac{3\mu_0 (\mu_r - 1) V N^2}{(\mu_r + 2) l^2}.$$

With $\mu_r \gg 1$, the result will be:

$$\Delta L = \frac{3\mu_0 V N^2}{l^2}.$$

References

- [1] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (John Wiley & Sons, New York, NY, 1998).