

A TUTORIAL ON PRINCIPAL COMPONENT ANALYSIS

DERIVATION, DISCUSSION AND SINGULAR VALUE DECOMPOSITION

Jon Shlens | jonshlens@ucsd.edu

25 March 2003 | Version 1

Principal component analysis (PCA) is a mainstay of modern data analysis - a black box that is widely used but poorly understood. The goal of this paper is to dispel the magic behind this black box. This tutorial focuses on building a solid intuition for how and why principal component analysis works; furthermore, it crystallizes this knowledge by deriving from first principles, the mathematics behind PCA. This tutorial does not shy away from explaining the ideas informally, nor does it shy away from the mathematics. The hope is that by addressing both aspects, readers of all levels will be able to gain a better understanding of the power of PCA as well as the when, the how and the why of applying this technique.

1 OVERVIEW

Principal component analysis (*PCA*) has been called one of the most valuable results from applied linear algebra. *PCA* is used abundantly in all forms of analysis - from neuroscience to computer graphics - because it is a simple, non-parametric method of extracting relevant information from confusing data sets. With minimal additional effort *PCA* provides a roadmap for how to reduce a complex data set to a lower dimension to reveal the sometimes hidden, simplified dynamics that often underlie it.

The goal of this tutorial is to provide both an intuitive feel for *PCA*, and a thorough discussion of this topic. We will begin with a simple example and provide an intuitive explanation of the goal of *PCA*. We will continue by adding mathematical rigor to place it within the framework of linear algebra and explicitly solve this problem. We will see how and why *PCA* is intimately related to the mathematical technique of singular value decomposition (*SVD*). This understanding will lead us to a prescription for how to apply *PCA* in the real world. We will discuss both

the assumptions behind this technique as well as possible extensions to overcome these limitations.

The discussion and explanations in this paper are informal in the spirit of a tutorial. The goal of this paper is to *educate*. Occasionally, rigorous mathematical proofs are necessary although relegated to the Appendix. Although not as vital to the tutorial, the proofs are presented for the adventurous reader who desires a more complete understanding of the math. The only assumption is that the reader has a working knowledge of linear algebra. Nothing more. Please feel free to contact me with any suggestions, corrections or comments.

2 MOTIVATION: A TOY EXAMPLE

Here is the perspective: we are an experimenter. We are trying to understand some phenomenon by measuring various quantities (e.g. spectra, voltages, velocities, etc.) in our system. Unfortunately, we can not figure out what is happening because the data appears clouded, unclear and even redundant. This is not a trivial problem, but rather a fundamental obstacle to experimental science. Examples abound from complex systems such as neuroscience, photo-science, meteorology and oceanography - the number of variables to measure can be unwieldy and at times even *deceptive*, because the underlying dynamics can often be quite simple.

Take for example a simple toy problem from physics diagrammed in Figure 1. Pretend we are studying the motion of the physicist's ideal spring. This system consists of a ball of mass m attached to a *massless, frictionless* spring. The ball is released a small distance away from equilibrium (i.e. the spring is stretched). Because the spring is "ideal," it oscillates indefinitely along the x -axis about its equilibrium at a set frequency.

This is a standard problem in physics in which the




Figure 1: A diagram of the toy example.

motion along the x direction is solved by an explicit function of time. In other words, the underlying dynamics can be expressed as a function of a single variable x .

However, being ignorant experimenters we do not know any of this. We do not know which, let alone how many, axes and dimensions are important to measure. Thus, we decide to measure the ball’s position in a three-dimensional space (since we live in a three dimensional world). Specifically, we place three movie cameras around our system of interest. At 200 Hz each movie camera records an image indicating a two dimensional position of the ball (a projection). Unfortunately, because of our ignorance, we do not even know what are the *real* “ x ”, “ y ” and “ z ” axes, so we choose three camera axes $\{\vec{a}, \vec{b}, \vec{c}\}$ at some arbitrary angles with respect to the system. The angles between our measurements might not even be 90° ! Now, we record with the cameras for 2 minutes. The big question remains: *how do we get from this data set to a simple equation of x ?*

We know a-priori that if we were smart experimenters, we would have just measured the position along the x -axis with one camera. But this is not what happens in the real world. We often do not know what measurements best reflect the dynamics of our system in question. Furthermore, we sometimes record more dimensions than we actually need!

Also, we have to deal with that pesky, real-world problem of *noise*. In the toy example this means

that we need to deal with air, imperfect cameras or even friction in a less-than-ideal spring. Noise contaminates our data set only serving to obfuscate the dynamics further. *This toy example is the challenge experimenters face everyday.* We will refer to this example as we delve further into abstract concepts. Hopefully, by the end of this paper we will have a good understanding of how to systematically extract x using principal component analysis.

3 FRAMEWORK: CHANGE OF BASIS

The Goal: Principal component analysis computes the most meaningful *basis* to re-express a noisy, garbled data set. The hope is that this new basis will filter out the noise and reveal hidden dynamics. In the example of the spring, the explicit goal of *PCA* is to determine: “the dynamics are along the x -axis.” In other words, the goal of *PCA* is to determine that \hat{x} - the unit basis vector along the x -axis - is the important dimension. Determining this fact allows an experimenter to discern which dynamics are important and which are just redundant.

3.1 A NAIVE BASIS

With a more precise definition of our goal, we need a more precise definition of our data as well. For each time sample (or experimental trial), an experimenter records a set of data consisting of multiple measurements (e.g. voltage, position, etc.). The number of measurement types is the *dimension* of the data set. In the case of the spring, this data set has 12,000 6-dimensional vectors, where each camera contributes a 2-dimensional projection of the ball’s position.

In general, each data sample is a vector in m -dimensional space, where m is the number of measurement types. Equivalently, every time sample is a vector that lies in an m -dimensional *vector space* spanned by an orthonormal basis. All measurement vectors in this space are a linear combination of this set of unit length basis vectors. A naive and simple

choice of a basis \mathbf{B} is the identity matrix \mathbf{I} .

$$\mathbf{B} = \begin{bmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \\ \vdots \\ \mathbf{b}_m \end{bmatrix} = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix} = \mathbf{I}$$

where each *row* is a basis vector \mathbf{b}_i with m components.

To summarize, at one point in time, camera A records a corresponding position $(x_A(t), y_A(t))$. Each trial can be expressed as a six dimensional column vector \vec{X} .

$$\vec{X} = \begin{bmatrix} x_A \\ y_A \\ x_B \\ y_B \\ x_C \\ y_C \end{bmatrix}$$

where each camera contributes two points and the entire vector \vec{X} is the set of coefficients in the naive basis \mathbf{B} .

3.2 CHANGE OF BASIS

With this rigor we may now state more precisely what *PCA* asks: *Is there another basis, which is a linear combination of the original basis, that best re-expresses our data set?*

A close reader might have noticed the conspicuous addition of the word *linear*. Indeed, *PCA* makes one stringent but powerful assumption: *linearity*. Linearity vastly simplifies the problem by (1) restricting the set of potential bases, and (2) formalizing the implicit assumption of continuity in a data set. *A subtle point it is, but we have already assumed linearity by implicitly stating that the data set even characterizes the dynamics of the system!* In other words, we are already relying on the superposition principal of linearity to believe that the data characterizes or provides an ability to interpolate between the individual data points¹.

¹To be sure, complex systems are almost always nonlinear and often their main qualitative features are a direct result of their nonlinearity. However, locally linear approximations usually provide a good approximation because non-linear, higher order terms vanish at the limit of small perturbations.

With this assumption *PCA* is now limited to re-expressing the data as a *linear combination* of its basis vectors. Let \mathbf{X} and \mathbf{Y} be $m \times n$ matrices related by a linear transformation \mathbf{P} . \mathbf{X} is the original recorded data set and \mathbf{Y} is a re-representation of that data set.

$$\mathbf{P}\mathbf{X} = \mathbf{Y} \quad (1)$$

Also let us define the following quantities².

- \mathbf{p}_i are the *rows* of \mathbf{P}
- \mathbf{x}_i are the *columns* of \mathbf{X}
- \mathbf{y}_i are the *columns* of \mathbf{Y} .

Equation 1 represents a *change of basis* and thus can have many interpretations.

1. \mathbf{P} is a matrix that transforms \mathbf{X} into \mathbf{Y} .
2. Geometrically, \mathbf{P} is a rotation and a stretch which again transforms \mathbf{X} into \mathbf{Y} .
3. The rows of \mathbf{P} , $\{\mathbf{p}_1, \dots, \mathbf{p}_m\}$, are a set of new basis vectors for expressing the *columns* of \mathbf{X} .

The latter interpretation is not obvious but can be seen by writing out the explicit dot products of $\mathbf{P}\mathbf{X}$.

$$\begin{aligned} \mathbf{P}\mathbf{X} &= \begin{bmatrix} \mathbf{p}_1 \\ \vdots \\ \mathbf{p}_m \end{bmatrix} \begin{bmatrix} \mathbf{x}_1 & \cdots & \mathbf{x}_n \end{bmatrix} \\ \mathbf{Y} &= \begin{bmatrix} \mathbf{p}_1 \cdot \mathbf{x}_1 & \cdots & \mathbf{p}_1 \cdot \mathbf{x}_n \\ \vdots & \ddots & \vdots \\ \mathbf{p}_m \cdot \mathbf{x}_1 & \cdots & \mathbf{p}_m \cdot \mathbf{x}_n \end{bmatrix} \end{aligned}$$

We can note the form of each column of \mathbf{Y} .

$$\mathbf{y}_i = \begin{bmatrix} \mathbf{p}_1 \cdot \mathbf{x}_i \\ \vdots \\ \mathbf{p}_m \cdot \mathbf{x}_i \end{bmatrix}$$

We recognize that each coefficient of \mathbf{y}_i is a dot-product of \mathbf{x}_i with the corresponding row in \mathbf{P} . In

²In this section \mathbf{x}_i and \mathbf{y}_i are *column* vectors, but be forewarned. In all other sections \mathbf{x}_i and \mathbf{y}_i are *row* vectors.

other words, the j^{th} coefficient of \mathbf{y}_i is a projection on to the j^{th} row of \mathbf{P} . This is in fact the very form of an equation where \mathbf{y}_i is a projection on to the basis of $\{\mathbf{p}_1, \dots, \mathbf{p}_m\}$. Therefore, the *rows* of \mathbf{P} are indeed a new set of basis vectors for representing of *columns* of \mathbf{X} .

3.3 QUESTIONS REMAINING

By assuming linearity the problem reduces to finding the appropriate *change of basis*. The row vectors $\{\mathbf{p}_1, \dots, \mathbf{p}_m\}$ in this transformation will become the *principal components* of \mathbf{X} . Several questions now arise.

- What is the best way to “re-express” \mathbf{X} ?
- What is a good choice of basis \mathbf{P} ?

These questions must be answered by next asking ourselves what features we would like \mathbf{Y} to exhibit. Evidently, additional assumptions beyond *linearity* are required to arrive at a reasonable result. The selection of these assumptions is the subject of the next section.

4 VARIANCE AND THE GOAL

Now comes the most important question: *what does “best express” the data mean?* This section will build up an intuitive answer to this question and along the way tack on additional assumptions. The end of this section will conclude with a mathematical goal for deciphering “garbled” data.

In a linear system “garbled” can refer to only two potential confounds: *noise* and *redundancy*. Let us deal with each situation individually.

4.1 NOISE

Noise in any data set must be low or - no matter the analysis technique - no information about a system can be extracted. There exists no absolute scale for noise but rather all noise is measured relative to the




Figure 2: A simulated plot of (x_A, y_A) for camera A . The signal and noise variances σ_{signal}^2 and σ_{noise}^2 are graphically represented.

measurement. A common measure is the *signal-to-noise ratio* (SNR), or a ratio of variances σ^2 .

$$SNR = \frac{\sigma_{signal}^2}{\sigma_{noise}^2} \quad (2)$$

A high SNR ($\gg 1$) indicates high precision data, while a low SNR indicates noise contaminated data.

Pretend we plotted all data from camera A from the spring in Figure 2. Any individual camera should record motion in a straight line. Therefore, any spread deviating from straight-line motion must be noise. The variance due to the signal and noise are indicated in the diagram graphically. The ratio of the two, the SNR , thus measures how “fat” the oval is - a range of possibilities include a thin line ($SNR \gg 1$), a perfect circle ($SNR = 1$) or even worse. In summary, we must assume that our measurement devices are reasonably good. Quantitatively, this corresponds to a high SNR .

4.2 REDUNDANCY

Redundancy is a more tricky issue. This issue is particularly evident in the example of the spring. In this case multiple sensors record the same dynamic




Figure 3: A spectrum of possible redundancies in data from the two separate recordings r_1 and r_2 (e.g. x_A, y_B). The best-fit line $r_2 = kr_1$ is indicated by the dashed line.

information. Consider Figure 3 as a range of possible plots between two arbitrary measurement types r_1 and r_2 . Panel (a) depicts two recordings with no redundancy. In other words, r_1 is entirely uncorrelated with r_2 . This situation could occur by plotting two variables such as $(x_A, \text{humidity})$.

However, in panel (c) both recordings appear to be strongly related. This extremity might be achieved by several means.

- A plot of (x_A, \tilde{x}_A) where x_A is in meters and \tilde{x}_A is in inches.
- A plot of (x_A, x_B) if cameras A and B are very nearby.

Clearly in panel (c) it would be more meaningful to just have recorded a single variable, the linear combination $r_2 - kr_1$, instead of two variables r_1 and r_2 separately. Recording solely the linear combination $r_2 - kr_1$ would both express the data more concisely and reduce the number of sensor recordings ($2 \rightarrow 1$ variables). Indeed, this is the very idea behind dimensional reduction.

4.3 COVARIANCE MATRIX

The *SNR* is solely determined by calculating variances. A corresponding but simple way to quantify the redundancy between individual recordings is to

calculate something like the variance. I say “something like” because the variance is the spread due to one variable but we are concerned with the spread between variables.

Consider two sets of simultaneous measurements with zero mean³.

$$A = \{a_1, a_2, \dots, a_n\}, \quad B = \{b_1, b_2, \dots, b_n\}$$

The variance of A and B are individually defined as follows.

$$\sigma_A^2 = \langle a_i a_i \rangle_i, \quad \sigma_B^2 = \langle b_i b_i \rangle_i$$

where the expectation⁴ is the average over n variables. The *covariance* between A and B is a straightforward generalization.

$$\text{covariance of } A \text{ and } B \equiv \sigma_{AB}^2 = \langle a_i b_i \rangle_i$$

Two important facts about the covariance.

- $\sigma_{AB}^2 = 0$ if and only if A and B are entirely uncorrelated.
- $\sigma_{AB}^2 = \sigma_A^2$ if $A = B$.

We can equivalently convert the sets of A and B into corresponding row vectors.

$$\begin{aligned} \mathbf{a} &= [a_1 \ a_2 \ \dots \ a_n] \\ \mathbf{b} &= [b_1 \ b_2 \ \dots \ b_n] \end{aligned}$$

so that we may now express the covariance as a dot product matrix computation.

$$\sigma_{\mathbf{ab}}^2 \equiv \frac{1}{n-1} \mathbf{ab}^T \quad (3)$$

where the beginning term is a constant for normalization⁵.

Finally, we can generalize from two vectors to an arbitrary number. We can rename the row vectors $\mathbf{x}_1 \equiv \mathbf{a}$, $\mathbf{x}_2 \equiv \mathbf{b}$ and consider additional indexed row

³These data sets are in *mean deviation form* because the means have been subtracted off or are zero.

⁴ $\langle \cdot \rangle_i$ denotes the average over values indexed by i .

⁵The simplest possible normalization is $\frac{1}{n}$. However, this provides a biased estimation of variance particularly for small n . It is beyond the scope of this paper to show that the proper normalization for an unbiased estimator is $\frac{1}{n-1}$.

vectors $\mathbf{x}_3, \dots, \mathbf{x}_m$. Now we can define a new $m \times n$ matrix \mathbf{X} .

$$\mathbf{X} = \begin{bmatrix} \mathbf{x}_1 \\ \vdots \\ \mathbf{x}_m \end{bmatrix} \quad (4)$$

One interpretation of \mathbf{X} is the following. Each *row* of \mathbf{X} corresponds to all measurements of a particular type (\mathbf{x}_i). Each *column* of \mathbf{X} corresponds to a set of measurements from one particular trial (this is \vec{X} from section 3.1). We now arrive at a definition for the *covariance matrix* $\mathbf{S}_\mathbf{X}$.

$$\mathbf{S}_\mathbf{X} \equiv \frac{1}{n-1} \mathbf{X} \mathbf{X}^T \quad (5)$$

Consider how the matrix form $\mathbf{X} \mathbf{X}^T$, in that order, computes the desired value for the ij^{th} element of $\mathbf{S}_\mathbf{X}$. Specifically, the ij^{th} element of the variance is the dot product between the vector of the i^{th} measurement type with the vector of the j^{th} measurement type.

- The ij^{th} value of $\mathbf{X} \mathbf{X}^T$ is equivalent to substituting \mathbf{x}_i and \mathbf{x}_j into Equation 3.
- $\mathbf{S}_\mathbf{X}$ is a square symmetric $m \times m$ matrix.
- The diagonal terms of $\mathbf{S}_\mathbf{X}$ are the *variance* of particular measurement types.
- The off-diagonal terms of $\mathbf{S}_\mathbf{X}$ are the *covariance* between measurement types.

Computing $\mathbf{S}_\mathbf{X}$ quantifies the correlations between all possible pairs of measurements. Between one pair of measurements, a large covariance corresponds to a situation like panel (c) in Figure 3, while zero covariance corresponds to entirely uncorrelated data as in panel (a).

$\mathbf{S}_\mathbf{X}$ is special. The covariance matrix describes all relationships between pairs of measurements in our data set. Pretend we have the option of manipulating $\mathbf{S}_\mathbf{X}$. We will suggestively define our manipulated covariance matrix $\mathbf{S}_\mathbf{Y}$. What features do we want to optimize in $\mathbf{S}_\mathbf{Y}$?

4.4 DIAGONALIZE THE COVARIANCE MATRIX

If our goal is to reduce redundancy, then we would like each variable to co-vary as little as possible with other variables. More precisely, to remove redundancy we would like the covariances between separate measurements to be zero. What would the optimized covariance matrix $\mathbf{S}_\mathbf{Y}$ look like? Evidently, in an “optimized” matrix all off-diagonal terms in $\mathbf{S}_\mathbf{Y}$ are zero. Therefore, removing redundancy diagonalizes $\mathbf{S}_\mathbf{Y}$.

There are many methods for diagonalizing $\mathbf{S}_\mathbf{Y}$. It is curious to note that *PCA* arguably selects the easiest method, perhaps accounting for its widespread application.

PCA assumes that all basis vectors $\{\mathbf{p}_1, \dots, \mathbf{p}_m\}$ are orthonormal (i.e. $\mathbf{p}_i \cdot \mathbf{p}_j = \delta_{ij}$). In the language of linear algebra, *PCA* assumes \mathbf{P} is an orthonormal matrix. Secondly *PCA* assumes the directions with the largest variances are the most “important” or in other words, most *principal*. Why are these assumptions easiest?

Envision how *PCA* might work. By the variance assumption *PCA* first selects a normalized direction in m -dimensional space along which the variance in \mathbf{X} is maximized - it saves this as \mathbf{p}_1 . Again it finds another direction along which variance is maximized, however, because of the orthonormality condition, it restricts its search to all directions perpendicular to all previous selected directions. This could continue until m directions are selected. The resulting ordered set of \mathbf{p} ’s are the *principal components*.

In principle this simple pseudo-algorithm works, however that would bely the true reason why the orthonormality assumption is particularly judicious. Namely, the true benefit to this assumption is that *it makes the solution amenable to linear algebra*. There exist decompositions that can provide efficient, explicit algebraic solutions.

Notice what we also gained with the second assumption. We have a method for judging the “importance” of the each principal direction. Namely, the variances associated with each direction \mathbf{p}_i quantify how principal each direction is. We could thus rank-order each basis vector \mathbf{p}_i according to the corresponding variances.

We will now pause to review the implications of all

the assumptions made to arrive at this mathematical goal.

4.5 SUMMARY OF ASSUMPTIONS AND LIMITS

This section provides an important context for understanding when *PCA* might perform poorly as well as a road map for understanding new extensions to *PCA*. The final section provides a more lengthy discussion about recent extensions.

I. Linearity

Linearity frames the problem as a change of basis. Several areas of research have explored how applying a nonlinearity prior to performing *PCA* could extend this algorithm - this has been termed *kernel PCA*.

II. Mean and variance are sufficient statistics.

The formalism of *sufficient statistics* captures the notion that the mean and the variance entirely describe a probability distribution. The only zero-mean probability distribution that is fully described by the variance is the Gaussian distribution.

In order for this assumption to hold, the probability distribution of \mathbf{x}_i must be Gaussian. Deviations from a Gaussian could invalidate this assumption⁶. On the flip side, this assumption formally guarantees that the *SNR* and the covariance matrix fully characterize the noise and redundancies.

III. Large variances have important dynamics.

This assumption also encompasses the belief

⁶**A sidebar question:** What if \mathbf{x}_i are not Gaussian distributed? Diagonalizing a covariance matrix might not produce satisfactory results. The most rigorous form of removing redundancy is statistical independence.

$$P(\mathbf{y}_1, \mathbf{y}_2) = P(\mathbf{y}_1)P(\mathbf{y}_2)$$

where $P(\cdot)$ denotes the probability density. The class of algorithms that attempt to find the $\mathbf{y}_1, \dots, \mathbf{y}_m$ that satisfy this statistical constraint are termed independent component analysis (*ICA*). In practice though, quite a lot of real world data are Gaussian distributed - thanks to the Central Limit Theorem - and *PCA* is usually a robust solution to slight deviations from this assumption.

that the data has a high *SNR*. Hence, principal components with larger associated variances represent interesting dynamics, while those with lower variances represent noise.

IV. The principal components are orthogonal.

This assumption provides an intuitive simplification that makes *PCA* soluble with linear algebra decomposition techniques. These techniques are highlighted in the two following sections.

We have discussed all aspects of deriving *PCA* - what remains is the linear algebra solutions. The first solution is somewhat straightforward while the second solution involves understanding an important decomposition.

5 SOLVING PCA: EIGENVECTORS OF COVARIANCE

We will derive our first algebraic solution to *PCA* using linear algebra. This solution is based on an important property of eigenvector decomposition. Once again, the data set is \mathbf{X} , an $m \times n$ matrix, where m is the number of measurement types and n is the number of data trials. The goal is summarized as follows.

Find some orthonormal matrix \mathbf{P} where $\mathbf{Y} = \mathbf{P}\mathbf{X}$ such that $\mathbf{S}_\mathbf{Y} \equiv \frac{1}{n-1}\mathbf{Y}\mathbf{Y}^T$ is diagonalized. The rows of \mathbf{P} are the *principal components* of \mathbf{X} .

We begin by rewriting $\mathbf{S}_\mathbf{Y}$ in terms of our variable of choice \mathbf{P} .

$$\begin{aligned} \mathbf{S}_\mathbf{Y} &= \frac{1}{n-1}\mathbf{Y}\mathbf{Y}^T \\ &= \frac{1}{n-1}(\mathbf{P}\mathbf{X})(\mathbf{P}\mathbf{X})^T \\ &= \frac{1}{n-1}\mathbf{P}\mathbf{X}\mathbf{X}^T\mathbf{P}^T \\ &= \frac{1}{n-1}\mathbf{P}(\mathbf{X}\mathbf{X}^T)\mathbf{P}^T \\ \mathbf{S}_\mathbf{Y} &= \frac{1}{n-1}\mathbf{P}\mathbf{A}\mathbf{P}^T \end{aligned}$$

Note that we defined a new matrix $\mathbf{A} \equiv \mathbf{X}\mathbf{X}^T$, where \mathbf{A} is *symmetric* (by Theorem 2 of Appendix A).

Our roadmap is to recognize that a symmetric matrix (\mathbf{A}) is diagonalized by an orthogonal matrix of its eigenvectors (by Theorems 3 and 4 from Appendix A). For a symmetric matrix \mathbf{A} Theorem 4 provides:

$$\mathbf{A} = \mathbf{E}\mathbf{D}\mathbf{E}^T \quad (6)$$

where \mathbf{D} is a diagonal matrix and \mathbf{E} is a matrix of eigenvectors of \mathbf{A} arranged as columns.

The matrix \mathbf{A} has $r \leq m$ orthonormal eigenvectors where r is the rank of the matrix. The rank of \mathbf{A} is less than m when \mathbf{A} is *degenerate* or all data occupy a subspace of dimension $r \leq m$. Maintaining the constraint of orthogonality, we can remedy this situation by selecting $(m - r)$ additional orthonormal vectors to “fill up” the matrix \mathbf{E} . These additional vectors do not effect the final solution because the variances associated with these directions are zero.

Now comes the trick. *We select the matrix \mathbf{P} to be a matrix where each row \mathbf{p}_i is an eigenvector of $\mathbf{X}\mathbf{X}^T$.* By this selection, $\mathbf{P} \equiv \mathbf{E}^T$. Substituting into Equation 6, we find $\mathbf{A} = \mathbf{P}^T\mathbf{D}\mathbf{P}$. With this relation and Theorem 1 of Appendix A ($\mathbf{P}^{-1} = \mathbf{P}^T$) we can finish evaluating \mathbf{S}_Y .

$$\begin{aligned} \mathbf{S}_Y &= \frac{1}{n-1} \mathbf{P}\mathbf{A}\mathbf{P}^T \\ &= \frac{1}{n-1} \mathbf{P}(\mathbf{P}^T\mathbf{D}\mathbf{P})\mathbf{P}^T \\ &= \frac{1}{n-1} (\mathbf{P}\mathbf{P}^T)\mathbf{D}(\mathbf{P}\mathbf{P}^T) \\ &= \frac{1}{n-1} (\mathbf{P}\mathbf{P}^{-1})\mathbf{D}(\mathbf{P}\mathbf{P}^{-1}) \\ \mathbf{S}_Y &= \frac{1}{n-1} \mathbf{D} \end{aligned}$$

It is evident that the choice of \mathbf{P} diagonalizes \mathbf{S}_Y . This was the goal for *PCA*. We can summarize the results of *PCA* in the matrices \mathbf{P} and \mathbf{S}_Y .

- The principal components of \mathbf{X} are the eigenvectors of $\mathbf{X}\mathbf{X}^T$; or the rows of \mathbf{P} .
- The i^{th} diagonal value of \mathbf{S}_Y is the variance of \mathbf{X} along \mathbf{p}_i .

In practice computing *PCA* of a data set \mathbf{X} entails (1) subtracting off the mean of each measurement type and (2) computing the eigenvectors of $\mathbf{X}\mathbf{X}^T$. This solution is encapsulated in demonstration Matlab code included in Appendix B.

6 A MORE GENERAL SOLUTION: SINGULAR VALUE DECOMPOSITION

This section is the most mathematically involved and can be skipped without much loss of continuity. It is presented solely for completeness. We derive another algebraic solution for *PCA* and in the process, find that *PCA* is closely related to singular value decomposition (*SVD*). In fact, the two are so intimately related that the names are often used interchangeably. What we will see though is that *SVD* is a more general method of understanding *change of basis*.

We begin by quickly deriving the decomposition. In the following section we interpret the decomposition and in the last section we relate these results to *PCA*.

6.1 SINGULAR VALUE DECOMPOSITION

Let \mathbf{X} be an arbitrary $n \times m$ matrix⁷ and $\mathbf{X}^T\mathbf{X}$ be a rank r , square, symmetric $n \times n$ matrix. In a seemingly unmotivated fashion, let us define all of the quantities of interest.

- $\{\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \dots, \hat{\mathbf{v}}_r\}$ is the set of *orthonormal* $m \times 1$ eigenvectors with associated eigenvalues $\{\lambda_1, \lambda_2, \dots, \lambda_r\}$ for the symmetric matrix $\mathbf{X}^T\mathbf{X}$.

$$(\mathbf{X}^T\mathbf{X})\hat{\mathbf{v}}_i = \lambda_i\hat{\mathbf{v}}_i$$

- $\sigma_i \equiv \sqrt{\lambda_i}$ are positive real and termed the *singular values*.

- $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_r\}$ is the set of *orthonormal* $n \times 1$ vectors defined by $\hat{\mathbf{u}}_i \equiv \frac{1}{\sigma_i} \mathbf{X}\hat{\mathbf{v}}_i$.

We obtain the final definition by referring to Theorem 5 of Appendix A. The final definition includes two new and unexpected properties.

⁷Notice that in this section only we are reversing convention from $m \times n$ to $n \times m$. The reason for this derivation will become clear in section 6.3.

- $\hat{\mathbf{u}}_i \cdot \hat{\mathbf{u}}_j = \delta_{ij}$
- $\|\mathbf{X}\hat{\mathbf{v}}_i\| = \sigma_i$

These properties are both proven in Theorem 5. We now have all of the pieces to construct the decomposition. The “value” version of singular value decomposition is just a restatement of the third definition.

$$\mathbf{X}\hat{\mathbf{v}}_i = \sigma_i\hat{\mathbf{u}}_i \quad (7)$$

This result says a quite a bit. \mathbf{X} multiplied by an eigenvector of $\mathbf{X}^T\mathbf{X}$ is equal to a scalar times another vector. The set of eigenvectors $\{\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \dots, \hat{\mathbf{v}}_r\}$ and the set of vectors $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_r\}$ are both orthonormal sets or bases in r -dimensional space.

We can summarize this result for all vectors in one matrix multiplication by following the prescribed construction in Figure 4. We start by constructing a new diagonal matrix Σ .

$$\Sigma \equiv \begin{bmatrix} \sigma_1 & & & & 0 \\ & \ddots & & & \\ & & \sigma_r & & \\ 0 & & & 0 & \\ & & & & \ddots \\ & & & & & 0 \end{bmatrix}$$

where $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r$ are the rank-ordered set of singular values. Likewise we construct accompanying orthogonal matrices \mathbf{V} and \mathbf{U} .

$$\begin{aligned} \mathbf{V} &= [\hat{\mathbf{v}}_1 \ \hat{\mathbf{v}}_2 \ \dots \ \hat{\mathbf{v}}_m] \\ \mathbf{U} &= [\hat{\mathbf{u}}_1 \ \hat{\mathbf{u}}_2 \ \dots \ \hat{\mathbf{u}}_n] \end{aligned}$$

where we have appended an additional $(m - r)$ and $(n - r)$ orthonormal vectors to “fill up” the matrices for \mathbf{V} and \mathbf{U} respectively⁸. Figure 4 provides a graphical representation of how all of the pieces fit together to form the matrix version of *SVD*.

$$\mathbf{X}\mathbf{V} = \mathbf{U}\Sigma \quad (8)$$

where each column of \mathbf{V} and \mathbf{U} perform the “value” version of the decomposition (Equation 7). Because

⁸This is the same procedure used to fix the degeneracy in the previous section.

\mathbf{V} is orthogonal, we can multiply both sides by $\mathbf{V}^{-1} = \mathbf{V}^T$ to arrive at the final form of the decomposition.

$$\mathbf{X} = \mathbf{U}\Sigma\mathbf{V}^T \quad (9)$$

Although it was derived without motivation, this decomposition is quite powerful. Equation 9 states that *any* arbitrary matrix \mathbf{X} can be converted to an orthogonal matrix, a diagonal matrix and another orthogonal matrix (or a rotation, a stretch and a second rotation). Making sense of Equation 9 is the subject of the next section.

6.2 INTERPRETING SVD

The final form of *SVD* (Equation 9) is a concise but thick statement to understand. Let us instead reinterpret Equation 7.

$$\mathbf{X}\mathbf{a} = k\mathbf{b}$$

where \mathbf{a} and \mathbf{b} are column vectors and k is a scalar constant. The set $\{\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \dots, \hat{\mathbf{v}}_m\}$ is analogous to \mathbf{a} and the set $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_n\}$ is analogous to \mathbf{b} . What is unique though is that $\{\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \dots, \hat{\mathbf{v}}_m\}$ and $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_n\}$ are orthonormal sets of vectors which *span* an m or n dimensional space, respectively. In particular, loosely speaking these sets appear to span all possible “inputs” (\mathbf{a}) and “outputs” (\mathbf{b}). Can we formalize the view that $\{\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \dots, \hat{\mathbf{v}}_n\}$ and $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_n\}$ span all possible “inputs” and “outputs”?

We can manipulate Equation 9 to make this fuzzy hypothesis more precise.

$$\begin{aligned} \mathbf{X} &= \mathbf{U}\Sigma\mathbf{V}^T \\ \mathbf{U}^T\mathbf{X} &= \Sigma\mathbf{V}^T \\ \mathbf{U}^T\mathbf{X} &= \mathbf{Z} \end{aligned}$$

where we have defined $\mathbf{Z} \equiv \Sigma\mathbf{V}^T$. Note that the previous columns $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_n\}$ are now rows in \mathbf{U}^T . Comparing this equation to Equation 1, $\{\hat{\mathbf{u}}_1, \hat{\mathbf{u}}_2, \dots, \hat{\mathbf{u}}_n\}$ perform the same role as $\{\hat{\mathbf{p}}_1, \hat{\mathbf{p}}_2, \dots, \hat{\mathbf{p}}_m\}$. Hence, \mathbf{U}^T is a *change of basis* from \mathbf{X} to \mathbf{Z} . Just as before, we were transforming column vectors, we can again infer that we are

The “value” form of SVD is expressed in equation 7.

$$\mathbf{X}\hat{\mathbf{v}}_i = \sigma_i\hat{\mathbf{u}}_i$$

The mathematical intuition behind the construction of the matrix form is that we want to express all n “value” equations in just one equation. It is easiest to understand this process graphically. Drawing the matrices of equation 7 looks like the following.

$$\begin{pmatrix} \text{---} m \text{---} \\ | \\ n \\ | \end{pmatrix} \times \begin{pmatrix} | \\ m \\ | \end{pmatrix} = \begin{pmatrix} \text{positive} \\ \text{number} \end{pmatrix} \begin{pmatrix} | \\ n \\ | \end{pmatrix}$$

We can construct three new matrices \mathbf{V} , \mathbf{U} and $\mathbf{\Sigma}$. All singular values are first rank-ordered $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r$, and the corresponding vectors are indexed in the same rank order. Each pair of associated vectors $\hat{\mathbf{v}}_i$ and $\hat{\mathbf{u}}_i$ is stacked in the i^{th} column along their respective matrices. The corresponding singular value σ_i is placed along the diagonal (the ii^{th} position) of $\mathbf{\Sigma}$. This generates the equation $\mathbf{XV} = \mathbf{U}\mathbf{\Sigma}$, which looks like the following.

$$\begin{pmatrix} \text{---} m \text{---} \\ | \\ n \\ | \end{pmatrix} \times \begin{pmatrix} \text{---} m \text{---} \\ | \\ m \\ | \end{pmatrix} = \begin{pmatrix} \text{---} n \text{---} \\ | \\ n \\ | \end{pmatrix} \times \begin{pmatrix} \text{---} n \times m \text{---} \\ \text{0} \\ \text{0} \end{pmatrix}$$

The matrices \mathbf{V} and \mathbf{U} are $m \times m$ and $n \times n$ matrices respectively and $\mathbf{\Sigma}$ is a diagonal matrix with a few non-zero values (represented by the checkerboard) along its diagonal. Solving this single matrix equation solves all n “value” form equations.

Figure 4: How to construct the matrix form of SVD from the “value” form.

transforming column vectors. The fact that the orthonormal basis \mathbf{U}^T (or \mathbf{P}) transforms column vectors means that \mathbf{U}^T is a basis that spans the columns of \mathbf{X} . Bases that span the columns are termed the *column space* of \mathbf{X} . The column space formalizes the notion of what are the possible “outputs” of any matrix.

There is a funny symmetry to *SVD* such that we can define a similar quantity - the *row space*.

$$\begin{aligned} \mathbf{XV} &= \mathbf{\Sigma U} \\ (\mathbf{XV})^T &= (\mathbf{\Sigma U})^T \\ \mathbf{V}^T \mathbf{X}^T &= \mathbf{U}^T \mathbf{\Sigma} \end{aligned}$$

$$\mathbf{V}^T \mathbf{X}^T = \mathbf{Z}$$

where we have defined $\mathbf{Z} \equiv \mathbf{U}^T \mathbf{\Sigma}$. Again the rows of \mathbf{V}^T (or the columns of \mathbf{V}) are an orthonormal basis for transforming \mathbf{X}^T into \mathbf{Z} . Because of the transpose on \mathbf{X} , it follows that \mathbf{V} is an orthonormal basis spanning the *row space* of \mathbf{X} . The row space likewise formalizes the notion of what are possible “inputs” into an arbitrary matrix.

We are only scratching the surface for understanding the full implications of *SVD*. For the purposes of this tutorial though, we have enough information to understand how *PCA* will fall within this framework.

6.3 SVD AND PCA

With similar computations it is evident that the two methods are intimately related. Let us return to the original $m \times n$ data matrix \mathbf{X} . We can define a new matrix \mathbf{Y} as an $n \times m$ matrix⁹.

$$\mathbf{Y} \equiv \frac{1}{\sqrt{n-1}} \mathbf{X}^T$$

where each *column* of \mathbf{Y} has zero mean. The definition of \mathbf{Y} becomes clear by analyzing $\mathbf{Y}^T \mathbf{Y}$.

$$\begin{aligned} \mathbf{Y}^T \mathbf{Y} &= \left(\frac{1}{\sqrt{n-1}} \mathbf{X}^T \right)^T \left(\frac{1}{\sqrt{n-1}} \mathbf{X}^T \right) \\ &= \frac{1}{n-1} \mathbf{X}^{TT} \mathbf{X}^T \\ &= \frac{1}{n-1} \mathbf{X} \mathbf{X}^T \\ \mathbf{Y}^T \mathbf{Y} &= \mathbf{S}_{\mathbf{X}} \end{aligned}$$

By construction $\mathbf{Y}^T \mathbf{Y}$ equals the covariance matrix of \mathbf{X} . From section 5 we know that the principal components of \mathbf{X} are the eigenvectors of $\mathbf{S}_{\mathbf{X}}$. If we calculate the *SVD* of \mathbf{Y} , the columns of matrix \mathbf{V} contain the eigenvectors of $\mathbf{Y}^T \mathbf{Y} = \mathbf{S}_{\mathbf{X}}$. *Therefore, the columns of \mathbf{V} are the principal components of \mathbf{X} .* This second algorithm is encapsulated in Matlab code included in Appendix B.

What does this mean? \mathbf{V} spans the row space of $\mathbf{Y} \equiv \frac{1}{\sqrt{n-1}} \mathbf{X}^T$. Therefore, \mathbf{V} must also span the column space of $\frac{1}{\sqrt{n-1}} \mathbf{X}$. We can conclude that finding the principal components¹⁰ amounts to finding an orthonormal basis that spans the *column space* of \mathbf{X} .

7 DISCUSSION AND CONCLUSIONS

7.1 QUICK SUMMARY

Performing *PCA* is quite simple in practice.

⁹ \mathbf{Y} is of the appropriate $n \times m$ dimensions laid out in the derivation of section 6.1. This is the reason for the “flipping” of dimensions in 6.1 and Figure 4.

¹⁰If the final goal is to find an orthonormal basis for the column space of \mathbf{X} then we can calculate it directly without constructing \mathbf{Y} . By symmetry the columns of \mathbf{U} produced by the *SVD* of $\frac{1}{\sqrt{n-1}} \mathbf{X}$ must also be the principal components.




Figure 5: Data points (black dots) tracking a person on a ferris wheel. The extracted principal components are $(\mathbf{p}_1, \mathbf{p}_2)$ and the phase is $\hat{\theta}$.

1. Organize a data set as an $m \times n$ matrix, where m is the number of measurement types and n is the number of trials.
2. Subtract off the mean for each measurement type or row \mathbf{x}_i .
3. Calculate the *SVD* or the eigenvectors of the covariance.

In several fields of literature, many authors refer to the individual measurement types \mathbf{x}_i as the *sources*. The data projected into the principal components $\mathbf{Y} = \mathbf{P}\mathbf{X}$ are termed the *signals*, because the projected data presumably represent the “true” underlying probability distributions.

7.2 DIMENSIONAL REDUCTION

One benefit of *PCA* is that we can examine the variances $\mathbf{S}_{\mathbf{Y}}$ associated with the principle components. Often one finds that large variances associated with

the first $k < m$ principal components, and then a precipitous drop-off. One can conclude that most interesting dynamics occur only in the first k dimensions.

In the example of the spring, $k = 1$. Likewise, in Figure 3 panel (c), we recognize $k = 1$ along the principal component of $r_2 = kr_1$. This process of throwing out the less important axes can help reveal hidden, simplified dynamics in high dimensional data. This process is aptly named *dimensional reduction*.

7.3 LIMITS AND EXTENSIONS OF PCA

Both the strength and weakness of *PCA* is that it is a *non-parametric* analysis. One only needs to make the assumptions outlined in section 4.5 and then calculate the corresponding answer. There are no parameters to tweak and no coefficients to adjust based on user experience - the answer is unique¹¹ and independent of the user.

This same strength can also be viewed as a weakness. If one knows a-priori some features of the dynamics of a system, then it makes sense to incorporate these assumptions into a *parametric* algorithm - or an algorithm with selected parameters.

Consider the recorded positions of a person on a ferris wheel over time in Figure 5. The probability distributions along the axes are approximately Gaussian and thus *PCA* finds $(\mathbf{p}_1, \mathbf{p}_2)$, however this answer might not be optimal. The most concise form of dimensional reduction is to recognize that the phase (or angle along the ferris wheel) contains all dynamic information. Thus, the appropriate parametric algorithm is to first convert the data to the appropriately centered polar coordinates and then compute *PCA*.

This prior *non-linear* transformation is sometimes termed a *kernel transformation* and the entire parametric algorithm is termed *kernel PCA*. Other common kernel transformations include Fourier and

¹¹To be absolutely precise, the principal components are **not** uniquely defined. One can always flip the direction by multiplying by -1 . In addition, eigenvectors beyond the rank of a matrix (i.e. $\sigma_i = 0$ for $i > \text{rank}$) can be selected almost at whim. However, these degrees of freedom do not effect the qualitative features of the solution nor a dimensional reduction.




Figure 6: Non-Gaussian distributed data causes *PCA* to fail. In exponentially distributed data the axes with the largest variance do not correspond to the underlying basis.

Gaussian transformations. This procedure is parametric because the user must incorporate prior knowledge of the dynamics in the selection of the kernel but it is also more optimal in the sense that the dynamics are more concisely described.

Sometimes though the assumptions themselves are too stringent. One might envision situations where the principal components need not be orthogonal. Furthermore, the distributions along each dimension (\mathbf{x}_i) need not be Gaussian. For instance, Figure 6 contains a 2-D exponentially distributed data set. The largest variances do not correspond to the meaningful axes of thus *PCA* fails.

This less constrained set of problems is not trivial and only recently has been solved adequately via *Independent Component Analysis (ICA)*. The formulation is equivalent.

Find a matrix \mathbf{P} where $\mathbf{Y} = \mathbf{PX}$ such that $\mathbf{S}_\mathbf{Y}$ is diagonalized.

however it abandons all assumptions except linearity, and attempts to find axes that satisfy the most formal form of redundancy reduction - *statistical independence*. Mathematically *ICA* finds a basis such that the joint probability distribution can be factor-

ized¹².

$$P(\mathbf{y}_i, \mathbf{y}_j) = P(\mathbf{y}_i)P(\mathbf{y}_j)$$

for all i and j , $i \neq j$. The downside of *ICA* is that it is a form of nonlinear optimization, making the solution difficult to calculate in practice and potentially not unique. However *ICA* has been shown a very practical and powerful algorithm for solving a whole new class of problems.

Writing this paper has been an extremely instructional experience for me. I hope that this paper helps to demystify the motivation and results of *PCA*, and the underlying assumptions behind this important analysis technique.

8 APPENDIX A: LINEAR ALGEBRA

This section proves a few unapparent theorems in linear algebra, which are crucial to this paper.

1. The inverse of an orthogonal matrix is its transpose.

The goal of this proof is to show that if \mathbf{A} is an orthogonal matrix, then $\mathbf{A}^{-1} = \mathbf{A}^T$.

Let \mathbf{A} be an $m \times n$ matrix.

$$\mathbf{A} = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$$

where \mathbf{a}_i is the i^{th} column vector. We now show that $\mathbf{A}^T \mathbf{A} = \mathbf{I}$ where \mathbf{I} is the identity matrix.

Let us examine the ij^{th} element of the matrix $\mathbf{A}^T \mathbf{A}$. The ij^{th} element of $\mathbf{A}^T \mathbf{A}$ is $(\mathbf{A}^T \mathbf{A})_{ij} = \mathbf{a}_i^T \mathbf{a}_j$.

Remember that the columns of an orthonormal matrix are orthonormal to each other. In other words, the dot product of any two columns is zero. The only exception is a dot product of one particular column with itself, which equals one.

$$(\mathbf{A}^T \mathbf{A})_{ij} = \mathbf{a}_i^T \mathbf{a}_j = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$$

$\mathbf{A}^T \mathbf{A}$ is the exact description of the identity matrix. The definition of \mathbf{A}^{-1} is $\mathbf{A}^{-1} \mathbf{A} = \mathbf{I}$. Therefore,

¹²Equivalently, in the language of *information theory* the goal is to find a basis \mathbf{P} such that the mutual information $I(\mathbf{y}_i, \mathbf{y}_j) = 0$ for $i \neq j$.

because $\mathbf{A}^T \mathbf{A} = \mathbf{I}$, it follows that $\mathbf{A}^{-1} = \mathbf{A}^T$.

2. If \mathbf{A} is any matrix, the matrices $\mathbf{A}^T \mathbf{A}$ and $\mathbf{A} \mathbf{A}^T$ are both symmetric.

Let's examine the transpose of each in turn.

$$\begin{aligned} (\mathbf{A} \mathbf{A}^T)^T &= \mathbf{A}^{TT} \mathbf{A}^T = \mathbf{A} \mathbf{A}^T \\ (\mathbf{A}^T \mathbf{A})^T &= \mathbf{A}^T \mathbf{A}^{TT} = \mathbf{A}^T \mathbf{A} \end{aligned}$$

The equality of the quantity with its transpose completes this proof.

3. A matrix is symmetric if and only if it is orthogonally diagonalizable.

Because this statement is bi-directional, it requires a two-part “if-and-only-if” proof. One needs to prove the forward and the backwards “if-then” cases.

Let us start with the forward case. If \mathbf{A} is orthogonally diagonalizable, then \mathbf{A} is a symmetric matrix. By hypothesis, orthogonally diagonalizable means that there exists some \mathbf{E} such that $\mathbf{A} = \mathbf{E} \mathbf{D} \mathbf{E}^T$, where \mathbf{D} is a diagonal matrix and \mathbf{E} is some special matrix which diagonalizes \mathbf{A} . Let us compute \mathbf{A}^T .

$$\mathbf{A}^T = (\mathbf{E} \mathbf{D} \mathbf{E}^T)^T = \mathbf{E}^{TT} \mathbf{D}^T \mathbf{E}^T = \mathbf{E} \mathbf{D} \mathbf{E}^T = \mathbf{A}$$

Evidently, if \mathbf{A} is orthogonally diagonalizable, it must also be symmetric.

The reverse case is more involved and less clean so it will be left to the reader. In lieu of this, hopefully the “forward” case is suggestive if not somewhat convincing.

4. A symmetric matrix is diagonalized by a matrix of its orthonormal eigenvectors.

Restated in math, let \mathbf{A} be a square $n \times n$ symmetric matrix with associated eigenvectors $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. Let $\mathbf{E} = [\mathbf{e}_1 \ \mathbf{e}_2 \ \dots \ \mathbf{e}_n]$ where the i^{th} column of \mathbf{E} is the eigenvector \mathbf{e}_i . This theorem asserts that there exists a diagonal matrix \mathbf{D} where $\mathbf{A} = \mathbf{E} \mathbf{D} \mathbf{E}^T$.

This theorem is an extension of the previous theorem 3. It provides a prescription for how to find the

matrix \mathbf{E} , the “diagonalizer” for a symmetric matrix. It says that the special diagonalizer is in fact a matrix of the original matrix’s eigenvectors.

This proof is in two parts. In the first part, we see that the any matrix can be orthogonally diagonalized if and only if it that matrix’s eigenvectors are all linearly independent. In the second part of the proof, we see that a symmetric matrix has the special property that all of its eigenvectors are not just linearly independent but also orthogonal, thus completing our proof.

In the first part of the proof, let \mathbf{A} be just some matrix, not necessarily symmetric, and let it have independent eigenvectors (i.e. no degeneracy). Furthermore, let $\mathbf{E} = [\mathbf{e}_1 \ \mathbf{e}_2 \ \dots \ \mathbf{e}_n]$ be the matrix of eigenvectors placed in the columns. Let \mathbf{D} be a diagonal matrix where the i^{th} eigenvalue is placed in the ii^{th} position.

We will now show that $\mathbf{AE} = \mathbf{ED}$. We can examine the columns of the right-hand and left-hand sides of the equation.

$$\begin{aligned} \text{Left hand side : } \mathbf{AE} &= [\mathbf{Ae}_1 \ \mathbf{Ae}_2 \ \dots \ \mathbf{Ae}_n] \\ \text{Right hand side : } \mathbf{ED} &= [\lambda_1 \mathbf{e}_1 \ \lambda_2 \mathbf{e}_2 \ \dots \ \lambda_n \mathbf{e}_n] \end{aligned}$$

Evidently, if $\mathbf{AE} = \mathbf{ED}$ then $\mathbf{Ae}_i = \lambda_i \mathbf{e}_i$ for all i . This equation is the definition of the eigenvalue equation. Therefore, it must be that $\mathbf{AE} = \mathbf{ED}$. A little rearrangement provides $\mathbf{A} = \mathbf{EDE}^{-1}$, completing the first part the proof.

For the second part of the proof, we show that a symmetric matrix always has orthogonal eigenvectors. For some symmetric matrix, let λ_1 and λ_2 be distinct eigenvalues for eigenvectors \mathbf{e}_1 and \mathbf{e}_2 .

$$\begin{aligned} \lambda_1 \mathbf{e}_1 \cdot \mathbf{e}_2 &= (\lambda_1 \mathbf{e}_1)^T \mathbf{e}_2 \\ &= (\mathbf{Ae}_1)^T \mathbf{e}_2 \\ &= \mathbf{e}_1^T \mathbf{A}^T \mathbf{e}_2 \\ &= \mathbf{e}_1^T \mathbf{Ae}_2 \\ &= \mathbf{e}_1^T (\lambda_2 \mathbf{e}_2) \\ \lambda_1 \mathbf{e}_1 \cdot \mathbf{e}_2 &= \lambda_2 \mathbf{e}_1 \cdot \mathbf{e}_2 \end{aligned}$$

By the last relation we can equate that $(\lambda_1 - \lambda_2) \mathbf{e}_1 \cdot \mathbf{e}_2 = 0$. Since we have conjectured that the eigenvalues are in fact unique, it must be

the case that $\mathbf{e}_1 \cdot \mathbf{e}_2 = 0$. Therefore, the eigenvectors of a symmetric matrix are orthogonal.

Let us back up now to our original postulate that \mathbf{A} is a symmetric matrix. By the second part of the proof, we know that the eigenvectors of \mathbf{A} are all orthonormal (we choose the eigenvectors to be normalized). This means that \mathbf{E} is an orthogonal matrix so by theorem 1, $\mathbf{E}^T = \mathbf{E}^{-1}$ and we can rewrite the final result.

$$\mathbf{A} = \mathbf{EDE}^T$$

Thus, a symmetric matrix is diagonalized by a matrix of its eigenvectors.

5. For any arbitrary $m \times n$ matrix \mathbf{X} , the symmetric matrix $\mathbf{X}^T \mathbf{X}$ has a set of orthonormal eigenvectors of $\{\hat{\mathbf{v}}_1, \hat{\mathbf{v}}_2, \dots, \hat{\mathbf{v}}_n\}$ and a set of associated eigenvalues $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$. The set of vectors $\{\mathbf{X}\hat{\mathbf{v}}_1, \mathbf{X}\hat{\mathbf{v}}_2, \dots, \mathbf{X}\hat{\mathbf{v}}_n\}$ then form an orthogonal basis, where each vector $\mathbf{X}\hat{\mathbf{v}}_i$ is of length $\sqrt{\lambda_i}$.

All of these properties arise from the dot product of any two vectors from this set.

$$\begin{aligned} (\mathbf{X}\hat{\mathbf{v}}_i) \cdot (\mathbf{X}\hat{\mathbf{v}}_j) &= (\mathbf{X}\hat{\mathbf{v}}_i)^T (\mathbf{X}\hat{\mathbf{v}}_j) \\ &= \hat{\mathbf{v}}_i^T \mathbf{X}^T \mathbf{X} \hat{\mathbf{v}}_j \\ &= \hat{\mathbf{v}}_i^T (\lambda_j \hat{\mathbf{v}}_j) \\ &= \lambda_j \hat{\mathbf{v}}_i \cdot \hat{\mathbf{v}}_j \\ (\mathbf{X}\hat{\mathbf{v}}_i) \cdot (\mathbf{X}\hat{\mathbf{v}}_j) &= \lambda_j \delta_{ij} \end{aligned}$$

The last relation arises because the set of eigenvectors of \mathbf{X} is orthogonal resulting in the Kronecker delta. In more simpler terms the last relation states:

$$(\mathbf{X}\hat{\mathbf{v}}_i) \cdot (\mathbf{X}\hat{\mathbf{v}}_j) = \begin{cases} \lambda_j & i = j \\ 0 & i \neq j \end{cases}$$

This equation states that any two vectors in the set are orthogonal.

The second property arises from the above equation by realizing that the length squared of each vector is defined as:

$$\|\mathbf{X}\hat{\mathbf{v}}_i\|^2 = (\mathbf{X}\hat{\mathbf{v}}_i) \cdot (\mathbf{X}\hat{\mathbf{v}}_i) = \lambda_i$$

9 APPENDIX B: CODE

This code is written for Matlab 6.5 (Release 13) from Mathworks¹³. The code is not computationally efficient but explanatory (terse comments begin with a %).

This first version follows Section 5 by examining the covariance of the data set.

```
function [signals,PC,V] = pca1(data)
% PCA1: Perform PCA using covariance.
%   data - MxN matrix of input data
%           (M dimensions, N trials)
%   signals - MxN matrix of projected data
%   PC - each column is a PC
%   V - Mx1 matrix of variances

[M,N] = size(data);

% subtract off the mean for each dimension
mn = mean(data,2);
data = data - repmat(mn,1,N);

% calculate the covariance matrix
covariance = 1 / (N-1) * data * data';

% find the eigenvectors and eigenvalues
[PC, V] = eig(covariance);

% extract diagonal of matrix as vector
V = diag(V);

% sort the variances in decreasing order
[junk, rindices] = sort(-1*V);
V = V(rindices);
PC = PC(:,rindices);

% project the original data set
signals = PC' * data;
```

This second version follows section 6 computing PCA through SVD.

```
function [signals,PC,V] = pca2(data)
```

```
% PCA2: Perform PCA using SVD.
%   data - MxN matrix of input data
%           (M dimensions, N trials)
%   signals - MxN matrix of projected data
%   PC - each column is a PC
%   V - Mx1 matrix of variances
```

```
[M,N] = size(data);

% subtract off the mean for each dimension
mn = mean(data,2);
data = data - repmat(mn,1,N);

% construct the matrix Y
Y = data' / sqrt(N-1);

% SVD does it all
[u,S,PC] = svd(Y);

% calculate the variances
S = diag(S);
V = S .* S;

% project the original data
signals = PC' * data;
```

10 REFERENCES

Bell, Anthony and Sejnowski, Terry. (1997) "The Independent Components of Natural Scenes are Edge Filters." *Vision Research* 37(23), 3327-3338.

[A paper from my field of research that surveys and explores different forms of decorrelating data sets. The authors examine the features of PCA and compare it with new ideas in redundancy reduction, namely Independent Component Analysis.]

Bishop, Christopher. (1996) *Neural Networks for Pattern Recognition*. Clarendon, Oxford, UK.

[A challenging but brilliant text on statistical pattern recognition (neural networks). Although the derivation of PCA is touch in section 8.6 (p.310-319), it does have a great discussion on potential extensions to the method and it puts PCA in context of other methods of dimensional

¹³<http://www.mathworks.com>

reduction. Also, I want to acknowledge this book for several ideas about the limitations of PCA.]

Lay, David. (2000). *Linear Algebra and It's Applications*. Addison-Wesley, New York.

[This is a beautiful text. Chapter 7 in the second edition (p. 441-486) has an exquisite, intuitive derivation and discussion of SVD and PCA. Extremely easy to follow and a must read.]

Mitra, Partha and Pesaran, Bijan. (1999) "Analysis of Dynamic Brain Imaging Data." *Biophysical Journal*. 76, 691-708.

[A comprehensive and spectacular paper from my field of research interest. It is dense but in two sections "Eigenmode Analysis: SVD" and "Space-frequency SVD" the authors discuss the benefits of performing a Fourier transform on the data before an SVD.]

Will, Todd (1999) "Introduction to the Singular Value Decomposition" Davidson College. www.davidson.edu/academic/math/will/svd/index.html

[A math professor wrote up a great web tutorial on SVD with tremendous intuitive explanations, graphics and animations. Although it avoids PCA directly, it gives a great intuitive feel for what SVD is doing mathematically. Also, it is the inspiration for my "spring" example.]