



GlobalTreeSearch: The first complete global database of tree species and country distributions

E. Beech, M. Rivers, S. Oldfield & P. P. Smith

To cite this article: E. Beech, M. Rivers, S. Oldfield & P. P. Smith (2017) GlobalTreeSearch: The first complete global database of tree species and country distributions, Journal of Sustainable Forestry, 36:5, 454-489, DOI: [10.1080/10549811.2017.1310049](https://doi.org/10.1080/10549811.2017.1310049)

To link to this article: <https://doi.org/10.1080/10549811.2017.1310049>



Accepted author version posted online: 23 Mar 2017.
Published online: 17 Apr 2017.



Submit your article to this journal [↗](#)



Article views: 10379



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 40 View citing articles [↗](#)

GlobalTreeSearch: The first complete global database of tree species and country distributions

E. Beech^a, M. Rivers^{a,b}, S. Oldfield^b, and P. P. Smith^a

^aBotanic Gardens Conservation International, Richmond, United Kingdom; ^bIUCN/SSC Global Tree Specialist Group

ABSTRACT

This article presents, for the first time, an overview of all known tree species by scientific name and country level distribution, and describes an online database—GlobalTreeSearch—that provides access to this information. Based on our comprehensive analysis of published data sources and expert input, the number of tree species currently known to science is 60,065, representing 20% of all angiosperm and gymnosperm plant species. Nearly half of all tree species (45%) are found in just 10 families, with the 3 most tree-rich families being Leguminosae, Rubiaceae, and Myrtaceae. Geographically, Brazil, Colombia, and Indonesia are the countries with the most tree species. The countries with the most country-endemic tree species reflect broader plant diversity trends (Brazil, Australia, China) or islands where isolation has resulted in speciation (Madagascar, Papua New Guinea, Indonesia). Nearly 58% of all tree species are single-country endemics. Our intention is for GlobalTreeSearch to be used as a tool for monitoring and managing tree species diversity, forests, and carbon stocks on a global, regional, and/or national level. It will also be used as the basis of the Global Tree Assessment, which aims to assess the conservation status of all of the world's tree species by 2020.

KEYWORDS

Global tree assessment; GlobalTreeSearch; tree database; tree distribution; tree diversity; tree endemism

Introduction

Trees first evolved over 300 million years ago, and “woodiness” has evolved since that time in plant families across the taxonomic array (Fitzjohn et al., 2014; Kenrick & Crane, 1997). Trees provide us with food, timber, and medicine, and are major components of the most biologically diverse ecosystems on earth, including woodlands and forests (FAO, 2014). Individual tree species play crucial roles in their ecosystems, supporting a multitude of other species due to their position at the base of trophic pyramid (e.g., Kennedy & Southwood, 1984). In the climate change negotiations in Paris in 2015, the conservation of trees, the cessation of deforestation, and the sustainable management of forests were all recognized as activities that address climate change (Poorter et al., 2015). However, effective forest conservation requires species-specific action, as individual tree species face threats that are unique to that species.

Despite the importance of trees and their ecosystem services, there has until now not been an authoritative global list of trees. In fact, our knowledge of the number of tree

CONTACT E. Beech  Emily.Beech@bgci.org  Botanic Gardens Conservation International

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/wjsf.

© 2017 Taylor & Francis

species has remained surprisingly sparse at the national and international levels. In 1753, in *Species Plantarum*, Linnaeus described 9,000 species of plants and marked up the woody species with the symbol of Saturn (Simpson, 2010). Since the time of Linnaeus, botanical exploration and discovery has hugely expanded, but the precise number of plant species still remains uncertain, with current estimates suggesting there are 370,495 seed plants (Lughadha et al., 2016). Estimates of the number of tree species have ranged from 45,000 to 100,000 (Fine & Ree, 2006; Oldfield, Lusty, & MacKinnon, 1998; Savolainen, 2000; Tudge, 2006), with reports suggesting there are 21,000 species in temperate regions (Hunt, 1996) and 40,000–53,000 in the tropics (Slik et al., 2015). Most of these figures are based on broad estimates or modeled numbers of trees; none of them have been derived from an authoritative global list of trees.

Here, for the first time, we present an overview of the complete list of all known tree species by scientific name and country level distribution, and we describe an online database, called GlobalTreeSearch (http://www.bgci.org/globaltree_search.php), that provides access to this information. GlobalTreeSearch has been designed as a tool for assessing, monitoring, and managing tree species diversity, forests, and carbon stocks on a global, regional, and/or national level.

Methodology

Definition of tree

As tree type growth habits have evolved many times across different plant families, there are many definitions of a tree. For the purposes of this project, we use the tree definition agreed by IUCN's Global Tree Specialist Group (GTSG): *a woody plant with usually a single stem growing to a height of at least two metres, or if multi-stemmed, then at least one vertical stem five centimetres in diameter at breast height*. We excluded cycads, tree ferns, tree-like Poaceae, Bromeliaceae, and Musaceae from our list. The World Checklist of Selected Plant Families (WCSP) and several other taxonomic databases use the Raunkiaer system for categorizing life forms, with trees most often being categorized as phanerophytes. However, this system excludes smaller trees and also includes some larger shrubs. Efforts have been made to include the smaller trees by incorporating country level and regional tree lists in addition to WCSP species. Shrubs were removed manually on a species-by-species basis. Some plant species have variable life-forms, but all species that are recorded as growing naturally as a tree somewhere have been included.

Data compilation and management

Tree species data were accessed from a range of sources including: the World Checklist of Selected Plant Families (WCSP, 2016), Flora of China (Wu Zhengyi, Raven, & Hong Deyuan 1994–2013), African Plant Database (Conservatoire et Jardin Botaniques Ville de Genève, 2017), FloraBase (Western Australian Herbarium, 1998), Plants of Southern Africa (SANBI, 2014), The PLANTS database (USDA, 2016), Brazilian Flora 2020 (Rio de Janeiro Botanical Garden, 2016), and Tropicos (Missouri Botanical Garden, 2016).

In addition to these major published sources, we extended the call for information to the botanical community and contacted over 80 experts in the GTSG and member institutions of

Botanic Gardens Conservation International (BGCI), receiving a wide range of regional, taxonomic, and country-specific tree checklists. Over 500 published sources were consulted, all of which are referenced in the Database Reference Sources listed in the Appendix. Collation and compilation of the data has taken over two years of full-time work.

Database description

The GlobalTreeSearch database was compiled in Microsoft Access and consists of four interlinked tables. The first table of taxon names contains all of the names received during the compilation of GlobalTreeSearch. This table currently contains 180,000 entries. Each unique tree taxon name links by ID number to the source of that name. This long list of tree names has been refined only to contain the names of accepted tree species names (i.e., the 60,065 final tree species). For this refined list of accepted tree species, we recorded associated geographic information. Each country occurrence for each species has its own unique source. This table currently contains over 195,500 entries.

Taxonomy

All taxon names received were recorded, but only taxonomically accepted names (or unresolved names) are included in the published GlobalTreeSearch. The current version of GlobalTreeSearch focuses on species names and does not contain infraspecific taxa. The taxonomic opinion follows the WCSP for its published families. Families that are not included in WCSP follow The Plant List (TPL, 2013). Other regional sources (Euro+Med, Reflora) and taxonomic sources (ILDIS) have been consulted when required. As The Plant List (TPL, 2013) was last updated in May 2012, any species published since then have been provisionally accepted on our list unless found to be synonymous with species from other more recent published sources, such as in a published monograph. In such cases, the source of the taxonomic status is recorded, allowing for updates as taxonomic opinion evolves.

Distribution information

The country-level distribution information was recorded following the ISO 3166 country names standard. Distribution data were sourced from various published references including the nomenclatural sources mentioned above, as well as additional monographs, checklists, and floras, some electronically and others through manual review. Distribution information has been recorded to country level for all taxa, with distribution in certain countries (Brazil, the United States, South Africa, Australia, and China) recorded to state or province. These countries were chosen as they are geographically expansive, and the distributions at sub-country level were generally easily obtained.

The country distributions were also matched to eight biomes (Oceania, Neotropical, Nearctic, Afrotropical, Antarctic, Palearctic, Australasia, Indo-Malay), following Olson et al. (2001). For the purpose of our analysis, Indonesia is considered entirely within the Indo-Malay biome and China within the Palearctic biome.

Expert review

GlobalTreeSearch has been reviewed by more than 30 experts chosen for their taxonomic or regional expertise. These experts reviewed the list of omitted taxa and amended distribution data where necessary. However, GlobalTreeSearch is not a static entity, which is to say that additions, taxonomic revisions, and deletions will continue to be carried out to ensure the list remains the most comprehensive and up-to-date list of the world's tree species. The database will be maintained and managed by BGCi through its website http://www.bgci.org/globaltree_search.php. An e-mail address will be set up to allow people to suggest revisions which will be verified through expert review before modification. In addition, if a suggested revision cannot easily be verified or resolved, then a comments field will record the nature of the query received.

Results

There are 60,065 tree species recorded worldwide. Based on current taxonomic knowledge and levels of botanical exploration, we believe that this compilation represents the best available record of the total number of tree species and a sound basis for conservation planning.

Over 45% (27,203) of tree species are found in just 10 families (Figure 1). The family with the most tree species is the Leguminosae with 5,405 tree species, followed by Rubiaceae (4,827), and then Myrtaceae (4,330).

The most diverse tree genera are *Syzygium* (1,069 species), *Eugenia* (884 species), and *Eucalyptus* (747 species), all in the family Myrtaceae (Figure 2). *Ficus* (Moraceae), *Diospyros* (Ebenaceae), and *Psychotria* (Rubiaceae) are the fourth, fifth, and sixth largest genera, respectively.

The country with the most diverse tree flora is Brazil, with 8,715 tree species, followed by Colombia (5,776 spp.) and Indonesia (5,142 spp.) (Figure 3). Nearly 58% of all tree species (34,575) are single country endemics. The countries with the most endemic trees




Figure 1. Top ten families with the most tree species.




Figure 2. Top ten genera with the most tree species.




Figure 3. Top ten countries with most tree species.

are Brazil (4,333 spp.), Madagascar (2,991 spp.), Australia (2,584 spp.), and China (2,149 spp.) (Figure 4).

Matching country distributions to biomes (Figure 5), the biome with the largest number of trees is the Neotropic biome with over 23,000 tree species, followed by the other tropical biomes, the Indo-Malay biome and the Afrotropic biome. There are no trees occurring in the Antarctic biome, and the next lowest diversity is in North America in the Nearctic biome, with fewer than 1,400 species. The most diverse tree genera reflect different centers of diversity. More than half of the species in the genus *Syzygium* (615) are found in the Indo-Malay biome, over 731 *Eugenia* species are found in the Neotropics, and almost all of the *Eucalyptus* genus is found in Australasia.




Figure 4. Top ten countries with the largest number of endemic species.




Figure 5. Tree species numbers by biome.

Discussion

The compilation of GlobalTreeSearch has been a multidisciplinary effort, with data contributed from botanic gardens, museums, academia, NGOs, forestry organizations, and agricultural institutions. GlobalTreeSearch asserts the number of tree species to be at the lower end of previous estimates. Nevertheless, the GlobalTreeSearch database contains over 375,500 plant names and country entries, an indication of the scale of the task. Having an accurate number of tree species and their country level distributions will encourage extensive applications by a wide range of users, from conservation planning and action to sustainable forestry practices.

The tropical hotspots of tree diversity identified by this analysis are unsurprising, with countries in the Neotropic biome coming out on top as reported in more general assessments of the distribution of biodiversity (Giam, Bradshaw, Tan, & Sodhi, 2010; Heywood & Watson, 1995; Myers, Mittermeier, Mittermeier, Da Fonseca, & Kent, 2000). Other estimates of tree diversity hotspots (such as Slik et al., 2015) have reported that the

Indo-Pacific region, comprising the Australasian, Oceania, and Indo-Malay biomes as well as Madagascar, is of comparable species diversity to the Neotropics. Again, our analysis reflects this assertion. Similarly, countries with the highest levels of tree endemism reflect broader measures of plant diversity (e.g., Brazil, China, Australia) or islands where isolation has resulted in speciation (e.g., Madagascar, Papua New Guinea, Indonesia). The preliminary analysis of tree distributions by country and biome represented in this article will be followed by more in-depth analyses in future publications.

Data availability was an important factor in developing this checklist. Several biodiverse countries have electronic online floras developed in response to the Global Strategy for Plant Conservation Target 1: An online flora of all known plants (Sharrock, Oldfield, & Wilson, 2014). For this reason, countries such as Colombia, Brazil, and China are likely to have more up-to-date data in the GlobalTreeSearch database. Other regions with diverse tree floras (e.g., Southeast Asia) have not been as extensively surveyed or have not compiled their data electronically. Although hard copy publications were used extensively in cases where floras are still incomplete (e.g., *Flora Malesiana*), there will still inevitably be data that are poorer for some areas. Projects are already under way to complete online floras on national, regional, and global scales, sources which will be utilized by GlobalTreeSearch when available.

The fact that Leguminosae is the most diverse tree family is not surprising given that Leguminosae is the third most diverse plant family after the (primarily herbaceous) families Asteraceae and Orchidaceae. Rubiaceae, the second most diverse tree family is the fourth most diverse plant family and is primarily tropical and woody. Myrtaceae is the third most diverse tree family and, interestingly, accounts for the top three tree genera *Syzygium*, *Eugenia*, and *Eucalyptus*. This reflects the fact that the family Myrtaceae is almost entirely woody, and it is characteristic of highly diverse subtropical and tropical zones.

GlobalTreeSearch is not a static database and it will incorporate both changes in taxonomy and increased availability of data. BGCI encourages submissions from regions where data may not be as readily available to improve the GlobalTreeSearch database. There is also scope for increasing the levels of regional data for countries other than those already included as new data become available. For example, island level data for countries such as Indonesia and the Philippines would be useful for conservation planning and forestry.

At the policy level, documenting trends in the conservation, sustainable use, and development of Forest Genetic Resources (FGR) has been undertaken by the Food and Agriculture Organization (FAO) of the United Nations. In 2014, FAO published the first State of the World's Forest Genetic Resources (SOWFGR) report (FAO, 2014). The purpose of this report was “[to build] the information and knowledge base required for action towards better conservation and sustainable management of FGR at national, regional, and international levels”. Based on the preliminary, draft findings of this report, FAO noted that “very few countries have detailed tree species checklists that include species characteristics allowing distinction between different life forms” (FAO, 2014). As a result, the inventories submitted by the 86 countries that contributed to the report were inconsistent and incomplete. The completion of GlobalTreeSearch will greatly support the FAO in achieving the strategic priorities outlined in their “Global Plan of Action for the Conservation, Sustainable Use, and

Development of Forest Genetic Resources” (FAO, 2013). Specifically, the GlobalTreeSearch database will enable country checklists to be routinely produced and will constitute a comprehensive data set that can be used to measure trends in the conservation and use of FGR over time. Furthermore, production of tree species distribution maps is an essential step in conservation and management of FGR, though few countries have the resources to produce such maps. With the GlobalTreeSearch database, the production of country-level and precise location distribution maps for all tree species is achievable.

Mapping species is also an important step in assessing the conservation status of all of the world’s tree species, an activity essential to ensure effective conservation and management of tree diversity. This is the aim of the Global Tree Assessment—a complete assessment of the conservation status of all trees by 2020 (Newton et al., 2015). The Global Tree Assessment is being made possible by the GlobalTreeSearch database, which will be used to prioritize species in need of conservation assessment. GlobalTreeSearch is used to develop a workflow to rapidly and accurately assess and map species not at risk of extinction (“Least Concern”), in conjunction with GBIF data, which in turn will enable subsequent activities to focus on the often less well-known species that are at risk of extinction. Tree red listing activities are already under way or completed for the Global Tree Assessment including assessment of European trees, assessment of various taxonomic groups (Magnoliaceae, oaks, Betulaceae, etc.), and initial assessment of the world’s commercial timbers. To date, conservation status assessments of only around 20,000 of the trees with scientifically accepted names have been recorded in global, regional, or national red lists; that is, about a third of known tree species (BGCI, ThreatSearch, unpublished data). BGCI will build on this work by carrying out a gap analysis bringing in data from its ThreatSearch database (a compilation of plant conservation assessments), PlantSearch database (taxa in botanic garden collections), and GardenSearch database (location of botanic gardens worldwide) to identify where capacity is needed to prioritize tree conservation through red listing and to conserve trees both *in situ* and *ex situ*.

The Global Tree Assessment supports the Global Strategy for Plant Conservation and the broader Aichi Targets of the Convention on Biological Diversity but is not simply a tool for tree conservation. GlobalTreeSearch and the Global Tree Assessment also support initiatives to improve sustainable forest management, for example through regulation of the international timber trade. FAO estimates that in 2006 the trade in timber products contributed some \$468 billion annually to global GDP (FAO, 2009). A recently compiled working list of internationally traded timbers documented 1,575 timber taxa (Mark, Newton, Oldfield, & Rivers, 2014). The conservation status of many of these species is unknown, and this is even more the case for the thousands of species that are traded locally or regionally. The national inventories and conservation assessments provided by GlobalTreeSearch and the GTA will greatly assist national authorities in regulating unsustainable trade in the species concerned. The Global Tree Assessment will also support implementation of the UNFCCC, which aims to tackle greenhouse gas emissions from deforestation and forest degradation through the “REDD+” program. Recognizing the potential for social and environmental risks and benefits from REDD+, the UNFCCC has agreed on a set of broad safeguards that countries should promote and support, specifically focusing on the conservation of natural forests and biodiversity (Visseren-Hamakers, McDermott, Marjanneke, & Cashore, 2012). Improved information on the

conservation status of tree species will help to focus REDD+ activities and enhance their conservation impacts.

Conclusions

It is perhaps surprising that it has taken until 2017 to compile GlobalTreeSearch, the first global, authoritative list of tree species. However, it is worth acknowledging that GlobalTreeSearch represents a huge scientific effort encompassing the discovery, collection, and description of tens of thousands of plant species. This is “big science” involving the work of thousands of botanists over a period of centuries, and the advent of digital checklists and databases over the past few decades has made the collation and refinement of so many data sources possible. Furthermore, GlobalTreeSearch is not a static list; it will continue to be refined, revised, and added to. Some geographic regions remain relatively unexplored botanically and, even where thorough fieldwork has been undertaken, there can be a substantial lag between discovery and publication. Similarly, plant taxonomy is constantly changing with new discoveries and taxonomic revisions augmenting our knowledge.

GlobalTreeSearch has both immediate value to a wide range of user communities and will allow for the assessment and monitoring of tree species diversity on a global, regional, and/or national level over time.

Acknowledgments

The authors would like to thank in particular the following organizations for providing data contributing to GlobalTreeSearch: African Plant Database; World Checklist of Selected Plant Families; IPNI; Flora of China; Brazilian Flora 2020; Missouri Botanical Gardens Tropicos; Plants of Southern Africa; FloraBase. A full list of the 500+ sources consulted for GlobalTreeList is available at the end of this article. We would also like to thank the following contributors and reviewers: Megan Barstow, Henk Beentje, Steven Brewer, Chien-Fan Chen, Lillian Chua, Alison Copeland, Iain Darbyshire, Neville Evans, Alfredo Fuentes, Zoe Irwin, Yvette Harvey-Brown, Thomas Heller, Alison Hutchinson, Ben Jones, David Kikodze, Esteban Martínez, Mónica Moraes R., Lydia Murphy, Sachiko Nishida, Naomi Rich, Marie-Stéphanie Samain, George Schatz, Anne-Cathrine Scheen, Ferry Slik, Harry Smith, Daniela Suarez, Jonathan Timberlake, Philip Thomas, Lex Thomson, Carmen Ulloa Ulloa, Salako K. Valère, Freddy Zenteno, and the following IUCN Species Survival Committee Specialist Groups: Global Tree Specialist Group, Cuban Plant Specialist Group, New Caledonia Plant Red List Authority, Madagascar Plant Specialist Group.

References

- Conservatoire et Jardin Botaniques Ville de Genève. (2017). *African plant database*. Retrieved April 1, 2016, from <http://www.ville-ge.ch/musinfo/bd/cjb/africa/>
- FAO. (2009). *State of the world's forests 2009*. Rome, Italy: FAO.
- FAO. (2013). *Global plan of action for the conservation, sustainable use and development of forest genetic resources*. Commission on Genetic Resources for Food and Agriculture, Rome, Italy: FAO.
- FAO. (2014). *The state of the world's forest genetic resources*. Rome, Italy: FAO.
- Fine, P. V., & Ree, R. H. (2006). Evidence for a time-integrated species-area effect on the latitudinal gradient in tree diversity. *The American Naturalist*, 168 (6), 796–804. doi:10.1086/508635

- Fitzjohn, R. G., Pennell, M. W., Zanne, A. E., Stevens, P. F., Tank, D. C., & Cornwell, W. K. (2014). How much of the world is woody? *Journal of Ecology*, 102 (5), 1266–1272. doi:10.1111/1365-2745.12260
- Giam, X., Bradshaw, C. J., Tan, H. T., & Sodhi, N. S. (2010). Future habitat loss and the conservation of plant biodiversity. *Biological Conservation*, 143 (7), 1594–1602. doi:10.1016/j.biocon.2010.04.019
- Heywood, V. H., & Watson, R. T. (1995). *Global biodiversity assessment* (Vol. 1140). Cambridge, UK: Cambridge University Press.
- Hunt, D. R. (1996). The genera of temperate broadleaved trees. *Broadleaves*, 2, 4–5.
- Kennedy, C. E. J., & Southwood, T. R. E. (1984). The number of species of insects associated with British trees: A re-analysis. *Journal of Animal Ecology*, 53, 455–478. doi:10.2307/4528
- Kenrick, P., & Crane, P. R. (1997). The origin and early evolution of plants on land. *Nature*, 389, 33–39. doi:10.1038/37918
- Lughadha, E. N., Govaerts, R., Belyaeva, I., Black, N., Lindon, H., Allkin, R., ... Nicolson, N. (2016). Counting counts: Revised estimates of numbers of accepted species of flowering plants, seed plants, vascular plants and land plants with a review of other recent estimates. *Phytotaxa*, 272 (1), 82–88. doi:10.11646/phytotaxa.272.1
- Mark, J., Newton, A., Oldfield, S., & Rivers, M. (2014). *The International timber trade: A working list of commercial timber tree species*. Richmond, UK: Botanic Gardens Conservation International.
- Missouri Botanical Garden (MBG). (2016). *Tropicos.org*. Retrieved December 7, 2016, from <http://www.tropicos.org>
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A. B., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858. doi:10.1038/35002501
- Newton, A., Oldfield, S., Rivers, M., Mark, J., Schatz, G. E., Garavito, N. T., ... Miles, L. (2015). Towards a Global Tree Assessment. *Oryx*, 49 (3), 410–415. doi:10.1017/S0030605315000137
- Oldfield, S., Lusty, C., & MacKinven, A. (1998). *The world list of threatened trees*. Cambridge: World Conservation Press. Cambridge, UK: UNEP-WCMC.
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., ... Kassem, K. R. (2001). Terrestrial ecoregions of the world: A new map of life on Earth. *Bioscience*, 51 (11), 933–938. doi:10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2
- Poorter, L., Van Der Sande, M. T., Thompson, J., Arets, E. J., Alarcón, A., Álvarez-Sánchez, J., ... Peña-Claros, M. (2015). Diversity enhances carbon storage in tropical forests. *Global Ecology Biogeography*, 24, 1314–1328. doi:10.1111/geb.12364
- Rio de Janeiro Botanical Garden. (2016). *Brazilian Flora 2020 in construction*. Retrieved February 1, 2017, from <http://floradobrasil.jbrj.gov.br>
- SANBI. (2014). *Plants of Southern Africa: An online checklist*. Retrieved September 01, 2016, from <http://posa.sanbi.org/searchspp.php>
- Savolainen, O. (2000). *Developing criteria and indicators for genetically sustainable forestry*. In Turok & Gerubek (Eds.) 2000. *International collaboration on forest genetic resources: The role of Europe*. Proceedings of the Second EUFORGEN Steering Committee meeting, Vienna, Austria, Rome: IPGRI. Retrieved from November 26–29, 1998.
- Sharrock, S., Oldfield, S., & Wilson, O. (2014). *Plant conservation report 2014: A review of progress in implementation of the global strategy for plant conservation 2011–2020*. Secretariat of the convention on biological diversity, Montreal, Canada, and Botanic Gardens Conservation International, Richmond, U.K. Technical Series No. 81.
- Simpson, N. (2010). Botanical symbols: A new symbol set for new images. *Botanical Journal of the Linnean Society*, 162, 117–129. doi:10.1111/boj.2010.162.issue-2
- Slik, J. W. F., Arroyo-Rodríguez, V., Aiba, S.-I., Alvarez-Loayza, P., Alves, L. F., Ashton, P., ... Venticinque, E. M. (2015). An estimate of the number of tropical tree species. *Proceedings of the National Academy of Sciences of the United States of America*, 112 (24), 7472–7477. doi:10.1073/pnas.1423147112
- The Plant List (TPL). (2013). *The plant list version 1.1*. Retrieved December 7, 2016, from <http://www.theplantlist.org/>
- Tudge, C. (2006). *The tree: A natural history of what trees are*. New York, NY: Crown Publishers.

- USDA, NRCS. (2016). *The PLANTS database*. Retrieved December 7, 2016, from <http://plants.usda.gov>
- Visseren-Hamakers, I. J., McDermott, C., Marjanneke, J. V., & Cashore, B. (2012). Trade-offs, co-benefits and safeguards: Current debates on the breadth of REDD+. *Current Opinion Environment Sustain*, 4 (6), 646–653. doi:10.1016/j.cosust.2012.10.005
- WCSP. (2016). *World checklist of selected plant families. Facilitated by the Royal Botanic Gardens, Kew*. Retrieved December 31, 2016, from <http://apps.kew.org/wcsp/qsearch.do>
- Western Australian Herbarium. (1998). *FloraBase—The Western Australian Flora. Department of parks and wildlife*. Retrieved September 1, 2016, from <https://florabase.dpaw.wa.gov.au/>
- Wu, Z., Raven, P. H., & Hong, D. (Eds.). (1994–2013). *Flora of China*. St Louis, MO: Science Press, Beijing & Missouri Botanical Garden Press.

Appendix

DATABASE REFERENCE SOURCES

- Abbott, J. R. (2009). *Phylogeny of the Polygalaceae and a revision of Badiera*. Doctoral dissertation, University of Florida, FL.
- Acevedo-Rodríguez, P. (2011). Allophylastrum: A new genus of Sapindaceae from northern South America. *PhytoKeys*, 5, 39–43.
- Acevedo-Rodríguez, P. & Strong, M. T. (2007). Catalogue of the seed plants of the West Indies. Retrieved 1 September 2014, from <http://botany.si.edu/Antilles/WestIndies/>.
- Acevedo-Rodríguez, P. & Strong, M. T. (2012). Catalogue of seed plants of the West Indies. *Smithsonian Contributions to Botany*, 98, 1–92.
- Acevedo-Rodríguez, P. & Brewer, S. W. (2016). *Spathelia belizensis*: A new species and first record for the genus in Central America (tribe Spathelieae, Rutaceae). *PhytoKeys*, 75, 145–151.
- Adema, F. & van der Ham, R. W. J. M. (1993). Cnesmocarpon (*Gen. Nov.*), Jagera and Trigonachras (Sapindaceae-Cupanieae): Phylogeny and systematics. *Blumea*, 38, 73–215.
- Agostini, G. & Fariñas, M. (1963). Holotype of Maytenus agostinii Steyerem. (Celastraceae). Caracas: Fundación Instituto Botánico de Venezuela.
- Akopian, S. S. (2007). On the Pyrus L. (Rosaceae) species in Armenia. *Flora, Vegetation and Plant Resources of Armenia*, 16, 15–26.
- Alejandro, G. J. D. & Meve, U. (2016). Rubovietnamia coronula *sp. nov.* (Rubiaceae: Gardenieae) from the Philippines. *Nordic Journal of Botany*, 34 (2), 385–389.
- Alemayehu, G., Asfaw, Z. & Kelbessa, E. (2016) Cordia africana (Boraginaceae) in Ethiopia: A review on its taxonomy, distribution, ethnobotany and conservation status. *International Journal of Botany Studies*, 1 (2), 38–46.
- Alfarhan, A. H., Al-Turki, T. A. & Basahy, A. Y. (2005). Flora of Jizan Region. Final Report Supported by King Abdulaziz City for Science and Technology. Riyadh, Saudi Arabia.
- Allen, C. K. (1942). Studies in the Lauraceae, IV: Preliminary study of the Paupasian species collected by the Archbold expeditions (concluded). *Journal of the Arnold Arboretum*, 23 (2), 133–155.
- Almeda, F., Marcela, A. & Mendoza, H. (2014). Two new species of Graffenrieda (Melastomataceae: Merianieae) from Colombia and Panama. *Phytotaxa*, 163 (1), 39–47.
- Amann, C., Amann, G., Arhel, R., Guiot, V. & Marquet, G. (2011). *Plantes de Mayotte*. Mamoudzou: Armen Factory.
- Amorim, B. S. & Alves, M. (2016). Taxonomic novelties in Gomidesia clade (Myrcia sl, Myrtaceae) from the Atlantic Forest of Brazil. *Phytotaxa*, 272 (4), 287–293.
- Anderson, C. (2016). Circumscription and nomenclature of Hiraea barclayana, H. reclinata, and H. ternifolia (Malpighiaceae), and of seven species misassigned to them. *Blumea—Biodiversity, Evolution and Biogeography of Plants*, 61 (2), 125–146.

- Andino, J. E. G., Fernandez-Fernandez, D. & Palacios, W. A. (2016). A new species of Sloanea (Elaeocarpaceae) subgenus *Quadrisejala* from Ecuadorian Amazonia. *Phytotaxa*. 263 (2), 147–153.
- Ang, W. F., Lok, A. F. S. L. & Tan, H. T. W. (2010). Rediscovery in Singapore of *Pinanga simplicifrons* (Miq.) Becc. (Arecaceae). *Nature in Singapore*, 3, 83–86.
- Annotated Checklist of the Flowering Plants of Nepal. (2017). Retrieved 1 February 2017, from http://www.efloras.org/flora_page.aspx?flora_id=110.
- ANPSA. (2017). Australia Native Plants Society. Retrieved 1 February 2017, from <http://www.anpsa.org.au>.
- APC. (2015). Australian Plant Census. Retrieved 1 February 2017, from <http://www.anbg.gov.au/chah/apc>.
- APNI. (2015) Australian Plant Name Index. Retrieved 1 February 2017, from <http://www.anbg.gov.au>.
- Applequist, W. L. (2016). A revision of the Malagasy species of *Homalium* sect. *Eumyriantheia* Warb. (Salicaceae). *Candollea*. 71 (1), 33–60.
- Arakaki, M., Ostolaza, C., Cáceres, F. & Roque, J. (2006). Cactaceae endémicas del Perú. *Revista Peruana de Biología*, 13 (2), 193–219.
- Arce, M. R., Gale, S. L. & Maxted, N. (2008). A taxonomic study of *Albizia* (Leguminosae: Mimosoideae: Ingeae) in Mexico and Central America. *Anales del Jardín Botánico de Madrid*. 65 (2), 255–305.
- Arnoldo, M. (2012). *Arnoldo's zakflora: wat in het wild groeit en bloeit op Aruba, Bonaire en Curaçao*. Zutphen: WalburgPers.
- Ashburner, K. & McAllister, H. A. (2013). *The Genus Betula. A taxonomic revision of birches*. London: Kew Publishing.
- Asprino, R. C. & Amorim, A. M. A. (2016). A new species of *Hirtella* (Chrysobalanaceae), and redescription of a closely related taxon, from the Atlantic Forest, Brazil. *Phytotaxa*. 265 (3), 259–266.
- Atha, D. E., Mitchell, J. D., Pell, S. K. & Camacho, F. R. (2011). A new species of *Comocladia* (Anacardiaceae) from Belize and Guatemala. *Brittonia*. 63 (3), 370–374.
- Atlas of Living Australia. (2017). *Atlas of living Australia*. Retrieved 1 February 2017, from <http://www.ala.org.au>.
- Australian Biological Resources Study. (2017). *Flora of Australia online*. Retrieved 1 February 2017, from <http://www.environment.gov.au/biodiversity/abrs/online-resources/flora/main/index.html>.
- Averyanov, L. V., Nguyen, K. S., Nguyen, H. T. & Harder, D.K. (2015). Preliminary assessment for conservation of *Pinus cernua* (Pinaceae) with a brief synopsis of related taxa in eastern Indochina. *Turczaninowia*. 18 (1), 738.
- Aymard, G. (1998). A new species of *Aldina* (Fabaceae: Caesalpinioideae) from Venezuelan Amazonia. *Novon a Journal for Botanical Nomenclature*. 8 (4), 330–331.
- Baba, Y. & Crayn, D. (2012). *Elaeocarpus hylobroma* (Elaeocarpaceae): A new species endemic to mountain tops in north-east Queensland, Australia. *Kew Bulletin*. 67 (4), 743–750.
- Baba, Y. (2013). *Evolution, systematics and taxonomy of Elaeocarpus (Elaeocarpaceae) in Australasia*. Doctoral dissertation, James Cook University, Queensland.
- Bacchetta, G., Fenu, G., Mattana, E., Zecca, G., Grassi, F., Casazza, G., et al. (2011). Genetic variability of the narrow endemic *Rhamnus persicifolia* Moris (Rhamnaceae) and its implications for conservation. *Biochemical Systematics and Ecology*. 39 (4), 477–484.
- Bajpai, O., Kumar, A., Srivastava, A. K., Kushwaha, A. K., Pandey, J. & Chaudhary, L.B. (2015). Tree species of the Himalayan Terai region of Uttar Pradesh, India: A checklist. *Check List*. 11 (4), 1718.
- Balachandran, N. & Rajendiran, K. (2016). *Cordia ramanujamii* (Cordiaceae): New species from Tamil Nadu, India. *Taiwania*. 61 (2), 74–77.
- Baldeón, S., Flores, M. & Roque, J. (2006). Fabaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 302–337.
- Ban, N. T., Ly, T. D. & Khoi, N. K. (2007). *Vietnam Red Data Book, Part II. Plants*. Hanoi: Natural Science and Technology Publishing.

- Bartholemew, B., Boufford, D. E., Chang, A. L., Cheng, Z., Dudley, T. R., He, S. A., et al. (1983). The 1980 Sino-American botanical expedition to western Hubei province, People's Republic of China. *Journal of the Arnold Arboretum*. 64 (1), 1–103.
- Bastin, J. F., Barbier, N., Réjou-Méchain, M., Fayolle, A., Gourlet-Fleury, S., Maniatis, D., et al. (2015). Seeing Central African forests through their largest trees. *Scientific reports*, 5, 13156.
- Beentje, H. J., & Smith, S. A. L. (Eds.). (2001). *Flora of tropical East Africa*. Crown agents for Overseas governments.
- Beltrán, H., Granda, A., León, B., Sagástegui, A., Sánchez, I. & Zapata, M. (2006). Asteraceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 64–164.
- Benitez de Rojas, C. & D'Arcy, W. G. (1998). The genera *Cestrum* and *Sessea* (Solanaceae: Cestreae) in Venezuela. *Annals of the Missouri Botanical Garden*. 85 (2), 273–351.
- Berendsohn, W. G., Gruber, A. K. & Monterrosa Salomón, J. (2012). Nova Silva Cuscatlanica. Árboles nativos e introducidos de El Salvador. Parte 2: Angiospermae—Familias M a P y Pteridophyta. *Englera*. 29 (2), 1–300.
- Berendsohn, W. G., Gruber, A. K. & Monterrosa Salomón, J. A. (2009) Nova Silva Cuscatlanica. Árboles nativos e introducidos de El Salvador. Parte 1: Angiospermae—Familias A a L. *Englera*. 29 (1), 1–438.
- Berg, C. C. (2002). *Ficus* subgenus *Pharmacosycea* section *Oreosycea* (Moraceae) in the Solomon Islands, Fiji, and the New Hebrides. *Blumea*. 47 (2), 299–314.
- Berg, C. C. (2004). Flora Malesiana precursor for the treatment of Moraceae 6: *Ficus* subgenus *Sycomorus*. *Blumea*. 49 (1), 155–200.
- Berg, C. C. (2007). New species of *Ficus* (Moraceae) from South America. *Blumea*. 52 (3), 569–594.
- Berg, C. C. (2007). Precursory taxonomic studies on *Ficus* (Moraceae) for the Flora of Thailand. *Thai Forest Bulletin (Botany)*, 35, 4–28.
- Bernal, R., Gradstein, S. R. & Celis, M. (Eds.). (2015). *Catálogo de plantas y líquenes de Colombia*. Retrieved 1 February 2017, from <http://catalogoplantasdecolombia.unal.edu.co/es/>.
- Beyra, A. & Lavin, M. (1999). Monograph of *Pictetia* (Leguminosae-Papilionoideae) and review of the *Aeschynomeneae*. *Systematic Botany Monographs*, 56, 1–93.
- Bhuinya, T. & Mukherjee, S. K. (2012). Status of *Litsea* Lam. (Lauraceae) in Andaman and Nicobar Islands, India. Paper Presented at the International Seminar on “Multidisciplinary Approaches in Angiosperm Systematics”. India: Department of Botany, University of Kalyani.
- Bhuinya, T., Singh, P. & Mukherjee, S. K. (2010). An account of the species of *Litsea* Lam. (Lauraceae) endemic to India. *Bangladesh Journal of Plant Taxonomy*. 17 (2), 183–191.
- Bhutan Biodiversity Portal (2017). *Bhutan Biodiversity Portal*. Retrieved 1 February 2017, from <http://www.biodiversity.bt>
- Bingham, M., Willemen, A., Ballings, P. & Hyde, M. A. (2017). *Flora of Zambia*. Retrieved 1 February 2017, from <http://www.zambiaflora.com/>
- Biodiversity Informatics and Co-Operation in Taxonomy for Interactive shared Knowledge base (2008). *Biotik*. Retrieved 1 February 2017, from www.biotik.org
- Biosecurity Queensland. (2017). *Weeds of Australia*. Biosecurity Queensland Edition. Retrieved 1 February 2017, from <http://keyserver.lucidcentral.org/weeds/data/media/Html/index.htm>.
- Birnbaum, P., Ibanez, T., Pouteau, R., Vandrot, H., Hequet, V., Blanchard, E., et al. (2015). Environmental correlates for tree occurrences, species distribution and richness on a high-elevation tropical island. *AoB Plants*, 7, 1–15.
- Bissiengou, P. & Sosef, M. S. M. (2008). Novitates Gabonenses 69. A new endemic species of and a new combination in *Campylopermum* (Ochnaceae). *Blumea*. 53 (3), 627–631.
- Borhidi, I. (1914). *The Flora of the Dutch West Indian Islands. II. The Flora of Curaçao, Aruba and Bonaire*. Leiden: Brill.
- Bollendorff, S. M., Van Welzen, P. C. & Slik, J. W. F. (2000). A taxonomic revision of *Mallotus* section *Polyadenii* (Euphorbiaceae). *Blumea*. 45 (2), 319–340.
- Borhidi, A. L., Oviedo-Prieto, R. & Fernández-Zequeira, M. (2016). Nuevos resultados de la revisión taxonómica de los géneros *Palicourea* y *Psychotria* (Rubiaceae, Psychotrieae) en Cuba. *Acta Botanica Hungarica*. 58 (1–2), 1–48.

- Bostock, P. D. & Holland, A. E. (2016) *Census of the Queensland Flora 2016*. Queensland Department of Science, Information Technology and Information: Brisbane. Retrieved 1 February 2017, from <http://www.data.qld.gov.au/dataset/census-of-the-queensland-flora-2016>.
- Botanical Survey of India. (2017). *eFlora of India*. Retrieved 1 February 2017, from <http://efloraindia.nic.in/efloraindia/homePage.action>.
- BRAHMS. (2017). *Plants of Namibia*. Retrieved 1 February 2017, from <http://herbaria.plants.ox.ac.uk/bol/namibia>.
- BRAHMS. (2017). *The National Herbarium of Trinidad and Tobago*. Available at: <http://herbaria.plants.ox.ac.uk/bol/trin>.
- Brambach, F., Coode, M., Biagioni, S. & Culmsee, H. (2016). *Elaeocarpus firdausii* (Elaeocarpaceae): A new species from tropical mountain forests of Sulawesi. *PhytoKeys*, 62, 1–14.
- Breitwieser I., Brownsey P. J., Nelson W. A. & Wilton A.D. (Eds.). (2016–2017). *Flora of New Zealand Online*. Retrieved 1 February 2017, from <http://www.nzflora.info>.
- Breteler, F. J. (2016). Novitates Gabonenses 84. A new species of Wilczekia (Celastraceae) from Gabon. *Plant Ecology and Evolution*. 149 (1), 131–134.
- Breteler, F. J. (Ed.). (1989). *The Connaraceae: A taxonomic study with emphasis on Africa*. Agricultural University: Michigan.
- Brummitt, N. A. & Utteridge, T. M. A. (2003). A second species of Distyliopsis (Hamamelidaceae) from New Guinea. *Kew Bulletin*. 58 (3), 727–731.
- Bünger, M. D. O., Mazine, F. F. & Stehmann, J. R. (2016). *Eugenia ruschiana* (Myrtaceae): A new species from southeastern Brazilian Atlantic forest. *Phytotaxa*. 266 (1), 40–44.
- Bunwong, S., Chantaranothai, P. & Keeley, S.C. (2014) Revisions and key to the Vernoniaceae (Compositae) of Thailand. *PhytoKeys*, 37, 25–101
- Bussmann, R. W., Gruhn, J. & Glenn, A. (2010). *Axinaea fernando-cabiesii* and *A. reginae* spp. nov. (Melastomataceae) from upper Amazonia of Peru, with notes on the conservation status of *A. flava*. *Nordic Journal of Botany*. 28 (5), 518–522.
- Byng, J. W., Barthelat, F., Snow, N. & Bernardini, B. (2016). Revision of *Eugenia* and *Syzygium* (Myrtaceae) from the Comoros archipelago. *Phytotaxa*. 252 (3), 163–184.
- Byng, J.W., Florens, F.V. & Baider, C. (2015). *Syzygium pyneei* (Myrtaceae): A new critically endangered endemic species from Mauritius. *PhytoKeys*, 46, 61–66.
- Byng, J. W., Barthelat, F., Snow, N. & Bernardini, B. (2016). Revision of *Eugenia* and *Syzygium* (Myrtaceae) from the Comoros archipelago. *Phytotaxa*. 252 (3), 163–184.
- CABI (2017) *CABI: Invasive Species Compendium*. Retrieved 1 February 2017, from <http://www.cabi.org/isc/>.
- Cabral, F. N., Bittrich, V. & Amaral, M. D. C. E. D. (2016). Two new species and one new record of *Caraipa* (Calophyllaceae) from Colombia. *Systematic Botany*. 41 (2), 348–353.
- The Calflora Database (2017). *Calflora: Information on California plants for education, research and conservation*. Retrieved 1 February 2017, from <http://calflora.org>.
- Caliari, C. P., Souza, V. C. & Mazine, F. F. (2016). Two new species of *Myrcia* (Myrtaceae) from Brazilian Atlantic Forest. *Phytotaxa*. 267 (3), 201–210.
- Callmänder, M. W., Lowry, P. P., Forest, F., Devey, D. S., Beentje, H. & Buerki, S. (2012). *Benstonea* Callm. & Buerki (Pandanaceae): Characterization, circumscription, and distribution of a new genus of screw-pines, with a synopsis of accepted species. *Candollea*. 67 (2), 323–345.
- Camargo, R. A. & de Azevedo Tozzi, A. M. G. (2014). A synopsis of the genus *Deguelia* (Leguminosae, Papilionoideae, Millettieae) in Brazil. *Brittonia*. 66 (1), 12–32.
- Campos-Ríos, M. & Chiang, F. (2010). *Bourreria tuxtlae* y *Bourreria veracruzana* (Boraginaceae) dos especies nuevas y clave para la identificación de las especies arbóreas del género, en México. *Polibotánica*, 30, 01–07.
- Cantley, J. T., Sporck-Koehler, M. J. & Chau, M. M. (2016). New and resurrected Hawaiian species of *pilo* (Coprosma, Rubiaceae) from the island of Maui. *PhytoKeys*, 60, 33.
- Cardoso, D., De Lima, H. C. & De Queiroz, L. P. (2013). *Staminodianthus*: A new neotropical Genistoid legume genus segregated from *Diploptropis*. *Phytotaxa*. 110 (1), 1–16.

- Carrington, S., Sabir, K., Broome, R. & Acevedo Rodriguez, P. (2007). *Plants of the Eastern Caribbean*. Barbados: The University of the West Indies. Retrieved 1 February 2017, from <http://ecflora.cavehill.uwi.edu/>
- Carter, S. & Wood, J. R. I. (1982). Two new succulent Euphorbia species from southwest Arabia. *Kew Bulletin*, 37 (1), 73–76.
- Castillo-Campos, G., García-Franco, J.G. & Martínez, M. L. (2013). *Spathacanthus magdalenae* sp. Nov. (Acanthaceae): A riparian forest species from Veracruz, Mexico. *Nordic Journal of Botany*, 31 (4), 449–452.
- Cayzer, L. W., Crisp, M. D. & Telford, I.R. (2000). *Auranticarpa*: A new genus of Pittosporaceae from northern Australia. *Australian Systematic Botany*, 13 (6), 903–917.
- Centre for Plant Biodiversity Research. (2017). *EUCLID*. Australian National Herbarium. Third Edition. Retrieved 1 February 2017, from <https://www.anbg.gov.au/cpbr/cd-keys/euclid3/>
- Chakrabarty, T., Lakra, G. S. & Diwakar, P. G. (2010). The Family Lauraceae in Andaman and Nicobar Islands. In Ramakrishna, C. Raghunathan & Sivaperuman, C. *Recent Trends in Biodiversity of Andaman and Nicobar Islands*. 179–193.
- Chanco, M., León, B. & Sánchez, I. (2006). Malvaceae endémicas del Perú. *Revista Peruana de Biología*, 13 (2), 413–425.
- Chang, C. & Kim, H. (2015). *The woody plants of Korea*. Retrieved 1 February 2017, from <http://florakorea.myspecies.info/en>.
- Chantaranothai, P. (2014). Three new records of Syzygium Gaertn. in Thailand and lectotypification of 19 taxa of Eugenia L. (Myrtaceae). *Thai Forest Bulletin (Botany)*, 42, 75–80.
- Chantaranothai, P. & Parnell, J. (1994). Revision of Acmena, Cleistocalyx, Eugenia s.s. and Syzygium (Myrtaceae) in Thailand. *Thai Forest Bulletin (Botany)*, 21, 1–123.
- Chaowasku, T. & Van Der Ham, R. W. (2013). Integrative systematics supports the establishment of Winitia: A new genus of Annonaceae (Malmeoideae, Miliuseae) allied to Stelechocarpus and Sageraea. *Systematics and Biodiversity*, 11 (2), 195–207.
- Chaowasku, T., Johnson, D. M., Van Der Ham, R. W. & Chatrou, L. W. (2015). Huberantha: A replacement name for Hubera (Annonaceae: Malmeoideae: Miliuseae). *Kew Bulletin*, 70 (2), 1–4.
- Cheek, M. & Haba, P. (2016). Spiny African Allophylus (Sapindaceae): A synopsis. *Kew Bulletin*, 71 (4), 57.
- Chileflora (2012). *Chileflora*. Retrieved 1 February 2017, from <http://www.chileflora.com/>.
- Chilebosque (2017). *Chilebosque*. Retrieved 1 February 2017, from <http://www.chilebosque.cl/>.
- Chong, K. Y., Tan, H. T. & Corlett, R. T. (2009). *A checklist of the total vascular plant flora of Singapore: native, naturalised and cultivated species*. Singapore: Raffles Museum of Biodiversity Research, National University of Singapore.
- Christensen, K. I. & Zieliński, J. (2008). Notes on the genus Crataegus (Rosaceae-Pyreae) in southern Europe the Crimea and western Asia. *Nordic Journal of Botany*, 26, 344–360.
- Chua, L. S. L., Suhaida, M., Hamidah, M. & Saw, L. G. (2010). *Malaysia Plant Red List, Peninsular Malaysian Diperocarpaceae*. Research Pamphlet No. 129. Forest Research Institute. Malaysia.
- Chung, R. C. K., Soepadmo, E. & Lim, A. L. (2005). A synopsis of the Bornean species of Microcos L. (Tiliaceae). *Gardens' Bulletin of Singapore*, 57, 101–130.
- Chung, R. C. K. (2001). Taxonomic notes on the Bornean Helicia and Heliciopsis (Proteaceae). *Journal of Tropical Forest Science*, 13 (3), 534–547.
- Chung, R.C.K., Tan, H.S. & Soepadmo, E. (2012). A remarkable new species of Jarandersonia (Malvaceae-Brownlowioideae) from Central Kalimantan, Borneo. *Systematic Botany*, 37 (1), 213–217.
- Cicuzza, D., Newton, A. & Oldfield, S. (2007). *The Red List of Magnoliaceae*. Cambridge: Fauna & Flora International.
- CIRAD. (2008). *Arbres, arbustes de la forêt réunionnaise*. Retrieved 1 February 2017, from <http://arbres-reunion.cirad.fr/>.
- Conn, B. & Damas, K. (2017) *Guide to trees of Papua New Guinea—PNGtrees descriptions*. Retrieved 1 February 2017, from <http://www.pngplants.org/PNGtrees/TreeDescriptions/>.
- Conn, B.J. (1995). *Handbooks of the flora of Papua New Guinea: Volume 3*. Melbourne: Melbourne University Press.

- Conservatoire et Jardin Botaniques Ville de Genève (2017). *African Plant Database*. Retrieved 1 April 2016, from <http://www.ville-ge.ch/musinfo/bd/cjb/africa/>.
- Cook Islands Government. (2007). *Cook Islands biodiversity database*. Retrieved 1 February 2017, from <http://cookislands.bishopmuseum.org/>.
- Cornejo, X. & Iltis, H. H. (2008). A revision of *Colicodendron* (Capparaceae). *Journal of the Botanical Research Institute of Texas*. 2 (1), 75–93.
- Cornejo, X. & Iltis, H. H. (2010). Three new species of *Capparidastrum* (Capparaceae) from Colombia, Venezuela, and Ecuador. *Harvard Papers in Botany*. 15 (1), 155–163.
- Cornejo, X., Lombardi, J. A. & Thomas, W. W. (2011). *Chionanthus parviflora*: A new species of Oleaceae Endemic to Northeastern Brazil. *Harvard Papers in Botany*. 16 (2), 421–423.
- Corner, E. J. (1970). New species of *Streblus* and *Ficus* (Moraceae). *Blumea*, 18, 393–411.
- Costion, C. M. & Lorence, D. H. (2012). The endemic plants of Micronesia: A geographical checklist and commentary. *Micronesia*. 43 (1), 51–100.
- Costion, C. M. & Plunkett, G. M. (2016). A revision of the genus *Osmoxylon* (Araliaceae) in Palau, including two new species. *PhytoKeys*, 58, 49–64.
- Costion, C. M., Kitalong, A. H. & Holm, T. (2009). Plant endemism, rarity, and threat in Palau, Micronesia: A geographical checklist and preliminary Red List assessment. *Micronesica*. 41 (1), 131–164.
- Cotton, E. & Meier, W. (2003). *Clidemia intonsa* and *Miconia chapensis* (Miconieae, Melastomataceae): Two new species endemic to cloud forest refuges in the Coastal Cordillera of Venezuela. *Willdenowia*. 33 (1), 197–203.
- Couvreux, T. L. & Niangadouma, R. (2016). New species of *Uvariopsis* (Annonaceae) and *Laccosperma* (Arecaceae/Palmae) from Monts de Cristal, Gabon. *PhytoKeys*. 68, 1–8.
- Couvreux, T. L., Niangadouma, R., Sonké, B. & Sauquet, H. (2015). *Sirdavidia*: An extraordinary new genus of Annonaceae from Gabon. *PhytoKeys*, 46, 1–19.
- Croizat, L. (1938). Notes on Chinese Euphorbiaceae. *Journal of the Arnold Arboretum*. 19 (2), 134–148.
- CSIRO. (2010) *Australian Tropical Rainforest Plants Edition 6*. Retrieved 1 February 2017, from <http://keys.trin.org.au/key-server/data/0e0f0504-0103-430d-8004-060d07080d04/media/Html/index.html>.
- Culmsee, H., Pitopang, R., Mangopo, H. & Sabir, S. (2011). Tree diversity and phytogeographical patterns of tropical high mountain rain forests in Central Sulawesi, Indonesia. *Biodiversity and Conservation*. 20 (5), 1103–1123.
- Daly, D. C. & Martínez-Habibe, M. C. (2016). Seven new species of *Dacryodes*. *Brittonia*, 68 (2), 120–137.
- Daly, D. C. (2005). *Protium aidanianum*: A new species from Western Amazonia. Studies in Neotropical Burseraceae XII. *Novon*. 15 (3), 410–413.
- Danin, A. (2003–2017) *Flora of Israel Online*. Retrieved 1 February 2017, from <http://flora.org.il/>
- Darbyshire, I. & Luke, Q. (2016). *Barleria mirabilis* (Acanthaceae): A remarkable new tree species from west Tanzania. *Kew Bulletin*. 71 (1), 1–6.
- Darbyshire, I., Kordofani, M., Farag, I., Candiga, R. and Pickering, H. (2015). The plants of Sudan and South Sudan: An annotated checklist. Richmond: Royal Botanic Gardens, Kew.
- Darwin, S. P. (2010). A taxonomic revision of *Timonius* subgen. *Pseudobobea* (Valeton) S.P. Darwin (Rubiaceae). *Candollea*. 65 (2), 217–240.
- Davila, N. & Kinoshita, L. S. (2016). A new species of *Platycarpum* (Rubiaceae, Henriquezieae) from Peruvian Amazon. *Phytotaxa*. 260 (3), 276–282.
- Dávila, N. & Vicentini, A. (2016). *Isertia psammophila* (Isertiaeae, Rubiaceae): A new species from white-sand areas of the Brazilian Amazon. *Phytotaxa*, 257 (2), 174–180.
- Dawson, S. E. (2016). Characteristics of the Madagascan genus *Canephora* (Rubiaceae: Octotropideae), and the description of two new species. *Kew Bulletin*, 71 (3), 1–10.
- de Carvalho, L. D. A. F. (1991). New Taxa of *Solanum* (Solanaceae) from Brazil, Colombia, Central America, and Venezuela. *Annals of the Missouri Botanical Garden*. 78 (1), 224–244.
- de Carvalho-Sobrinho, J.G., Machado, M.C. & de Queiroz, L.P. (2012). *Spirotheca elegans* (Malvaceae: Bombacoideae): A new species from Bahia, Brazil. *Systematic Botany*. 37 (4), 978–982.

- de Faria, J. E. Q., Mazine, F.F. & Proença, C. E. B. (2015). Two new species of *Eugenia* (Myrtaceae) from the Cabo Frio Center of Plant Diversity, Rio de Janeiro, Brazil. *Phytotaxa*. 208 (3), 201–208.
- de Granville, J. J. (2007). A new species of *Bactris* (Palmae) from French Guiana. *Brittonia*. 59 (4), 354–356.
- de Kok, R. (2013). The genus *Premna* L. (Lamiaceae) in the Flora Malesiana area. *Kew Bulletin*. 68 (1), 55–84.
- de Kok, R. P., Rusea, G. & Latiff, A. (2009). The genus *Teijsmanniodendron* Koord. (Lamiaceae). *Kew Bulletin*. 64 (4), 587–625.
- de Kok, R. P. J. (2015). A revision of *Cryptocarya* (Lauraceae) from Thailand and Indo-China. *Garden's Bulletin Singapore*. 67 (2), 309–350.
- de Kok, R. P. J. (2016). A revision of *Beilschmiedia* (Lauraceae) of peninsular Malaysia. *Blumea*. 61 (2), 147–164.
- de Kok, R. P. J. (2016). A revision of *Cryptocarya* R. Br. (Lauraceae) of Peninsular Malaysia. *Kew Bulletin*. 71 (1), 1–26.
- De Lima, L. F. G., Baumgratz, J. F. A., Lughadha, E. N. & Dos Santos, J. U. M. (2016). Two new species of *Graffenrieda* (Melastomataceae, Merianieae) from the Amazon Rainforest. *Phytotaxa*. 267 (1), 77–83.
- De Queiroz, L.P., Lewis, G.P. & Allkin, R. (1999). A revision of the genus *Moldenhawera* Schrad. (Leguminosae-Caesalpinioideae). *Kew Bulletin*. 54 (4), 817–852.
- de Salas, M.F. & Baker, M.L. (2016). *A census of the vascular plants of Tasmania, including Macquarie Island. (Tasmanian Herbarium, Tasmanian Museum and Art Gallery. Hobart)* Retrieved 1 February 2017, from <http://www.tmag.tas.gov.au>.
- De Sanctis, M., Adeeb, A., Farcomeni, A., Patriarca, C., Saed, A. & Attorre, F. (2012). Classification and distribution patterns of plant communities on Socotra Island, Yemen. *Applied Vegetation Science*, 16, 148–165.
- de Stapf, M.N.S. & dos Santos Silva, T. (2013). Typifications in *Cordia* (Cordiaceae), with an assessment of the status of *Cordia blanchetii*. *Kew Bulletin*. 68 (2), 355–359.
- De Stefano, R.D., Janovec, J.P. & Can, L.L. (2013). Three decades to connect the sexes: *Calatola microcarpa* (Icacinaeae): A new species from the Southwestern Amazon. *Phytotaxa*. 124 (1), 43–49.
- De Wilde, W. J. J. O. (2002). Additions to Asian Myristicaceae: *Endocomia*, *Gymnacranthera*, *Horsfieldia*, *Knema*. *Blumea*. 47 (2), 347–362.
- Demaio, P., Karlin, U. O. & Medina, M. (2015). *Árboles Nativos de Argentina*. Argentina: Ecoval Ediciones.
- Deomurari, A., Jani, M., Matieda, K. & KaPatel, J. (2014) *IBIS-Flora Beta Version—Angiosperm Flora of India*. Retrieved 1 February 2017, from <http://www.flora.indianbiodiversity.org/flora/angiosperms>.
- Devecchi, M. F., Thomas, W. W. & Pirani, J. R. (2016). *Simaba arenaria* (Simaroubaceae): A new species from sandy coastal plains in Northeastern Brazil, with notes on seedling morphology. *Systematic Botany*. 41 (2), 401–407.
- Dludlu, M. N., Muasya, A. M., Chimphango, S. B. & Stirton, C. H. (2015). Taxonomy of the Southern African *Psoralea aphylla* complex (Psoraleae, Leguminosae). *South African Journal of Botany*, 97, 77–100.
- Dombo, A., Da Costa, E. & Neto, G. (2002). *Zambia Plant Red Data List*. Retrieved 1 February 2017, from <http://www.nationalredlist.org/zambia-plant-red-data-list-2002/>.
- Donnell, A. (2012). *A systematic revision of Bakeridesia Hochr. (Malvaceae)*. Doctoral dissertation, Ohio University, United States.
- Dressler, S. (1999). Revision of the genus *Cleistanthus* (Euphorbiaceae) in the Philippines. *Blumea*. 44(1), 109–148.
- East African Plant Red List Authority (EAPRLA). (2015). Tree assessments from EAPRLA workshops I-X. Nairobi, Kenya.
- Eastwood, A. (2005). *Globally threatened trees of the Caucasus. A report on the Caucasus regional tree Red Listing workshop held in Tbilisi, Georgia, 26–28 September 2005*. Cambridge: Fauna & Flora International.

- Eastwood, A., Lazkov, G. & Newton, A. (2009) *The Red List of trees of Central Asia*. Cambridge: Fauna & Flora International.
- Endemia. (2016). *Endemia: Faune et Flore de Nouvelle-Caledonie*. Retrieved 1 February 2017, from <http://endemia.nc>.
- Esser, H. & Saw, L. G. (2015). A new species of *Polyosma* (Escalloniaceae/Polyosmaceae) from Thailand and new synonyms. *Phytotaxa*. 221 (1), 89–92.
- Euro+Med (2006–). Euro+Med PlantBase: The information resource for Euro-Mediterranean plant diversity. Retrieved 1 February 2017, from <http://ww2.bgbm.org/EuroPlusMed/>.
- Ewango, C. E., Kenfack, D., Sainge, M. N., Thomas, D. W. & van der Burgt, X. M. (2016). *Gambeya korupensis* (Sapotaceae: Chrysophylloideae): A new rain forest tree species from the Southwest Region in Cameroon. *Kew Bulletin*, 71 (2), 1–6.
- Falanruw, M. V. C. (2015). *Trees of Yap: A field guide*. California: U.S. Department of Agriculture—Forest Service, Pacific Southwest Research Station.
- Farjon, A. (2001). *A world checklist and bibliography of conifers* (2nd ed.). Richmond: Royal Botanic Gardens Kew.
- Farjon, A. & Filer, D. (2013). *An atlas of the world's conifers*. Leiden: Brill.
- Fauna & Flora of Liberia (2017) *Plant atlas of Liberia*. Retrieved 1 February 2017, from <http://www.liberianfaunaflora.org/plant-atlas-of-liberia>.
- Fauset, S., Johnson, M. O., Gloor, M., Baker, T. R., Monteagudo, A., Brien, R. J., et al. (2015). Hyperdominance in Amazonian forest carbon cycling. *Nature Communications*, 6, 6857.
- Fedde, F. (1921). *Repertorium specierum novarum regni vegetabilis*. Fasciculus XVII. Berlin: Selbstverlag des Herausgebers.
- Fernández-Alonso, J. L. & Cogollo-Pacheco, Á. A. (2016). *Chionanthus megistocarpus* (Oleaceae): A new species from the Western Cordillera of Colombia. *Phytotaxa*. 269 (1), 14–20.
- Fernandez-Fernandez, D., Ulloa, C. U. & Penneys, D. S. (2016). A new species of *Blakea* (Blakeaceae, Melastomataceae) from Ecuador. *Phytotaxa*. 284 (1), 69–74.
- Fernando, E. S., Co, L. C., Lagunzad, D. A., Gruezo, W. S., Barcelona, J. F., Madulid, D. A., et al. (2008). Threatened plants of the Philippines: A preliminary assessment. *Asia Life Sciences, Supplement 3*, 1–52.
- Feulner, G. (1997). First observations of *Olea cf. europaea* (the Wild Olive) and *Ehretia obtusifolia* in the United Arab Emirates. *Tribulus*. 7 (1), 12–14.
- Fiaschi, P. (2016). A new species of *Dendropanax* (Araliaceae) from the Brazilian Atlantic Forest. *Brittonia*. 68 (2), 103–110.
- Fiaschi, P. & Plunkett, G. M. (2016). Two new species of the *Didymopanax* clade of *Schefflera* (Araliaceae) from the Brazilian Amazon. *Phytotaxa*. 245 (2), 153–160.
- Field Museum (1999–2016). *Amostras de Herbário da Neotrópica*. Retrieved 1 February 2017, from <http://fm1.fieldmuseum.org/vrrc/index.php?PHPSESSID=5c08cd33d8686913db18e82fbc1b5555>.
- Figueiredo, E. (2010). Unpublished assessments of trees of S. Tomé and Príncipe (draft). Centro de Ecologia, Funcional, University of Coimbra. Portugal.
- Figueiredo, E. & Smith, G. F. (2008). *Plants of Angola/Plantas de Angola*. Strelitzia 22. South African National Biodiversity Institute, Pretoria.
- Figueiredo, E., Paiva, J., Stévant, T., Oliveira, F. & Smith, G. F. (2011). Annotated catalogue of the flowering plants of São Tomé and Príncipe. *Bothalia*. 41 (1), 41–82.
- Filipe de Portugal, S. T., Lewis, G. P. & Hawkins, J. A. (2010). A revision of the South American genus *Apuleia* (Leguminosae, Cassieae). *Kew Bulletin*. 65 (2), 225–232.
- Filipowicz, N., Nee, M. & Renner, S. (2012). Description and molecular diagnosis of a new species of *Brunfelsia* (Solanaceae) from the Bolivian and Argentinean Andes. *PhytoKeys*, 10, 83–94.
- Flora Malesiana (2017) *Flora Malesiana*. Retrieved 1 February 2017, from <http://portal.cybertaxon.org/flora-malesiana/>.
- Flora of North America Editorial Committee (Eds.). (1993+). *Flora of North America North of Mexico*. 20+ vols. New York and Oxford.
- Flores, R., Ibanez, A., & Correa, M. D. (2016). *Eugenia veraguensis* (Myrtaceae): A new species from Golfo de Chiriquí in Veraguas Province, Panama, with notes on *Eugenia rhombea*. *Phytotaxa*. 270(3), 217–222.

- FOI (2017). *Flowers of India*. Retrieved 1 February 2017, from <http://www.flowersofindia.net/>.
- Forero, E. (1983). Connaraceae. *Flora Neotropica*, 36, 1–207.
- Forest Research Institute Malaysia. (2015). *Malaysia Flora Database*. Retrieved 1 February 2015, <http://www.chm.frim.gov.my/Bio-Diversity-Databases/Flora-Database.aspx>
- Fosberg, F. R., Sachet, M. H., & Oliver, R. (1979). Geographical checklist of the Micronesian Dicotyledonae. *Micronesica*, 15 (1–2).
- Freire, S.E., Katinas, L. & Sancho, G. (2002). Gochnatia (Asteraceae, Mutisieae) and the Gochnatia complex: Taxonomic implications from morphology. *Annals of the Missouri Botanical Garden*, 89 (4), 524–550.
- Friedmann, F. (1994). Flore des Seychelles: dicotylédones. IRD Editions, Collection Faune et Flore tropicales 44. Marseille, France.
- Fritsch, P. W., & Al, F. (2015). A taxonomic revision of Antillean Symplocos (Symplocaceae). *Phytotaxa*, 194 (1), 1–67.
- Funk, V. A., Berry, P., Alexander, S., Hollowell, T.H. & Kelloff, C. L. (2007). *Checklist of the plants of the Guiana Shield (Venezuela: Amazonas, Bolivar, Delta Amacuro; Guyana, Surinam, French Guiana)*. Washington, DC: National Museum of Natural History.
- Gagnidze R. I. (Ed.) (1985–2011). *Saqartvelos flora (Flora of Georgia)* (2nd ed.). Tbilisi: Metsniereba.
- Gagnon, E., Bruneau, A., Hughes, C. E., de Queiroz, L. & Lewis, G.P. (2016). A new generic system for the pantropical Caesalpinia group (Leguminosae). *PhytoKeys*, 71, 1–160.
- Gahagen, B., Terbush, M. & Ballard, H. (2016). A taxonomic synthesis of the Tovomita weddelliana (Clusiaceae) species complex. *Systematic Botany*, 40 (4), 968–988.
- Gaikwad, K. N. & Mali, M. V. (2012). Tree flora of Nashik city (Maharashtra). *International Journal of Life Science & Pharma Research*, 2 (2), 94–101.
- Gamba-Moreno, D. L. and Almeda, F. (2014). Systematics of the Octopleura clade of Miconia (Melastomataceae: Miconieae) in tropical America. *Phytotaxa*, 179 (1), 1–174.
- Gann, G. D., J. C. Trejo-Torres and C. G. Stocking. (2015–2016). Plantas de la Isla de Puerto Rico/ Plants of the Island of Puerto Rico. Florida: The Institute for Regional Conservation.
- Gao, L. M. & Li, D. Z. (2009). Rhododendron qiaojiaense (Ericaceae): A new species from Yunnan, China. *Annales Botanici Fennici*, 46 (1), 67–70.
- Garavito, N. T., Dávila, E. A., Caro, S. A., Murakami, A. A., Baldeón, S., Beltrán, H., et al. (2014) *A regional Red List of Montane tree species of the tropical Andes: Trees at the top of the world*. Richmond: Botanic Gardens Conservation International.
- GBIF. (2017). Global Biodiversity Information Facility. Retrieved 1 February 2016, from <http://www.gbif.org/>
- Germishuizen, G., Meyer, N. L., Steenkamp, Y. & Keith, M. (Eds.) (2006). *A checklist of South African plants*. Southern African Botanical Diversity Network Report No. 41. SABONET, Pretoria, Republic of South Africa.
- Gibbs, D. & Chen, Y. (2009). *The Red List of Maples*. Cambridge: Fauna & Flora International.
- Gibbs, D., Chamberlain, D. & Argent, G. (2011). *The Red List of rhododendrons*. Richmond: Botanic Gardens Conservation International.
- Gibbs, P. & Semir, J. (2003). A taxonomic revision of the genus Ceiba Mill. (Bombacaceae). *Anales del Jardín Botánico de Madrid*, 60(2), 259–300.
- Gibbs, P. E. & Alverson, W. S. (2006). How many species of Spirotheca (Malvaceae sl, Bombacoideae)? *Brittonia*, 58(3), 245–258.
- Gilles, D., Zaiss, R., Blach-Overgaard, A., Catarino, L., Damen, T., Deblauwe, V. et al. (2016). RAINBIO: A mega-database of tropical African vascular plants distributions. *PhytoKeys*, 74, 1–18.
- Giriraj, A., Irfan-Ullah, M., Ramesh, B. R., Karunakaran, P. V., Jentsch, A. & Murthy, M. S. R. (2008). Mapping the potential distribution of Rhododendron arboreum Sm. ssp. nilagiricum (Zenker) Tagg (Ericaceae): An endemic plant using ecological niche modelling. *Current Science*, 94 (12), 1605–1612.
- GISD. (2017). *The Global Invasive Species Database*. IUCN. Retrieved 1 February 2017, from <http://www.iucngisd.org/gisd/>.

- National Museum of Nature and Science. (2017). *Global Red List of Japanese threatened plants*. Retrieved 1 February 2017, from https://www.kahaku.go.jp/english/research/db/botany/redlist/list/list_div.html.
- Goldenberg, R. & Meirelles, J. (2011). A new species of Graffenrieda (Merianieae, Melastomataceae) with a calyptrate calyx. *Systematic Botany*. 36 (1), 119–123.
- Goldenberg, R., Almeda, F., Caddah, M. K., Martins, A. B., Meirelles, J., Michelangeli, F. A. & Weiss, M. (2013). Nomenclator botanicus for the neotropical genus Miconia (Melastomataceae: Miconieae). *Phytotaxa*. 106 (1), 1–171.
- Goldenberg, R., Michelangeli, F. A., Aona, L. Y., & Amorim, A. M. (2016). Angiosperms and the Linnean shortfall: Three new species from three lineages of Melastomataceae at one spot at the Atlantic Forest. *PeerJ*, 4, e1824.
- Gómez, J. R. D. S., & Michelangeli, F. A. (2016). Miconia portogallensis (Melastomataceae): A new species from the Sierra Madre del Sur, Guerrero, Mexico. *Phytotaxa*. 278 (2), 132–140.
- González Gutiérrez, P. A. (2014). *Evolution and biogeography of Buxus L. (Buxaceae) in Cuba and the Caribbean*. Doctoral dissertation, Freie Universität Berlin, Germany.
- González-Espinosa, M., Meave, J. A., Lorea-Hernández, F. G., Ibarra-Manríquez, G. & Newton, A. C. (2011). *The Red List of Mexican cloud forest trees*. Cambridge: Fauna & Flora International.
- Gordon, J. (2005). *Red Listing of Mesoamerican dry forest tree species*. Cambridge: Fauna & Flora International.
- Gosline, G. & Malécot, V. (2011). A monograph of Octoknema (Octoknemaceae—Olacaceae s.l.). *Kew Bulletin*. 66 (3), 367–404.
- Gottschling, M. & Miller, J. S., (2007). A revision of Bourreria (Boraginales, Ehretiaceae) in South America. *Annals of the Missouri Botanical Garden*. 94 (4), 734–744.
- Government of India. (2017). Botanical survey of India. Retrieved 1 February 2017, from <http://bsi.gov.in/Index.aspx>.
- Grandtner, M. M. (2005). *Elsevier's dictionary of trees. Volume 1: North America*. Amsterdam: Elsevier Science.
- Grandtner, M.M. & Chevrete, J. (2014). *Elsevier's dictionary of trees. Volume 2: South America*. Cambridge, United States: Academic Press.
- Graveson, R. (2015). Plants of Saint Lucia. Retrieved 1 September 2015, from <http://saintlucianplants.com/>.
- Grimshaw, J. (2003). Notes on the temperate species of Juglans. *IDS Yearbook*. 107–130.
- Grossheim, A. A. (1939–1967). *Flora Kavkaza, Vol. I-II, Baku, Azerbaijan, Izdatel'stvo AzΦAH, Vol. III, Baku, Azerbaijan, Izdatel'stvo Akademii Nauk Azerbajjanskoi SSR, Vol. IV-VI. Leningrad, Izdatel'stvo Akademii Nauk SSSR, Vol. VII. Leningrad, Russian Federation: Nauka*.
- Grossheim, A. A. (1949). *Opređelitel' Pactenii Kavkaz*. Moscow: Sovetckaia Nauka.
- Guerrero, A., Judd, W. S. & Morris, A. B. (2004). A new species of Illicium subsection Parviflora (Illiciaceae) from the Massif de la Hotte, Haiti. *Brittonia*. 56 (4), 346–352.
- Guillaumin, A. N. D. R. (1931). Contribution to the Flora of the New Hebrides: Plants collected by SF Kajewski in 1928 and 1929. *Journal of the Arnold Arboretum*, 12 (4), 221–264.
- Guo, W., Fan, Q., Jing, H., Hu, X., & Liao, W. (2016). Sorbus calcicola (Rosaceae): A new species from Guangxi Province in China. *Phytotaxa*. 261 (3), 260–266.
- Gustafsson, M. H., & Borchsenius, F. (2016). Clusia nubium (Clusiaceae): A new species from cloud-forests of southwestern Ecuador. *Phytotaxa*. 253 (3), 219–222.
- Hackel, E. (1908). Notes on Philippine Gramineae, III. *The Philippine Journal of Science*, 3 (C), 172.
- Haegens, R. M. A. P. (2000). *Taxonomy, phylogeny, and biogeography of Baccaurea, Distichirhops, and Nothobaccaurea (Euphorbiaceae)*. Doctoral dissertation, Universiteit Leiden, The Netherlands.
- Hahn, N. (2016). Senegalia montis-salinarum: A new species of Fabaceae: Mimosoideae endemic to the Soutpansberg, South Africa. *Phytotaxa*. 244 (2), 174–180.
- Haretche, F., Mai, P. & Brazeiro, A. (2012). Woody flora of Uruguay: Inventory and implication within the Pampean region. *Acta Botanica Brasilica*. 26 (3), 537–552.
- Harriet, G.J. & Kerry, W.S. (1998). *1997 IUCN Red list of threatened plants*. Gland: IUCN.

- Hartley, T.G. (1974). A revision of the genus *Acronychia* (Rutaceae). *Journal of the Arnold Arboretum*. 55 (3), 469–523.
- Harvard University (2017). Harvard University Herbaria and Libraries. Retrieved 1 February 2017, from <http://huh.harvard.edu/>.
- Harwood, B. & Tathana, N. (2011) Two new *Casearia* (Salicaceae) species from Thailand, and notes on *Casearia grewiaefolia* var. *gelonioides*. *Thai Forest Bulletin (Botany)*, 39, 23–27.
- Hassler, M. (2017). World plants: Synonymic checklists of the vascular plants of the world (version Nov 2016). In: Roskov, Y., Abucay, L., Orrell, T., Nicolson, D., Bailly, N., Kirk, P. et al. (Eds.). (2017). *Species 2000 & ITIS catalogue of life*. Retrieved 1 February 2017, from <http://www.catalogueoflife.org>.
- Heatubun, C. D., Baker, W. J., Mogea, J. P., Harley, M. M., Tjitrosoedirdjo, S. S. & Dransfield, J. (2009). A monograph of *Cyrtostachys* (Arecaceae). *Kew Bulletin*. 64 (1), 67–94.
- Hendrian. (2004). Revision of *Ochrosia* (Apocynaceae) in Malaysia. *Blumea*. 49 (1), 101–128.
- Hernandez, F.G.L. (1997). On *Cinnamomum* (Lauraceae) in Mexico. *Acta Botánica Mexicana*, 40, 1–18.
- Hind, D. J. N. (2011). *An annotated preliminary checklist of the Compositae of Bolivia. Version 2*. Richmond: Royal Botanic Gardens, Kew
- Hoogland, R. D. (1979). Studies in the Cunoniaceae II. The Genera *Caldcluvia*, *Pullea*, *Acsmithia*, and *Spiraeanthemum*. *Blumea*. 25 (2), 481–505.
- Hopkins, H. C. F. & Pillon, Y. (2011). Further new endemic taxa of Cunoniaceae from New Caledonia. *Kew Bulletin*. 66 (3), 405–423.
- Huamantupa, I., Vásquez, R., Foster, R., Cuba, M. & Calatayud, G. (2014). Adiciones de angiospermas a la Flora del Perú procedentes de los bosques Andino Amazónicos del sur peruano. *Revista Peruana de Biología*. 21 (2), 163–170.
- Editorial Committee of the Flora of Taiwan. (Ed.). (1975–). *Flora of Taiwan*. Taipei: Department of Botany, National Taiwan University.
- Hughes, M. (2013). *Memecylon pseudomegacarpum* M. Hughes (Melastomataceae): A new species of tree from Peninsular Malaysia. *European Journal of Taxonomy*, (56), 1–9.
- Hunter, J. T. (2001). *Eucalyptus saxicola* (Myrtaceae): A new species from the Central Tablelands of New South Wales (section *Maidenaria* series *Bridgesianae*). *Telopea*. 9 (2), 403–407.
- Hyde, M. A., Wursten, B. T., Ballings, P. & Coates Palgrave, M. (2017). Flora of Zimbabwe. Retrieved 1 February 2017, from <http://zimbabweflora.co.zw/>.
- IDigBio. (2017). *Integrated digitized biocollections*. National Science Foundation. Retrieved 1 February 2017, from <https://www.idigbio.org/>.
- India Biodiversity Portal (2013). *India biodiversity portal*. Retrieved 1 September 2016, from <http://indiabiodiversity.org/>.
- International Legume Database & Information Service. (2006–2013). *ILDIS World Database of Legumes Version 10.38*. Retrieved 1 September 2016, from <http://www.legumes-online.net/>.
- IPNI. (2012). *The International Plant Names Index (IPNI)*. Retrieved 1 September 2016, from http://www.ipni.org/how_to_cite_us.html
- IUCN France, MNHN and DIREN Polynésie Française. (2015). Liste rouge de la Flore vasculaire endémique de Polynésie française (2015). Available at: https://inpn.mnhn.fr/espece/listerouge/FR/Flore_Vasculaire_Polynesie_2015
- IUCN. (2016). *The IUCN Red List of threatened species. Version 2016–3*. Retrieved 16 December 2016, from <http://www.iucnredlist.org>.
- Jacks, B. R. (2005). Revision of *Myrsine* (Myrsinaceae) in Australia. *Australian Systematic Botany*. 18 (5), 399–438.
- Jiménez, J. E., Juárez, P., & Chaves-Fallas, J. M. (2016). *Cupania moralesii* (Sapindaceae): A new endemic tree species from the premontane forest of Costa Rica. *Phytotaxa*. 275 (1), 69–74.
- Jimenez-Ramirez, J., González, C. A., & Rios-Carrasco, S. (2016). *Triumfetta acahuizotlanensis* (Malvaceae): A new tree species endemic of Guerrero, Mexico. *Phytotaxa*. 265 (3), 273–278.
- Jing, H., Fan, Q., Guo, W., & Liao, W. (2016). *Sorbus prunifolia* (Rosaceae): A new species from Yunnan Province, China. *Phytotaxa*. 258 (2), 171–178.
- JSTOR. (2017). Global Plants. ITHAKA. Retrieved 1 February 2017, from <http://plants.jstor.org/>.

- Judkevich, M. D., Salas, R. M., & Keller, H. (2016). *Randia brevītuba* (Rubiaceae): A new species from the Southern Cone of America and comments on *Randia armata*. *Systematic Botany*, 41 (1), 238–244.
- Kalema, J. & Beentje, H. (2012). *Conservation checklist of the trees of Uganda*. Richmond: Kew Publishing.
- Kariagin, I. I. (Ed.). (1950, 1953–1957). *Flora Azerbaijana (Flora of Azerbaijan)*. Vol. I, IV–VII. Baku, Azerbaijan: Izdatel'stvo Akademii Nauk Azerbajanskoy SSR.
- Kawasaki, M. L., & Pérez, Á. J. (2016). New species of Myrtaceae from Yasuní National Park, Ecuador. *Harvard Papers in Botany*, 21(1), 55–58.
- Kelly, L. M., Almeda, F., & Fritsch, P. W. (2016). A taxonomic revision of Mexican and Central American Symplocos (Symplocaceae). *Phytotaxa*, 264(1), 1–115.
- Keng, H. (1984). *Flora malesianae* precursores. LVIII, part 2: The genus *Gordonia* (Theaceae) in Malesia. *Garden's Bulletin Singapore*, 37, 1–47.
- Ketskhoveli, N. (Ed.-in-Chief) (1971–1984). *Saqartvelos Flora* (2nd ed.), Vol. I–IX. Tbilisi: Metsniereba.
- Khumchompoo, S. & Thongpukdee, A. (2005). *Miliusa longiflora* (Hook. f. & Thomson) Baill. ex Finet & Gagnep. (Annonaceae): A new record for Thailand. *Thai Forest Bulletin (Botany)*, 33, 32–34.
- Kiet, L. C., Kessler, P. J. & Eurlings, M. (2005). A new species of *Aquilaria* (Thymelaeaceae) from Vietnam. *Blumea*, 50 (1), 135–141.
- Knapp, S. (2009). Synopsis and lectotypification of *Solanum* (Solanaceae) species endemic in the West Indies. *Anales del Jardín Botánico de Madrid*, 66 (1), 65–84.
- Knapp, S., Spooner, D. M. & León, B. (2006). Solanaceae endémicas del Perú. *Revista Peruana de Biología*, 13 (2), 612–643.
- Koenen, E. J. & de Wilde, J. J. (2012). A taxonomic revision of the reinstated genus *Leplaea* and the newly recognized genus *Neoguarea* (Meliaceae, Sapindales): The exclusion of *Guarea* from Africa. *Plant Ecology and Evolution*, 145(2), 209–241.
- Köhler, E. (2006). Three new *Buxus* species (Buxaceae) from eastern Cuba. *Willdenowia*, 36(1), 479–489.
- Köhler, E., Fernández, R. & Zamudio, S. (1993). *Buxus moctezumae* Köhler, Fernández et Zamudio (Buxaceae) una especie nueva de Estado de Querétaro, México Con 37 Figuras. *Feddes Repertorium*, 104(5–6), 295–305.
- Komarov, V. L. (Ed.) (1934–1964). *Flora SSSR*, Vol I–IV, Leningrad, Russian Federation: Vol. VII–XXX. Moskva-Leningrad, Russian Federation: Izdatel'stvo Akademii Nauk SSSR.
- Kostermans, A. (1959). A monograph of the genus *Heritiera* Aiton (Sterculiaceae) (including *Argyrodendron* F. v. M. and *Tarrietia* Bl.). *Reinwardtia*, 4 (4), 465–583.
- Kostermans, A. (1960). Miscellaneous botanical notes 2. *Reinwardtia*, 5 (4), 375–411.
- Kostermans, A. (1969). Materials for a revision of Lauraceae II. *Reinwardtia*, 7 (1), 451–536.
- Kostermans, A. J. (1965). A monograph of the genus *Parinari* Aubl. (Rosaceae-Chrysobalanoideae) in Asia and the Pacific region. *Reinwardtia*, 7 (2), 147–213.
- Kress, W. J., DeFilipps, R. A., Farr, E., Kyi, D. Y. Y., Angrist, J. D., Pischke, J. S. et al. (2003). *A checklist of the trees, shrubs, herbs, and climbers of Myanmar*. Washington DC, United States: National Museum of Natural History.
- Kriebel, R. (2016). A monograph of *Conostegia* (Melastomataceae, Miconieae). *PhytoKeys*, 67, 1–326.
- Krishnamani, R., Kumar, A. & Harte, J. (2004). Estimating species richness at large spatial scales using data from small discrete plots. *Ecography*, 27 (5), 637–642.
- Kulju, K. K. M., Sierra, S. E. C. and Van Welzen, P. C. (2007). Re-shaping *Mallotus* [part 2]: Inclusion of *Neotrewia*, *Octospermum* and *Trewia* in *Mallotus* s.s. (Euphorbiaceae s.s.). *Blumea*, 52 (1), 115–136.
- Lachashvili, N. J., Eradze, N. V. and Khetsuriani, L. D. (2016): Conspectus of trees and shrubs of Tbilisi environs (East Georgia, South Caucasus). *Annals of Agrarian Science*, 14.
- Lachenaud, O. & Séné, O. O. (2012). Un nouveau *Multidentia* (Rubiaceae) d'Afrique centrale. *Plant Ecology and Evolution*, 145 (1), 132–137.

- Lara, L. J. C., Berrazaín Iturralde, R., Leiva Sánchez, A. T. & Oldfield S. (2005). *Memorias del primer Taller para la Categorización de Árboles Cubanos. Grupo de Especialistas en Plantas de Cuba, Flora & Fauna International*. Cuba: Jardín Botánico Nacional, La Habana.
- Larsen, K. & Ming, H. C. (1991). New taxa of Myrsinaceae from Thailand. *Nordic Journal of Botany*. 11 (1), 61–78.
- Leenhouts, P. W. (1986). A taxonomic revision of Nephelium (Sapindaceae). *Blumea*. 31 (2), 373–436.
- Leenhouts, P. W. & Vente, M. (1982). A taxonomic revision of Harpullia (Sapindaceae). *Blumea*. 28 (1), 1–51.
- León, B. (2006). Clusiaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 261–264.
- León, B. (2006). Ericaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 285–293.
- León, B. (2006). Flacourtiaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 338.
- León, B. (2006). Lauraceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 380–388.
- León, B. (2006). Myrsinaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 458–462.
- León, B. (2006). Sapindaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 605–607.
- León, B. (2006). Symplocaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 647–649.
- León, B. (2006). Tiliaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 655.
- León, B. & Monsalve, C. (2006). Annonaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 35–41.
- León, B. & Monsalve, C. (2006). Apocynaceae endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 46–48.
- León-Yáñez, S., Valencia, R., Pitman, N., Endara, L., Ulloa Ulloa C. & Navarrete H., (Eds.) (2011). *Libro Rojo de las Plantas Endémicas del Ecuador. 2nd edición*. Quito, Ecuador: Publicaciones del Herbario QCA,
- Lim, C. K., Dransfield, J., Kiew, R. & Saw, L. G. (1998). Four new Pinanga Blume (Palmae) species from Peninsular Malaysia. *Garden's Bulletin Singapore*. 50 (1), 99–114.
- Lim, T. K. (2012). *Edible medicinal and non-medicinal plants. Volume 1. Fruits*. New York, United States: Springer.
- Linares, J. & Sousa, M. (2007). Nuevas especies de Dalbergia (Leguminosae: Papilionoideae: Dalbergieae) en México y Centroamérica. *Ceiba*. 48(1–2), 61–82.
- Liu, B., & Qin, H. N. (2013). Taxonomic revision of the Symplocos nakaharae complex (Symplocaceae) with special reference to fruit morphology. *Journal of Systematics and Evolution*, 51 (1), 94–114.
- Loeuille, B., & Pirani, J. R. (2016). Two new syncephalous species of Eremanthus (Asteraceae: Vernonieae) from southeastern Brazil. *Phytotaxa*. 243 (2), 128–136.
- Loizeau, P. & Barriera, G. (2007). *Aquifoliaceae néotropicales: Descriptions, Illustrations, Identification, et Recherche d'Information*. Retrieved 1 September 2016, from <http://www.ville-ge.ch/cjb/>.
- Loots, S. (2005). *A Red Data Book of Namibian Plants*. SABONET Report No. 38. Pretoria.
- Loots, S. (2010). *Unpublished assessments of the Namibian Botanical Research*. Institute of Namibia. Namibia.
- Lorence, D. H. & Wood, K. R. (2012). Psychotria kosraensis (Rubiaceae): A new species from Kosrae, Caroline Islands, Micronesia. *Novon: A Journal for Botanical Nomenclature*. 22 (1), 51–55.
- Lozada-Pérez y, L., & Rojas Gutiérrez, J. (2016). Una Nueva Especie, Omiltemia guerrerensis (Rubiaceae) de Guerrero, México. *Novon: A Journal for Botanical Nomenclature*. 24(4), 365–368.
- Lozano, P. & Klitgaard, B. B. (2006). The genus Machaerium (Leguminosae: Papilionoideae: Dalbergieae) in Ecuador. *Brittonia*. 58(2), 124–150.
- Luoma-aho, T. (2004). Forest Genetic Resources Conservation and Management: Proceedings of the Asia Pacific Forest Genetic Resources Programme (APFORGEN) Inception Workshop, Kepong, Kuala Lumpur, Malaysia, 15–18 July, 2003. Bioversity International.
- Luong, V. D., Luu, H. T., Nguyen, T. Q. T. & Nguyen, Q. D. (2016) Camellia luteopallida (Theaceae): A new species from Vietnam. *Annales Botanici Fennici*. 53(1–2), 135–138.
- M. B. Thomas, J. F. Rock Herbarium, University of Hawaii at Manoa. (2015). *Consortium of Pacific Herbaria Database (CPH)*. Retrieved 1 September 2016, from <http://www.pacificherbaria.org/>

- Maas, P. J., Westra, L. Y. T. & Chatrou, L. W. (2003). *Duguetia* (Annonaceae). *Flora Neotropica*, 88, 1–274.
- Maas, P. J. M., Westra, L. Y. T., Guerrero, S. A., Lobão, A. Q., Scharf, U., Zamora, N.A. et al. (2015). Confronting a morphological nightmare: Revision of the Neotropical genus *Guatteria* (Annonaceae). *Blumea*, 60(1–2), 1–219.
- Mackinder, B. A. & Clark, R. (2012). *Baphia rosa* (Leguminosae: Papilionoideae): A new species from the Kabompo District of Zambia and a review of the classification of *B. chrysophylla*, *B. claessensii* and *B. gillettii* from the Democratic Republic of the Congo. *Kew Bulletin*, 67(3), 413–420.
- Mackinder, B. A. & Harris, D. J. (2006). A synopsis of the genus *Berlinia* (Leguminosae–Caesalpinioideae). *Edinburgh Journal of Botany*, 63(2–3), 161–182.
- Mackinder, B. A. & van der Burgt, X.M. (2009). *Berlinia korupensis* (Leguminosae–Caesalpinioideae): A new tree species from Cameroon. *Kew Bulletin*, 64(1), 129–134.
- Mackinder, B. A. & Wieringa, J. J. (2007). Novitates Gabonenses 58: Two new species of *Berlinia* (Leguminosae–Caesalpinioideae: Detarieae). *Kew Bulletin*, 62(1), 159–164.
- Mackinder, B. A. & Wieringa, J.J. (2013). *Hymenostegia viridiflora* (Detarieae, Caesalpinioideae, Leguminosae): A new tree species from Cameroon. *Blumea*, 58(1), 13–17.
- Mackinder, B. A., Wieringa, J. J. & van der Burgt, X. M. (2010). A revision of the genus *Talbotiella* Baker f.(Caesalpinioideae: Leguminosae). *Kew Bulletin*, 65(3), 401–420.
- Madriñán, S. (2004). *Rhodostemonodaphne* (Lauraceae). *Flora Neotropica*, 92, 1–102.
- Makashvili, A. (1952–3). *Tbilisi Midamoebis Flora. Volume 1–2*. Tbilisi: Publishing House of the Stalin Tbilisi State University.
- Marinho, L. C., Gahagen, B., & Amorim, A. M. (2016). Description of *Tovomita clarkii* (Clusiaceae): An endemic species from Venezuela. *Phytotaxa*, 261 (1), 87–91.
- Marinho, L. C., Martins, M. V., Amorim, A. M., & Bittrich, V. (2016). *Vismia atlantica* (Hypericaceae): A new species previously thought to be well-known from the Brazilian Atlantic Forest 1. *The Journal of the Torrey Botanical Society*, 143 (3), 330–337.
- Mark, J., Newton, A., Oldfield, S. & Rivers, M. (2014). The International Timber Trade: A working list of commercial timber tree species. Richmond: BGCI.
- Marques, C. P., Fonseca, T., Ferreira, M. & Laranjeira, P. (2010). First forest inventory of Timor-Leste. Districts of Bobonaro and Covalima (2008|2009). Final Report. Universidade de Trás-os-Montes e Alto Douro, Portugal Google Scholar.
- Marshall, A. R., Couvreur, T. L., Summers, A. L., Deere, N. J., Luke, W. Q., Ndangalasi, H. J et al. (2016). A new species in the tree genus *Polyceratocarpus* (Annonaceae) from the Udzungwa Mountains of Tanzania. *PhytoKeys*, 63, 63–76.
- Marshall, A. J. & Beehler B. M. (2007). *The Ecology of Indonesian Papua, Part Two*. Tuttle Publishing.
- Martinelli, G. & Moraes M. A. (2013). *Livro vermelho da flora do Brasil*. Rio de Janeiro, Brazil: Instituto de Pesquisas Jardim Botânico do Rio de Janeiro, Centro Nacional de Conservação da Flora.
- Masri, T., Nuwayhid, R. Y., Masri, N. & Fakhry, K. (2009). Autoecology of *Castanea sativa* Mill. in Lebanon. *Acta Hort.*, 815, 83–94.
- Maxwell, J. F. & Veldkamp, J. F. (1990). Notes on the Astronieae (Melastomataceae)—I. *Astrocalyx*, *Astronia*. *Blumea*, 35, 71–114.
- Mazine, F. F., & Souza, V. C. (2015). Lectotypifications and a new combination in *Eugenia* sect. *Racemosae* (Myrtaceae). *Phytotaxa*, 205 (3), 157–167.
- Mazine, F.F. & Faria, J.E.Q. (2013). A new species of *Eugenia* (Myrtaceae) from South America. *Phytotaxa*, 151 (1), 53–57.
- Meerts, P. (2016). An annotated checklist to the trees and shrubs of the Upper Katanga (DR Congo). *Phytotaxa*, 258 (3), 201–250.
- Meireles, J. E. & Tozzi, A. M. G. A. (2015). *Limadendron*: A new genus of Leguminosae (Papilionoideae, Brongniartieae) from South America. *Plant Systematics and Evolution*, 301 (2), 701–707.
- Meirelles, J., Lima, D. F., & Goldenberg, R. (2016). *Miconia astrocalyx* (Melastomataceae, Miconieae): A new species from Brazilian Cerrado. *Phytotaxa*, 257(2), 187–192.

- Meléndez González, P. A. (2009). Sinopsis del Genero *Machaerium* (Leguminosae Papilionoideae) en Venezuela. *Acta Botánica Venezuelica*. 32(2), 363–416.
- Mendoza, W., Rodríguez, M., Rodríguez, E., Weigend, M., Chanco, M., Ortiz, R., et al. (2006). El libro rojo de las plantas endémicas del Perú. *Revista Peruana de Biología*. 13 (2), 64–164.
- Merrill, E. D. (1909). New or Noteworthy Philippine Plants, VII. *The Philippine Journal of Science*, 4, C.
- Michelangeli, F.A. & Goldenberg, R. (2014). A new species of Graffenrieda (Melastomataceae) from the northern Amazon basin. *Brittonia*. 66 (2), 170–173.
- Ministerio de Medio Ambiente y Agua. (2012). *Libro Rojo de la Flora amenazada de Bolivia*. Vol. I. *Zona Andina*. La Paz, Bolivia: Ministerio de Medio Ambiente y Agua.
- Missouri Botanical Garden. (2017). *Pakistan plant database*. Retrieved 1 February 2017, from <http://www.tropicos.org/project/Pakistan>.
- Missouri Botanical Garden. (2017). *A world checklist of Thymelaeaceae*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/Thymelaeaceae>.
- Missouri Botanical Garden. (2017). *Bolivia catalogue*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/BC>.
- Missouri Botanical Garden. (2017). *Catálogo de las Plantas Vasculares de la Selva Central, Peru*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/SCP>.
- Missouri Botanical Garden. (2017). *Catalogue of the vascular plants of Ecuador*. Retrieved 1 February 2017, from <http://tropicos.org/Project/CE>.
- Missouri Botanical Garden. (2017). *Catalogue of the vascular plants of Madagascar*. Available at: <http://www.tropicos.org/Project/MADA>
- Missouri Botanical Garden. (2017). *Catalogue of the vascular plants of the Department of Antioquia (Colombia)*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/CV>.
- Missouri Botanical Garden. (2017). *eFloras.org*. Retrieved 1 February 2017, from <http://www.efloras.org/>.
- Missouri Botanical Garden. (2017). *Flora de Nicaragua*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/FN>
- Missouri Botanical Garden. (2017). *Flora Mesoamericana*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/FM>.
- Missouri Botanical Garden. (2017). *Flora of Panama*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/FOPWFO>.
- Missouri Botanical Garden. (2017). *Index to the climate change vulnerability of Andean plants*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/AndeanClimateChange>.
- Missouri Botanical Garden. (2017). *Listado de la Flora del Parque Nacional Madidi, Bolivia*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/MDICHK>.
- Missouri Botanical Garden. (2017). *Manual de Plantas de Costa Rica*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/Costa%20Rica>.
- Missouri Botanical Garden. (2017). *Panama checklist*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/PAC>.
- Missouri Botanical Garden. (2017). *Paraguay checklist*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/Paraguay>.
- Missouri Botanical Garden. (2017). *Peru checklist*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/PEC>.
- Missouri Botanical Garden. (2017). *Selected Rubiaceae Tribes and Genera*. Retrieved 1 February 2017, from <http://www.tropicos.org/Project/Rubiaceae>.
- Missouri Botanical Garden. (2017). *Tropicos*. Retrieved 1 February 2017, from <http://www.tropicos.org/Home.aspx>.
- Missouri Botanical Garden. (2017). *WBN flora database*. Available at: <http://www.tropicos.org/Project/Weda%20Bay>.
- Mitchell, J. D. & Daly, D. C. (2015). A revision of *Spondias* L. (Anacardiaceae) in the Neotropics. *PhytoKeys*, (55), 1–92.
- Moffett, R. O. (2007). Name changes in the Old World *Rhus* and recognition of *Searsia* (Anacardiaceae). *Bothalia*. 37 (2), 165–175.

- Monk, K., De Fretes, Y. & Reksodiharjo-Lilley, G. (1997). Ecology of Nusa Tenggara and Maluka. The Ecology of Indonesia Series vol 5. Dalhousie University, Singapore.
- Morales, J. (2006). Novedades en las Proteaceae de Costa Rica. *Darwiniana, nueva serie*, 44 (2), 490–492.
- Morales, M., & Fortunato, R. H. (2016). A new xerophytic species of Mimosa (Mimosoideae, Leguminosae) from Madagascar. *Phytotaxa*, 270 (4), 277–285.
- Mori, S.A. & Cornejo, X. (2013). Two new species (*Gustavia johnclarkii* and *G. hubbardiorum*) and other contributions to the systematics of *Gustavia* (Lecythidaceae). *Brittonia*, 65 (3), 330–341.
- MSBP. (2003). Darwin Initiative Research Exercise on Community Tree Seeds, UK Workshop 2003.
- Müller, J. (2006). Systematics of *Baccharis* (Compositae-Astereae) in Bolivia, including an overview of the genus. *Systematic Botany Monographs*, 76, 1–341.
- Munzinger, J., Lowry, P. P., Buerki, S., & Callmander, M. W. (2016). A taxonomic revision of the endemic new Caledonian Genus *Storthocalyx* (Sapindaceae). *Systematic Botany*, 41 (2), 387–400.
- Muscarella, R., Uriarte, M., Erickson, D. L., Swenson, N. G., Zimmerman, J. K., et al. (2014). A well-resolved phylogeny of the trees of Puerto Rico based on DNA barcode sequence data. *PLoS ONE*, 9 (11), e112843.
- Muséum national d'Histoire naturelle. (2017). *Inventaire National du Patrimoine Nature (INPN)*. Retrieved 1 February 2017, from <https://inpn.mnhn.fr/accueil/index?lg=en>.
- MyBIS. (2015). Malaysia Biodiversity Information System (MyBIS). Retrieved 1 February 2017, from <http://www.mybis.gov.my>.
- Nagamasu, H., Rueangruea, S., Suddee, S. & Tagane, S. (2015). *Prunus kaengkrahanensis* (Roseaceae), a new species from Southwestern Thailand. *Thai Forest Bulletin (Botany)*, 43, 43–45.
- National Red List. (2016). *The National Red List*. Retrieved 1 September 2016, from <http://www.nationalredlist.org>.
- Naturalis Biodiversity Center. (2017). *BioPortal*. Retrieved 1 February 2017, from <http://bioportal.naturalis.nl/>.
- NatureFiji. (2017). *NatureFiji—MareqetiViti-MareqetiViti*. Retrieved 1 February 2017, from <https://naturefiji.org/>.
- New Zealand Plant Conservation Network. (2017). *New Zealand Plant Conservation Network*. Available at: <http://www.nzpcn.org.nz/>.
- Newman, M., Ketphanh, S., Svengsuksa, B., Thomas, P., Sengdala, K., Lamxay, V. et al. (2007). *A checklist of the vascular plants of Lao PDR*. Scotland: Royal Botanic Garden Edinburgh.
- Ng, F .S. P. (2005). Taxonomic notes on Bornean *Litsea*, *Lindera*, *Neolitsea* and *Iteadaphne* (Lauraceae). *Garden's Bulletin Singapore*, 57, 217–246.
- Ngearnsaengsarua, C., Middleton, D. J. & Chayamarit, K. (2011). A revision of the genus *Litsea* Lam.(Lauraceae) in Thailand. *Thai Forest Bulletin (Botany)*, 39, 40–119.
- Nishida, S. (2008). Taxonomic revision of *Beilschmiedia* (Lauraceae) in Borneo. *Blumea*, 53 (2), 345–383.
- Nishida, S., (1999). Revision of *Beilschmiedia* (Lauraceae) in the neotropics. *Annals of the Missouri Botanical Garden*, 86 (3), 657–701.
- Noblick, L.R. & Lorenzi, H. (2010). New *Syagrus* species from Brazil. *Palms*, 54 (1), 18–42.
- Ohwi, J., (1984). *Flora of Japan. A combined much revised, and extended translation by the author of his Flora of Japan (1953) and Flora of Japan: Pteridophyta (1957)*. Washington, DC: Smithsonian Institution.
- Oldfield, S. & Eastwood, A. (2007). *The Red List of Oaks*. Cambridge: Fauna & Flora International.
- Ordas, J. A. D., Taradji, A. R. J., Valdez Jr, M. B., Banag, C. I., & Alejandro, G. J. D. (2016). *Neonauclea conicalycina*: A new myrmecophytic species of Naucleaeae (Rubiaceae) from Cebu, Philippines. *Phytotaxa*, 273 (2), 127–132.
- Ortiz-Díaz, J. J., Arnelas, I., & Pinzon, J. P. (2016). *Coccoloba tunii* (Polygonaceae): A new species from Chiapas (Mexico). *Phytotaxa*, 275 (1), 75–80.
- Ortiz-Rodriguez, A. E., Escobar-Castellanos, M. A., & Pérez-Farrera, M. A. (2016). Phylogenetic analyses and morphological characteristics support the description of a second species of *Tridimeris* (Annonaceae). *PhytoKeys*, 74, 79.

- Ortiz-Rodriguez, A. E., Ramos, C. M. B., & Gomez-Dominguez, H. (2016). A new species of *Amphitecna* (Bignoniaceae) endemic to Chiapas, Mexico. *PhytoKeys*, 65, 15–23.
- Orwa, C., Mutua, A., Kindt, R., Jamnadass, R. & Simons, A. (2009). *Macaranga tanarius* Muell. Arg. (Euphorbiaceae). *Agroforestry Database 4.0*. Retrieved 1 September 2016, from http://www.worldagroforestry.org/treedb/AFTPDFS/Macaranga_tanarius.pdf
- Österreichische Bundesforste. (2014). *Forest carbon inventory in proposed central Suau REDD+ Area, Milne Bay Province, Papua New Guinea*. Suva, Fiji: SPC/GIZ Regional REDD+ Project.
- Ottens-Treurniet, M. A. D., & van Welzen, P. C. (2016). A revision of the Malesian genus *Blumeodendron* (Euphorbiaceae). *Blumea-Biodiversity, Evolution and Biogeography of Plants*, 61 (1), 64–82.
- Page, N. (2016). New tree species described from the Western Ghats, India. *Oryx*, 50 (4), 582–582.
- Palmpedia. (2015). *Palmpedia: A Palm grower's guide*. Retrieved 1 September 2016, from <http://www.palmpedia.net/>.
- Palmweb. (2017). *Palmweb: Palms of the world online*. Retrieved 1 February 2017, from <http://palmweb.org/>.
- Pannell, C. M. (2008). A key to *Aglaia* (Meliaceae) in Australia, with a description of a new species, *A. cooperae*, from Cape York Peninsula, Queensland. *Journal of the Adelaide Botanic Garden*, 22, 67–71.
- Pathak, M. K., Bhaumik, M. & Krishna, G. (2014). A new species of *Adinandra* (Pentaphylacaceae) from India. *Edinburgh Journal of Botany*, 71 (03), 379–383.
- Pelissari, G. & Neto, S. R. (2014). *Ficus tubulosa* (Moraceae): A new Amazonian species and the re-establishment of *Ficus trachelosyce*. *Phytotaxa*, 170 (3), 207–212.
- Pendry, C. A. (2004). Monograph of *Ruprechtia* (Polygonaceae). *Systematic Botany Monographs*, 67, 1–113.
- Pennington, R. T., Daza, A., Reynel, C. and Lavin, M. (2011). *Poissonia eriantha* (Leguminosae) from Cuzco, Peru: An overlooked species underscores a pattern of narrow endemism common to seasonally dry neotropical vegetation. *Systematic Botany*, 36 (1), 59–68.
- Pennington, T. D. (2016). Systematic treatment of American *Trichilia* (Meliaceae). *Phytotaxa*, 259 (1), 18–162.
- Pennington, T. D., & Clarkson, J. J. (2013). A revision of *Guarea* (Meliaceae). *Edinburgh Journal of Botany*, 70 (2), 179.
- Pennington, T. D. & Clarkson, J. J. (2013). A revision of *Guarea* (Meliaceae). *Edinburgh Journal of Botany*, 70 (2), 179–362.
- Pennington, T. D. & Clarkson, J. J. (2016). A revision of American *Trichilia* (Meliaceae). *Phytotaxa*, 259 (1), 1–2.
- Pennington, T. D. & Mori, S. A. (1993). *Guarea michel-moddei* (Meliaceae): A new species from central French Guiana. *Brittonia*, 45 (3), 231–234.
- PFAF. (2017). *Plants for a future*. Retrieved 1 February 2017, from <http://pfaf.org>.
- Philipson, W. R. (1976). A synopsis of the Malesian species of *Osmoxylon* (including *Boerlagiodendron*), Araliaceae. *Blumea*, 23 (1), 99–119.
- Phillipson, P. B., & Allorge, L. (2016). A remarkable new species of *Clerodendrum* L. (Lamiaceae) from Madagascar. *Candollea*, 71 (1), 117–126.
- Pickering, H. & Patzelt, A. (2008). *Field guide to the wild plants of Oman*. Richmond: Royal Botanic Gardens, Kew.
- Pipoly III, J. J. (1992). Notes on the genus *Myrsine* (Myrsinaceae) in Peru. *Novon*, 2 (4), 392–407.
- Pipoly III, J. J. (1995). *Clusia fabiolae*: A new species, with a synopsis of *Clusia* section *Anandrogryne* (Clusiaceae) in Guayana. *SIDA, Contributions to Botany*, 16 (4), 737–755.
- Pipoly III, J.J. & Takeuchi, W. (2004). New species of *Tapeinosperma* and *Discocalyx* (Myrsinaceae) from Morobe Province, Papua New Guinea. *Harvard Papers in Botany*, 8 (2), 153–159.
- Pipoly III, J.J., Kearns, D. M. & Berry, P. E. (2000). *Key to the species and sections of Clusia in the Venezuelan Guayana*. Retrieved 1 February 2017, from <http://www.mobot.org/MOBOT/research/ven-guayana/clusiaceae/clusia.html>.
- Plantes et botanique. (2017). *Plantes & botanique*. Retrieved 1 February 2017, from <http://www.plantes-botanique.org>.

- PlantNET. (2017). *New South Wales Flora Online*. Retrieved 1 February 2017, from <http://plantnet.rbgsyd.nsw.gov.au/>.
- PNGplants. (2017). *PNGplants database: Plant collections from Papua New Guinea*. Retrieved 1 February 2017, from <http://www.pngplants.org/PNGdatabase>.
- Poncy, O. (2007). A new species of *Inga* Mill. (Fabaceae, Mimosoideae) from the Guianas. *Adansonia*. 3 (29), 249–254.
- Pool, A. (2013). New species, combinations, and lectotypifications in neotropical and Northern Mexican *Frangula* (Rhamnaceae). *Novon: A Journal for Botanical Nomenclature*. 22 (4), 447–467.
- Poorum, K. & Ranghoo-Sanmukhiya, V. M. (2009). Phylogeny of *Trochetia* species based on morphological and molecular markers. *University of Mauritius Research Journal*. 15(1), 560–576.
- Prance, G. T. (1984). Chrysobalanaceae. *Flora Malesiana-Series 1, Spermatophyta*. 10 (1), 635–678.
- Prance, G. T., Plana, V., Edwards, K. S., & Pennington, R. T. (2007). *Flora neotropica. Monograph 100. Proteaceae*. New York: New York Botanical Garden, 8–9.
- Prance, G. T. (2012). A revision of *Barringtonia* (Lecythidaceae). *Allertonia*, 12, 1–164.
- Prance, G. T. & Jongkind, C. C. (2015). A revision of African Lecythidaceae. *Kew Bulletin*. 70 (1), 1–68.
- Prance, G. T., Plana, V., Edwards, K. S. & Pennington, R. T. (2007). *Proteaceae (Flora Neotropica Monograph 100)*. New York: The New York Botanical Garden Press. New York, United States.
- Prata, E. M. B., De Carvalho, R. B., & Vicentini, A. (2016). *Pagamea spruceana* (Rubiaceae, Gaertnereae): A new species from flooded white-sand forests in the Upper Rio Negro region, Brazil. *Phytotaxa*, 269 (3), 186–192.
- Press, J. R., Shrestha, K. K. & Sutton, D.A. (2000). Annotated checklist of the flowering plants of Nepal. London: Natural History Museum Publications.
- Priyanti, P., Chikmawati, T., Sobir, S., & Hartana, A. (2016). *Durio connatus* (Malvaceae): A new species from Kalimantan, Indonesia. *Phytotaxa*. 272 (3), 215–219.
- PROSEA. (2017). *PROSEA: Plant resources of South-East Asia*. Retrieved 1 February 2017, from <http://proseanet.org/prosea/>.
- PROTA. (2017). *Plant resources of Tropical Africa*. Retrieved 1 February 2017, from <http://www.prota.org/>.
- PVNH. (2017). *The herbarium of Vanuatu (PVNH)*. Retrieved 1 February 2017, from http://publish.plantnet-project.org/project/vanuaflora_en
- Raimondo, D., Staden, L. V., Foden, W., Victor, J. E., Helme, N. A., Turner, R. C. et al. (2009). *Red List of South African plants 2009*. Pretoria, South African National Biodiversity Institute.
- Rainer, H. (2001). Nomenclatural and taxonomic notes on *Annona* (Annonaceae). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 103, 513–524.
- Rainer, H. (2006). Monographic studies in the genus *Annona* L.(Annonaceae): Inclusion of the genus *Rollinia* A. St.-Hil. *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 108, 191–205.
- Ramesh, B.R., Pascal, J.P. & Nouguier, C. (1997). *Atlas of endemics of the Western Ghats (India): Distribution of tree species in the evergreen and semi-evergreen forests*. India: Institut français de Pondichéry.
- Real Jardín Botánico de Madrid. (2017). *Flora de Guinea: Base de datos de Flora de Guinea. Consejo Superior de Investigaciones Científicas*. Retrieved 1 February 2017, from <http://www.floradeguinea.com/herbario/>.
- Rejmánek, M. & Richardson, D. M. (2013). Trees and shrubs as invasive alien species—2013 update of the global database. *Diversity and Distributions*, 19 (8), 1093–1094.
- Reksodiharjo, W. S. (1960). The genus *Coelostegia* Benth. (Bombac.). *Reinwardtia*, 5 (3), 269–91.
- Retief, E. & Van Wyk, A. E. (2001). The genus *Ehretia* (Boraginaceae: Ehretioideae) in southern Africa. *Bothalia*. 31 (1), 9–23.
- Rich, T. C. G., Houston, L., Robertson, A. & Proctor, M. C. F. (2010). *Whitebeams, Rowans and Service trees of Britain and Ireland: A monograph of British and Irish Sorbus, L.* London: Botanical Society of the British Isles.

- Richard, P. S. S. & Muthukumar, S. A. (2012). Arborescent angiosperms of Mundanthurai range in the Kalakad-Mundanthurai Tiger Reserve (KMTR) of the southern Western Ghats, India. *Check List*. 8 (5), 951–962.
- Ricker, M., Hernández, H. M., Sousa, M. & Ochoterena, H. (2013). Especies arbóreas y arborescentes de México: Asteraceae, Leguminosae y Rubiaceae. *Revista Mexicana de Biodiversidad*. 84 (2), 439–470.
- Ricketson, J. M. and Pipoly III, J. J. (2013). Nomenclatural transfers in the pantropical genus *Myrsine* (Myrsinaceae). *Novon: A Journal for Botanical Nomenclature*. 22 (4), 468–472.
- Rico Arce, M. D. L. & Griffiths, P. (2002). Two little known *Acacia* species from Mexico. *Curtis's Botanical Magazine*. 19 (3), 152–159.
- Rio de Janeiro Botanical Garden (2016). *Brazilian Flora 2020 in construction*. Retrieved 1 February 2017, from <http://floradobrasil.jbrj.gov.br>.
- Risna, R. A. (2009). *Autekologi dan studi populasi Myristica teijsmannii* Miq. (Myristicaceae) di Cagar Alam Pulau Sempu, Jawa Timur. Masters dissertation, Bogor Agricultural University, West Java, Indonesia.
- Ritter, M. W. & Naugle, C. M. (1999). Population characteristics, germination and proposed management of *Elaeocarpus joga* Merr. on Guam: A regionally endemic tree. *Micronesica*. 31 (2), 275–281.
- Robi, A. J. (2015). *A taxonomic revision of the family Lauraceae from South India*. Doctoral dissertation, Kannur University, Kerala, India.
- Romkham, N. (2013). *Wrightia calcicola* (Apocynaceae: Apocynoideae): A new species from Thailand. *Thai Forest Bulletin (Botany)*, 41, 81–84.
- Royal Botanic Gardens and Domain Trust. (2017.) *PlantNET (The NSW Plant Information Network System)*. Retrieved 1 February 2017, from <http://plantnet.rbgsyd.nsw.gov.au>.
- Royal Botanic Gardens Edinburgh. (2017). *Edinburgh Rhododendron Monographs*. Retrieved 1 February 2017, from http://data.rbge.org.uk/service/factsheets/Edinburgh_Rhododendron_Monographs.xhtml.
- Royal Botanic Gardens, Kew. (1999). *Survey of Economic Plants for Arid and Semi-Arid Lands (SEPASAL) database*. Retrieved 1 February 2017, from <http://apps.kew.org/sepasalweb/sepaweb>.
- Royal Botanic Gardens, Kew. (2014). *Kew Gardens—Conservation Assessment Tracker*. Retrieved 1 August 2015, from www.google.com/fusiontables/DataSource?docid=14VS4gADo50vHpLa4loT4B6K-ZX-6G20YP19il7A#rows:id=1.
- Royal Botanic Gardens, Kew. (2017). *Discover plants and fungi*. Retrieved 1 February 2017, from <http://www.kew.org/science-conservation/plants-fungi>.
- Royal Botanic Gardens, Kew. (2017). *Flora Zambesiaca*. Retrieved 1 February 2017, from <http://apps.kew.org/efloras/search.do>.
- Royal Botanic Gardens, Kew. (2017). *The Herbarium Catalogue*. Retrieved 1 February 2017, from <http://apps.kew.org/herbcat/navigator.do>.
- Rueangruea, S., Tagane, S., Suddee, S. & Tetsana, N. (2015). *Toona calcicola*: A new species and *Reinwardtiadendron humile*, a new record to Thailand. *Thai Forest Bulletin (Botany)*, 43, 79–86.
- Rzedowski, J., Lemos, R. M. & de Rzedowski, G. C. (2005). Inventario del conocimiento taxonómico, así como de la diversidad y del endemismo regionales de las especies mexicanas de *Bursera* (Burseraceae). *Acta Botanica Mexicana*, 70, 85–111.
- SANBI (2014). *Plants of Southern Africa—an online checklist*. Retrieved 1 September 2016, from <http://posa.sanbi.org/searchspp.php>.
- SANBI (2014). *Red List of South African Plants version 2014.1*. Retrieved 1 September 2016, from <http://redlist.sanbi.org/>.
- Sánchez-Chávez, E., & Zamudio, S. (2016). Tres especies nuevas de Myrtaceae de la Sierra Madre Oriental, México. *Acta botánica mexicana*, 115, 51–64.
- Sanoja, E. (2004). Diagnósis y observaciones sobre la biología de *Catostemma lemense*, nueva Bombacaceae de Venezuela. *Acta Botánica Venezolánica*. 27 (2), 83–94.
- Santamaria-Aguilar, D., & Ortiz, R. D. C. (2016). *Freziera dasycarpa* (Pentaphylacaceae): A new species from the montane forest of Antioquia, Colombia. *Phytotaxa*. 263 (3), 279–285.
- Santos, L. L., Sales, M. F., & Sobral, M. (2016). A new Brazilian species of *Myrcia* (Myrtaceae) and a new synonym for *Psidium amplexicaule*. *Phytotaxa*. 257 (2), 122–132.

- Sarma, J., Shameer, P. S., & Mohanan, N. N. (2016). A new species of *Garcinia* (Clusiaceae) from Assam, North East India. *Phytotaxa*. 252 (1), 73–76.
- Sasidharan, N. & Swarupanandan, K. (1992). A new species of *Cassine* (Celastraceae) from India. *Reinwardtia*. 11 (1), 29–32.
- Saunders, R. M. & Chalermglin, P. (2008). A synopsis of *Goniothalamus* species (Annonaceae) in Thailand, with descriptions of three new species. *Botanical Journal of the Linnean Society*. 156 (3), 355–384.
- Schatz, G. E. & Maas, P. J. M. (2010). Synoptic revision of *Stenanona* (Annonaceae). *Blumea*. 55 (3), 205–223.
- Schatz, G. E., Ramos A, C.A., Ortiz, O. O. & McPherson, G. (2015). A new, restricted range species of *Annona* (Annonaceae) endemic to the Caribbean Slope of Panama. *Novon*. 24 (2), 203–208.
- Schatz, G. E., Shulkina, T. & Solomon, J. C. (2014). *Red List of the endemic plants of the Caucasus: Armenia, Azerbaijan, Georgia, Iran, Russia, and Turkey*. St. Louis: Missouri Botanical Garden Press.
- Schmidt, L. H. & Mbor, A. (2008). *Commiphora africana* (A. Rich.) Eng. *Seed Leaflet*, 138.
- Schot, A. M. (2004). Systematics of *Aporosa* (Euphorbiaceae). *Blumea Supplement No. 17*. The Netherlands: Nationaal Herbarium Nederland, Universiteit Leiden.
- Sebola, R. J., & Balkwill, K. (2009). A monograph of the Oliniaceae. *South African Journal of Botany*.
- Secco, R. D. S. (2008). *Alchornea lojaensis*: A new species of Euphorbiaceae for the Flora of Ecuador. *Kew Bulletin*. 63 (3), 511–513.
- SEMARNAT (2010). NOM-059-SEMARNAT-2010.
- Sevilla, S. & Welzen, P. C. (2001). Revision and phylogeny of *Agrostistachys* and *Chondrostylis* (Euphorbiaceae). *Blumea*. 46 (1), 71–97.
- Shaw, K., Stritch, L., Rivers, M., Roy, S., Wilson, B. & Govaerts, R. (2014). The Red List of Betulaceae. Richmond: BGCI.
- Shimizu, G. H., Goncalves, D. J. P., Franca, F., Simoes, A. O., & Yamamoto, K. (2016). A remarkable new species of *Qualea* (Vochysiaceae) from Piauí state, Brazil. *Phytotaxa*. 273 (4), 262–268.
- Sidyasa, K., Uhaedi, S., Gusti M. T. I., Whitmore, T.C. & Sidiyaka, K. (1997). Tree flora of Indonesia: Check list for Irian Jaya. Bogor: Ministry of Forestry, Forestry Research and Development Agency, Forest & Nature Conservation Research & Development Centre.
- Sierra, S. E. C. & Van Welzen, P. C. (2005). A taxonomic revision of *Mallotus* section *Mallotus* (Euphorbiaceae) in Malesia. *Blumea*. 50 (2), 249–274.
- Sierra, S. E. C., Aparicio, M., Gebraad, M. J. H., Kulju, K. K. M. & Van Welzen, P. C. (2007). The morphological range in *Mallotus* (Euphorbiaceae) and a taxonomic revision of its section *Rottleropsis* (including *Axenfeldia*) in Malesia, Thailand and Africa. *Blumea*. 52 (1), 21–113.
- Sierra, S. E. C., Van Welzen, P. C. & Slik, J. W. F. (2005). A taxonomic revision of *Mallotus* section *Philippinenses* (former section *Rottlera*–Euphorbiaceae) in Malesia and Thailand. *Blumea*. 50 (2), 221–248.
- Siniscalchi, C. M., Loeuille, B., Semir, J., & Pirani, J. R. (2016). *Lychnophora spiciformis* (Asteraceae: Vernoniaeae): A new species from Bahia, Brazil. *Phytotaxa*. 253 (1), 48–56.
- Sistema Nacional de Áreas de Conservación (Sinac) y Programa REDD-CCAD-GIZ. (2014). *Protocolo de campo para la identificación de especies arbóreas: Información taxonómica y dendrológica de las especies arbóreas de Costa Rica*. 2014. San José, Costa Rica.
- Sleumer, H. (1969). Materials towards the knowledge of the Icacinaceae of Asia, Malesia, and adjacent areas. *Blumea*. 17 (1), 181–264.
- Sleumer, H. (1988). The genera *Discocalyx* Mez, *Fittingia* Mez, *Loheria* Merr. and *Tapeinosperma* Hook.f. (Myrsinaceae) in New Guinea. *Blumea*. 33 (1), 81–107.
- Sleumer, H. O. (1980). Flacourtiaceae. *Flora Neotropica*, 22, 1–499.
- Slik, J. W. F. (2009). *Plants of Southeast Asia*. Retrieved 1 September 2016, from <https://www.asianplant.net>.
- Slik, J. W. F. & Van Welzen, P. C. (2001). A taxonomic revision of *Mallotus* sections *Hancea* and *Stylanthus* (Euphorbiaceae). *Blumea*. 46 (1), 3–66.
- Smith, A. C. (1971). Studies of pacific island plants, XXIV. The genus *Terminalia* (Combretaceae) in Fiji, Samoa, and Tonga. *Brittonia*. 23 (4), 394–412.

- Smithsonian Institution. (2017). *Floras of the West Indies: Catalogue of Seed Plants of the West Indies*. Retrieved 1 September 2016, from <http://botany.si.edu/Antilles/WestIndies/catalog.htm>.
- Smithsonian Tropical Research Institute. (2017). *Panama*. Retrieved 1 September 2016, from <http://biogeodb.stri.si.edu/biodiversity/>.
- Snow, N. (2004). Systematics of *Pilidiostigma* (Myrtaceae). *Systematic Botany*, 29(2), 393–406.
- Sobral, M. A. R. C. O. S., Mazine, F. F., Leoni, L., Souza, M. C., & A. D. E. (2016). Five new southeastern Brazilian Myrtaceae. *Phytotaxa*, 253 (1), 57–70.
- Sobral, M., Caliari, C. P., Gressler, E., Mazine, F. F., Magenta, M., & Viana, P. L. (2016). Seven new southeastern Brazilian species of *Myrcia* (Myrtaceae). *Phytotaxa*, 247 (1), 27–44.
- Soejarto, D. D. (1980). Revision of South American *Saurauia* (Actinidiaceae). *Fieldiana: Bot. N.S.*, 2, 1–141.
- Soejima, A., Tagane, S., Van, N. N., Duy, C. N., Huong, N. T. T., & Yahara, T. (2016). *Callicarpa bachmaensis* Soejima & Tagane (Lamiaceae): A new species from Bach Ma National Park in Thua Thien Hue Province, Central Vietnam. *PhytoKeys*, 62, 33–39.
- Soepadmo, E. & Wong, K. M. (1995). *Tree flora of Sabah and Sarawak: Volume One*. Sandakan: Sabah Forestry Department.
- Soepadmo, E., Wong, K. M. & Saw, L. G. (1996). *Tree flora of Sabah and Sarawak: Volume Two*. Sandakan: Sabah Forestry Department.
- Soepadmo, E. & Saw, L. G. (2000). *Tree flora of Sabah and Sarawak: Volume Three*. Sandakan: Sabah Forestry Department.
- Soepadmo, E., Saw, L. G. & Chung, R. C. K. (2002). *Tree flora of Sabah and Sarawak: Volume Four*. Sandakan: Sabah Forestry Department.
- Soepadmo, E., Saw, L. G. & Chung, R. C. K. (2004). *Tree flora of Sabah and Sarawak: Volume Five*. Sandakan: Sabah Forestry Department.
- Soepadmo, E., Saw, L. G., Chung, R. C. K. & Kiew, R. (2007). *Tree flora of Sabah and Sarawak: Volume Six*. Sandakan: Sabah Forestry Department.
- Soepadmo, E., Saw, L. G., Chung, R. C. K. & Kiew, R. (2011). *Tree flora of Sabah and Sarawak: Volume Seven*. Sandakan: Sabah Forestry Department.
- Soepadmo, E., Saw, L. G., Chung, R. C. K. & Kiew, R. (2014). *Tree flora of Sabah and Sarawak: Volume Eight*. Sandakan: Sabah Forestry Department.
- Solano, J. (2016). *Passiflora chimuensis*, una nueva especie y primer registro de la sección *Pseudostrophea* (Subgénero *Astrophea*, Passifloraceae) en Mesoamérica. *Journal of the Botanical Research Institute of Texas*, 10 (1).
- Sosef, M. S. (2013). The genus *Idertia* (Ochnaceae). *Plant Ecology and Evolution*, 146 (3), 351–359.
- Sosef, M. S. & Dauby, G. (2012). Contribution to the taxonomy of *Garcinia* (Clusiaceae) in Africa, including two new species from Gabon and a key to the Lower Guinean species. *PhytoKeys*, 17, 41–62.
- Sosnovsky, D. I. (Ed.) (1952a, 1952b). *Flora Azerbaijana. Vol. II, III*. Baku, Azerbaijan: Izdatel'stvo Akademii Nauk Azerbajanskoy SSR.
- Sothers, C., Prance, G. T., Buerki, S., De Kok, R. & Chase, M. W. (2014). Taxonomic novelties in Neotropical Chrysobalanaceae: Towards a monophyletic *Couepia*. *Phytotaxa*, 172 (2), 176–200.
- Sotuyo, S. & Lewis, G. P. (2007). A new species of *Caesalpinia* from the Río Balsas Depression, Mexico, and an updated taxonomic circumscription of the *Caesalpinia hintonii* complex (Leguminosae: Caesalpinioideae: Caesalpinieae: Poincianella group). *Brittonia*, 59 (1), 33–36.
- Sousa, M. & Sotuyo, S. (2012). El género *Muellera* L. f. en Mesoamérica y norte de Sudamérica. *Acta Botánica Mexicana*, 100, 15–40.
- Sousa, S. (2009). *Standleyi* una nueva sección del género *Lonchocarpus* (Leguminosae, Papilionoideae, Millettieae), nuevas especies y subespecies para Mesoamérica y Sudamérica. *Acta Botánica Mexicana*, 86, 39–69.
- Sousa, S. (2011). Especies nuevas y una lectotipificación en *Lonchocarpus* (Leguminosae) para Mesoamérica. *Revista Mexicana de Biodiversidad*, 82 (4), 1108–1122.
- South African National Biodiversity Institute. (2016). *Botanical database of Southern Africa (BODATSA)*. Retrieved 1 September 2016, from <http://newposa.sanbi.org/>.

- South Australian Seed Conservation Centre. (2017). *Seeds of South Australia*. Retrieved 1 February 2016, from <http://www.saseedbank.com.au/>.
- Stergios, B. D. (2001). Contributions to South American Caesalpiniaceae. Vi. *Campsiandra felipeana* (Caesalpinieae): A dwarf swamp-forest endemic of the lower Orinoco watershed. *Acta Botánica Venezuelica*. 24 (2), 85–91.
- Stergios, B.D. & Aymard, G. A. (2008). A striking new species of *Aldina* (Fabaceae-Swartzieae-Aldininae) from the Venezuelan Guayana highlands. *Harvard Papers in Botany*. 13 (1), 29–33.
- Stirton, C. H. & Muasya, A. M. (2016). Seven new species and notes on the genus *Aspalathus* (Crotalariaeae, Fabaceae). *South African Journal of Botany*, 104, 35–46
- Stone, R.D. & Luke, Q. (2015). *Lijndenia udzungwarum* (Melastomataceae–Olisbeoideae): A new, endemic species from the Udzungwa Mountains of southern Tanzania. *Phytotaxa*. 226 (2), 169–176.
- Stone, R. D. & Ntetha, N. A. (2013). *Warneckea parvifolia* (Melastomataceae–Olisbeoideae): A new “sand-forest” endemic from northeastern KwaZulu-Natal (South Africa) and southernmost Mozambique, and a phylogenetic analysis of eastern and southern African representatives of *W.* section *Warneckea*. *South African Journal of Botany*, 88, 317–325.
- Stoops E. & Welzen P.C. van. (2013). A revision of *Ptychopyxis* (Euphorbiaceae) in southeast Asia. *Nordic Journal of Botany*, 31, 94–112.
- Strijk, J.S., Sirimongkol, S., Rueangruea, S., Ritphet, N. & Chamchumroon, V. (2014). *Lithocarpus orbicarpus* (Fagaceae): A new species of Stone Oak from Phang Nga province, Thailand. *PhytoKeys*, 34, 33–45.
- Swanepoel, W. (2005). *Commiphora kaokoensis* (Burseraceae): A new species from Namibia, with notes on *C. dinteri* and *C. namaensis*. *Bothalia*. 35 (1), 47–53.
- Swanepoel, W. (2006). Two new species of *Commiphora* (Burseraceae) from southern Africa. *Bothalia*. 36 (1), 45–56.
- Swanepoel, W. (2011). *Commiphora buruxa* (Burseraceae): A new species from southern Namibia. *South African Journal of Botany*. 77 (3), 608–612.
- Swanepoel, W. (2014). *Commiphora namibensis* (Burseraceae): a new species from Angola. *Phytotaxa*. 178 (3), 211–216.
- Swenson, U., & Munzinger, J. (2016). Five new species and a systematic synopsis of *Pycnandra* (Sapotaceae), the largest endemic genus in New Caledonia. *Australian Systematic Botany*. 29 (1), 1–40.
- Swenson, U., Nylinder, S. & Munzinger, J. (2013). Towards a natural classification of Sapotaceae subfamily Chrysophylloideae in Oceania and Southeast Asia based on nuclear sequence data. *Taxon*. 62 (4), 746–770.
- Tagane, S., Nguyen, V. H., Van Ngoc, N., Son, H. T., Toyama, H., Yang, C. J., et al. (2016). *Homalium glandulosum* (Salicaceae): A new species from Vu Quang National Park, North Central Vietnam. *PhytoKeys*, 58, 97.
- Takeuchi, W. (2007). Notes on *Acronychia* (Rutaceae) from the Kaijende Highlands of Papua New Guinea. *Harvard Papers in Botany*. 11 (2), 203–206.
- Takeuchi, W. (2014). A distinctive addition to the tree flora of Papua New Guinea: *Helicia woxvoldiana* sp. nov. (Proteaceae), a large-flowered myrmecophyte from the upper Sepik. *Phytotaxa*. 172 (2), 94–100.
- Takhtajan, A. L. (Ed.) (1954–2001). *Flora Armenii Vol 1–9*. Yerevan, Armenia: Izdatel'stvo Akademii Nauk Armianskoy SSR, Vol 10. Liechtenstein: A.R.G. Gantner Verlag KG
- Takhtajan, A. L. (Ed.) (2003, 2006, 2008). *Caucasian flora conspectus, Vol. I-II*. Saint-Petersburg: Saint-Petersburg University Press.
- Takhtajan, A. L. (Ed.) (2003, 2006, 2008). *Caucasian flora conspectus, Vol. III*. Saint-Petersburg—Moscow: KMK Scientific Press.
- Talybov, T. G. & Ibrahimov A. M. (2009). Wild species of Pears (*Pyrus* L.) in The Flora of Nakhichevan Autonomous Republic of Azerbaijan. *Turczaninowia*. 12 (3–4), 82–87.
- Tan, H. S., Chung, R. C. K. & Soepadmo, E. (2011). A synopsis of *Jarandersonia* (Malvaceae—Brownlowioideae). *Garden's Bulletin Singapore*. 63 (1–2), 137–144.

- Tange, C. (2013). A revision of the genus *Greenea* (Rubiaceae). *Thai Forest Bulletin (Botany)*, 41, 64–80.
- Taylor, C. M. (2016). Rubiacearum Americanarum Magna Hama Pars XXXI: More new neotropical species and morphological notes for Psychotria (Psychotrieae). *Novon: A Journal for Botanical Nomenclature*. 24 (4), 413–434.
- Ter Steege, H., Pitman, N. C., Killeen, T. J., Laurance, W. F., Peres, C. A., Guevara, J. E., et al. (2015). Estimating the global conservation status of more than 15,000 Amazonian tree species. *Science Advances*. 1(10), 1500936.
- Ter Steege, H., Vaessen, R. W., Cárdenas-López, D., Sabatier, D., Antonelli, A., De Oliveira, S. M., Pitman, et al. (2016). The discovery of the Amazonian tree flora with an updated checklist of all known tree taxa. *Scientific Reports*, 6, 29549.
- The Plant List. (2013). Version 1.1. Retrieved 1 October 2015, from <http://www.theplantlist.org>.
- Thomas, W. W. (2011). Nothotalisia: A new genus of Picramniaceae from tropical America. *Brittonia*. 63(1), 51–61.
- Timyan, J. (1996). *Bwa yo: Important trees of Haiti*. Washington D.C.: South-East Consortium for International Development.
- Todzia, C. A. (1992). A reevaluation of the genus *Phyllostylon* (Ulmaceae). *SIDA, Contributions to Botany*, 15, 263–270.
- Tomas, W. M., Ishii, I. H., Urbanetz, C., Tomas, M. A., Camilo, A. R. & Cano, Á. (2015). The Carandilla Palm (*Trithrinax schizophylla* Drude, Arecaceae) is not extinct in Brazil: First primary records from the Chaco region of Mato Grosso do Sul. *Check List*. 11(4), 1669.
- Tong, Y., Pang, K. S. & Xia N. (2014). *Carpinus insulares* (Betulaceae): A new species from Hong Kong, China. *Journal of Tropical and Subtropical Botany*. 22(2), 121–124.
- Torres-Leite, F., Bruniera, C. P., Zappi, D. C., & Carrijo, T. T. (2016). True axillary inflorescences in *Rudgea* (Palicoureeae, Rubiaceae): A newly reported characteristic of two new Brazilian species, *R. quisquiliae* and *R. axilliflora*. *Phytotaxa*. 272(3), 191–200.
- Toyama, H., Tagane, S., Chhang, P., Nagamasu, H., & Yahara, T. (2016). Flora of Bokor National Park, Cambodia III: A new species, *Garcinia bokorensis* (Clusiaceae). *Acta Phytotaxonomica et Geobotanica*. 67(1), 47–53.
- Toyama, H., Tagane, S., Van Son Dang, H. T., Nagamasu, H., Naiki, A., & Yahara, T. (2016). A new species of *Eustigma* (Hamamelidaceae) from Hon Ba Nature Reserve, Vietnam. *PhytoKeys*, 65, 47–55.
- Trainor, C. R. (2011). New island records of ‘*Eucalyptus alba sensu lato*’ for Damar and Romang, Lesser Sundas, Indonesia. *Northern Territory Naturalist*. 23(1), 45–53.
- Trinidad and Tobago Biodiversity (2012). Retrieved 1 October 2015, from <http://www.biodiversity.gov.tt>.
- Trudgen, M. S. & Baker, W. J. (2008). A revision of the *Heterospathe elegans* (Arecaceae) complex in New Guinea. *Kew Bulletin*. 63(4), 639–647.
- Turner, I. M. (2012). A new combination in *Monocarpia* (Annonaceae). *Edinburgh Journal of Botany*. 69(1), 95–98.
- Ulloa, C. U., Penneys, D. S., Neill, D. A. & Fernandez, D. M. (2015). A new species of *Chalybea* (Blakeeae, Melastomataceae) from the Ecuador-Peru border. *Phytotaxa*. 212(4), 264–270.
- Ulloa, C.U., Ståhl, B., Minga, D. & Ansaloni, R. (2015). A new species of *Symplocos* (Symplocaceae) from southern Ecuador. *PhytoKey*, 56, 41–46.
- UNEP-WCMC. (2015). *Overview of Dalbergia spp. from South and Central America: A basic review*. Cambridge: UNEP-World Conservation Monitoring Centre.
- University of Florida Herbarium. (2017). University of Florida Herbarium (FLAS). Retrieved 1 February 2017, from <https://www.flmnh.ufl.edu/herbarium/>
- USDA, NRCS. (2017). The PLANTS Database. National Plant Data Team. Retrieved 1 February 2017, from <https://plants.usda.gov/java/citePlants>
- Useful Tropical Plants Database. (2017). Useful Tropical Plants Database. Retrieved 1 February 2017, from <http://tropical.theferns.info/>
- Utteridge, T. M. & Davis, A. P. (2009). Two new combinations in *Pyrostria* (Rubiaceae, Vanguerieae) from Thailand. *Kew Bulletin*. 64(4), 751–752.

- Valdés, C. P. & Rodríguez, I. V. (2000). *Wallenia maestrensis* (Myrsinaceae): A new species from eastern Cuba. *Willdenowia*, 30(1), 141–145.
- Valencia, S., Rosales, S. J. L. S., & Arellano, O. J. S. (2016). A new species of *Quercus*, section *Lobatae* (Fagaceae) from the Sierra Madre Oriental, Mexico. *Phytotax*, 269(2), 120–126.
- Van den Eynden, V., Oatham, M. P. & Johnson, W. (2008). How free access internet resources benefit biodiversity and conservation research: Trinidad and Tobago's endemic plants and their conservation status. *Oryx*, 42(3), 400–407.
- van der Burgt, X. M. (2016). *Didelotia korupensis* and *Tessmannia korupensis* (Leguminosae, Caesalpinioideae): Two new tree species from Korup National Park in Cameroon. *Blumea-Biodiversity, Evolution and Biogeography of Plants*, 61(1), 51–58.
- van der Burgt, X. M., Mackinder, B. A., Wieringa, J. J. & de la Estrella, M. (2015). The *Gilbertiodendron ogoouense* species complex (Leguminosae: Caesalpinioideae), Central Africa. *Kew Bulletin*, 70(2), 1–42.
- Van der Maesen, L. J. G. (2013). Novitates Gabonenses 85: Leguminosae-Papilionoideae in the flora of Gabon: New species and nomenclatural notes. *Webbia*, 68(2), 97–101.
- Van der Meijden, R. (1982). *Systematics and evolution of xanthophyllum: (Polygalaceae)*. *Leiden Botanical Series, Volume 7*. Leiden: Leiden University Press.
- Van Dung, L., Son, H. T., Tran, N. I. N. H., & Nhan, P. H. (2016). *Camellia quangcuongii* (Theaceae): A new species from Vietnam. *The Journal of Japanese Botany*, 91, 226–230.
- Van Ngoc, N., Van Dung, L., Tagane, S., Binh, H. T., Son, H. T., Trung, V. Q., et al. (2016). *Lithocarpus dahuoaiensis* (Fagaceae): A new species from Lam Dong Province, Vietnam. *PhytoKeys*, 69, 23.
- Van Welzen, P. C. & Sierra, S. E. C. (2006). The *Mallotus wrayi* complex (Euphorbiaceae). *Blumea*, 51(2), 373–388.
- Vargas, O. M. (2011). Anomenclator of *Diplostephium* (Asteraceae: Astereae): A list of species with their synonyms and distribution by country. *Lundellia*, 14, 32–51.
- Velayos, M., Cabezas, F., Barberá, P., de la Estrella, M., Aedo, C., Morales, R., et al. (2013). Preliminary checklist of vascular plants of Bioko Island (Equatorial Guinea). *Botanica Complutensis*, 37, 109–133.
- Villaseñor, J. L. (2016). Checklist of the native vascular plants of Mexico. *Revista Mexicana de Biodiversidad*, 87(3), 559–902.
- Vink, W. (1985). The Winteraceae of the old world. V. *Exospermum* links *Bubbia* to *Zygogynum*. *Blumea*, 31(1), 39–55.
- Vink, W. (2016). The Winteraceae of the Old World. VIII. Some *Zygogynum* species from New Guinea. *Blumea-Biodiversity, Evolution and Biogeography of Plants*, 61(1), 41–50.
- Vivero, J. L., Kelbessa, E. & Demissew, S. (2005). *The Red List of trees and shrubs of Ethiopia and Eritrea*. Cambridge: Fauna & Flora International.
- Vivero, J. L., Szejner, M., Gordon, J. and Magin, G. (2006). *The Red List of trees of Guatemala*. Cambridge: Fauna & Flora International
- von Adolf Engler, B. (1994). *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*. Stuttgart: Schweizerbart Science Publishers.
- Wagner, M. R., Cobbinah, J. R. & Bosu, P. P. (2008). Forest entomology in West Tropical Africa: forest insects of Ghana. Dordrecht: Springer Science & Business Media.
- Wagner, W. L. & Lorence, D. H. (2002–2017). *Flora of the Marquesas Islands*. Retrieved 1 February 2017, from: <http://botany.si.edu/pacificislandbiodiversity/marquesasflora/index.htm>.
- Wagner, W. L., Herbst, D. R. & Lorence, D. H. (2005–2017). *Flora of the Hawaiian Islands*. Retrieved 1 February 2017, from: <http://botany.si.edu/pacificislandbiodiversity/hawaiianflora/index.htm>.
- Wagner, W. L. & Lorence, D. H. (2011). A nomenclator of Pacific oceanic island *Phyllanthus* (Phyllanthaceae), including *Glochidion*. *PhytoKeys*, 4, 67–94.
- Wahlert, G. A., Phillipson, P. B., Mabberley, D. J., & Lowry, P. P. (2016). *Grewia hispidissima* Wahlert, Phillipson & Mabb., sp. nov. (Malvaceae, Grewioideae): A new species of restricted range from northwestern Madagascar. *Adansonia*, 38 (1), 117–121.
- WCSP (2016). *World checklist of selected plant families. Facilitated by the Royal Botanic Gardens, Kew*. Retrieved 1 February 2017, from <http://apps.kew.org/wcsp/>.

- Wearn, J. A. & Mabberley, D. J. (2011). *Clerodendrum* (Lamiaceae) in Borneo. *Systematic Botany*. 36(4), 1050–1061.
- Weerasinghe, A., Puvendran, S., Wickramasinghe, A., Karunaratne, D.N., Wijesundara, S. & Karunaratne, V. (2013). Potent bioactivities of the endemic Annonaceae heightens its dire conservation status. *Journal of the National Science Foundation of Sri Lanka*. 41 (4), 345–350.
- Weerasooriya, A. D. & Saunders, R. M. (2010). Monograph of *Mitrephora* (Annonaceae). *Systematic Botany Monographs*, 90, 1–167.
- Welzen, P. C. van (Ed.). (2017). *Flora Malesiana Euphorbiaceae*. Retrieved 1 February 2017, from <http://www.nationaalherbarium.nl/euphorbs>.
- Welzen, P. C. van & Chayamarit, K. (2017). *Flora of Thailand Euphorbiaceae*. Retrieved 1 February 2017, <http://www.nationaalherbarium.nl/thaieuph>.
- Werff, H. V. D. (2013). A revision of the genus *Ocotea* Aubl.(Lauraceae) in Madagascar and the Comoro Islands. *Adansonia*. 35 (2), 235–279.
- West, J. (2010). *Australian tropical rainforest plants*. Retrieved 1 February 2017, <http://keys.trin.org.au/key-server/data/0e0f0504-0103-430d-8004-060d07080d04/media/Html/index.html>.
- Western Australian Herbarium. (1998–). *FloraBase: The Western Australian Flora*. Department of Parks and Wildlife. Retrieved 1 September 2016, from <https://florabase.dpaw.wa.gov.au/>
- Whistler, W. A. (1986). A revision of *Psychotria* (Rubiaceae) in Samoa. *Journal of the Arnold Arboretum*. 67 (3), 341–370.
- Whistler, W. A. (2004). Rainforest trees of Samoa: A guide to the common native and naturalized lowland and foothill forest trees of the Samoan Archipelago. Honolulu: Isle Botanica.
- Whitford, H. N. (1911). The Forest of the Philippines: Part II. The Principal Forest Trees. Manila: Department of the Interior, Bureau of Forestry.
- Wieringa, J. J. & Mackinder, B. A. (2012). Novitates Gabonensis 79: *Hymenostegia elegans* and *H. robusta* spp. nov. (Leguminosae–Caesalpinioideae) from Gabon. *Nordic Journal of Botany*. 30 (2), 144–152.
- Wieringa, J. J., Mackinder, B. A. & Van Proosdij, A. S. (2013). *Gabonius* gen. nov. (Leguminosae, Caesalpinioideae, Detarieae): A distant cousin of *Hymenostegia* endemic to Gabon. *Phytotaxa*. 142 (1), 15–24.
- Wilkie, P. (2007). A new species of *Pterospermum* (Dombeyoideae, Malvaceae/Sterculiaceae) from Cambodia and Vietnam. *Edinburgh Journal of Botany*. 64 (02), 179–183.
- Wilmot-Dear, C. M., & Friis, I. (1998). *Cypholophus decipiens* (Urticaceae): Taxonomy and range of a species often misplaced in *Boehmeria*. *Kew Bulletin*. 53 (4), 919–927.
- Wilmot-Dear, C. M. & Friis, I. (2013). The old world species of *Boehmeria* (Urticaceae, tribus Boehmerieae). A taxonomic revision. *Blumea*. 58 (2), 85–216.
- Wilson, P. G. (2009). Conspectus of the genus *Eugenia* (Myrtaceae) in the Philippines. *Gardens' Bulletin Singapore*. 60 (2), 399–410.
- Womersley, J. S. (1978). *Handbooks of the Flora of Papua New Guinea. Volume I*. Melbourne: Melbourne University Press.
- Wong, K. M., Ariffin, A. M., & Joffre, A. A. (2016). Novitates Bruneienses, 5. *Polyalthia watui* (Annonaceae): A new tree species from Brunei, Borneo. *Gardens' Bulletin Singapore*. 68 (1), 71–76.
- Wood, K. R., Appelhans, M. S., & Wagner, W. L. (2016). *Melicope oppenheimeri*, section *Pelea* (Rutaceae): A new species from West Maui, Hawaiian Islands: With notes on its ecology, conservation, and phylogenetic placement. *PhytoKeys*. 25 (69), 51–64.
- Wood, K., Lorence, D. H. & Kiehn, M. (2016). *Coprosma kawaikiniensis* (Rubiaceae): A new species from the Dubautia-Sadleria shrubland-fernland community on Kaua'i, Hawaiian Islands. *PhytoKeys*, 60, 21–32.
- WorldWideWattle. (2017) *World Wide Wattle*. Retrieved 1 February 2017, from [s](#)
- Wu Zhengyi, Raven, P. H. & Hong Deyuan (Eds.). (1994–2013). *Flora of China*. St Louis: Science Press, Beijing & Missouri Botanical Garden Press.
- Xue, B., Su, Y. C., Thomas, D. C. & Saunders, R. M. (2012). Pruning the polyphyletic genus *Polyalthia* (Annonaceae) and resurrecting the genus *Monoon*. *Taxon*. 61 (5), 1021–1039.

- Ya Sokolov, S. (Ed.). (2013). *Derev'ya i Kustarniki SSSR*. Volumes 1–7. Russia: Book On Demand Ltd.
- Yahara, T., Tagane, S., Mase, K., Chhang, P. & Toyama, H. (2016). Flora of Bokor National Park V: Two new species of *Machilus* (Lauraceae), *M. bokorensis* and *M. brevipaniculata*. *PhytoKeys*, 65, 35–46.
- Yii, P. C. & Chai, P. P. K. (2001). New combinations, new names and new species of *Madhuca* (Sapotaceae) from Sabah and Sarawak, Borneo. *Gardens' Bulletin Singapore*, 53, 343–356.
- Zamora, N. (2014). Protocolo de campo para la identificación de especies arbóreas: Información taxonómica y dendrológica de las especies arbóreas de Costa Rica. Sistema Nacional de Áreas de Conservación (Sinac) y Programa REDD-CCAD-GIZ. Costa Rica: Ministerio de Ambiente y Energía.
- Zarones, L. (2014). Native plants of the Mariana Islands for wildlife and ornamental use. Mariana Islands: Laolao Bay Pride Campaign.
- Zhu, H., Roos, M. C. & Ridsdale, C. E. (2012). A taxonomic revision of the Malesian species of *Lasianthus* (Rubiaceae). *Blumea*, 57 (1), 1–102.