

Table of contents

Plenary

Calculus

Complex Analysis

Ordinary Differential Equations

Nonlinear Optimization, Part I: Unconstrained Optimization

Nonlinear Optimization, Part II: Constrained Optimization

Algorithm Design

Plenary

All the proofs and results I don't want to write twice.

- 1.x.x) Real analysis
- 2.x.x) Calculus
- 3.x.x) One-Liner Definitions
- 4.x.x) Linear Algebra
- 5.x.x) Discrete Mathematics
- 9.x.x) General Stuff

Definition 1.1.1 (Supremum/Infimum) Let $X \subset \mathbb{R}$ be a non-empty. The supremum of X is a real number $M =: \sup X$ that satisfies

- (i) M is an upper bound of X , and
- (ii) if M' is an upper bound of X , then $M' \geq M$

that is, M is the least upper bound of X . The infimum of X is the greatest lower bound of X .

Definition 1.1.2 (Subsequential limit) Let $\{x_n \in \mathbb{R}\}$ sequence. $\bar{x} \in \mathbb{R}$ is called a **subsequential limit** of $\{x_n\}$ if $\{x_n\}$ has a subsequence $\{x_{n_k}\}$ which converges to \bar{x} .

Definition 1.1.3 (Limit superior/limit inferior) Let $\{x_n \in \mathbb{R}\}$ sequence, and let $S(x_n)$ be the set of all subsequential limits of $\{x_n\}$.

Then we define the **limit superior** of $\{x_n\}$ to be

$$\limsup x_n := \sup S(x_n)$$

and the **limit inferior** of $\{x_n\}$ to be

$$\liminf x_n := \inf S(x_n)$$

Alternatively, we can also define them by

$$\begin{aligned}\limsup x_n &:= \lim_{n \rightarrow \infty} \sup \{x_k \mid k \geq n\} \\ \liminf x_n &:= \lim_{n \rightarrow \infty} \inf \{x_k \mid k \geq n\}\end{aligned}$$

Definition 1.1.4 (Cluster point) Let S be a subset of a topological space X . A point x in X is a cluster point of the set S if every neighborhood of x contains at least one point of S different from x itself.

A cluster point is also called a limit point or accumulation point.

In real analysis, $c \in \mathbb{R}$ is a cluster point of a non-empty set $A \subseteq \mathbb{R}$ if for every $\varepsilon > 0$ there exists a point $x \in A \setminus \{c\}$ such that $x \in (c - \varepsilon, c + \varepsilon)$.

In complex analysis, $c \in \mathbb{C}$ is a cluster point of a non-empty set $A \subseteq \mathbb{C}$ if for every $\varepsilon > 0$ there exists a point $z \in A \setminus \{c\}$ such that $z \in B_\varepsilon(c)$.

Definition 1.1.5 (Dense) Informally, a subset A of a topological space X is said to be **dense** in X if every point of X either belongs to A or else is arbitrarily “close” to a member of A .

A subset A of a topological space X is said to be a dense subset of X if any of the following equivalent conditions are satisfied:

- (i) The smallest **closed subset** of X containing A is X itself.
- (ii) The closure of A in X is equal to X . ($\text{cl}_X A = X$).
- (iii) Every point in X either belongs to A or is a **cluster point** of A .

Definition 1.1.6 (Point of closure) For S as a subset of a Euclidean space, x is a point of closure of S if every open ball centered at x contains a point of S (this point can be x itself).

Definition 1.1.7 (Closure) The closure of a subset S of points in a topological space can be defined using any of the following equivalent definitions:

- (i) $\text{cl } S$ is the set of all **points of closure** of S .
- (ii) $\text{cl } S$ is the set S together with all of its **limit points**.
- (iii) $\text{cl } S$ is the intersection of all closed sets containing S .
- (iv) $\text{cl } S$ is the smallest closed set containing S .
- (v) $\text{cl } S$ is the union of S and its boundary ∂S .

Definition 1.1.8 (Open sets) A subset U of a metric space (M, d) is called open if for any point x in U , there exists a real number $\varepsilon > 0$ such that any point $y \in M$ satisfying $d(x, y) < \varepsilon$ belongs to U .

Equivalently, U is open if every point U has a neighborhood contained in U .

An example of a metric space is $(\mathbb{R}^2, \|\cdot\|)$.

Definition 1.1.9 (Closed sets) A subset A of a topological space (X, τ) is closed if its complement $X \setminus A$ is an **open** subset of (X, τ) .

A set A is closed in X if and only if it is equal to its closure $\text{cl } A$ in X .

Yet another equivalent definition is that a set is closed if and only if it contains all of its boundary points.

Definition 1.2.1 (Monotone sequences) A sequence $\{x_n\}$ is said to be **increasing** if $x_0 \leq x_1 \leq x_2 \leq \dots$ and **decreasing** if $x_0 \geq x_1 \geq x_2 \geq \dots$ and **monotone** if it is either increasing or decreasing.

Theorem 1.2.2 (Monotone convergence theorem) If $\{x_n\}$ is monotone and bounded, then $\{x_n\}$ converges.

$$\lim_{n \rightarrow \infty} = \begin{cases} \sup\{x_n : n \in \mathbb{N}\} & \text{if } \{x_n\} \text{ is increasing} \\ \inf\{x_n : n \in \mathbb{N}\} & \text{if } \{x_n\} \text{ is decreasing} \end{cases}$$

Theorem 1.2.3 (Monotone subsequence theorem) Every sequence has a monotone subsequence.

Proof. Let $\{x_n\}$ be a sequence. We call a term x_p a **peak term** of $\{x_n\}$ if

$$x_p \geq x_n \quad (\forall n \geq p)$$

That is, all terms after x_p never go above x_p again. Then there are only two cases:

Case 1: $\{x_n\}$ has infinitely many peak terms.

Then the subsequence formed by all the peak terms form a decreasing subsequence of $\{x_n\}$.

Case 2: $\{x_n\}$ has finitely many peak terms.

Let $x_{p_1}, x_{p_2}, \dots, x_{p_j}$ be **all** the peak terms.

Let $n_1 = p_j + 1$ be the first term after the last peak term.

Since x_{n_1} is not a peak term. $\implies \exists n_2 > n_1$ such that $x_{n_1} < x_{n_2}$.

Since x_{n_2} is not a peak term, $\implies \exists n_3 > n_2$ such that $x_{n_2} < x_{n_3}$.

Continuing indefinitely, we can form an increasing subsequence $\{x_{n_k}\}$. □

Theorem 1.2.4 (Bolzano-Weierstrass Theorem) Every bounded sequence has a convergent subsequence.

Proof. Let $\{x_n\}$ be a bounded sequence. By the monotone subsequence theorem, $\{x_n\}$ has a monotone subsequence $\{x_{n_k}\}$.

Since $\{x_n\}$ is bounded, so is $\{x_{n_k}\}$.

Since $\{x_{n_k}\}$ is both monotone and bounded, it follows from the **monotone convergence theorem** that $\{x_{n_k}\}$ converges. □

Theorem 1.2.5 (Monotone seq. with a convergent subseq. is convergent) Let $\{x_n\}$ be a monotone sequence with a subsequence $\{x_{n_k}\}$ that converges to L . Then $\{x_n\}$ converges to L .

Proof. WLOG, assume that $\{x_n\}$ is decreasing. Given any $\varepsilon > 0$, we want to find a $N_\varepsilon \in \mathbb{N}$ such that

$$|x_n - L| < \varepsilon \quad \forall (n \geq N_\varepsilon)$$

Since $\{x_{n_k}\}$ is decreasing and converges to L , we can find (and fix) a k_ε such that

$$0 < x_{n_k} - L < \varepsilon \quad \forall (k \geq k_\varepsilon) \tag{*}$$

So we take $N_\varepsilon = n_{k_\varepsilon}$. Then since $\{x_n\}$ is decreasing,

$$x_n \leq x_{N_\varepsilon} = x_{n_{k_\varepsilon}} \quad \forall (n \geq N_\varepsilon)$$

Moreover, $L \leq x_n \leq x_{n_{k_\varepsilon}}$, and hence

$$0 \leq x_n - L \leq x_{n_{k_\varepsilon}} - L$$

and from (*), we have that this entire inequality $< \varepsilon$, and hence

$$0 \leq x_n - L < \varepsilon$$

and finally

$$|x_n - L| < \varepsilon$$

□

Theorem 1.2.6 (Mean value theorem) Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on the $[a, b]$, and differentiable on (a, b) . Then there exists $c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Generalized to multiple variables, the mean value theorem can be written as:

Let $f : [a, b] \rightarrow \mathbb{R}$, where $a, b \in \mathbb{R}^n$, and $[a, b]$ refers to the line segment connecting a and b , namely

$$[a, b] := \{\lambda a + (1 - \lambda)b \mid \lambda \in [0, 1]\}$$

Suppose f is continuous on $[a, b]$ and differentiable on (a, b) . Then there exists $c \in [a, b]$ such that

$$\nabla f(c)^T(b - a) = f(b) - f(a)$$

In some arguments, we use $f : [x, x + td] \rightarrow \mathbb{R}$ and write that there exists $\eta \in [x, x + td]$ such that

$$\nabla f(\eta)^T d = \frac{f(x + td) - f(x)}{t}$$

Result 1.2.7 (Preprocessed limits) Let $k, \ell \in \mathbb{N}$ and $a, b, c \in \mathbb{R}$ be fixed.

(a) $\lim_{n \rightarrow \infty} \frac{1}{n^k} = 0$

(b) $\lim_{n \rightarrow \infty} b^n = 0$ if $|b| < 1$

(c) $\lim_{n \rightarrow \infty} c^{\frac{1}{n}} = 1$ if $c > 0$

(d) $\lim_{n \rightarrow \infty} n^{\frac{1}{n}} = 1$

(e) $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$

(f) $\lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^n = \frac{1}{e}$

(g) $\lim_{n \rightarrow \infty} \frac{n^k}{c^n} = 0$ if $c > 0$

if $k < \ell$ and $1 < a < b$, we have

$$n^k \ll n^\ell \ll a^n \ll b^n \ll n!$$

Theorem 1.2.8 (Bernoulli's inequality)

$$(1+x)^r \geq 1+rx$$

This holds under any of the following conditions:

- $r \in \mathbb{Z}, r \geq 1$ and $x \in \mathbb{R}, x \geq -1$ (inequality is strict if $x \neq 0$ and $r \geq 2$)
- $r \in \mathbb{Z}, r \geq 0$ and $x \in \mathbb{R}, x \geq -2$
- $r \in \mathbb{Z}, r$ is even and $x \in \mathbb{R}$
- $r \in \mathbb{R}, r \geq 1$ and $x \in \mathbb{R}, x \geq -1$ (inequality is strict if $x \neq 0$ and $r \neq 1$)

and separately,

$$(1+x)^r \leq 1+rx$$

for every $r \in \mathbb{R}, 0 \leq r \leq 1$ and $x \geq -1$.

Result 1.2.9 (Limit to infinity of a rational function) Let P, Q be polynomial functions, where Q is of a higher degree. Then

$$\lim_{x \rightarrow \infty} \frac{P(x)}{Q(x)} = 0$$

Compute. Consider the example of

$$\lim_{x \rightarrow \infty} \frac{x^2 - 3x}{x^3 + 2x + 5}$$

We can divide both numerator and denominator by x^2 to obtain

$$\lim_{x \rightarrow \infty} \frac{1 - \frac{3}{x}}{x + \frac{2}{x} + \frac{5}{x^2}}$$

And we can see that the numerator $\rightarrow 1$ while the denominator $\rightarrow \infty$. □

Result 1.2.10 (Limit of $\frac{e^x}{x}$ as $x \rightarrow \infty$)

$$\lim_{x \rightarrow \infty} \frac{e^x}{x} = \infty$$

Proof. Since e^x can be written as a Taylor series

$$e^x = 1 + x + \frac{x^2}{2} + \dots$$

We have $e^x \geq 1 + x + x^2$ and hence

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{e^x}{x} &\geq \lim_{x \rightarrow \infty} \frac{1 + x + \frac{x^2}{2}}{x} \\ &= \lim_{x \rightarrow \infty} \frac{1}{x} + 1 + \frac{x}{2} \\ &= \infty \end{aligned}$$

□

Result 1.2.11 (Limit of $\frac{\ln x}{x}$ as $x \rightarrow \infty$)

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x} = 0$$

Proof. Given any ε , we have to find a $N \in \mathbb{N}$ such that

$$n \geq N \implies \frac{\ln x}{x} < \varepsilon$$

But, if you've been paying attention,

$$\frac{\ln x}{x} < \varepsilon \iff \frac{e^{\varepsilon x}}{\varepsilon x} > \frac{1}{\varepsilon}$$

And since $\varepsilon x \rightarrow +\infty$, using [Result 1.2.10](#) with εx as the limiting variable tells us that indeed there exists such an N , hence completing the proof. \square

Result 1.2.12 (Limit of a polynomial divided by an exponential) Let $a, b \in \mathbb{R}$ be fixed, with $b > 1$. Then we have

$$\lim_{x \rightarrow \infty} \frac{x^a}{b^x} = 0$$

Proof. Given any ε we want to find a $N \in \mathbb{N}$ such that

$$n \geq N \implies \frac{x^a}{b^x} < \varepsilon$$

But this is equivalent to

$$a \ln x - x \ln b < \ln \varepsilon$$

So it suffices to prove that

$$a \ln x - x \ln b \rightarrow -\infty.$$

Rewriting, we have

$$\begin{aligned} a \ln x - x \ln b &= x \left(a \cdot \frac{\ln x}{x} - \ln b \right) \\ &= \infty(-\ln b) \quad \because \frac{\ln x}{x} \rightarrow 0 \\ &= -\infty \end{aligned}$$

This completes the proof. \square

Definition 1.2.13 (Norm properties) Given a vector space X over a subfield F of the complex numbers \mathbb{C} , a **norm** on X is a real-valued function $p : X \rightarrow \mathbb{R}$ with the following properties, where $|k|$ denotes the absolute value of a scalar k .

(N1) (*Positive definiteness*) For all $x \in X$, if $p(x) = 0$ then $x = 0$.

(N2) (*Absolute homogeneity*) $p(kx) = |k|p(x)$ for all $x \in X$ and scalars k .

(N3) (*Subadditivity/Triangle inequality*) $p(x + y) \leq p(x) + p(y)$ for all $x, y \in X$

Theorem 1.2.14 (Limit and limit superior/inferior) Let $\{x_n\}$ be a bounded sequence. Then $\{x_n\}$ converges to \bar{x} if and only if

$$\limsup x_n = \liminf x_n = \bar{x}$$

In short,

$$\lim_{n \rightarrow \infty} x_n = \bar{x} \text{ (exists)} \iff \limsup x_n = \liminf x_n = \bar{x}$$

Sidenote: all convergent sequences are bounded, so the boundedness can be taken for free once convergence is established.

Result 1.2.15 (Limit of a polynomial divided by its successor) Let P be a polynomial. Show that

$$\lim_{x \rightarrow \infty} \frac{P(x)}{P(x+1)} = 1$$

Proof. We will write $P(x)$ as

$$P(x) := \sum_{i=0}^n a_i x^i$$

where n is the degree of polynomial P .

$$\begin{aligned} P(x+1) &= P\left(x\left(1 + \frac{1}{x}\right)\right) \\ &= a_0 + a_1 x \left(1 + \frac{1}{x}\right) + a_2 x^2 \left(1 + \frac{1}{x}\right)^2 + \dots + a_n x^n \left(1 + \frac{1}{x}\right)^n \\ &= x^n \left[\frac{a_0}{x^n} + \frac{a_1}{x^{n-1}} \left(1 + \frac{1}{x}\right) + \frac{a_2}{x^{n-2}} \left(1 + \frac{1}{x}\right)^2 + \dots + a_n \left(1 + \frac{1}{x}\right)^n \right] \\ P(x) &= x^n \left(\frac{a_0}{x^n} + \frac{a_1}{x^{n-1}} + \frac{a_2}{x^{n-2}} + \dots + a_n \right) \\ \implies \frac{P(x)}{P(x+1)} &= \frac{\frac{a_0}{x^n} + \frac{a_1}{x^{n-1}} \left(1 + \frac{1}{x}\right) + \frac{a_2}{x^{n-2}} \left(1 + \frac{1}{x}\right)^2 + \dots + a_n \left(1 + \frac{1}{x}\right)^n}{\frac{a_0}{x^n} + \frac{a_1}{x^{n-1}} + \frac{a_2}{x^{n-2}} + \dots + a_n} \\ \implies \lim_{x \rightarrow \infty} \frac{P(x)}{P(x+1)} &= \frac{a_n}{a_n} = 1 \end{aligned}$$

□

Theorem 2.1.1 (Fundamental theorem of calculus)

First part Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Let $F : [a, b] \rightarrow \mathbb{R}$ be defined by

$$F(x) = \int_a^x f(t) dt$$

Then F is uniformly continuous on $[a, b]$ and differentiable on (a, b) , and

$$F'(x) = f(x)$$

on (a, b) so F is an antiderivative of f .

Corollary

$$\int_a^b f(t) dt = F(b) - F(a)$$

Second part Let $f : [a, b] \rightarrow \mathbb{R}$. Let $F : [a, b] \rightarrow \mathbb{R}$ be continuous and also the antiderivative of f in (a, b) . If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(t) dt = F(b) - F(a)$$

This is stronger than the corollary because it does not assume that f is continuous.

Definition 3.1.1 (Affine functions) An affine function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is of the form

$$f(x) = Ax - b \quad (A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m)$$

Definition 3.1.2 (Coercive functions) A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is coercive if

$$\lim_{\|x\| \rightarrow \infty} f(x) = +\infty$$

Definition 3.1.3 (Supercoercive functions) A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is supercoercive if

$$\lim_{\|x\| \rightarrow \infty} \frac{f(x)}{\|x\|} = +\infty$$

Remark 4.1.1 (Thinking about matrix dimensions) Let $A \in \mathbb{R}^{m \times n}$, $x \in \mathbb{R}^n$, and $b \in \mathbb{R}^m$. We can validly write

$$Ax = b$$

So A is a gadget that takes a n -dim vector and returns a m -dim vector.

(A has m rows and n columns)

Definition 4.1.2 (Positive (semi)definiteness) A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is positive definite if

$$x^T A x > 0 \quad \forall (x \in \mathbb{R}^n)$$

and positive semidefinite if

$$x^T A x \geq 0 \quad \forall (x \in \mathbb{R}^n)$$

Definition 4.1.3 (Inner product space) An inner product space is a vector space V over the field F together with an *inner product*.

An inner product is a map

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow F$$

that satisfies the following for all $x, y, z \in V$ and $a, b \in F$:

(I1) (Positive definiteness) If x is non-zero, then

$$\langle x, x \rangle > 0$$

(I2) (*Linearity in the first argument*)

$$\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$$

(I3) (*Conjugate symmetry*)

$$\langle x, y \rangle = \overline{\langle y, x \rangle}$$

Definition 5.0.0 (Absolute basics of boolean algebra)

- (a) Literal: a boolean variable x or $\neg x$ (or \bar{x})
- (b) Conjunction: \wedge (and)
- (c) Disjunction: \vee (or)
- (d) Clause: a disjunction of **distinct** literals

Definition 5.1.1 (Conjunctive normal form) This is a **conjunction** of one or more **clauses**.

$$(A \vee B) \wedge (C \vee D \vee E)$$

Definition 5.1.2 (Disjunctive normal form) This is a **disjunction** of one or more **conjunctions**.

$$(A \wedge B) \vee (C \wedge D \wedge E)$$

Proposition 5.1.3 (Extending a CNF to 3 variables) Given a 1-variable or 2-variable **CNF**, we want to write a logically equivalent 3-variable CNF. (Useful for 3-SAT problems). Here's how:

2-var CNF. Say we have the expression $(x \vee y)$. This is logically equivalent to

$$(x \vee y \vee z) \wedge (x \vee y \vee \bar{z})$$

Notice that if z is TRUE then we can drop the left branch because it's true and hence

$$(x \vee y \vee z) \wedge (x \vee y \vee \bar{z}) \equiv (x \vee y \vee \bar{z}) \equiv (x \vee y)$$

Similarly if z is FALSE then we drop the right branch and get

$$(x \vee y \vee z) \wedge (x \vee y \vee \bar{z}) \equiv (x \vee y \vee z) \equiv (x \vee y)$$

1-var CNF. Now consider the expression x . Instead of adding just one variable we now add two and get the logically equivalent expression

$$(x \vee y \vee z) \wedge (x \vee y \vee \bar{z}) \wedge (x \vee \bar{y} \vee z) \wedge (x \vee \bar{y} \vee \bar{z})$$

If $(y, z) = (\text{TRUE}, \text{TRUE})$ we can drop all clauses containing y or z , leaving us with

$$(x \vee \bar{y} \vee \bar{z})$$

but then $(\bar{y}, \bar{z}) = (\text{FALSE}, \text{FALSE})$ and hence it is logically equivalent to just x . Repeating this logic for all combinations of (y, z) , we can see that $(*)$ is logically equivalent to x .

Definition 9.1.1 (Gamma function) The gamma function is defined via a convergent improper integral:

$$\Gamma(z) := \int_0^{\infty} e^{-t} t^{z-1} dt \quad (\text{Re}(z) > 0)$$

Note that “ \int_0^{∞} ” is a shorthand for “ $\lim_{k \rightarrow \infty} \int_0^k$ ”.

Observe that $\Gamma(1) = 1$.

$$\int_0^{\infty} e^{-t} dt = \left[-e^{-t} \right]_0^{\infty} = 1$$

And that $\Gamma(n+1) = n\Gamma(n)$.

$$\begin{aligned} \int_0^{\infty} e^{-t} t^n dt &= \left[-e^{-t} \cdot t^n \right]_0^{\infty} - \int_0^{\infty} -e^{-t} \cdot nt^{n-1} dt \\ &= 0 + \int_0^{\infty} e^{-t} \cdot nt^{n-1} dt \quad (\text{by Result 1.2.12}) \\ &= n\Gamma(n) \end{aligned}$$

Definition 9.1.2 (Language reductions) If problem A is reducible to problem B , we write $A \leq B$.

Reducing A to B by a **Many-one reduction** is to find a function f which converts inputs x of A into inputs $f(x)$ of B , such that $A(x) = B(f(x))$ under all values of x .

Reducing A to B by a **Turing reduction** is to find a function which mimics the behavior of A using an oracle of B . i.e., $A(x) = \text{TRUE} \iff B(f(x)) = \text{TRUE}$.

A being reducible to B means solving A cannot be harder than the combined difficulty of solving B and executing the reduction. In particular, if the reduction runs in constant-time, A cannot be harder than B . In other words, \leq is referring to hardness.

Definition 9.1.3 (Everything P-, NP-related) This is a compilation of everything P- and NP-related. For in-depth definitions, refer to each link below.

A problem L is in P if it runs in polynomial time.

A problem L is in NP if it has a polynomial-time verifier.

We say that $L_1 \leq_P L_2$ if there is a polynomial-time **reduction** from L_1 to L_2 .

A problem L is NP-complete when $L \in \text{NP}$, and every problem L' in NP has a polynomial-time reduction to it:

$$\forall L' \in \text{NP} : L' \leq_P L$$

A problem H is NP-hard when for every $L \in \text{NP}$, there is a polynomial-time reduction from L to H :

$$\forall L \in \text{NP} : L \leq_P H \quad (*)$$

The only difference between NP-complete and NP-hard is that NP-complete has the extra constraint of having to be in NP.

(*), based on a **previous remark**, also implies that H is at least as hard as the hardest problem in NP.

Theorem 9.1.4 (Cauchy-Schwarz inequality) For all vectors u and v of an inner product space,

$$|\langle u, v \rangle|^2 \leq \langle u, u \rangle \cdot \langle v, v \rangle$$

This gives the following corollaries:

(a) Let $u_i, v_i \in \mathbb{R}$ for $i = 1, \dots, n$ for any integer n . Then

$$\left(\sum u_i v_i \right)^2 \leq \left(\sum u_i^2 \right) \left(\sum v_i^2 \right)$$

(b) Let $u_k, v_k \in \mathbb{C}$ for $k = 1, \dots, n$ for any integer n . Then

$$\left| \sum u_i v_i \right|^2 \leq \left(\sum |u_i|^2 \right) \left(\sum |v_i|^2 \right)$$

Proof. To prove (a), we observe that \mathbb{R}^n equipped with the standard dot product is an **inner product space**. We can build vectors $u, v \in \mathbb{R}^n$ by arranging u_i for $i = 1, \dots, n$ into a column vector and do the same for v_i to get v .

Then applying the Cauchy-Schwarz inequality with the standard dot product, we have

$$|u \cdot v|^2 \leq (u \cdot u)(v \cdot v)$$

Which gives the statement in (a) exactly.

To prove (b), instead of the **inner product space** constructed from \mathbb{R}^n and the standard dot product, we use \mathbb{C}^n and the complex inner product defined by

$$\langle u, w \rangle := u_1 \bar{w}_1 + \dots + u_n \bar{w}_n$$

Then by the Cauchy-Schwarz inequality, for all $u, w \in \mathbb{C}^n$,

$$\begin{aligned} |\langle u, w \rangle|^2 &= \left| \sum u_k \bar{w}_k \right|^2 \\ &\leq \langle u, u \rangle \cdot \langle w, w \rangle \\ &= \left(\sum u_k \bar{u}_k \right) \left(\sum w_k \bar{w}_k \right) \\ &= \left(\sum |u_k|^2 \right) \left(\sum |w_k|^2 \right) \end{aligned}$$

That is

$$|u_1 \bar{w}_1 + \dots + u_n \bar{w}_n|^2 \leq \left(|u_1|^2 + \dots + |u_n|^2 \right) \left(|\bar{w}_1|^2 + \dots + |\bar{w}_n|^2 \right)$$

But since $|z|^2 = |\bar{z}|^2$ for all $z \in \mathbb{C}$, we can define a collection v_1, \dots, v_n such that $v_k = \bar{w}_k$, then we can rewrite the above inequality as

$$|u_1 v_1 + \dots + u_n v_n|^2 \leq \left(|u_1|^2 + \dots + |u_n|^2 \right) \left(|v_1|^2 + \dots + |v_n|^2 \right)$$

And finally since the collection w_k were arbitrarily chosen, so can the collection v_k . □

Calculus

Remark 1.1.1 (Trigonometric identities)

$$\begin{aligned}\sin 2x &= 2 \sin x \cos x & \sin^2 x &= \frac{1 - \cos 2x}{2} \\ \cos 2x &= 1 - 2 \sin^2 x & \cos^2 x &= \frac{\cos 2x + 1}{2} \\ &= 2 \cos^2 x - 1\end{aligned}$$

$$\begin{aligned}\sin(A \pm B) &= \sin A \cos B \pm \cos A \sin B \\ \cos(A \pm B) &= \cos A \cos B \mp \sin A \sin B\end{aligned}$$

$$\begin{array}{l|l}\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) & 2 \sin A \cos B = \sin\left(\frac{A+B}{2}\right) + \sin\left(\frac{A-B}{2}\right) \\ \sin A - \sin B = 2 \cos\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right) & 2 \cos A \sin B = \sin\left(\frac{A+B}{2}\right) - \sin\left(\frac{A-B}{2}\right) \\ \cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right) & 2 \cos A \cos B = \cos\left(\frac{A+B}{2}\right) + \cos\left(\frac{A-B}{2}\right) \\ \cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right) & -2 \sin A \sin B = \cos\left(\frac{A+B}{2}\right) - \cos\left(\frac{A-B}{2}\right)\end{array}$$

Result 1.1.2 (Basic trigonometric constants) Trigonometric preprocessing to finish homework in $O(1)$ time.

	0	$\pi/6$	$\pi/4$	$\pi/3$	$\pi/2$
sin	0	$1/2$	$1/\sqrt{2}$	$\sqrt{3}/2$	1
cos	1	$\sqrt{3}/2$	$1/\sqrt{2}$	$1/2$	0
tan	0	$1/\sqrt{3}$	1	$\sqrt{3}$	-

$$\begin{aligned}\sin(-x) &= -\sin(x) \\ \cos(-x) &= \cos(x) \\ \tan(-x) &= -\tan(x)\end{aligned}$$

Remark 2.1.1 (Differentiation identities)

Product rule $(uv)' = u'v + uv'$

Quotient rule $\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$

(can be derived from product rule using u and $\frac{1}{v}$)

Remark 2.1.2 (Integration identities)

$f(x)$	$\int f(x) dx$
$\frac{1}{x^2 + a^2}$	$\frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right)$
$\frac{1}{\sqrt{a^2 - x^2}}$	$\sin^{-1} \left(\frac{x}{a} \right) \quad (x < a)$
$\frac{1}{x^2 - a^2}$	$\frac{1}{2a} \ln \left(\frac{x - a}{x + a} \right) \quad (x > a)$
$\frac{1}{a^2 - x^2}$	$\frac{1}{2a} \ln \left(\frac{a + x}{a - x} \right) \quad (x < a)$
$\tan x$	$\ln(\sec x) \quad (x < \frac{\pi}{2})$
$\cot x$	$\ln(\sin x) \quad (0 < x < \pi)$
$\sec x$	$-\ln(\sec x + \tan x) \quad (x < \frac{\pi}{2})$
$\csc x$	$-\ln(\csc x + \cot x) \quad (0 < x < \pi)$

Remark 2.1.3 (Chain rule) In all the following scenarios, let $h := f \circ g$.

When f takes a scalar Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$. Then $h : \mathbb{R} \rightarrow \mathbb{R}$ and we have

$$h'(t) = f'(g(t)) \cdot g'(t)$$

And $f', g' : \mathbb{R} \rightarrow \mathbb{R}$.

When f takes a vector Let $g : \mathbb{R} \rightarrow \mathbb{R}^n$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Then $h : \mathbb{R} \rightarrow \mathbb{R}$ and we have

$$h'(t) = \nabla f(g(t))^T g'(t)$$

Note that $\nabla f(g(t)) \in \mathbb{R}^n$ and $g'(t) \in \mathbb{R}^n$.

When f takes a complex number Let $g : \mathbb{R} \rightarrow \mathbb{C}$ and $f : \mathbb{C} \rightarrow \mathbb{R}$. Then $h : \mathbb{R} \rightarrow \mathbb{R}$.

In particular, we write $g(t) = g_1(t) + ig_2(t)$ and $f : x + iy \mapsto f(x + iy)$.

Interestingly, we still have

$$(f \circ g)'(t) = f_x(g(t)) \cdot g_1'(t) + f_y(g(t)) \cdot g_2'(t)$$

Note the lack of i terms on the term with g_2' . This is intentional.

Remember anyway that $f \circ g : \mathbb{R} \rightarrow \mathbb{R}$, and so we must have $(f \circ g)' : \mathbb{R} \rightarrow \mathbb{R}$.

Definition 2.1.4 (Differentiability) In single-variable calculus, $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable if

$$\lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

exists (and if so, is denoted as $f'(a)$).

In multivariable calculus, $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable if there exists a **linear map** $J : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - J(h)}{\|h\|} = 0$$

Then from this perspective, differentiability of single-variable complex functions can be written as: $f : \mathbb{C} \rightarrow \mathbb{C}$ is differentiable if there is a linear map $J : \mathbb{C} \rightarrow \mathbb{C}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z+h) - f(z) - J(h)|}{|h|} = 0$$

Comment All of these cases are equivalent to saying that there exists a $k \in \mathbb{R}$ such that.

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = k$$

Essentially, that there exists a local **linearization** to the function.

Definition 2.1.5 (Directional derivative) Let $D \subset \mathbb{R}^n$ be open. $f : D \rightarrow \mathbb{R}$ is directionally differentiable at $\bar{x} \in \mathbb{R}^n$ in the direction $d \in \mathbb{R}^n$ if

$$\lim_{t \downarrow 0} \frac{f(\bar{x} + td) - f(\bar{x})}{t}$$

exists. This limit is denoted by $f'(\bar{x}; d)$ and is called the directional derivative of f at \bar{x} in the direction d .

If f is directionally differentiable at \bar{x} in every direction $d \in \mathbb{R}^n$, we call f directionally differentiable at \bar{x} .

If f is directionally differentiable at every $\bar{x} \in \mathbb{R}^n$, we call it directionally differentiable.

Remark 2.1.6 (Gradient) Only scalar-valued functions can have gradients.

The **gradient** of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is the vector-valued function ∇f whose value at p gives the direction and rate of fastest increase. Further, ∇f can be written as

$$\nabla f(p) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(p) \\ \vdots \\ \frac{\partial f}{\partial x_n}(p) \end{bmatrix}$$

The gradient of f is defined as the unique vector field whose dot product with any vector d at each point x is the directional derivative of f along d . That is,

$$\nabla f(x)^T d = f'(x; d)$$

Remark 2.1.7 (Jacobian) The Jacobian of a vector-valued function in several variables generalizes the **gradient** of a scalar-valued function in several variables.

In other word, the Jacobian of a scalar-valued function is its gradient.

So let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. In particular,

$$f(x) := \begin{bmatrix} f_1(x) \\ \vdots \\ f_m(x) \end{bmatrix} \quad f_i : \mathbb{R}^n \rightarrow \mathbb{R}$$

Then the Jacobian of f is an $m \times n$ matrix

$$\mathbf{J} = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \cdots & \frac{\partial f}{\partial x_n} \end{bmatrix} = \begin{bmatrix} \nabla^T f_1 \\ \vdots \\ \nabla^T f_m \end{bmatrix} = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix} = \nabla f(x)^T$$

Note that by definition of the **gradient**, we have

$$\nabla f(x)^T = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \cdots & \frac{\partial f}{\partial x_n} \end{bmatrix} = \mathbf{J}$$

and hence we say that the Jacobian is the transpose of the gradient.

Algorithm Design

Definition 0.0.1 (Vertex cover) Let $G = (V, E)$ be an undirected graph. A vertex cover $U \subseteq V$ satisfies

$$(u, v) \in E \implies u \in U \vee v \in U.$$

In other words, every edge in E has at least one endpoint in the vertex cover U . Such a set is said to *cover* the edges of G .

Definition 1.1 (Flow network) A *flow network* is a directed graph $G = (V, E)$ with a single *source node* s and a single *target node* t , as well as a positive number $c(e)$ for each edge $e \in E$, called the capacity of e .

Definition 1.3 (Flow) Let G be a flow network. A *flow* on G is given by a positive number $f(e)$ for each edge e in G satisfying the following two constraints:

- **Capacity constraints.** For each edge $e \in E$, we have $0 \leq f(e) \leq c(e)$
- **Flow conservation.** For each vertex $v \in V$ that is not the source or target vertex,

$$\sum_{e \text{ into } v} f(e) = \sum_{e \text{ leaving } v} f(e)$$

The *value* of flow f is all of the flow leaving s :

$$\text{val}(f) := \sum_{e \text{ leaving } s} f(e)$$

where s is the source node of G .

Problem 1 (Max Flow) Input: A flow network G with source s , target t , and positive edge capacities $c(e)$ for $e \in E$.

Output: A flow f with the maximum value.

Definition 1.4 (Residual graph) Let G be a flow network and let f be a flow on G . The *residual graph* of G and f , denoted by G_f , is the directed graph defined as follows:

The vertices of G_f are the same as the vertices of G .

For each edge $e = (u, v)$ in G , if $f(e) < c(e)$ then we add the edge (u, v) to G_f , labelled with the number $c(e) - f(e)$. If $f(e) > 0$, then we also add the edge (v, u) to G_f , labelled with the number $f(e)$.

All paths from s to t in the residual graph correspond to a sequence where flow can be re-routed to increase its value.

Definition 3.0 (Binary representation) Let

$$\{0, 1\}^* := \{\varepsilon, 0, 1, 00, 01, 10, 11, 100, 101, \dots\}$$

be the set of all finite binary strings. (where ε is the empty string.)

Definition 3.1 (Decision problem) A decision problem L is a subset of $\{0, 1\}^*$. The computational task corresponding to L is “Given a string $x \in \{0, 1\}^*$, is $x \in L$?”

Problem 8 (L -membership problem) **Input:** A boolean string x .

Output: Decide if $x \in L$.

Example 3.1.1 (Rewriting problems as L -membership problems)

- *Graph Connectivity.* Given a graph $G = (V, E)$, is it connected?

$$\begin{aligned} L &= \{x \in \{0, 1\}^* \mid x \text{ encodes a connected graph}\} \\ &= \{G \mid G \text{ is a connected graph}\} \end{aligned}$$

- *Max Flow (Decision Version).* Given a flow network G and a positive integer k , does the max flow on G have value $\geq k$?

$$\begin{aligned} L &= \{x \in \{0, 1\}^* \mid x \text{ encodes a } (G, k) \text{ such that } \text{val}(G) \geq k\} \\ &= \{(G, k) \mid \text{val}(G) \geq k\} \end{aligned}$$

- *Sum.* Given $a, b, c \in \mathbb{Z}$, does $a + b = c$?

$$\begin{aligned} L &= \{x \in \{0, 1\}^* \mid x \text{ encodes } (a, b, c) \text{ such that } a + b = c\} \\ &= \{(a, b, c) \mid a + b = c\} \end{aligned}$$

Choice of encoding is important and definitely affects runtime. However, for discussion we will assume that the most natural and succinct encoding is chosen.

Definition 3.2 (Polynomial-time algorithms) An algorithm A runs in *polynomial time* if $\exists c \in \mathbb{R}$ s.t. $\forall x \in \{0, 1\}^*$, A terminates after $O(|x|^c)$ computation steps.

A decision problem L is *polynomial-time computable* if there exists a polynomial-time A s.t. $\forall x \in \{0, 1\}^*$, $x \in L \iff A(x) = \text{Yes}$.

We define

$$P := \{L \subseteq \{0, 1\}^* \mid L \text{ is polynomial-time computable}\}$$

The complexity class P is our proxy for *efficiently computable languages*. (“language” is another way to refer to L , in addition to “problem”.)

Definition 3.3.1 (Vertex Cover Problem)

$$VC := \{(G, k) \mid G \text{ is a graph with a **vertex cover** of size } \leq k\}$$

Definition 3.3.2 (Satisfiability Problem)

$$\text{FORMSAT} := \{F \mid F \text{ is a satisfiable boolean formula}\}$$

Definition 3.3 (Nondeterministic polynomial-time algorithms) A decision problem L has a polynomial-time verifier if there is a polynomial time algorithm B taking two strings x, y as input, and a polynomial $p(n)$ such that

$$x \in L \iff \exists y \in \{0, 1\}^*, |y| \leq p(|x|) : B(x, y) = \text{Yes}$$

The complexity class

$$NP := \{L \subseteq \{0, 1\}^* \mid L \text{ has a polynomial-time verifier}\}$$

- A polynomial-time verifier for VC would take a graph (G, k) and a proposed vertex cover U and check if $|U| \leq k$ and that U is a vertex cover.
- A polynomial-time verifier for SAT would take in a boolean formula F and a proposed assignment x and check if $F(x) = \text{Yes}$.

Observe that $P \subseteq NP$. If L has a polynomial-time algorithm, then it also has a polynomial-time verifier.

Proof. If $L \in P$, then by definition there exists a polynomial-time algorithm A with

$$x \in L \iff A(x) = \text{Yes}$$

Then, following the definition of NP, we need to find a polynomial-time verifier B such that

$$x \in L \iff \exists y \in \{0, 1\}^*, |y| \leq p(|x|) : B(x, y) = \text{Yes}$$

But we can simply use $B(x, y) := A(x)$. □

Definition 3.4 (Complement of a decision problem) The complement of a decision problem L is defined as

$$\bar{L} = \{0, 1\}^* \setminus L := \{x \in \{0, 1\}^* \mid x \notin L\}$$

Note that $\{0, 1\}^* = L \cup \bar{L}$ for any decision problem L .

Exercise 3.5 Prove that if $L \in P$ then $\bar{L} \in P$.

Proof. If $L \in P$, then by definition there exists a polynomial-time algorithm A with

$$x \in L \iff A(x) = \text{Yes}$$

Then, following the definition of P, we need to find a polynomial-time algorithm B such that

$$x \in \bar{L} \iff B(x) = \text{Yes}$$

But we can simply use $B(x) := \neg A(x)$. □

Definition 3.6 (coNP) The complexity class coNP is defined as

$$\text{coNP} := \{L \mid \bar{L} \in \text{NP}\}$$

For example, recall that $\text{SAT} = \{F \mid F \text{ is a satisfiable boolean formula}\}$. Then

$$\overline{\text{SAT}} = \left\{ x \in \{0, 1\}^* \left| \begin{array}{l} x \text{ is an invalid encoding of a formula, or} \\ x \text{ encodes an unsatisfiable boolean formula} \end{array} \right. \right\}$$

But given $x \in \{0, 1\}^*$, it is easy to test its validity as a boolean formula, hence we focus on the second constraint:

$$\text{CONT} := \{F \mid F \text{ is an unsatisfiable boolean formula}\}$$

Note that since $\overline{\text{CONT}} = \text{SAT} \in \text{NP}$, we have that $\overline{\text{CONT}} \in \text{NP}$.

However, is $\text{CONT} \in \text{NP}$? Observe that $F \in \text{CONT}$ if and only if for every assignment x to the variables of F , we have that $F(x) = \text{No}$. Since there are 2^n assignments to check, it is not clear how to encode this checking procedure into a single polynomial-sized certificate. For this reason, many researchers conjecture that $\text{NP} \neq \text{coNP}$.

Remark 3.6.1 (The complexity class $\text{NP} \cap \text{coNP}$) We can show that $\text{P} \subseteq \text{coNP}$ by

$$L \in \text{P} \implies \bar{L} \in \text{P} \implies \bar{L} \in \text{NP} \implies L \in \text{coNP}$$

And since $\text{P} \subseteq \text{NP}$, we have

$$\text{P} \subseteq \text{NP} \cap \text{coNP}$$

Complex Analysis

Definition 0.0.0 (General terminology)

Entire function is a complex-valued function that is holomorphic on \mathbb{C} .

A **real-valued** function is any function $f : X \rightarrow \mathbb{R}$.

A **complex-valued** function is any function $f : X \rightarrow \mathbb{C}$.

A subset of \mathbb{R}^n or \mathbb{C}^n is called **compact** if it is closed and bounded.

The C^n notation:

- C^0 : continuous
- C^1 : continuously differentiable
- C^2 : twice continuously differentiable

Theorem 0.0.1 (Conventional notation)

For this chapter on Complex Analysis.

Let $U \subseteq \mathbb{C}$ be an open set.

Let $D(P, r)$ be the open disc centered at P with radius r . Then

- (i) $\partial D(P, r)$ is the (closed) curve at the border of $D(P, r)$
- (ii) $\overline{D}(P, r)$ is the closed disc centered at P with radius r .

Result 0.0.2 (Basic complex arithmetic)

$$\begin{aligned} |z|^2 &= z\bar{z} \\ |zw|^2 &= |z|^2|w|^2 \\ |z + w|^2 &= |z|^2 + |w|^2 + \operatorname{Re}(z \cdot \bar{w}) \\ |z + w|^2 + |z - w|^2 &= 2|z|^2 + 2|w|^2 \end{aligned}$$

Proof. Let $z := x + iy$, and $w := u + iv$.

$$|z|^2 = x^2 + y^2 = (x + iy)(x - iy) = z\bar{z}$$

$$\begin{aligned} |zw|^2 &= (xu - yv)^2 + (xv + yu)^2 \\ &= (x^2u^2 - 2xyvu + y^2v^2) + (x^2v^2 + 2xyvu + y^2u^2) \\ &= x^2u^2 + y^2v^2 + x^2v^2 + y^2u^2 \\ &= (x^2 + y^2)(u^2 + v^2) \\ &= |z|^2|w|^2 \end{aligned}$$

□

Theorem 0.0.3 (Complex differentiability) A complex function $f(z) := u(z) + iv(z)$ is complex-differentiable at z_0 if and only if u and v satisfy the **Cauchy-Riemann Equations** at z_0 .

To say a function is **holomorphic** is much stronger, since a holomorphic function is complex-differentiable at every point of some open subset of the complex plane \mathbb{C} .

Definition 1.1.3 (Complex Partial)

$$\frac{\partial f}{\partial z} \equiv \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) f$$

$$\frac{\partial f}{\partial \bar{z}} \equiv \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) f$$

Definition 1.4.1 (Holomorphic functions) Let $U \subset \mathbb{C}$ be open. Let $f : U \rightarrow \mathbb{C}$ be in $C^1(U)$. f is said to be *holomorphic* if

$$\frac{\partial f}{\partial \bar{z}} = 0$$

Properties of holomorphic functions If f and g are holomorphic in a domain U , then so are $f + g$, $f - g$, fg , and $f \circ g$.

Additionally, if g has no zeros in U , then f/g is holomorphic too.

Examples of holomorphic functions Here are some building blocks to get started (remember that you can use these with the properties above to show that other more complicated functions are holomorphic too):

- (i) $f(z) = 1/z$ on $\mathbb{C} \setminus \{0\}$
- (ii) $f(z) = 1/p(z)$ on \mathbb{C} where $p(z) \neq 0$
- (iii) $f(z) = z$ on \mathbb{C}

All these can be proved using a destructuring of $z := x + iy$ and using [Definition 1.1.3](#).

Here are some functions that are not holomorphic:

- (i) $f(z) = \bar{z}$
- (ii) $f(z) = |z|$

Showing that a function is holomorphic If we can write $f \equiv u + iv$, and u and v have **continuous** first partial derivatives and satisfy the [Cauchy-Riemann equations](#), then f is holomorphic.

Definition 1.4.2 (Cauchy-Riemann Equations) If $f(z) = u(z) + iv(z)$ is [holomorphic](#), then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Proposition 1.4.3 If $f : U \rightarrow \mathbb{C}$ is C^1 and f satisfies the Cauchy-Riemann equations, then

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial x} = -i \frac{\partial f}{\partial y}$$

Definition 1.4.4 (Harmonic functions) Let $U \subset \mathbb{C}$ be open. Let $f : U \rightarrow \mathbb{C}$ be in $C^2(U)$. f is said to be *harmonic* if

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0$$

The operator

$$\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$$

is called the *Laplace operator*, or *Laplacian*, and is denoted by Δ . We write

$$\Delta f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$$

Theorem 1.5.1 Let $f, g \in C^1(U)$ where

$$U := \{(x, y) \in \mathbb{R}^2 : |x - a| < \delta, |y - b| < \varepsilon\}$$

and let $\frac{\partial f}{\partial y} = \frac{\partial g}{\partial x}$ on U . Then there exists a function $h \in C^2(U)$ such that

$$\frac{\partial h}{\partial x} = f \quad \text{and} \quad \frac{\partial h}{\partial y} = g$$

on U . If f and g are real-valued, then we may take h to be real-valued also.

Theorem 1.5.3 Let $U \subset \mathbb{C}$ be either an open rectangle or open disc, and let F be holomorphic on U . Then there exists a holomorphic function H on U such that

$$\frac{\partial H}{\partial z} = F$$

on U .

Definition 2.1.1 (Bounded C^1 functions) A function $\phi : [a, b] \rightarrow \mathbb{R}$ is continuously differentiable (and we write $\phi \in C^1([a, b])$) if

- (a) ϕ is continuous on $[a, b]$
- (b) ϕ' exists on (a, b)
- (c) ϕ' has a continuous extension to $[a, b]$

In other words, for (c) we require that

$$\lim_{t \rightarrow a^+} \phi'(t) \quad \text{and} \quad \lim_{t \rightarrow b^-} \phi'(t)$$

both exist.

The motivation for this definition is so if $\phi \in C^1([a, b])$, then we have

$$\begin{aligned} \phi(b) - \phi(a) &= \lim_{\varepsilon \rightarrow 0^+} (\phi(b - \varepsilon) - \phi(a + \varepsilon)) \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{a+\varepsilon}^{b-\varepsilon} \phi'(t) dt \\ &= \int_a^b \phi'(t) dt \end{aligned}$$

and hence have the **fundamental theorem of calculus** hold for $\phi \in C^1([a, b])$.

Definition 2.1.2 (Continuous complex curve) Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be defined by $\gamma(t) := \gamma_1(t) + i\gamma_2(t)$.

Then γ is said to be continuous on $[a, b]$ if both γ_1 and γ_2 are.

The curve γ is $C^1([a, b])$ if γ_1 and γ_2 are continuously differentiable on $[a, b]$. Under these circumstances we will write

$$\gamma'(t) = \frac{d\gamma}{dt} = \frac{d\gamma_1}{dt} + i \frac{d\gamma_2}{dt}$$

Definition 2.1.3 (Complex integration) Let $\psi : [a, b] \rightarrow \mathbb{C}$ be continuous on $[a, b]$. Write $\psi(t) = \psi_1(t) + i\psi_2(t)$. Then we define

$$\int_a^b \psi(t) dt := \int_a^b \psi_1(t) dt + i \int_a^b \psi_2(t) dt$$

Using this definition along with Definitions 2.1.1 and 2.1.2, we have that if $\gamma \in C^1([a, b])$ is complex-valued, then

$$\gamma(b) - \gamma(a) = \int_a^b \gamma'(t) dt$$

Proposition 2.1.4 Let $U \subseteq \mathbb{C}$ be open and let $\gamma : [a, b] \rightarrow U$ be a C^1 curve. If $f : U \rightarrow \mathbb{R}$ and $f \in C^1(U)$ and we write

$$\begin{aligned} f : x + iy &\mapsto f(x + iy) \\ \gamma(t) &= \gamma_1(t) + i\gamma_2(t) \end{aligned}$$

then

$$\begin{aligned} f(\gamma(b)) - f(\gamma(a)) &= \int_a^b (f \circ \gamma)'(t) dt \\ &= \int_a^b \left(\frac{\partial f}{\partial x}(\gamma(t)) \cdot \frac{d\gamma_1}{dt} + \frac{\partial f}{\partial y}(\gamma(t)) \cdot \frac{d\gamma_2}{dt} \right) dt \\ &= \int_a^b f_x(\gamma(t)) \cdot \gamma_1'(t) + f_y(\gamma(t)) \cdot \gamma_2'(t) dt \end{aligned}$$

This follows from Definition 2.1.3 and the chain rule.

(the lack of an i term is intentional. Remember that $f \circ \gamma : \mathbb{R} \rightarrow \mathbb{R}$)

Definition 2.1.5 (Complex line integral) Let $U \subseteq \mathbb{C}$ open, $F : U \rightarrow \mathbb{C}$ continuous on U , and let $\gamma : [a, b] \rightarrow U$ be a C^1 curve. Then we define the complex line integral

$$\oint_{\gamma} F(z) dz := \int_a^b F(\gamma(t)) \cdot \frac{d\gamma}{dt} dt$$

Proposition 2.1.6 (Holomorphic line integral) Let $U \subseteq \mathbb{C}$ open, $F : U \rightarrow \mathbb{C}$ continuous on U , and let $\gamma : [a, b] \rightarrow U$ be a C^1 curve. If f is a holomorphic function on U , then

$$f(\gamma(b)) - f(\gamma(a)) = \oint_{\gamma} \frac{\partial f}{\partial z}(z) dz$$

Definition 2.1.6a (Complex antiderivative) A function f has an antiderivative F if and only if, for every $\gamma : [a, b] \rightarrow \mathbb{C}$,

$$\oint_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a))$$

This comes from using a holomorphic function on [Proposition 2.1.4](#), and then applying [Definition 2.1.5](#).

Proposition 2.1.7 (Moving $\|$ into integral) Let $\phi : [a, b] \rightarrow \mathbb{C}$ be continuous. Then

$$\left| \int_a^b \phi(t) dt \right| \leq \int_a^b |\phi(t)| dt$$

Proposition 2.1.8 (Upper bound of line integral) Let $U \subseteq \mathbb{C}$ be open and $f \in C^0(U)$. Let $\gamma : [a, b] \rightarrow U$ be a C^1 curve, and let $\ell(\gamma)$ be given by

$$\ell(\gamma) := \int_a^b \left| \frac{d\gamma}{dt}(t) \right| dt$$

Then we have

$$\left| \oint_{\gamma} f(z) dz \right| \leq \left(\sup_{t \in [a, b]} |f(\gamma(t))| \right) \cdot \ell(\gamma)$$

(Note that $\ell(\gamma)$ is the length of γ .)

Proposition 2.1.9 (Parameterization-independence of line integrals) Let $U \subseteq \mathbb{C}$ be open and $f : U \rightarrow \mathbb{C}$ be a continuous function. Let $\gamma : [a, b] \rightarrow U$ be a C^1 curve. Suppose that $\phi : [c, d] \rightarrow [a, b]$ is a bijective increasing C^1 with a C^1 inverse.

Let $\tilde{\gamma} = \gamma \circ \phi$. Then

$$\oint_{\tilde{\gamma}} f(z) dz = \oint_{\gamma} f(z) dz$$

The proof involves the standard change of variable formula from calculus.

Theorem 2.2.1 (Existence of f' on holomorphic f) Let $U \subseteq \mathbb{C}$ be open and let f be holomorphic on U . Then f' exists at each point of U and

$$f'(z) = \frac{\partial f}{\partial z}$$

for all $z \in U$.

As a result of this theorem, we often will write $f' = \frac{\partial f}{\partial z}$ when f is holomorphic.

Theorem 2.2.2 (Holomorphic by existence of derivative) Let $U \subseteq \mathbb{C}$ be open. If $f \in C^1(U)$ and f has a complex derivative at each point of U , then f is holomorphic on U .

In other words, if a continuous, complex-valued function f on U has a complex derivative at each point and if f' is continuous on U , then f is holomorphic on U .

Theorem 2.2.3 (Holomorphism and directional derivatives) Let f be holomorphic in a neighborhood $P \in \mathbb{C}$. Let $w_1, w_2 \in \mathbb{C}$ have unit modulus. Consider the directional derivatives

$$D_{w_1}f(P) := \lim_{t \rightarrow 0} \frac{f(P + tw_1) - f(P)}{t}$$

$$D_{w_2}f(P) := \lim_{t \rightarrow 0} \frac{f(P + tw_2) - f(P)}{t}$$

Then

- (a) $|D_{w_1}f(P)| = |D_{w_2}f(P)|$
- (b) if $|f'(P)| \neq 0$, then the directed angle from w_1 to w_2 equals the directed angle from $D_{w_1}f(P)$ to $D_{w_2}f(P)$.

Note:

- 2.2.3(a) alone implies that f is holomorphic.
- 2.2.3(b) alone implies that f is holomorphic.

Lemma 2.3.1 Let $(\alpha, \beta) \subseteq \mathbb{R}$ be an open interval and let $H, F : (\alpha, \beta) \rightarrow \mathbb{R}$ be continuous functions. Let $p \in (\alpha, \beta)$ and suppose that dH/dx exists and equals $F(x)$ for all $x \in (\alpha, \beta) \setminus \{p\}$. Then $(dH/dx)(p)$ exists and $(dH/dx)(x) = F(x)$ for all $x \in (\alpha, \beta)$.

$$\forall x \in (\alpha, \beta) \setminus \{p\} : \frac{dH}{dx}(x) = F(x) \implies \forall x \in (\alpha, \beta) : \frac{dH}{dx}(x) = F(x)$$

It's as if the continuity fills in the gap at p .

Theorem 2.3.2 Let $U \subseteq \mathbb{C}$ be either an open rectangle or an open disc and let $P \in U$. Let f and g be continuous, real-valued functions on U which are continuously differentiable on $U \setminus \{P\}$. Suppose further that

$$\frac{\partial f}{\partial y} = \frac{\partial g}{\partial x} \quad \text{on } U \setminus \{P\}$$

Then there exists a C^1 function $h : U \rightarrow \mathbb{R}$ such that

$$\frac{\partial h}{\partial x} = f, \quad \frac{\partial h}{\partial y} = g$$

at every point of U (including P).

Theorem 2.3.3 (Existence of holomorphic antiderivative) Let $U \subseteq \mathbb{C}$ be either an open rectangle or an open disc. Let $P \in U$ be fixed. Suppose that F is continuous on U and holomorphic on $U \setminus \{P\}$. Then there is a holomorphic H on U such that $\partial H / \partial z = F$.

Note that since H is holomorphic, by [Theorem 2.2.1](#), we can write $H' = F$.

Lemma 2.4.1 Let γ be the boundary of a disc $D(z_0, r)$ in the complex plane, equipped with the counterclockwise orientation. Let z be a point inside the circle $\partial D(z_0, r)$. Then

$$\frac{1}{2\pi i} \oint_{\gamma} \frac{1}{\zeta - z} d\zeta = 1$$

The proof involves considering the function

$$I(z) := \oint_{\gamma} \frac{1}{\zeta - z} d\zeta$$

and showing that $I(z)$ is independent of z , and that $I(z_0) = 2\pi i$.

Theorem 2.4.2 (Cauchy integral formula) Suppose that $U \subseteq \mathbb{C}$ is open and that f is a holomorphic function on U . Let $z_0 \in U$ and let $r > 0$ such that $\overline{D}(z_0, r) \subseteq U$. Let $\gamma : [0, 1] \rightarrow \mathbb{C}$ be the C^1 curve $\gamma(t) = z_0 + r \cos(2\pi t) + ir \sin(2\pi t)$. Then, for each $z \in D(z_0, r)$,

$$f(z) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$$

The converse of this theorem is true too: if f is given by the Cauchy integral formula, then f is holomorphic.

Example 2.4.2a (Examples with Cauchy integral formula) Here's some ground-truth computations to get started. (Almost all problems in MATH 466 can be re-routed back to these)

$$\oint_{\gamma} \zeta^k d\zeta = \begin{cases} 0 & \text{if } k \neq -1 \\ 2\pi i & \text{if } k = -1 \end{cases} \quad (k \in \mathbb{Z})$$

Theorem 2.4.3 (Cauchy integral theorem) If f is a holomorphic function on an open disc $U \subseteq \mathbb{C}$, and if $\gamma : [a, b] \rightarrow U$ is a C^1 curve in U with $\gamma(a) = \gamma(b)$, then

$$\oint_{\gamma} f(z) dz = 0$$

Note that this implies that the **Cauchy integral formula** gives a zero whenever z does not lie in the contour γ , since the integrand is holomorphic. (Integrand is holomorphic because numerator is assumed to be holomorphic, and the denominator is never zero.)

Proof. By **Theorem 1.5.3**, there is a holomorphic function $G : U \rightarrow \mathbb{C}$ with $G' = f$ on U . Since $\gamma(a) = \gamma(b)$, we have that

$$0 = G(\gamma(b)) - G(\gamma(a))$$

By **Proposition 2.1.6**, this equals

$$\oint_{\gamma} G'(z) dz = \oint_{\gamma} f(z) dz$$

(Reminder that since G is holomorphic, $G' = \frac{\partial G}{\partial z}$ by **Theorem 2.2.1**) □

Definition 2.6.1 (Piecewise C^1 curve) A piecewise C^1 curve $\gamma : [a, b] \rightarrow \mathbb{C}$ is a continuous function such that there exists a finite set of numbers $a_1 \leq a_2 \leq \dots \leq a_k$ satisfying $a_1 = a$ and $a_k = b$, and with the property that for every $i \leq j \leq k-1$, $\gamma|_{[a_j, a_{j+1}]}$ is a C^1 curve.

γ is a piecewise C^1 curve in an open set U if $\gamma([a, b]) \subseteq U$.

Note that while joining $C^1(\mathbb{R})$ curves may not lead to a piecewise $C^1(\mathbb{R})$ curve, doing it in \mathbb{C} somehow works.

Definition 2.6.2 (Integrating over a piecewise C^1 curve) If $U \subseteq \mathbb{C}$ is open and $\gamma : [a, b] \rightarrow U$ is a piecewise C^1 curve in U and if $f : U \rightarrow \mathbb{C}$ is a continuous function on U , then

$$\oint_{\gamma} f(z) dz := \sum_{j=1}^k \oint_{\gamma|_{[a_j, a_{j+1}]}} f(z) dz$$

where a_1, a_2, \dots, a_k are as in [Definition 2.6.1](#).

Lemma 2.6.3 Let $U \subseteq \mathbb{C}$ be open. Let $\gamma : [a, b] \rightarrow U$ be a piecewise C^1 curve. Let $\phi : [c, d] \rightarrow [a, b]$ be a piecewise C^1 strictly monotone increasing function with $\phi(c) = a$ and $\phi(d) = b$. Let $f : U \rightarrow \mathbb{C}$ be a continuous function on U . Then the function $\gamma \circ \phi : [c, d] \rightarrow U$ is a piecewise C^1 curve and

$$\oint_{\gamma} f(z) dz = \oint_{\gamma \circ \phi} f(z) dz$$

(Really, $\{\gamma(t) \mid t \in [a, b]\} = \{(\gamma \circ \phi)(s) \mid s \in [c, d]\}$, and there are no added crossovers on the parameterization of $\gamma \circ \phi$ because ϕ is strictly monotone increasing.)

Lemma 2.6.4 Let $U \subseteq \mathbb{C}$ be open, $f : U \rightarrow \mathbb{C}$ a holomorphic function and $\gamma : [a, b] \rightarrow U$ a piecewise C^1 curve. Then

$$f(\gamma(b)) - f(\gamma(a)) = \oint_{\gamma} f'(z) dz$$

(This is really just [Proposition 2.1.6](#) restated with a piecewise C^1 version of γ)

Proposition 2.6.5 If $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ is a holomorphic function, and if γ_r describes the circle of radius r around 0, traversed once around counterclockwise, then, for any two positive numbers $r_1 < r_2$,

$$\oint_{\gamma_{r_1}} f(z) dz = \oint_{\gamma_{r_2}} f(z) dz$$

Proposition 2.6.6 Let $0 < r < R < \infty$ and define the annulus

$\mathcal{A} := \{z \in \mathbb{C} : r < |z| < R\}$. Let $f : \mathcal{A} \rightarrow \mathbb{C}$ be a holomorphic function. If $r < r_1 < r_2 < R$ and if for each j the curve γ_{r_j} describes the circle of radius r_j around 0, traversed once counterclockwise, then we have

$$\oint_{\gamma_{r_1}} f(z) dz = \oint_{\gamma_{r_2}} f(z) dz$$

(On this annulus (donut), integrating a holomorphic f along any two circles centered at zero will yield the same value.)

Theorem 2.6.7 (Cauchy integral formula and theorem: general form) Let $U \subseteq \mathbb{C}$ be open. Let $f : U \rightarrow \mathbb{C}$ be holomorphic. Then

$$\oint_{\gamma} f(z) dz = 0$$

for any piecewise C^1 closed curve γ in U that can be deformed in U through closed curves to a closed curve lying entirely in a disc contained in U .

In addition, suppose that $\overline{D}(z, r) \subseteq U$. Then

$$\frac{1}{2\pi i} \oint_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = f(z)$$

for any piecewise C^1 closed curve γ in $U \setminus \{z\}$ that can be continuously deformed in $U \setminus \{z\}$ to $\partial D(z, r)$ equipped with counterclockwise orientation.

Theorem 3.1.1 (Analyticity of holomorphic functions) Let $U \subseteq \mathbb{C}$ be open and let f be a holomorphic on U . Then $f \in C^\infty(U)$. Moreover, if $\overline{D}(P, r) \subseteq U$ and $z \in D(P, r)$, then

$$\left(\frac{\partial}{\partial z} \right)^k f(z) = \frac{k!}{2\pi i} \oint_{|\zeta - P|=r} \frac{f(\zeta)}{(\zeta - z)^{k+1}} d\zeta$$

for all $k \in \mathbb{N}_0$.

Corollary 3.1.2 (Derivative of a holomorphic function is holomorphic) Let $U \subseteq \mathbb{C}$ be open. If $f : U \rightarrow \mathbb{C}$ is holomorphic, then $f' : U \rightarrow \mathbb{C}$ is holomorphic.

Theorem 3.1.3 If ϕ is a continuous function on $\{\zeta : |\zeta - P| = r\}$, then the function f given by

$$f(z) := \frac{1}{2\pi i} \oint_{|\zeta - P|=r} \frac{\phi(\zeta)}{\zeta - z} d\zeta$$

is defined and holomorphic on $D(P, r)$.

This theorem induces a very strong way to create a holomorphic function. Instead of differentiability, we only need a continuous ϕ to build a holomorphic f .

Theorem 3.1.4 (Morera's Theorem) Let $U \subseteq \mathbb{C}$ be open. Let $f : U \rightarrow \mathbb{C}$ be a continuous function on a connected open subset U of \mathbb{C} . Suppose that for every closed, piecewise C^1 curve $\gamma : [0, 1] \rightarrow U$, with $\gamma(0) = \gamma(1)$, we have

$$\oint_{\gamma} f(\zeta) d\zeta = 0$$

Then f is holomorphic on U .

Lemma 3.2.1 The sequence $\{a_k \in \mathbb{C}\}$ converges to a limit if and only if for each $\varepsilon > 0$ there is an N_0 such that $j, k \geq N_0$ implies that $|a_j - a_k| < \varepsilon$.

Definition 3.2.2 (Complex power series) Let $P \in \mathbb{C}$ be fixed. A *complex power series* (centered at P) is an expression of the form

$$\sum_{k=0}^{\infty} a_k (z - P)^k$$

where a_k for $k = 0, \dots, \infty$ are complex constants.

Note that this power series expansion is only a formal expression. It may or may not converge.

A *necessary* condition for $\sum a_k (z - P)^k$ to converge is that $a_k (z - P)^k \rightarrow 0$.

Lemma 3.2.3 (Abel's Theorem) If $\sum_{k=0}^{\infty} a_k (z - P)^k$ converges at some z , then the series converges at each $w \in D(P, r)$, where $r = |z - P|$.

Definition 3.2.4 (Radius of convergence of power series) Let $\sum_{k=0}^{\infty} a_k (z - P)^k$ be a power series. Then

$$r := \sup \left\{ |w - P| \mid \sum_{k=0}^{\infty} a_k (w - P)^k \text{ converges} \right\}$$

is called the *radius of convergence* of the power series. We will call $D(P, r)$ the disc of convergence.

Lemma 3.2.5 If $\sum_{k=0}^{\infty} a_k (z - P)^k$ is a power series with radius of convergence r , then the series converges for each $w \in D(P, r)$ and diverges for each w such that $|w - P| > r$.

Note that the convergence or divergence question for $|w - P| = r$ is left open.

Lemma 3.2.6 (Computing radius of convergence) Using

$$\ell : \limsup_{k \rightarrow +\infty} |a_k|^{1/k},$$

the radius of convergence r of the power series $\sum_{k=0}^{\infty} a_k (z - P)^k$ is given by

$$r = \begin{cases} 1/\ell & \text{if } \ell > 0 \\ +\infty & \text{if } \ell = 0 \end{cases}$$

Definition 3.2.7 (Uniform convergence of complex functions) A series $\sum_{k=0}^{\infty} f_k(z)$ of functions $f_k(z)$ converges uniformly on a set E to the function $g(z)$ if for each $\varepsilon > 0$ there is an N_0 such that if $N \geq N_0$, then

$$\left| g(z) - \sum_{k=0}^N f_k(z) \right| < \varepsilon \quad \forall z \in E$$

Definition 3.2.8 (Uniformly Cauchy series) Let $\sum_{k=0}^{\infty} f_k(z)$ be a series of functions on a set E . The series is said to be *uniformly Cauchy* if, for any $\varepsilon > 0$, there is a positive integer N_0 such that if $m \geq j \geq N_0$, then

$$\left| \sum_{k=j}^m f_k(z) \right| < \varepsilon \quad \forall z \in E$$

Proposition 3.2.9 Let $\sum_{k=0}^{\infty} a_k(z-P)^k$ be a power series with radius of convergence r . Then, for any number R with $0 \leq R < r$, the series $\sum_{k=0}^{\infty} |a_k(z-P)^k|$ **converges uniformly** on $\overline{D}(P, R)$.

In particular, the series $\sum_{k=0}^{+\infty} a_k(z-P)^k$ converges uniformly and absolutely on $\overline{D}(P, R)$.

Lemma 3.2.10 If a power series

$$\sum_{j=0}^{\infty} a_j(z-P)^j \quad (*)$$

has a radius of convergence $r > 0$, then the series defines a C^∞ function $f(z)$ on $D(P, r)$. The function f is holomorphic on $D(P, r)$. The series obtained by termwise differentiation k times of $(*)$,

$$\sum_{j=k}^{\infty} [j(j-1)\dots(j-k+1)] a_j(z-P)^{j-k}$$

converges on $D(P, r)$, and its sum is $[\partial/\partial z]^k f(z)$ for each $z \in D(P, r)$.

Proposition 3.2.11 If both series $\sum_{j=0}^{\infty} a_j(z-P)^j$ and $\sum_{j=0}^{\infty} b_j(z-P)^j$ converge on a disc $D(P, r)$, $r > 0$, and if

$$\sum_{j=0}^{\infty} a_j(z-P)^j = \sum_{j=0}^{\infty} b_j(z-P)^j$$

on $D(P, r)$, then $a_j = b_j$ for every j .

Theorem 3.3.1 Let $U \subseteq \mathbb{C}$ be open and let f be holomorphic on U . Let $P \in U$ and suppose that $D(P, r) \subseteq U$. Then the complex power series

$$\sum_{k=0}^{\infty} \frac{(\partial^k f / \partial z^k)(P)}{k!} (z-P)^k$$

has radius of convergence at least r . It converges to $f(z)$ on $D(P, r)$.

Theorem 3.4.1 (The Cauchy estimates) Let $U \subseteq \mathbb{C}$ be open and $f : U \rightarrow \mathbb{C}$ be holomorphic. Let $P \in U$ and assume that the closed disc $\overline{D}(P, r)$, $r > 0$, is contained in U . Set

$$M := \sup_{z \in \overline{D}(P, r)} |f(z)|.$$

Then for $k = 1, 2, 3, \dots$ we have

$$\left| \frac{\partial^k f}{\partial z^k}(P) \right| \leq \frac{Mk!}{r^k}$$

Lemma 3.4.2 Let $U \subseteq \mathbb{C}$ be open and connected and $f : U \rightarrow \mathbb{C}$ be holomorphic. If $\partial f / \partial \bar{z} = 0$ on U , then f is constant on U .

Proof. Since f is holomorphic, $\partial f / \partial \bar{z} = 0$. But we have assumed that $\partial f / \partial z = 0$. Thus $\partial f / \partial x = \partial f / \partial y = 0$. So f is constant. □

Theorem 3.4.3 (Liouville's Theorem) A bounded **entire** function is constant.

Proof. Let f be entire and assume that $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Fix a $P \in \mathbb{C}$ and let $r > 0$. We apply the **Cauchy estimate** for $k = 1$ on $\overline{D}(P, r)$. The result is

$$\left| \frac{\partial f}{\partial z}(P) \right| \leq \frac{M}{r}$$

Since this inequality holds for all $r > 0$, we can blow it up to $+\infty$ and conclude that

$$\frac{\partial f}{\partial z}(P) = 0$$

But since P is arbitrary, we conclude that

$$\frac{\partial f}{\partial z} \equiv 0$$

By **Lemma 3.4.2**, the proof is complete. □

Theorem 3.4.4 If $f : \mathbb{C} \rightarrow \mathbb{C}$ is an entire function and if for some real number C and some positive integer k it holds that

$$|f(z)| \leq C|z|^k$$

for all $z \in \mathbb{C}$ with $|z| > 1$, then f is a polynomial in z of degree at most k .

Theorem 3.4.5 Let $p(z)$ be a non-constant (holomorphic) polynomial. Then p has a root. That is, there exists an $\alpha \in \mathbb{C}$ such that $p(\alpha) = 0$.

This is in fact the fundamental theorem of algebra, and one of the most elegant applications of **Liouville's Theorem**.

Proof. Suppose there isn't an $\alpha \in \mathbb{C}$ such that $p(\alpha) = 0$. Then

$$g(z) := \frac{1}{p(z)}$$

is entire. Notice that as $|z| \rightarrow \infty$, $|p(z)| \rightarrow +\infty$. Thus $1/|p(z)| \rightarrow 0$ as $|z| \rightarrow \infty$ and hence g is bounded. By Liouville's Theorem, g is constant; hence p is constant. Contradiction! □

Corollary 3.4.6 If $p(z)$ is a holomorphic polynomial of degree k , then there are k complex numbers $\alpha_1, \dots, \alpha_k$ (not necessarily distinct) and a non-zero constant C such that

$$p(z) = C(z - \alpha_1) \dots (z - \alpha_k)$$

Theorem 3.5.1 Let $U \subseteq \mathbb{C}$ be an open set. Let $f_j : U \rightarrow \mathbb{C}$, $j = 1, 2, 3 \dots$ be a sequence of holomorphic functions. Suppose that there is a function $f : U \rightarrow \mathbb{C}$ such that, for each compact subset E of U , the sequence $f_j|_E$ converges uniformly to $f|_E$. Then f is holomorphic on U . (In particular, $f \in C^\infty(U)$)

Corollary 3.5.2 If f_j, f, U are as defined in [Theorem 3.5.1](#), then for any integer $k \in \{0, 1, 2, \dots\}$ we have

$$\left(\frac{\partial}{\partial z}\right)^k f_j(z) \rightarrow \left(\frac{\partial}{\partial z}\right)^k f(z)$$

uniformly on compact sets.

Theorem 3.6.1 Let $U \subseteq \mathbb{C}$ be a connected open set and let $f : U \rightarrow \mathbb{C}$ be holomorphic. Let $\mathbf{Z} := \{z \in U \mid f(z) = 0\}$. If there are a $z_0 \in \mathbf{Z}$ and a sequence $\{z_j\} \subseteq \mathbf{Z} \setminus \{z_0\}$ such that $z_j \rightarrow z_0$, then $f \equiv 0$.

Corollary 3.6.2 Let $U \subseteq \mathbb{C}$ be connected and open, and $D(P, r) \subseteq U$. If f is holomorphic on U and $f|_{D(P, r)} \equiv 0$, then $f \equiv 0$ on U .

Note the strength of this statement. As long as f is holomorphic, if it's zero on just a tiny $D(P, r)$, then it is zero on the entire domain.

Corollary 3.6.3 Let $U \subseteq \mathbb{C}$ be connected and open. Let f, g be holomorphic on U . If $\{z \in U \mid f(z) = g(z)\}$ has an [accumulation point](#) in U , then $f \equiv g$.

Corollary 3.6.4 Let $U \subseteq \mathbb{C}$ be connected and open and let f, g be holomorphic on U . If $f \cdot g \equiv 0$ on U , then either $f \equiv 0$ on U or $g \equiv 0$ on U .

Corollary 3.6.5 Let $U \subseteq \mathbb{C}$ be connected and open and let f be holomorphic on U . If there is a $P \in U$ such that

$$\left(\frac{\partial}{\partial z}\right)^j f(P) = 0$$

for every j , then $f \equiv 0$.

Corollary 3.6.6 If f and g are entire holomorphic functions and if $f(x) = g(x)$ for all $x \in \mathbb{R} \subseteq \mathbb{C}$, then $f \equiv g$.

Definition 4.1.0 (Types of singularities) Let $U \subseteq \mathbb{C}$ be open and $P \in U$. Suppose that $f : U \setminus \{P\} \rightarrow \mathbb{C}$ is holomorphic.

There are three possibilities for the behavior of f near P :

- (i) (*Removable singularity*) $|f(z)|$ is bounded on $D(P, r) \setminus \{P\}$ for some $r > 0$ with $D(P, r) \subseteq U$.
- (ii) (*Pole*) $\lim_{z \rightarrow P} |f(z)| = +\infty$.
- (iii) (*Essential singularity*) Neither (i) nor (ii) applies.

Theorem 4.1.1 (The Riemann removable singularities theorem) Let $f : D(P, r) \setminus \{P\} \rightarrow \mathbb{C}$ be holomorphic and bounded. Then

- (a) $\lim_{z \rightarrow P} f(z)$ exists

(b) the function $\hat{f} : D(P, r) \rightarrow \mathbb{C}$ defined by

$$\hat{f}(z) = \begin{cases} f(z) & \text{if } z \neq P \\ \lim_{\zeta \rightarrow P} f(\zeta) & \text{if } z = P \end{cases}$$

is holomorphic

Notice that, a priori, it is not even clear that $\lim_{z \rightarrow P} f(z)$ exists, or, even if it does, that the function \hat{f} has any regularity at P beyond just continuity.

Theorem 4.1.4 (Casorati-Weierstrass) If $f : D(P, r_0) \setminus \{P\} \rightarrow \mathbb{C}$ is holomorphic and P is an essential singularity of f , then $f(D(P, r) \setminus \{P\})$ is **dense** in \mathbb{C} for any $0 < r < r_0$.

Definition 4.2.0 (Laurent series) A Laurent series on $D(P, r)$ is a (formal) expression of the form

$$\sum_{j=-\infty}^{+\infty} a_j (z - P)^j$$

where j are integer indices.

We say that the infinite series $\sum_{j=-\infty}^{+\infty} \alpha_j$ converges if $\sum_{j=0}^{+\infty} \alpha_j$ and $\sum_{j=1}^{+\infty} \alpha_{-j}$ converge. In this case, we set

$$\sum_{j=-\infty}^{+\infty} \alpha_j = \left(\sum_{j=0}^{+\infty} \alpha_j \right) + \left(\sum_{j=1}^{+\infty} \alpha_{-j} \right)$$

This doubly infinite series converges to a complex number σ if and only if for each $\varepsilon > 0$ there is an $N > 0$ such that if $\ell \geq N$ and $k \geq N$, then $|\sigma - \sum_{j=-k}^{\ell} \alpha_j| < \varepsilon$.

It is important to realize that ℓ and k are independent here. In particular, the existence of the limit $\lim_{k \rightarrow +\infty} \sum_{j=-k}^{+k} \alpha_j$ does not imply in general that $\sum_{j=-\infty}^{+\infty} \alpha_j$ converges.

Lemma 4.2.1 This is the analogue for Laurent series for **Lemma 3.2.3**.

If $\sum_{j=-\infty}^{+\infty} a_j (z - P)^j$ converges at $z_1 \neq P$ and at $z_2 \neq P$ and if $|z_1 - P| < |z_2 - P|$, then the series converges for all z with $|z_1 - P| < |z - P| < |z_2 - P|$.

Lemma 4.2.2 This is the analogue for Laurent series for **Lemma 3.2.5**. Let

$$\sum_{j=-\infty}^{+\infty} a_j (z - P)^j$$

converge at (at least) one point z_0 . There are unique non-negative numbers r_1 and r_2 (r_1 or r_2 may be $+\infty$) such that the series converges absolutely for all z with

$$r_1 < |z - P| < r_2$$

and diverges for all z with

$$|z - P| < r_1 \quad \text{or} \quad r_2 < |z - P|$$

Proposition 4.2.4 (Uniqueness of Laurent expansion) Let $0 \leq r_1 < r_2 \leq \infty$. If the Laurent series $\sum_{j=-\infty}^{+\infty} a_j(z-P)^j$ converges on $D(P, r_2) \setminus \overline{D}(P, r_1)$ to a function f , then, for any $r \in (r_1, r_2)$ and each $j \in \mathbb{Z}$, we have

$$a_j = \frac{1}{2\pi i} \oint_{|\zeta-P|=r} \frac{f(\zeta)}{(\zeta-P)^{j+1}} d\zeta$$

In particular, the a_j 's are uniquely determined by f .

Theorem 4.3.1 (The Cauchy integral formula for an annulus) Suppose that $0 \leq r_1 < r_2 \leq +\infty$ and that $f : D(P, r_2) \setminus \overline{D}(P, r_1) \rightarrow \mathbb{C}$ is holomorphic. Then, for each s_1, s_2 such that $r_1 < s_1 < s_2 < r_2$ and each $z \in D(P, s_2) \setminus \overline{D}(P, s_1)$, it holds that

$$f(z) = \frac{1}{2\pi i} \oint_{|\zeta-P|=s_2} \frac{f(\zeta)}{\zeta-z} d\zeta - \frac{1}{2\pi i} \oint_{|\zeta-P|=s_1} \frac{f(\zeta)}{\zeta-z} d\zeta$$

Theorem 4.3.2 (The existence of Laurent expansions) If $0 \leq r_1 < r_2 \leq +\infty$ and $f : D(P, r_2) \setminus \overline{D}(P, r_1) \rightarrow \mathbb{C}$ is holomorphic, then there exist complex numbers a_j such that

$$\sum_{j=-\infty}^{+\infty} a_j(z-P)^j$$

converges on $D(P, r_2) \setminus \overline{D}(P, r_1)$ to f . If $r_1 < s_1 < s_2 < r_2$, then the series converges absolutely and uniformly on $D(P, s_2) \setminus \overline{D}(P, s_1)$.

Proposition 4.3.3 (Laurent expansion of holomorphic functions) If $f : D(P, r) \setminus \{P\} \rightarrow \mathbb{C}$ is holomorphic, then f has a unique Laurent series expansion.

$$f(z) = \sum_{j=-\infty}^{+\infty} a_j(z-P)^j$$

which converges absolutely for $z \in D(P, r) \setminus \{P\}$. The convergence is uniform on compact subsets of $D(P, r) \setminus \{P\}$. The coefficients are given by

$$a_j = \frac{1}{2\pi i} \oint_{\partial D(P, s)} \frac{f(\zeta)}{(\zeta-P)^{j+1}} d\zeta$$

for any $0 < s < r$.

There are three mutually exclusive possibilities for the Laurent series of this proposition:

- (i) $a_j = 0$ for all $j < 0$;
- (ii) for some $k > 0$, $a_j = 0$ for all $-\infty < j < -k$
- (iii) neither (i) nor (ii) applies

These three cases correspond exactly to the **three types of isolated singularities**: (i) $\iff P$ is a removable singularity; (ii) $\iff P$ is a pole; (iii) $\iff P$ is an essential singularity.

Proposition 4.4.1 Let f be holomorphic on $D(P, r) \setminus \{P\}$ and suppose that f has a pole of order k at P . Then the Laurent series coefficients a_j of f expanded about P , for $j = -k, -k + 1, -k + 2, \dots$, are given by the formula

$$a_j = \frac{1}{(k+j)!} \left(\frac{\partial}{\partial z} \right)^{k+j} \left((z-P)^k \cdot f \right) \Big|_{z=P}$$

Definition 4.5.1 (Holomorphically simply connected (HSC)) An open set $U \subseteq \mathbb{C}$ is holomorphically simply connected if U is connected and if, for each holomorphic function $f : U \rightarrow \mathbb{C}$, there is a holomorphic function $F : U \rightarrow \mathbb{C}$ such that $F' \equiv f$.

Lemma 4.5.2 A connected open set U is holomorphically simply connected if and only if for each holomorphic function $f : U \rightarrow \mathbb{C}$ and each piecewise C^1 closed curve γ in U ,

$$\oint_{\gamma} f(z) dz = 0.$$

Definition 4.5.3a (Residue of a function at a point) The **residue** of a function f at point P is denoted by $\text{Res}_f(P)$, and is the coefficient of $(z-P)^{-1}$ in the **Laurent expansion** of f about P .

In particular, if f is holomorphic, then $\text{Res}_f(P)$ is given by

$$\text{Res}_f(P) = \frac{1}{2\pi i} \oint_{\gamma} f(\zeta) d\zeta$$

where γ is a counterclockwise simply closed curve around P and not including any other singularities inside the curve.

Theorem 4.5.3 (Residue theorem) Suppose that $U \subseteq \mathbb{C}$ is a **HSC** open set, and that P_1, \dots, P_n are distinct points of U . Suppose that $f : U \setminus \{P_1, \dots, P_n\} \rightarrow \mathbb{C}$ is a holomorphic function and γ is a closed, piecewise C^1 curve in $U \setminus \{P_1, \dots, P_n\}$.

Set R_j to be the coefficient of $(z-P_j)^{-1}$ in the Laurent expansion of f about P_j .

Then

$$\oint_{\gamma} f(z) dz = \sum_{j=1}^n R_j \cdot \left(\oint_{\gamma} \frac{1}{\zeta - P_j} d\zeta \right)$$

Using the notation of **Res_f** and **Ind_γ**, we can rewrite this as

$$\oint_{\gamma} f(z) dz = 2\pi i \sum_{j=1}^n \text{Res}_f(P_j) \cdot \text{Ind}_{\gamma}(P_j)$$

Definition 4.5.4 (Index of a curve) Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a piecewise C^1 closed curve. Suppose $P \notin \gamma([a, b])$. Then the **index** of γ with respect to P , is defined as

$$\text{Ind}_{\gamma}(P) := \frac{1}{2\pi i} \oint_{\gamma} \frac{1}{\zeta - P} d\zeta$$

The index is also sometimes called the “winding number of γ about P ”. As we will see later, $\text{Ind}_{\gamma}(P)$ coincides with the number of times γ winds about P , counting orientation.

Lemma 4.5.5 (Index of a curve is an integer) Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a piecewise C^1 closed curve. Suppose P is a point not on the image of that curve, then

$$\frac{1}{2\pi i} \oint_{\gamma} \frac{1}{\zeta - P} d\zeta \equiv \frac{1}{2\pi i} \int_a^b \frac{\gamma'(t)}{\gamma(t) - P} dt$$

is an integer.

Proposition 4.5.6 Let f be a function with a pole of order k at P . Then

$$\operatorname{Res}_f(P) = \frac{1}{(k-1)!} \left(\frac{\partial}{\partial z} \right)^{k-1} \left((z-P)^k f(z) \right) \Big|_{z=P}$$

Proof. This is the case $j = -1$ of [Proposition 4.4.1](#). □

Nonlinear Optimization, Part I: Unconstrained Optimization

Definition 1.1.2 (The argmin set) The set which minimizes values of f over a domain X is denoted by

$$\operatorname{argmin}_{x \in X} f := \left\{ x \in X \mid f(x) = \inf_{x \in X} f \right\}$$

Definition 1.1.3 (Local vs. global minima) Let $X \subset \mathbb{R}^n$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Then $\bar{x} \in X$ is called a

- global minimizer of f over X if $\bar{x} \in \operatorname{argmin}_X f$, i.e. $\forall x \in X : f(\bar{x}) \leq f(x)$
- local minimizer of f over X if $\exists \varepsilon > 0$ such that $\forall x \in X \cap B_\varepsilon(\bar{x}) : f(\bar{x}) \leq f(x)$

For strict global/local minimizers, the above conditions hold with strict inequality.

Definition 1.1.4 (Level sets and Lower level sets) For $f : \mathbb{R}^n \rightarrow \mathbb{R}$ the level set for the level $c \in \mathbb{R}$ is given by

$$f^{-1}(\{c\}) = \{x \in \mathbb{R}^n \mid f(x) = c\}$$

The lower level set (or *sublevel* set) of f to the level $c \in \mathbb{R}$ is

$$\operatorname{lev}_c f := f^{-1}((-\infty, c]) = \{x \in \mathbb{R}^n \mid f(x) \leq c\}$$

Proposition 1.1.5 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous. If $\exists c \in \mathbb{R}$ such that $\operatorname{lev}_c f$ is non-empty and bounded then f takes its minimum over \mathbb{R}^n .

Definition 1.2.1 (Convex sets) A set $C \subset \mathbb{R}^n$ is called convex if

$$\lambda x + (1 - \lambda)y \in C \quad \forall (x, y \in C, \lambda \in (0, 1))$$

or simply a set which contains all connecting lines of points from the set.

Definition 1.2.3 (Convex functions) Let $C \subset \mathbb{R}^n$ be convex. Let $\lambda \in (0, 1)$ and $x, y \in C$ and let

Then $f : C \rightarrow \mathbb{R}$ is said to be

- convex on C if

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$$

- strictly convex on C if

$$f(\lambda x + (1 - \lambda)y) < \lambda f(x) + (1 - \lambda)f(y)$$

- strongly convex on C if $\exists \mu > 0$ such that

$$f(\lambda x + (1 - \lambda)y) + \frac{\mu}{2} \lambda(1 - \lambda) \|x - y\|^2 \leq \lambda f(x) + (1 - \lambda)f(y)$$

This has parallels [here](#) and [here](#).

Example 1.2.5 (Convex functions)

- (a) $\exp : \mathbb{R} \rightarrow \mathbb{R}$ and $-\log : (0, \infty) \rightarrow \mathbb{R}$ are convex.
(b) (*Affine functions*) $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ of the form

$$f(x) = Ax - b \quad (A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m)$$

is called **affine** (linear). All affine functions, hence all linear functions ($b = 0$) $\mathbb{R}^n \rightarrow \mathbb{R}$ are convex.

- (c) (*Norms*) Any norm $\|\cdot\|$ on \mathbb{R}^n is convex.

Proposition 1.2.6 (Convexity preserving operations)

- (Positive combinations) For $i = 1, \dots, n$ let $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and $\lambda_i \geq 0$. Then $\sum_{i=1}^n \lambda_i f_i$ is convex.
- (Composition with affine mapping) $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and $g : \mathbb{R}^m \rightarrow \mathbb{R}^n$ affine. Then $f \circ g$ is convex.

Theorem 1.2.7 (Taylor's Theorem, $k = 2$) Let $D \subset \mathbb{R}^n$ be open, let $f : D \rightarrow \mathbb{R}$ be twice continuously differentiable, and $x, y \in D$ such that $[x, y] \subset D$. Then there exists $\eta \in [x, y]$ such that

$$f(y) - f(x) = \nabla f(x)^T (y - x) + \frac{1}{2} (y - x)^T \nabla^2 f(\eta) (y - x)$$

Theorem 1.2.8 (Schwarz's Theorem) Let $D \subset \mathbb{R}^n$ be open and $f : D \rightarrow \mathbb{R}$ be twice continuously differentiable at $x \in D$. Then $\nabla^2 f(x)$ is symmetric.

Definition 2.1.1 (Directional derivative) Let $D \subset \mathbb{R}^n$ be open. $f : D \rightarrow \mathbb{R}$ is directionally differentiable at $\bar{x} \in \mathbb{R}^n$ in the direction $d \in \mathbb{R}^n$ if

$$\lim_{t \downarrow 0} \frac{f(\bar{x} + td) - f(\bar{x})}{t}$$

exists. This limit is denoted by $f'(x; d)$ and is called the directional derivative of f at \bar{x} in the direction d .

If f is directionally differentiable at \bar{x} in every direction $d \in \mathbb{R}^n$, we call f directionally differentiable at \bar{x} .

If f is directionally differentiable at every $\bar{x} \in \mathbb{R}^n$, we call it directionally differentiable.

Lemma 2.1.2 (Directional derivative and gradient) Let $D \subset \mathbb{R}^n$ be open and $f : D \rightarrow \mathbb{R}$ differentiable at $x \in D$. Then f is directionally differentiable at x with

$$f'(x; d) = \nabla f(x)^T d \quad \forall (d \in \mathbb{R}^n)$$

Where $f'(x; d)$ is the **directional derivative** of f at x in the direction d .

Lemma 2.1.4 (Basic optimality condition) Let $X \subset \mathbb{R}^n$ be open and $f : X \rightarrow \mathbb{R}$. If \bar{x} is a local minimizer of f over X and f is **directionally differentiable** at \bar{x} then

$$f'(x; d) \geq 0 \quad \forall (d \in \mathbb{R}^n)$$

Theorem 2.1.5 (Fermat's rule) Let $X \subset \mathbb{R}^n$ be open and $f : X \rightarrow \mathbb{R}$ differentiable at $\bar{x} \in X$. If \bar{x} is a local minimizer (or maximizer) of f over X then $\nabla f(\bar{x}) = 0$.

Theorem 2.1.6 (Second-order necessary condition) Let $X \subset \mathbb{R}^n$ be open and $f : X \rightarrow \mathbb{R}$ twice continuously differentiable. If \bar{x} is a local minimizer (maximizer) of f over X then $\nabla^2 f(\bar{x})$ is positive (negative) semidefinite.

Proof. We only prove the case in which \bar{x} is a local minimizer. The maximum case follows from it by substituting f for $-f$.

Assume on the contrary that $\nabla^2 f(\bar{x})$ were not positive semidefinite. Then there exists $d \in \mathbb{R}^n$ such that $d^T \nabla^2 f(\bar{x}) d < 0$. By **Taylor's Theorem**, for $t > 0$ there exists $\eta_t \in [\bar{x}, \bar{x} + td]$ such that

$$f(\bar{x} + td) = f(\bar{x}) + t \nabla f(\bar{x})^T d + \frac{t^2}{2} d^T \nabla^2 f(\eta_t) d$$

But since \bar{x} is a local minimizer, by **Fermat's rule** we have $\nabla f(\bar{x}) = 0$ and hence

$$f(\bar{x} + td) = f(\bar{x}) + \frac{t^2}{2} d^T \nabla^2 f(\eta_t) d$$

As $t \downarrow 0$, $\eta_t \rightarrow \bar{x}$ and hence for some $t > 0$ sufficiently small, we have

$$\frac{t^2}{2} d^T \nabla^2 f(\eta_t) d < 0$$

as $\nabla^2 f$ is continuous by assumption. This yields $f(\bar{x} + td) < f(\bar{x})$ for all $t > 0$ sufficiently small, which contradicts the fact that \bar{x} is a local minimizer of f over X . Hence, $\nabla^2 f(\bar{x})$ must be positive semidefinite. \square

Lemma 2.1.7 Let $X \subset \mathbb{R}^n$ be open and $f : X \rightarrow \mathbb{R}$ twice continuously differentiable. If $\bar{x} \in \mathbb{R}^n$ is such that $\nabla^2 f(\bar{x})$ positive definite then $\exists \mu, \varepsilon > 0$ such that $B_\varepsilon(\bar{x}) \subset X$ and

$$d^T \nabla^2 f(x) d \geq \mu \|d\|_2^2 \quad \forall (d \in \mathbb{R}^n, x \in B_\varepsilon(\bar{x}))$$

Theorem 2.1.8 (Sufficient optimality condition) Let $X \subset \mathbb{R}^n$ be open and $f : X \rightarrow \mathbb{R}$ twice continuously differentiable. Moreover, let \bar{x} be a stationary point of f such that $\nabla^2 f(\bar{x})$ is positive definite. Then \bar{x} is a strict local minimizer of f .

Theorem 2.2.1 (First-order characterizations) Let $C \subset \mathbb{R}^n$ be open and convex and let $f : C \rightarrow \mathbb{R}$ be differentiable on C . Then the following hold for all $x, \bar{x} \in C$:

(a) f is convex on C iff

$$f(x) \geq f(\bar{x}) + \nabla f(\bar{x})^T (x - \bar{x}) \quad (*)$$

(b) f is strictly convex on C iff $(*)$ holds with strict inequality whenever $x \neq \bar{x}$.

(c) f is strongly convex with modulus $\mu > 0$ on C iff

$$f(x) \geq \left[f(\bar{x}) + \nabla f(\bar{x})^T (x - \bar{x}) \right] + \frac{\mu}{2} \|x - \bar{x}\|^2$$

This has parallels [here](#) and [here](#).

Corollary 2.2.2 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Then the following hold:

(a) (*Affine minorization*) There exists an [affine](#) function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ which minorizes f everywhere, i.e.

$$g(x) \leq f(x) \quad (x \in \mathbb{R}^n)$$

(b) If f is strongly convex then it is strictly convex and coercive (level-bounded).

Corollary 2.2.3 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Then the following are equivalent:

- (i) \bar{x} is a global minimizer of f , i.e. $\bar{x} \in \operatorname{argmin} f$;
- (ii) \bar{x} is a local minimizer;
- (iii) \bar{x} is a stationary point of f .

Proof. (i) \implies (ii) is obvious. (ii) \implies (iii) follows from [Fermat's Theorem](#). (iii) \implies (i) follows from [Theorem 2.2.1 \(a\)](#). □

Corollary 2.2.4 (Monotonocity of gradient mappings) Let $C \subset \mathbb{R}^n$ be open and convex and let $f : C \rightarrow \mathbb{R}$ be differentiable on C . Then the following hold for all $x, y \in C$

(a) f is convex on C iff

$$\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq 0 \tag{*}$$

(b) f is strictly convex on C iff (*) holds with a strict inequality whenever $x \neq y$.

(c) f is strongly convex with modulus $\mu > 0$ on C iff

$$\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq \mu \|x - y\|^2$$

This has parallels [here](#) and [here](#).

Theorem 2.2.5 (Twice differentiable convex functions) Let $C \subset \mathbb{R}^n$ be open and convex and let $f : C \rightarrow \mathbb{R}$ be twice continuously differentiable on C . Then the following hold:

- (a) f is convex on C iff $\nabla^2 f(x)$ is positive semidefinite $\forall x \in C$.
- (b) If $\nabla^2 f(x)$ is positive definite $\forall x \in C$ then f is strictly convex on C .
- (c) f is strongly convex with modulus $\mu > 0$ on C iff, $\forall x \in C$, the smallest eigenvalue of $\nabla^2 f(x)$ is bounded by μ from below.

Corollary 2.2.6 (Convexity of quadratic functions) Let $A \in \mathbb{R}^{n \times n}$ be symmetric and $b \in \mathbb{R}^n$, $\gamma \in \mathbb{R}$, and define $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$f(x) = \frac{1}{2}x^T A x + b^T x + \gamma$$

Then the following hold:

- (a) f is convex if and only if A is positive semidefinite
- (b) f is strongly convex if and only if A is positive definite

Proof. In view of [Theorem 2.2.5](#), it suffices to show that f is twice continuously differentiable.

$$\begin{aligned}\nabla f(x) &= Ax + b \\ \nabla^2 f(x) &= A\end{aligned}$$

and we are done. □

Theorem 2.2.7 (Convex optimization) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $X \subset \mathbb{R}^n$ be a non-empty convex set. Consider the convex optimization problem

$$\min f(x) \quad \text{s.t.} \quad x \in X \tag{*}$$

Then the following hold:

- (a) A point \bar{x} is a global minimizer of $(*)$ if and only if it is a local minimizer of $(*)$.
- (b) The solution set $\operatorname{argmin}_X f$ of $(*)$ is convex (possibly empty).
- (c) If f is strictly convex, then the solution set has at most one element.
- (d) If f is strongly convex and differentiable and X is closed, then $(*)$ has exactly one solution ($\operatorname{argmin}_X f$ is a singleton).

Proposition 2.3.1 (Operator norms) Let $\|\cdot\|_*$ be a (vector) norm on \mathbb{R}^n and \mathbb{R}^m , respectively. Then for $A \in \mathbb{R}^{m \times n}$,

$$\|A\|_* := \sup_{x \neq 0} \frac{\|Ax\|_*}{\|x\|_*}$$

is a norm on $\mathbb{R}^{m \times n}$ with

$$\|A\|_* = \sup_{\|x\|_*=1} \|Ax\|_* = \sup_{\|x\|_* \leq 1} \|Ax\|_*$$

Proposition 2.3.2 Let $A \in \mathbb{R}^{m \times n}$. Then we have

$$\|A\|_1 = \max_{j=1, \dots, n} \sum_{i=1}^m |a_{ij}| \quad (\text{maximum absolute column sum})$$

$$\|A\|_2 = \sqrt{\lambda_{\max}(A^T A)} \quad (\text{spectral norm})$$

$$\|A\|_\infty = \max_{i=1, \dots, m} \sum_{j=1}^n |a_{ij}| \quad (\text{maximum absolute row sum})$$

Proposition 2.3.3 Let $\|\cdot\|_*$ be a norm on \mathbb{R}^n , \mathbb{R}^m , and \mathbb{R}^p respectively. Then for all $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$ the following hold:

- $\|Ax\|_* \leq \|A\|_* \|x\|_*$ for all $x \in \mathbb{R}^n$ (*compatibility*)
- $\|AB\|_* \leq \|A\|_* \|B\|_*$ (*submultiplicativity*)

Proposition 2.3.4 (Banach Lemma) Let $C \in \mathbb{R}^{n \times n}$ with $\|C\| < 1$ where $\|\cdot\|$ is a submultiplicative matrix norm. Then $I + C$ is invertible and we have

$$\|(I + C)^{-1}\| \leq \frac{1}{1 - \|C\|}$$

Corollary 2.3.5 Let $A, B \in \mathbb{R}^{n \times n}$ with $\|I - BA\| < 1$ for some submultiplicative norm $\|\cdot\|$. Then A and B are invertible with

$$\|B^{-1}\| \leq \frac{\|A\|}{1 - \|I - BA\|}$$

Definition 3.1.1 (Descent direction) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $x \in \mathbb{R}^n$. A vector $d \in \mathbb{R}^n$ is said to be a *descent direction* of f at x if there exists $\ell > 0$ such that

$$f(x + td) < f(x) \quad (t \in (0, \ell])$$

Proposition 3.1.2 (Sufficient condition for descent direction) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be *directionally differentiable* at $x \in \mathbb{R}^n$ in the direction $d \in \mathbb{R}^n$ with

$$f'(x; d) < 0$$

Then d is a *descent direction* of f at x . In particular, this is true if f is differentiable at x with

$$\nabla f(x)^T d < 0$$

Proof. The first statement follows immediately from the definition of the *directional derivative*. The second one uses [Lemma 2.1.2](#). □

Corollary 3.1.3 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable, $B \in \mathbb{R}^{n \times n}$ be positive definite and $x \in \mathbb{R}^n$ with $\nabla f(x) \neq 0$. Then $-B\nabla f(x)$ is a descent direction of f at x .

Proof. This result follows almost immediately from the definition of a *descent direction* and the definition of a *positive definite* matrix. □

Definition 3.1.4 (Step-size rule) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable and let $\mathcal{A}_f := \{(x, d) \mid \nabla f(x)^T d < 0\}$. A set-valued mapping

$$T : (x, d) \in \mathcal{A}_f \mapsto T(x, d) \subset \mathbb{R}_{++}$$

is called a step-size rule for f .

We call it well-defined for f if $T(x, d) \neq \emptyset$ for all $(x, d) \in \mathcal{A}_f$.

If the step-size rule is well-defined for all continuously differentiable functions $\mathbb{R}^n \rightarrow \mathbb{R}$, we simply call it well-defined.

Definition 3.1.5 (Efficient step-size) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable. The **step-size rule** T is called efficient for f if there exists $\theta > 0$ such that

$$f(x + td) \leq f(x) - \theta \left(\frac{\nabla f(x)^T d}{\|d\|} \right)^2$$

Theorem 3.1.6 (Global convergence of general descent method) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable, and let $\{x^k\}$, $\{d^k\}$, $\{t_k\}$ be generated by **Algorithm 3.1.1**. Moreover, assume that the following hold:

(i) (*Angle condition*) There exists $c > 0$ such that

$$-\frac{\nabla f(x^k)^T d^k}{\|\nabla f(x^k)\| \cdot \|d^k\|} \geq c \quad \forall k \in \mathbb{N}$$

i.e. the angle between the gradient vector and the descent direction is at most 90° .

(ii) (*Efficient step-size*) There exists $\theta > 0$ such that

$$f(x^k + t_k d^k) \leq f(x^k) - \theta \left(\frac{\nabla f(x^k)^T d^k}{\|d^k\|} \right)^2 \quad \forall k \in \mathbb{N}$$

Then every cluster point of $\{x^k\}$ is a stationary point of f .

Proof. By (ii), there exists $\theta > 0$ such that

$$f(x^{k+1}) = f(x^k + t_k d^k) \leq f(x^k) - \theta \left(\frac{\nabla f(x^k)^T d^k}{\|d^k\|} \right)^2 \quad \forall k \in \mathbb{N}$$

Putting $k := c^2 \theta$, the angle condition implies

$$f(x^{k+1}) \leq f(x^k) - k \|\nabla f(x^k)\|^2 \quad (*)$$

Let \bar{x} be a cluster point of $\{x^k\}$. As $\{f(x^k)\}$ is monotonically decreasing and convergent to $f(\bar{x})$ on a subsequence (since $\{x^k\} \rightarrow \bar{x}$ on a subsequence and f is continuous), **this implies** that the whole sequence $\{f(x^k)\}$ converges to $f(\bar{x})$.

In particular, we have

$$f(x^{k+1}) - f(x^k) \rightarrow 0$$

Therefore, $(*)$ implies $\|\nabla f(x^k)\| \rightarrow 0$ by squeezing. \square

Definition 3.2.0 (Armijo rule and sufficient decrease) Choose $\beta, \sigma \in (0, 1)$. For $x, d \in \mathcal{A}_f$ the Armijo rule T_A is defined by

$$T_A(x, d) = \max_{l \in \mathbb{N}_0} \{ \beta^l \mid f(x^k + \beta^l d^k) \leq f(x^k) + \beta^l \sigma \nabla f(x^k)^T d^k \}$$

The inequality

$$f(x^k + \beta^l d^k) \leq f(x^k) + \beta^l \sigma \nabla f(x^k)^T d^k$$

is called the *Armijo condition*. It ensures a **sufficient decrease** on the objective function.

Example 3.2.1 (Insufficient decrease) Consider the function $f(x) = (x - 1)^2 - 1$ with optimal value $f^* = -1$.

The sequence $\{x_k\}$ with $x_k := -\frac{1}{k}$ has $f(x_k) = \frac{1+2k}{k^2}$ and

$$f(x_{k+1}) - f(x_k) = \frac{2k^2 + 4k + 1}{k^2(k+1)^2} < 0$$

Hence we've found a case where the objective value decreases, but $f(x_k)$ converges to a non-optimal value. ($f(x_k) \rightarrow 0$, but we want $f(x_k) \rightarrow -1$)

Lemma 3.2.3 (Convergence to gradient) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable. Moreover, let $\{x^k \in \mathbb{R}^n\} \rightarrow x$, $\{d^k \in \mathbb{R}^n\} \rightarrow d$ and $\{t_k > 0\} \downarrow 0$. Then

$$\lim_{k \rightarrow \infty} \frac{f(x^k + t_k d^k) - f(x^k)}{t_k} = \nabla f(x)^T d$$

Proof. By the **mean value theorem**, for all $k \in \mathbb{N}$, there exists $\eta^k \in [x^k, x^k + t_k d^k]$ such that

$$f(x^k + t_k d^k) - f(x^k) = t_k \nabla f(\eta^k)^T d^k$$

Clearly, $\eta^k \rightarrow x$ and hence the continuity of ∇f yields

$$\nabla f(\eta^k)^T d^k \rightarrow \nabla f(x)^T d$$

This readily implies

$$\lim_{k \rightarrow \infty} \frac{f(x^k + t_k d^k) - f(x^k)}{t_k} = \lim_{k \rightarrow \infty} \nabla f(\eta^k)^T d^k = \nabla f(x)^T d$$

□

Theorem 3.2.4 (Global convergence of the gradient method) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable. Then every cluster point of a sequence generated by the **Gradient method with Armijo rule** is a stationary point of f .

Proof. Assume on the contrary that $\nabla f(\bar{x}) \neq 0$.

Let \bar{x} be a cluster point of the generated sequence $\{x^k\}$, and let $\{x^k\}_K$ be a subsequence converging to \bar{x} . By the continuity of f , $\{f(x^k)\}_K \rightarrow f(\bar{x})$.

As $\{f(x^k)\}$ is monotonically decreasing by the Armijo condition and converges on a subsequence to $f(\bar{x})$, **by inspection**, $\{f(x^k)\}_{\mathbb{N}}$ converges to $f(\bar{x})$.

In particular, we have

$$f(x^k) - f(x^{k+1}) \rightarrow 0$$

Substituting $t_k = \beta^l$ and $x^{k+1} = x^k + \beta^l d^k$ into steps **(S2)** and **(S3)** of the algorithm, we have

$$0 \leq t_k \|\nabla f(x^k)\|^2 = -t_k \nabla f(x^k)^T d^k \leq \frac{f(x^k) - f(x^{k+1})}{\sigma} \rightarrow 0$$

Since $\{\nabla f(x^k)\}_K \rightarrow \nabla f(\bar{x}) \neq 0$ (by continuity of ∇f), by squeeze theorem on the above inequality, this implies that $\{t_k\}_K \rightarrow 0$. Due to **(S3)**, for all $k \in K$ sufficiently large, we have

$$f(x^k + \beta^{l_k-1} d^k) - f(x^k) > \beta^{l_k-1} \sigma \nabla f(x^k)^T d^k \quad (*)$$

where $\beta^{l_k} = t_k$ and $l_k \in \mathbb{N}$ is the exponent *uniquely* determined by the Armijo rule in **(S3)**. Note that l_k is the smallest value of l that satisfies the **Armijo condition**, and hence $l_k - 1$ does *not* satisfy the Armijo condition, hence (*).

Passing to the limit on K and using **Lemma 3.2.3** gives

$$-\|\nabla f(\bar{x})\|^2 \geq -\sigma \|\nabla f(\bar{x})\|^2$$

Which is a contradiction because $\sigma \in (0, 1)$ and $\nabla f(\bar{x}) \neq 0$ by assumption. Hence, \bar{x} is indeed a stationary point of f , completing the proof. \square

Proposition 3.2.7 (Kantorovich inequality) Let $A \in \mathbb{R}^{n \times n}$ symmetric positive definite. Then

$$\frac{4\lambda_{\min}\lambda_{\max}}{(\lambda_{\min} + \lambda_{\max})^2} \leq \frac{\|d\|^4}{(d^T A d)(d^T A^{-1} d)} \quad \forall (d \in \mathbb{R}^n \setminus \{0\})$$

Theorem 3.2.8 (Gradient method for strongly convex quadratics) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be given by

$$f(x) = \frac{1}{2} x^T A x + b^T x$$

where $A \in \mathbb{R}^{n \times n}$ is symmetric positive definite and $b \in \mathbb{R}^n$. Let $\bar{x} := -A^{-1}b$ be the (unique) global minimizer of f . Assume that $\{x^k\}$ is generated by the gradient method from **Algorithm 3.2.1**. Then the following hold.

(a) (*Convergence of function values*)

$$f(x^{k+1}) - f(\bar{x}) \leq \left(\frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}} \right)^2 (f(x^k) - f(\bar{x})) \quad \forall (k \in \mathbb{N})$$

i.e. the sequence $\{f(x^k)\}$ converges linearly to $f(\bar{x})$.

(b) (*Convergence of variables*)

$$\|x^k - \bar{x}\| \leq \sqrt{\frac{\lambda_{\max}}{\lambda_{\min}}} \left(\frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}} \right)^k \|x^0 - \bar{x}\| \quad \forall (k \in \mathbb{N})$$

i.e. $\{x^k\}$ converges to \bar{x} for any starting point x^0 .

Definition 3.2.9 (Condition number) For a symmetric positive definite matrix A , its *condition number* is given by

$$\text{cond}(A) := \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)}$$

The condition number of the matrix influences the convergence rate in **Theorem 3.2.8**. If $\text{cond}(A)$ is very large then convergence can be very slow.

Definition 3.3.1 (Convergence rates) Let $\{x^k \in \mathbb{R}^n\} \rightarrow \bar{x}$ and $\|\cdot\|$ be an arbitrary norm on \mathbb{R}^n . Then $\{x^k\}$ converges (at least)

(i) **linearly** to \bar{x} if there exists $c \in (0, 1)$ such that

$$\|x^{k+1} - \bar{x}\| \leq c \|x^k - \bar{x}\| \quad (k \in \mathbb{N} \text{ sufficiently large})$$

(ii) **superlinearly** to \bar{x} if

$$\lim_{k \rightarrow \infty} \frac{\|x^{k+1} - \bar{x}\|}{\|x^k - \bar{x}\|} = 0$$

(iii) **quadratically** to \bar{x} if there exists $C > 0$ such that

$$\|x^{k+1} - \bar{x}\| \leq C \|x^k - \bar{x}\|^2 \quad \forall (k \in \mathbb{N})$$

Definition 3.3.2 (Landau symbols) Let $\{a_k > 0\}, \{b_k > 0\} \downarrow 0$. Then we define

$$\begin{aligned} a_k = o(b_k) &\iff \lim_{k \rightarrow \infty} \frac{a_k}{b_k} = 0 \\ a_k = O(b_k) &\iff \exists C > 0, \forall k \in \mathbb{N} : a_k \leq C b_k \end{aligned}$$

Rewriting [Definition 3.3.1](#) using Landau notation, we say $\{x^k\} \rightarrow \bar{x}$ converges

(i) superlinearly if and only if

$$\|x^{k+1} - \bar{x}\| = o(\|x^k - \bar{x}\|)$$

(ii) quadratically if and only if

$$\|x^{k+1} - \bar{x}\| = O(\|x^k - \bar{x}\|^2)$$

Remark 3.3.3a (Newton's method) Our goal is to effectively solve

$$F(x) = 0$$

where $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is assumed to be continuously differentiable. The method we are going to study is called *Newton's method* and its basic idea is shockingly simple and relies on *linearization*, one of the most basic principles in mathematics:

Suppose \bar{x} is a root of F and x^k is our current approximation of it. Then consider a local, linear approximation

$$x \mapsto F_k(x) := F(x^k) + F'(x^k)(x - x^k)$$

of F at x^k . Now, compute x^{k+1} as a root of F_k , and we should move closer to \bar{x} .

If $F'(x^k) \in \mathbb{R}^{n \times n}$ is invertible we can write

$$x^{k+1} = x^k - F'(x^k)^{-1} F(x^k)$$

But for numerical reasons one does not explicitly invert a matrix, but instead will compute the *Newton direction* d^k as solution to the *Newton equation*

$$F'(x^k)d = -F(x^k)$$

and then update $x^{k+1} := x^k + d^k$. This yields [Algorithm 3.3.1](#).

Lemma 3.3.3 (Local invertibility) Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be continuously differentiable and $\bar{x} \in \mathbb{R}^n$ such that $F'(\bar{x})$ is invertible. Then there exists $\varepsilon > 0$ such that $F'(x)$ is invertible for all $x \in B_\varepsilon(\bar{x})$. Moreover, there exists $c > 0$ such that

$$\|F'(x)^{-1}\| \leq c \quad (x \in B_\varepsilon(\bar{x}))$$

Remark 3.3.3b (Differentiability in Landau notation) Using the Landau notation, we can express the fact that F is **differentiable** at $\bar{x} \in \mathbb{R}^n$ if and only if

$$\|F(x^k) - F(\bar{x}) - F'(\bar{x})(x^k - \bar{x})\| = o(\|x^k - \bar{x}\|)$$

for all sequences $\{x^k\} \rightarrow \bar{x}$.

Here's how it's expanded:

$$\lim_{x^k \rightarrow \bar{x}} \frac{\|F(x^k) - F(\bar{x}) - F'(\bar{x})(x^k - \bar{x})\|}{\|x^k - \bar{x}\|} = 0$$

Definition 3.3.4 (Local Lipschitz) We say that $G : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is locally Lipschitz (continuous) at $\bar{x} \in \mathbb{R}^n$ if there exists $L = L(\bar{x}) > 0$ such that

$$\|G(x) - G(y)\| \leq L \|x - y\| \quad (x, y \in B_\varepsilon(\bar{x}))$$

Lemma 3.3.5 Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be continuously differentiable and $\{x^k\}$ such that $\{x^k\} \rightarrow \bar{x}$. Then

$$\|F(x^k) - F(\bar{x}) - F'(\bar{x})(x^k - \bar{x})\| = o(\|x^k - \bar{x}\|)$$

If F' is, in addition, **locally Lipschitz continuous**, we also have

$$\|F(x^k) - F(\bar{x}) - F'(\bar{x})(x^k - \bar{x})\| = O(\|x^k - \bar{x}\|^2)$$

Theorem 3.3.6 (Convergence of local Newton's method for equations) Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be continuously differentiable and let \bar{x} be a root of F such that $F'(\bar{x})$ is invertible. Then there exists $\varepsilon > 0$ such that for every $x^0 \in B_\varepsilon(\bar{x})$, the following hold:

- (a) The local Newton method from **Algorithm 3.3.1** is well-defined and generates a sequence $\{x^k\}$ convergent to \bar{x} .
- (b) The rate of convergence is at least superlinear.
- (c) If in addition F' is **locally Lipschitz continuous** at \bar{x} the rate of convergence is quadratic.

Remark 3.3.6a (Newton's method in optimization) We now want to exploit our study of Newton's method for solving smooth, nonlinear equations to tackle unconstrained optimization problems of the form

$$\min_{x \in \mathbb{R}^n} f(x)$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is at least twice continuously differentiable. Recall that a necessary condition for \bar{x} to be a local minimizer of f is

$$\nabla f(\bar{x}) = 0$$

So we can put $F := \nabla f$ and we have $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ continuously differentiable, and all local minimizers of f are among its roots.

This yields **Algorithm 3.3.2**.

Theorem 3.3.7 (Convergence of local Newton's method for optimization) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable and let \bar{x} be a stationary point of f such that $\nabla^2 f(\bar{x})$ is invertible. Then there exists $\varepsilon > 0$ such that for every $x^0 \in B_\varepsilon(\bar{x})$, the following hold:

- (a) **Algorithm 3.3.2** is well-defined and generates a sequence $\{x^k\}$ convergent to \bar{x} .
- (b) The rate of convergence is at least superlinear.
- (c) If in addition $\nabla^2 f$ is **locally Lipschitz continuous** at \bar{x} the rate of convergence is quadratic.

Theorem 3.3.9 (Global convergence of **Algorithm 3.3.3)** Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable. Then every cluster point of a sequence generated by **Algorithm 3.3.3** is a stationary point of f .

Lemma 3.3.10 (Moré and Sorensen) Let \bar{x} be an isolated cluster point of $\{x^k \in \mathbb{R}^n\}$ and assume that $\{\|x^{k+1} - x^k\|\}_K \rightarrow 0$ for every subsequence $\{x^k\}_K \rightarrow \bar{x}$. Then the whole sequence $\{x^k\}$ converges to \bar{x} .

Corollary 3.3.11 Let \bar{x} be an isolated cluster point of a sequence $\{x^k\}$ generated by **Algorithm 3.3.3**. Then the whole sequence $\{x^k\}$ converges to \bar{x} .

Proposition 3.3.12 (Acceptance of full step-size in globalized Newton's method) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable, and $\bar{x} \in \mathbb{R}^n$ such that $\nabla f(\bar{x}) = 0$ and $\nabla^2 f(\bar{x})$ is positive definite. Assume that $\{x^k\} \rightarrow \bar{x}$ and that d^k is given by

$$d^k = -\nabla^2 f(x^k)^{-1} \nabla f(x^k)$$

(**Newton Equation**)

Then there exists $k_0 \in \mathbb{N}$ such that $\forall \sigma \in (0, \frac{1}{2})$, $k \geq k_0$:

$$f(x^k + d^k) \leq f(x^k) + \sigma \nabla f(x^k)^T d^k$$

The significance of this last equation being that f experiences **sufficient decrease**.

Theorem 3.3.13 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable and let $\{x^k\}$ be generated by **Algorithm 3.3.3**. If \bar{x} is a cluster point of $\{x^k\}$ such that $\nabla^2 f(\bar{x})$ is positive definite then the following hold:

- (a) The whole sequence $\{x^k\}$ converges to \bar{x} and \bar{x} is a strict local minimizer of f .
- (b) For all $k \in \mathbb{N}$ sufficiently large, the search direction d^k will be determined through the **Newton equation**.
- (c) For all $k \in \mathbb{N}$ sufficiently large, the full step-size $t_k = 1$ will be accepted
- (d) $\{x^k\}$ converges superlinearly to \bar{x}
- (e) If $\nabla^2 f$ is locally Lipschitz then $\{x^k\}$ converges to \bar{x} quadratically.

Definition 3.3.14a (Quasi-Newton equation) In the context of iterating over $\{x_k\}$ and $\{H_k\}$ where H_k is the approximation of $\nabla^2 f(x^k)$ at x^k , H_{k+1} satisfies the quasi-Newton equation if

$$H_{k+1}s^k = y^k,$$

where $s^k := x^{k+1} - x^k$ and $y^k := \nabla f(x^{k+1}) - \nabla f(x^k)$.

Remark 3.3.14b (Direct Quasi-Newton methods) In order to devise a strategy of how to approximate the Hessian matrix of the underlying function f the current iterate x^k we first need to agree on which properties we would like it to have. To this end, let H_k be an approximation of $\nabla^2 f(x^k)$. We would like for H_{k+1} to satisfy the following criteria:

- I. $H_{k+1} = H_{k+1}^T$ is symmetric.
- II. H_{k+1} satisfies the **quasi-Newton equation**.
- III. H_{k+1} can be obtained efficiently from H_k .
- IV. The resulting method has strong local convergence properties.

Remark 3.3.14 Let x^k be a current iterate for minimizing $f : \mathbb{R}^n \rightarrow \mathbb{R}$ that is twice continuously differentiable.

- (a) The Hessian $\nabla^2 f(x^k)$ of f at x^k does not necessarily satisfy the **quasi-Newton equation**.
- (b) Condition I. is motivated by **Schwarz's Theorem**.
- (c) The quasi-Newton equation can be motivated by the Mean-Value Theorem in integral form, which yields

$$\begin{aligned}\nabla f(x^{k+1}) - \nabla f(x^k) &= \int_0^1 \nabla^2 f(x^k + t(x^{k+1} - x^k)) dt \cdot (x^{k+1} - x^k) \\ y^k &= \int_0^1 \nabla^2 f(x^k + t(x^{k+1} - x^k)) dt \cdot s^k\end{aligned}$$

The term $\int_0^1 \nabla^2 f(x^k + t(x^{k+1} - x^k)) dt$ can be interpreted as some sort of “averaged Hessian”.

Remark x.x.x (Collection of unconstrained minimization methods)

- 1. Gradient method
- 2. Globalized Newton's method
- 3. Globalized BFGS method
- 4. Globalized inexact Newton's method

Algorithm 3.1.1 (General line-search descent algorithm) Goal is to solve

$$\min_{x \in \mathbb{R}^n} f(x).$$

- (S0) *Initialization*: Choose $x^0 \in \mathbb{R}^n$ and put $k := 0$.
- (S1) *Termination*: If x^k satisfies a termination criterion: STOP.
- (S2) *Search direction*: Determine d^k such that $\nabla f(x^k)^T d^k < 0$.
- (S3) *Step-size*: Determine $t_k > 0$ such that $f(x^k + t_k d^k) < f(x^k)$.
- (S4) *Update*: Put $x^{k+1} := x^k + t_k d^k$, $k \leftarrow k + 1$ and go to (S1).

Algorithm 3.2.1 (Gradient method with Armijo rule) Goal is to solve

$$\min_{x \in \mathbb{R}^n} f(x).$$

- (S0) Choose $x^0 \in \mathbb{R}^n$, $\sigma, \beta \in (0, 1)$, $\varepsilon \geq 0$ and put $k := 0$.
- (S1) If $\|\nabla f(x^k)\| \leq \varepsilon$, STOP.
- (S2) Put $d^k := -\nabla f(x^k)$.
- (S3) Determine $t_k > 0$ by

$$t_k := \max_{l \in \mathbb{N}_0} \{\beta^l \mid f(x^k + \beta^l d^k) \leq f(x^k) + \beta^l \sigma \nabla f(x^k)^T d^k\}$$

- (S4) Put $x^{k+1} := x^k + t_k d^k$, $k \leftarrow k + 1$ and go to (S1).

Algorithm 3.3.1 (Local Newton's method for equations) Goal is to solve

$$F(x) = 0$$

where $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and F is assumed to be continuously differentiable.

- (S0) Choose $x^0 \in \mathbb{R}^n$, $\varepsilon \geq 0$ and put $k := 0$.
- (S1) If $\|F(x^k)\| \leq \varepsilon$, STOP.
- (S2) Compute d^k as a solution of
$$F'(x^k)d = -F(x^k)$$
- (S3) Put $x^{k+1} := x^k + d^k$, $k \leftarrow k + 1$ and go to (S1).

Algorithm 3.3.2 (Local Newton's method for unconstrained optimization)
Goal is to solve

$$\min_{x \in \mathbb{R}^n} f(x).$$

The following is obtained by substituting F for ∇f in the **local Newton's method for equations**.

- (S0) Choose $x^0 \in \mathbb{R}^n$, $\varepsilon \geq 0$ and put $k := 0$.

(S1) If $\|\nabla f(x^k)\| \leq \varepsilon$, STOP.

(S2) Compute d^k as a solution of

$$\nabla^2 f(x^k)d = -\nabla f(x^k)$$

(S3) Put $x^{k+1} := x^k + d^k$, $k \leftarrow k + 1$ and go to (S1).

Algorithm 3.3.3 (Globalized Newton's method for unconstrained optimization) Goal is to solve

$$\min_{x \in \mathbb{R}^n} f(x).$$

The following is obtained by substituting F for ∇f in the **local Newton's method for equations**.

(S0) Choose $x^0 \in \mathbb{R}^n$, $\rho > 0$, $p > 2$, $\beta \in (0, 1)$, $\sigma \in (0, \frac{1}{2})$, $\varepsilon \geq 0$ and put $k := 0$.

(S1) If $\|\nabla f(x^k)\| \leq \varepsilon$, STOP.

(S2) Try to compute d^k as a solution of

$$\nabla^2 f(x^k)d = -\nabla f(x^k)$$

If no solution can be found or if

$$\nabla f(x^k)^T d^k > -\rho \|d^k\|^p$$

(insufficient decrease)

then fall back to $d^k := -\nabla f(x^k)$

(S3) Determine t_k by

$$t_k := \max_{l \in \mathbb{N}_0} \{\beta^l \mid f(x^k + \beta^l d^k) \leq f(x^k) + \beta^l \sigma \nabla f(x^k)^T d^k\}$$

(S4) Put $x^{k+1} := x^k + t_k d^k$, $k \leftarrow k + 1$ and go to (S1).

Nonlinear Optimization, Part II: Constrained Optimization

Definition 5.0.0 (Standard Nonlinear Program)

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad & g_i(x) \leq 0 \quad \forall i \in I \\ & h_j(x) = 0 \quad \forall j \in J \end{aligned} \quad (5.1)$$

Where $f, g_i, h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ are continuously differentiable.

We call (5.1) a nonlinear program (NLP) in standard form.

By convention, we let the feasible set of (5.1) be denoted by X , with

$$X := \left\{ x \in \mathbb{R}^n \mid \begin{array}{l} g_i(x) \leq 0 \quad \forall i \in I \\ h_j(x) = 0 \quad \forall j \in J \end{array} \right\} \quad (5.2)$$

By the continuity of the constraint functions, X is closed.

We will use $I := \{1, \dots, m\}$ and $J := \{1, \dots, p\}$, and define the **active set** at $\bar{x} \in X$ as

$$I(\bar{x}) := \{i \in I \mid g_i(\bar{x}) = 0\}$$

Definition 5.1.1 (Cones) A non-empty set $K \subset \mathbb{R}^n$ is said to be a cone if

$$\lambda v \in K \quad (\lambda \geq 0, v \in K)$$

i.e. K is a cone if and only if it is closed under multiplication with non-negative scalars.

Example 5.1.2 (Examples of cones)

- (a) (*Non-negative Orthant*) For all $n \in \mathbb{N}$, the non-negative orthant \mathbb{R}_+^n is a convex cone, which is also a polyhedron as

$$\mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid (-e_i)^T x \leq 0, \forall i = 1, \dots, n\}$$

- (b) (*Cone complimentary constraints*) Let $K \subset \mathbb{R}^n$ be a cone. Then the set

$$\Lambda := \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n \mid x, y \in K, \langle x, y \rangle = 0\}$$

is a cone. A prominent example is $K = \mathbb{R}^n$, in which case Λ is called the complementary constraint set.

- (c) (*Positive semidefinite matrices*) For $n \in \mathbb{N}$, the set of positive semidefinite $n \times n$ matrices is a convex cone in the space of $n \times n$ symmetric matrices.

Definition 5.1.3 (Tangent cone) Let $S \subset \mathbb{R}^n$ and $\bar{x} \in S$. Then the set

$$T_S(\bar{x}) := \left\{ d \in \mathbb{R}^n \mid \exists \{x^k \in S\} \rightarrow \bar{x}, \{t_k\} \downarrow 0 : \frac{x^k - \bar{x}}{t_k} \rightarrow d \right\}$$

is called the (Bouligand) tangent cone of S at \bar{x} .

Proposition 5.1.4 Let $S \subset \mathbb{R}^n$ and $x \in S$. Then $T_S(x)$ is a closed cone.

Theorem 5.1.5 (Basic first-order optimality condition) Let \bar{x} be a local minimizer of $f \in C^1$ over S . Then the following hold:

(a) $\nabla f(\bar{x})^T d \geq 0 \quad (d \in T_S(\bar{x}))$

(b) If S is convex then

$$\nabla f(\bar{x})^T (x - \bar{x}) \geq 0 \quad (x \in S)$$

Definition 5.1.6 (Projection on a set) Let $S \subset \mathbb{R}^n$ be non-empty and $x \in \mathbb{R}^n$. Then we define the projection of x on S by

$$P_S(x) := \operatorname{argmin}_{y \in S} \|x - y\|$$

Proposition 5.1.7 (Projection on a closed convex set) Let $x \in \mathbb{R}^n$ and $S \subset \mathbb{R}^n$ be a non-empty, closed, and convex set. Then the following hold:

(a) $P_S(x)$ has exactly one element, i.e. P is a function $\mathbb{R}^n \rightarrow S$

(b) $P_S(x) = x$ if and only if $x \in S$

(c) $(P_S(x) - x)^T (y - P_S(x)) \geq 0 \quad (y \in S)$

Proof. (a) follows immediately from [Theorem 2.2.7\(d\)](#) since the function $y \mapsto \frac{1}{2} \|x - y\|^2$ is strongly convex.

(b) is obvious.

(c) follows from [Theorem 5.1.5\(b\)](#) applied to $f : y \mapsto \frac{1}{2} \|x - y\|^2$. □

Lemma 5.1.8 Let $B \in \mathbb{R}^{l \times n}$. Then

$$K := \{B^T x \mid x \geq 0\}$$

is a (non-empty) closed, convex cone.

Theorem 5.1.9 (Farkas' Lemma) Let $B \in \mathbb{R}^{l \times n}$ and $h \in \mathbb{R}^n$. Then the system

$$B^T x = h \quad (x \in \mathbb{R}^l, x \geq 0)$$

has a solution if and only if $h^T d \geq 0$ for all $d \in \mathbb{R}^n$ such that $Bd \geq 0$.

Proof. Proving that (1) implies (2):

Let $x \geq 0$ such that $B^T x = h$. Then for any d such that $Bd \geq 0$, we have

$$h^T d = (B^T x)^T d = x^T Bd$$

But $x^T Bd \geq 0$ because $x \geq 0$ and $Bd \geq 0$.

Proving that (2) implies (1) by contrapositive.

Assume that (1) is false. Then

$$h \notin \{B^T x \mid x \geq 0\} =: K$$

By [Lemma 5.1.8](#), K is a closed convex cone.

Set $\bar{s} := P_K(h)$ and $\bar{d} := \bar{s} - h$. Note that $\bar{s} \in K$, and $h \notin K$, and hence $\bar{d} \neq 0$.

By [Proposition 5.1.7\(c\)](#),

$$\bar{d}^T(s - \bar{s}) \geq 0 \quad \forall (s \in K) \quad (*)$$

Substituting $s := 0$ and $s := 2\bar{s}$, we obtain two simultaneous inequalities

$$\bar{d}^T \bar{s} \leq 0 \quad \text{and} \quad \bar{d}^T \bar{s} \geq 0$$

And hence $\bar{d}^T \bar{s} = 0$. Using this with $(*)$ gives

$$\bar{d}^T s \geq 0$$

Then by definition of cone K , for all $x \geq 0$,

$$\begin{aligned} \bar{d}^T B^T x &\geq 0 \\ \Rightarrow (B\bar{d})^T x &\geq 0 \end{aligned}$$

Inserting $x := e_i$ (where e_i is the i^{th} component vector) for $i = 1, \dots, n$ implies $(B\bar{d})^T \geq 0$.

On the other hand (recall $\bar{d}^T \bar{s} = 0$ from above)

$$\begin{aligned} h^T \bar{d} &= (\bar{s} - \bar{d})^T \bar{d} \\ &= \bar{s}^T \bar{d} - \|\bar{d}\|^2 \\ &= -\|\bar{d}\|^2 \\ &\leq 0 \end{aligned}$$

But since $\bar{d} \neq 0$, we have the strict inequality $h^T \bar{d} < 0$.

Therefore, $B\bar{d} \geq 0$, but $h^T \bar{d} < 0$, i.e. (2) does not hold. \square

Definition 5.1.10 (Karush-Kuhn-Tucker conditions) Consider the standard NLP in [\(5.1\)](#). and let X be the feasible set of [\(5.1\)](#).

1. The function $L : \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} L(x, \lambda, \mu) &= f(x) + \sum_{i=1}^m \lambda_i g_i(x) + \sum_{j=1}^p \mu_j h_j(x) \\ &= f(x) + \lambda^T g(x) + \mu^T h(x) \end{aligned}$$

is called the Lagrangian (function) of [\(5.1\)](#).

2. The set of conditions

$$\begin{aligned} \nabla_x L(x, \lambda, \mu) &= 0, \\ h(x) &= 0, \\ \lambda \geq 0, \quad g(x) \leq 0, \quad \lambda^T g(x) &= 0 \end{aligned}$$

are called the Karush-Kuhn-Tucker conditions for [\(5.1\)](#), where

$$\begin{aligned} \nabla_x L(x, \lambda, \mu) &= \nabla f(x) + \sum_{i=1}^m \lambda_i \nabla g_i(x) + \sum_{j=1}^p \mu_j \nabla h_j(x) \\ &= \nabla f(x) + \lambda^T \nabla g(x) + \mu^T \nabla h(x) \end{aligned}$$

3. A triple $(\bar{x}, \bar{\lambda}, \bar{\mu})$ that satisfies the KKT conditions is called a KKT point.
4. Given \bar{x} , a feasible point for (5.1), we define

$$M(\bar{x}) := \{(\lambda, \mu) \mid (\bar{x}, \lambda, \mu) \text{ is a KKT point of (5.1)}\}$$

as the set of all KKT multipliers (possibly empty) at \bar{x} .

Definition 5.1.10a (Constraint qualification (CQ)) We define a condition about the feasible set X of a standard NLP (5.1) that guarantees that the KKT conditions hold at a local minimizer as a constraint qualification.

If a CQ holds on $\bar{x} \in X$, then KKT is necessary for \bar{x} to be a local minimizer.

If a CQ holds on $\bar{x} \in X$, then \bar{x} being a local minimizer implies that there exists a $(\bar{\lambda}, \bar{\mu})$ such that $(\bar{x}, \bar{\lambda}, \bar{\mu})$ is a KKT point.

Definition 5.1.11 (Linearized cone) Let X be the feasible set of (5.1). The linearized cone (of X) at $\bar{x} \in X$ is defined by

$$L_X(\bar{x}) := \left\{ d \in \mathbb{R}^n \mid \begin{array}{ll} \nabla g_i(\bar{x})^T d \leq 0 & \forall i = I(\bar{x}) \\ \nabla h_j(\bar{x})^T d = 0 & \forall j = 1, \dots, p \end{array} \right\}$$

Definition 5.1.12 (Abadie constraint qualification (ACQ)) We say that the ACQ holds at $\bar{x} \in X$ if

$$T_X(\bar{x}) = L_X(\bar{x})$$

That is, the tangent cone is exactly the linearized cone.

Theorem 5.1.13 (KKT conditions under ACQ) Let $\bar{x} \in X$ be a local minimizer of (5.1) such that ACQ holds at \bar{x} . Then there exists $(\bar{\lambda}, \bar{\mu}) \in \mathbb{R}^m \times \mathbb{R}^p$ such that $(\bar{x}, \bar{\lambda}, \bar{\mu})$ is a KKT point of (5.1).

Proof. By Theorem 5.1.5,

$$\nabla f(\bar{x})^T d \geq 0 \quad \forall (d \in T_X(\bar{x})) \quad (*)$$

Set

$$B := \begin{pmatrix} -\nabla g_i(\bar{x})^T & (i = 1, \dots, m) \\ -\nabla h_j(\bar{x})^T & (j = 1, \dots, p) \\ \nabla h_j(\bar{x})^T & (j = 1, \dots, p) \end{pmatrix} \in \mathbb{R}^{(m+2p) \times n}$$

Then purely from the definition of a linearized cone and how we set B , we have

$$d \in L_X(\bar{x}) \iff Bd \geq 0$$

By ACQ, we have $d \in T_X(\bar{x}) \iff Bd \geq 0$

Combined with (*), we have

$$\nabla f(\bar{x})^T d \geq 0 \quad \forall (d : Bd \geq 0)$$

(Think $h = \nabla f(\bar{x})$ and apply the Farkas Lemma.)

By the Farkas Lemma,

$$\exists y = \begin{pmatrix} y^1 \in \mathbb{R}^m \\ y^2 \in \mathbb{R}^p \\ y^3 \in \mathbb{R}^p \end{pmatrix}$$

such that $y \geq 0$, and $B^T y = \nabla f(\bar{x})$

Define $\bar{\lambda} \in \mathbb{R}^n, \bar{\mu} \in \mathbb{R}^p$ by

$$\bar{\lambda}_i = \begin{cases} y_i^1 & \text{if } i = 1, \dots, m \\ 0 & \text{otherwise} \end{cases}$$

and

$$\bar{\mu}_i = \begin{cases} y_j^2 - y_j^3 & \text{if } j = m+1, \dots, m+2p \\ 0 & \text{otherwise} \end{cases}$$

Then $(\bar{x}, \bar{\lambda}, \bar{\mu})$ is a KKT point.

MORE NOTES

$$\begin{aligned} 0 &= \nabla f(\bar{x}) + \sum_{i=0}^m y_i^1 \nabla g_i(\bar{x}) + \sum_{j=m+1}^{m+2p} (y_j^2 - y_j^3) \nabla h_j(\bar{x}) \\ &= \nabla f(\bar{x}) + \sum_{i=0}^m \bar{\lambda}_i \nabla g_i(\bar{x}) + \sum_{j=m+1}^{m+2p} \bar{\mu}_j \nabla h_j(\bar{x}) \end{aligned}$$

and then there is a line with a tick/check next to it:

$$\bar{\lambda}^T g(\bar{x}) = \sum_{i=0}^m \bar{\lambda}_i g_i(\bar{x}) = 0$$

□

Definition 5.1.14 (Constraint qualifications) A condition on X (i.e. on g and h) that ensures that the KKT conditions hold at a local minimizer is called a **constraint qualification**.

Definition 5.1.15 (LICQ and MFCQ) Let \bar{x} be feasible for (1). We say that

- (a) **(LICQ)** the linear independence constraint qualification holds at \bar{x} if the gradients

$$\begin{aligned} \nabla g_i(\bar{x}) & \quad (i \in I(\bar{x})), \\ \nabla h_j(\bar{x}) & \quad (j \in J) \end{aligned}$$

are linearly independent.

- (b) **(MFCQ)** the Mangasarian-Fromovitz constraint qualification holds at \bar{x} if the gradients

$$\nabla h_j(\bar{x}) \quad (j \in J)$$

are linearly independent, and $\exists d \in \mathbb{R}^n$ such that

$$\begin{aligned} \nabla g_i(\bar{x})^T d &< 0 \quad (i \in I(\bar{x})) \\ \nabla h_j(\bar{x})^T d &= 0 \quad (j \in J) \end{aligned}$$

Proposition 5.1.16 (LICQ implies MFCQ) Let \bar{x} be feasible for (1) such that **LICQ** holds at \bar{x} . Then **MFCQ** holds.

With $I := 1, \dots, m$ and $J := 1, \dots, p$, the Standard NLP is

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad & g_i(x) \leq 0 \quad \forall i \in I \\ & h_j(x) = 0 \quad \forall j \in J \end{aligned} \quad (5.1)$$

We define the active set $I(\bar{x})$ as

$$I(\bar{x}) := \{i \in I \mid g_i(\bar{x}) = 0\}$$

Let X be the feasible set of the NLP.

TANGENT CONE:

$$T_X(\bar{x}) := \left\{ d \in \mathbb{R}^n \mid \exists \{x^k \in X\} \rightarrow \bar{x}, \{t_k\} \downarrow 0 : \frac{x^k - \bar{x}}{t_k} \rightarrow d \right\}$$

LINEARIZED CONE:

$$L_X(\bar{x}) := \left\{ d \in \mathbb{R}^n \mid \begin{array}{l} \nabla g_i(\bar{x})^T d \leq 0 \quad \forall i \in I(\bar{x}) \\ \nabla h_j(\bar{x})^T d = 0 \quad \forall j \in J \end{array} \right\}$$

Definition 5.1.22 We say that the affine constraint qualification (Affine CQ) holds for (5.1) if all constraints are affine. That is, there exists

$$\begin{aligned} a_i &\in \mathbb{R}^n & \alpha_i &\in \mathbb{R} & \forall i \in I \\ b_j &\in \mathbb{R}^n & \beta_j &\in \mathbb{R} & \forall j \in J \end{aligned}$$

such that for all $x \in \mathbb{R}^n$,

$$\begin{aligned} g_i(x) &= a_i^T x - \alpha_i & \forall i \in I \\ h_j(x) &= b_j^T x - \beta_j & \forall j \in J \end{aligned}$$

Proposition 5.1.23 Let the Affine CQ hold for (5.1). Then ACQ holds at every feasible point.

Proof. Let $\bar{x} \in X$. In view of Lemma 5.1.18 we only have to show that $L_X(\bar{x}) \subset T_X(\bar{x})$. Let $d \in L_X(\bar{x})$. Then by defn of the linearized cone we have

$$\begin{aligned} \alpha_i^T d &\leq 0 & \forall i \in I \\ \beta_j^T d &= 0 & \forall j \in J \end{aligned}$$

Now, let $\{t_k\} \downarrow 0$ and put $x^k := \bar{x} + t_k d$. Then

$$x^k \rightarrow \bar{x} \quad \text{and} \quad \frac{x^k - \bar{x}}{t_k} \rightarrow d$$

Hence, we still need to show that $x^k \in X$ ($\forall k$ suff. large)

For $i \notin I(\bar{x})$ Then $a_i^T \bar{x} < \alpha_i$, hence by continuity

$$a_i^T x^k < \alpha_i \quad \forall (x \text{ suff. large})$$

For $i \in I(\bar{x})$ Then

$$a_i^T x^k = a_i^T \bar{x} + t_k a_i^T d \leq a_i^T \bar{x} = \alpha_i$$

because $a_i^T d \leq 0$ by definition of $L_X(\bar{x})$.

For $j \in J$ Then

$$b_j^T x^k = b_j^T \bar{x} + t_k b_j^T d = \beta_j$$

because $b_j^T d = 0$ by definition of $L_X(\bar{x})$.

These three together show that $x^k \in X$ for k sufficiently large. This completes the proof. \square

5.1.5 CONVEX PROBLEMS

Consider

$$\begin{aligned} \min f(x) \quad \text{s.t.} \quad & g_i(x) \leq 0 & \forall i \in I \\ & h_j(x) = b_j^T x - \beta_j = 0 & \forall j \in J \end{aligned} \quad (5.2)$$

where $f, g_i, h_j \in \mathbb{R}^n \rightarrow \mathbb{R}$ are cont. diff **and convex**, and $b_j \in \mathbb{R}^n, \beta_j \in \mathbb{R}$. Then

$$X = \left\{ x \in \mathbb{R}^n \mid \begin{array}{ll} g_i(x) \leq 0 & \forall i \in I \\ h_j(x) = 0 & \forall j \in J \end{array} \right\}$$

is convex (see Midterms).

Theorem 5.1.24 Let \bar{x} be feasible for (5.2), and consider the following statements:

- (a) There exists $(\bar{\lambda}, \bar{\mu}) \in M(\bar{x})$
- (b) \bar{x} is a global minimizer of (5.2)

Then (a) implies (b). Hence, if a CQ holds at \bar{x} , then (a) iff (b).

Proof. Let $(\bar{x}, \bar{\lambda}, \bar{\mu})$ be a KKT point of (5.2), and let $\bar{x} \in X$.

Then, by Theorem 2.2.1,

$$f(x) \geq f(\bar{x}) + \nabla f(\bar{x})^T (x - \bar{x})$$

But with the KKT conditions, we can rewrite $\nabla f(\bar{x})$ as

$$\nabla f(\bar{x}) = - \sum_{i \in I(\bar{x})}^m \bar{\lambda}_i \nabla g_i(\bar{x}) - \sum_{j=1}^p \bar{\mu}_j \nabla h_j(\bar{x})$$

So then

$$f(x) \geq f(\bar{x}) - \sum_{i \in I(\bar{x})}^m \bar{\lambda}_i \nabla g_i(\bar{x})^T (x - \bar{x}) - \sum_{j=1}^p \bar{\mu}_j \nabla h_j(\bar{x})^T (x - \bar{x}) \quad (*)$$

Again by Theorem 2.2.1, we have

$$\nabla g_i(\bar{x})^T (x - \bar{x}) \leq g_i(x) - g_i(\bar{x})$$

Separately,

$$\nabla h_j(\bar{x})^T (x - \bar{x}) = 0$$

Putting everything back to (*) and noting that $g_i(\bar{x}) = 0$ by definition of an active set,

$$f(x) \geq f(\bar{x}) - \sum_{i \in I(\bar{x})}^m \bar{\lambda}_i g_i(x)$$

But since $\bar{\lambda}_i \geq 0$ and $g_i(x) \leq 0$, we have

$$f(x) \geq f(\bar{x})$$

and hence \bar{x} is a global minimum of (5.2). Hence shown that (a) implies (b). The converse direction is the definition of a CQ.

And hence (a) iff (b). □

Definition 5.1.25 (Slater constraint qualification) We say that Slater CQ holds for (5.2) if there exists \hat{x} such that

$$\begin{aligned} g_i(\hat{x}) &< 0 & \forall i \in I \\ h_j(\hat{x}) &= 0 & \forall j \in J \end{aligned}$$

We call such an \hat{x} a Slater point.

Proposition 5.1.26 Let SCQ hold for (5.2). Then ACQ holds at every feasible point.

Proof. Let $\bar{x} \in X$ and set

$$F(\bar{x}) := \left\{ d \in \mathbb{R}^n \mid \begin{array}{ll} \nabla g_i(\bar{x})^T \leq 0 & \forall i \in I(\bar{x}) \\ b_j^T d = 0 & \forall j \in J \end{array} \right\}$$

Lemma ($F(\bar{x}) \subset T_X(\bar{x})$) Let $d \in F(\bar{x})$, take $\{t_k\} \downarrow 0$. Set $x^k := \bar{x} + t_k d$. Then $\frac{x^k - \bar{x}}{t_k} \rightarrow d$. Moreover,

For $i \notin I(\bar{x})$, $g_i(x^k) < 0$ for k sufficiently large.

For $i \in I(\bar{x})$, $\frac{g_i(x^k) - g_i(\bar{x})}{t_k} \rightarrow \nabla g_i(\bar{x})^T d < 0$, and $g_i(\bar{x}) = 0$, so we have $g_i(x^k) < 0$ for k sufficiently large.

For $j \in J$, $h_j(x^k) = b_j^T x^k - \beta_j = t_k \nabla h_j(\bar{x})^T d - \beta_j$

$$\begin{aligned} h_j(x^k) &= b_j^T x^k - \beta_j \\ &= t_k \nabla h_j(\bar{x})^T d + \nabla h_j(\bar{x}) \beta_j = 0 \end{aligned}$$

Thus

$$clF(\bar{x}) \subset clT_X(\bar{x}) = T_X(\bar{x})$$

We now show $L_X(\bar{x}) \subset dF(\bar{x})$. To this end, let $d \in L_X(\bar{x})$, and let \hat{x} be a Slater point.

Set \hat{d} to be $\hat{x} - \bar{x}$. Then by Theorem 2.2.1,

$$\nabla g_i(\bar{x})^T \hat{d} \leq g_i(\hat{x}) - g_i(\bar{x}) < 0 \quad \forall (i \in I(\bar{x})) \quad (*)$$

< 0 because of definitions of Slater and Active Set.

Moreover, by the affine-ness of h_j , we have

$$\nabla h_j(\bar{x})^T \hat{d} = h_j(\hat{x}) - h_j(\bar{x}) = 0 \quad \forall (j \in J) \quad (**)$$

because both $h_j(\hat{x})$ and $h_j(\bar{x})$ are zero.

Now we take a small pertubation of d using \hat{d} :

$$d(\delta) := d + \delta \hat{d} \quad (\delta > 0)$$

Then $d(\delta) \in F(\bar{x})$, since

$$\nabla g_i(\bar{x})^T d(\delta) = \nabla g_i(\bar{x})^T d + \delta \nabla g_i(\bar{x})^T \hat{d} < 0 \quad \forall (i \in I(\bar{x}))$$

$$\nabla h_j(\bar{x})^T d(\delta) = \nabla h_j(\bar{x})^T d + \delta \nabla h_j(\bar{x})^T \hat{d} = 0 \quad \forall (j \in J)$$

< 0 because the 1st term ≤ 0 and 2nd term < 0 , because d is in the linearized cone, and $\hat{d} \dots?$

And $= 0$ because both terms $= 0$.

Hence,

$$d = \lim_{\delta \downarrow 0} d(\delta) \subset cl F(\bar{x})$$

□

Consider the standard NLP:

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad & g_i(x) \leq 0 \quad \forall i \in I \\ & h_j(x) = 0 \quad \forall j \in J \end{aligned} \quad (1)$$

But now assume that $f, g_i, h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ with no smoothness.

The Lagrangian of (1) is

$$\begin{aligned} L(x, \lambda, \mu) &= f(x) + \sum_{i \in I} \lambda_i g_i(x) + \sum_{j \in J} \mu_j h_j(x) \\ &= f(x) + \lambda^T g(x) + \mu^T h(x) \end{aligned}$$

The Dual Problem

Observe that if x is a feasible point,

$$\sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} = \begin{cases} f(x) & \text{if } x \in X, \\ +\infty & \text{if } x \notin X \end{cases}$$

Therefore the primal problem (1) is equivalent to

$$\min_{x \in \mathbb{R}^n} \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(x, \lambda, \mu)$$

Q. When can we switch min(inf) and sup?

Definition 6.1.1 (Lagrangian dual) The Lagrangian dual of (1) is given by

$$\max d(\lambda, \mu) \quad \text{s.t.} \quad \lambda \geq 0$$

where the dual objective is given by $d : \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R} \cup \{+\infty\}$ and

$$d(\lambda, \mu) := \inf_{x \in \mathbb{R}^n} L(x, \lambda, \mu)$$

The function $p : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ given by

$$p(x) := \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(x, \lambda, \mu)$$

is called the **primal objective**.

Example 6.1.2 (LP duality) Consider the standard linear program (LP):

$$\begin{aligned} \min_{x \in \mathbb{R}^n} c^T x \quad \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned}$$

with $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, $c \in \mathbb{R}^n$.

The Lagrangian reads

$$\begin{aligned} L(x, \lambda, \mu) &= c^T x - \lambda^T x + \mu^T (b - Ax) \\ &= (c - \lambda - A^T \mu)^T x + b^T \mu \end{aligned} \tag{*}$$

So then

$$\nabla_x L(x, \lambda, \mu) = c - \lambda - A^T \mu$$

The function that takes $x \mapsto L(x, \lambda, \mu)$ is affine (from (*)), and in particular it is convex. And hence it takes its minimum if and only if $\nabla_x L(x, \lambda, \mu) = 0$, in which case,

$$\inf_{x \in \mathbb{R}^n} L(x, \lambda, \mu) = b^T \mu$$

otherwise if $\nabla_x L(x, \lambda, \mu) \neq 0$, the infimum must be $-\infty$.

So then

$$d(\lambda, \mu) = \begin{cases} b^T \mu & \text{if } c = A^T \mu + \lambda, \\ -\infty & \text{otherwise} \end{cases}$$

Therefore, the dual problem reads

$$\max_{\lambda, \mu} d(\lambda, \mu) \quad \text{s.t.} \quad \lambda \geq 0$$

Which is the same as

$$\max_{\lambda, \mu} b^T \mu \quad \text{s.t.} \quad \lambda \geq 0, A^T \mu + \lambda = c$$

and again,

$$\max_{\mu} b^T \mu \quad \text{s.t.} \quad A^T \mu \leq c$$

Theorem 6.2.1 (Weak duality) Let \hat{x} be feasible for (P) and $(\hat{\lambda}, \hat{\mu})$ be feasible for (D). Then

$$p(\hat{x}) \geq d(\hat{\lambda}, \hat{\mu})$$

Proof. We have

$$p(\hat{x}) = f(\hat{x}) \quad (\hat{x} \in X)$$

and hence

$$\begin{aligned} p(\hat{x}) &\geq f(\hat{x}) + \hat{\lambda}^T g(\hat{x}) + \hat{\mu}^T h(\hat{x}) \\ &= L(\hat{x}, \hat{\lambda}, \hat{\mu}) \\ &\geq \inf_{x \in \mathbb{R}^n} L(x, \hat{\lambda}, \hat{\mu}) \\ &= d(\hat{\lambda}, \hat{\mu}) \end{aligned}$$

Remark If $p(\hat{x}) = d(\hat{\lambda}, \hat{\mu})$, then \hat{x} solves (P), and $(\hat{\lambda}, \hat{\mu})$ solves (D). □

From weak duality, if we define

$$\bar{p} := \inf_{x \in \mathbb{R}^n} p(x) \geq \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} d(\lambda, \mu) =: \bar{d}$$

Then $\bar{p} - \bar{d} \geq 0$

Example 6.2.2 (Non-zero duality gap) Consider the following objective function:

$$\min f(x) := \begin{cases} x^2 - 2x & x \geq 0 \\ x & \text{otherwise} \end{cases} \quad \text{s.t.} \quad g(x) := -x \leq 0$$

The Lagrangian reads

$$L(x, \lambda) = \begin{cases} x^2 - (2 + \lambda)x & \text{if } x \geq 0 \\ (1 - \lambda)x & \text{otherwise} \end{cases}$$

A short computation shows that

$$d(\lambda) = \begin{cases} -\frac{(2+\lambda)^2}{4} & \text{if } \lambda \geq 1 \\ -\infty & \text{otherwise} \end{cases}$$

Therefore,

$$\bar{d} = d(1) = -\frac{9}{4} < 1 = \bar{p}$$

Hence the duality gap

$$\bar{p} - \bar{d} = \frac{5}{4} > 0$$

Definition 6.3.1 (Saddle point) The triple $(\bar{x}, \bar{\lambda}, \bar{\mu}) \in \mathbb{R}^n \times \mathbb{R}_+^m \times \mathbb{R}^p$ is called a saddle point of the Lagrangian L of (P) if

$$L(\bar{x}, \lambda, \mu) \leq L(\bar{x}, \bar{\lambda}, \bar{\mu}) \leq L(x, \bar{\lambda}, \bar{\mu})$$

Theorem 6.3.2 The following are equivalent:

- (i) $(\bar{x}, \bar{\lambda}, \bar{\mu})$ is a saddle point of (P)
- (ii) \bar{x} solves (P); $(\bar{\lambda}, \bar{\mu})$ solves (D)

Proof.

(i) \implies (ii): If $(\bar{x}, \bar{\lambda}, \bar{\mu})$ is a saddle point of (P), then

$$\begin{aligned}
 L(\bar{x}, \bar{\lambda}, \bar{\mu}) &\stackrel{\text{S.P.}}{\leq} \inf_x L(x, \bar{\lambda}, \bar{\mu}) \\
 &\leq \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} \inf_x L(x, \lambda, \mu) \\
 &\leq \inf_x \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(x, \lambda, \mu) \\
 &\leq \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(\bar{x}, \lambda, \mu) \\
 &\stackrel{\text{S.P.}}{\leq} L(\bar{x}, \bar{\lambda}, \bar{\mu})
 \end{aligned} \tag{*}$$

Then,

$$\begin{aligned}
 d(\bar{\lambda}, \bar{\mu}) &= \inf_x L(x, \bar{\lambda}, \bar{\mu}) \\
 &\stackrel{(*)}{=} \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(\bar{x}, \lambda, \mu) \\
 &= p(\bar{x}) = \bar{p} < +\infty
 \end{aligned}$$

Hence if $x \in X$, and by weak duality, \bar{x} solves (P), and $(\bar{\lambda}, \bar{\mu})$ solves (D).

(ii) \implies (i): Observe that

$$\begin{aligned}
 L(\bar{x}, \bar{\lambda}, \bar{\mu}) &\stackrel{\bar{x} \in X}{\leq} f(\bar{x}) \\
 &\stackrel{\bar{x} \in X}{=} p(\bar{x}) \\
 &\stackrel{\text{defn. of } p}{=} \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(\bar{x}, \lambda, \mu) \\
 &= d(\bar{\lambda}, \bar{\mu}) \\
 &= \inf_x L(x, \bar{\lambda}, \bar{\mu}) \\
 &\leq L(\bar{x}, \bar{\lambda}, \bar{\mu})
 \end{aligned}$$

But that's just the original LHS value, and hence all lines are equal. Hence

$$\begin{aligned}
 L(\bar{x}, \bar{\lambda}, \bar{\mu}) &= \inf_x L(x, \bar{\lambda}, \bar{\mu}) \\
 &= \sup_{\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}^p} L(\bar{x}, \lambda, \mu)
 \end{aligned}$$

And hence $(\bar{x}, \bar{\lambda}, \bar{\mu})$ is a saddle point. □

Consider again

$$\min f(x) \quad \text{s.t.} \quad x \in X \tag{1}$$

and a penalty function

$$P_\alpha^r := f + \alpha r, \quad \alpha > 0 \tag{2}$$

with $r \geq 0$, $r(x) = 0 \iff x \in X^r$

Definition 7.2.1 The penalty function P_α^r is called exact at a local min \bar{x} of (1) if there exists $\bar{\alpha}$ such that \bar{x} is a local min of P_α^r for all $\alpha > \bar{\alpha}$

Consider now the standard NLP

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad & g_i(x) \leq 0 \quad \forall i \in I \\ & h_j(x) = 0 \quad \forall j \in J \end{aligned} \quad (3)$$

with $f, g_i, h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ at least cont. diff.

A whole class of penalty functions in the sense of (2) for problem (3) is defined via

$$r_q(x) := \|(\max\{g(x), 0\}, h(x))\|_q$$

Where the max is interpreted component-wise, and then we're taking the q -norm. The value in (\cdot) is a vector with the first i elements being $\max\{g_i(x), 0\}$ and the last j elements being $h_j(x)$.

and

$$\|z\|_q = \begin{cases} \left(\sum_{i=1}^{\ell} (z_i)^q \right)^{\frac{1}{q}} & \text{if } q \in [1, \infty) \\ \max_{i=1, \dots, \ell} |z_i| & \text{if } q = +\infty \end{cases}$$

we focus on $q = 1$:

$$P_\alpha^1(x) = f(x) + \alpha \sum_{j=1}^p |h_j(x)| + \alpha \sum_{i=1}^m \max\{g_i(x), 0\}$$

Theorem 7.2.2 Let $(\bar{x}, \bar{\lambda}, \bar{\mu})$ be a KKT point of the convex NLP

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) \quad \text{s.t.} \quad & g_i(x) \leq 0 \quad \forall i \in I \\ & h_j(x) = 0 \quad \forall j \in J \end{aligned} \quad (4)$$

with $f, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ convex and cont. diff, and $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ affine (and hence convex).

Then $\bar{x} \in \operatorname{argmin}_X P_\alpha^1(x)$, for all $\alpha \geq \|(\bar{\lambda}, \bar{\mu})\|_\infty$.

In particular, P_α^1 is exact at \bar{x} if a CQ holds.

Proof. By Theorem 5.1.23 (KKT for convex problems), \bar{x} is a global minimizer of (4). Therefore, by Theorem 6.3.2 (Saddle point theorem), \bar{x} is a global minimizer of the Lagrangian $L(\cdot, \bar{\lambda}, \bar{\mu})$.

Therefore, for all $x \in \mathbb{R}^n$ and for all $\alpha \geq \|(\bar{\lambda}, \bar{\mu})\|_\infty$, we have

$$P_\alpha^1(\bar{x}) = f(\bar{x}) + \alpha \sum_{j=1}^p |h_j(\bar{x})| + \alpha \sum_{i=1}^m \max\{g_i(\bar{x}), 0\}$$

For reference,

$$\|(\bar{\lambda}, \bar{\mu})\|_\infty := \max\{|\bar{\lambda}_1|, \dots, |\bar{\lambda}_m|, |\bar{\mu}_1|, \dots, |\bar{\mu}_p|\}$$

But remember that since the point is feasible, the second and third terms are both zero and hence

$$\begin{aligned}
P_\alpha^1(\bar{x}) &= f(\bar{x}) \\
&\stackrel{\text{KKT}}{=} f(\bar{x}) + \bar{\lambda}^T g(\bar{x}) + \bar{\mu}^T h(\bar{x}) \\
&= L(\bar{x}, \bar{\lambda}, \bar{\mu}) \\
&\stackrel{\text{Thm 6.3.2}}{\leq} L(x, \bar{\lambda}, \bar{\mu}) \\
&= f(x) + \sum_{i=1}^m \bar{\lambda}_i g_i(x) + \sum_{j=1}^p \bar{\mu}_j h_j(x) \\
&\leq f(x) + \sum_{i=1}^m \bar{\lambda}_i \max\{g_i(x), 0\} + \sum_{j=1}^p |\bar{\mu}_j| \cdot |h_j(x)| \\
&\leq f(x) + \alpha \sum_{i=0}^m \max\{g_i(x), 0\} + \alpha \sum_{j=1}^p |h_j(x)| \\
&= P_\alpha^1(x)
\end{aligned}$$

Hence \bar{x} is the global minimizer of P_α^1 , that is for any $\alpha \geq \|(\bar{\lambda}, \bar{\mu})\|_\infty$

$$\bar{x} \in \underset{X}{\operatorname{argmin}} P_\alpha^1$$

□

SQP Methods

Consider

$$\min f(x) \quad \text{s.t.} \quad h_j(x) = 0 \quad \forall j \in J \quad (1)$$

with $f, h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ twice cont diff.

Define $\Phi : \mathbb{R}^n \times \mathbb{R}^p \rightarrow \mathbb{R}^n \times \mathbb{R}^p$ by

$$\Phi(x, \mu) := \begin{bmatrix} \nabla_x L(x, \mu) \\ h(x) \end{bmatrix}$$

where $L : \mathbb{R}^n \times \mathbb{R}^p \rightarrow \mathbb{R}$ is the Lagrangian of (1). Then

$$(x, \mu) \text{ is a KKT point of (1)} \iff \Phi(x, \mu) = 0 \quad (*)$$

where Φ is C^1 .

Idea: Apply Newton's method to $(*)$.

Algorithm 8.1.1 (Lagrange-Newton method)

(S0) Choose $(x^0, \mu^0) \in \mathbb{R}^n \times \mathbb{R}^p$ and set $k := 0$.

(S1) If $\Phi(x^k, \mu^k) = 0$: STOP

(S2) Determine $(\Delta x^k, \Delta \mu^k)$ as solution of

$$\Phi'(x^k, \mu^k) \begin{pmatrix} \Delta x \\ \Delta \mu \end{pmatrix} = -\Phi(x^k, \mu^k)$$

(S3) Put $(x^{k+1}, \mu^{k+1}) := (x^k, \mu^k) + (\Delta x^k, \Delta \mu^k)$, $k \leftarrow k + 1$, and go to (S1).

“Hessian is the Jacobian of the gradient”

Crucial part for well-definedness is to have Φ' be invertible at (x^k, μ^k)

Theorem 8.2.1 Let $(\bar{x}, \bar{\mu}) \in \mathbb{R}^n \times \mathbb{R}^p$ be a KKT point of (1), i.e. a root of the function $\Phi(x, \mu) = 0$ such that:

- (i) (LICQ) The vectors $\nabla h_j(x)$ are linearly independent, for $j \in J$
- (ii) (2nd order sufficient condition) We have

$$d^T \nabla_{xx}^2 L(\bar{x}, \bar{\mu}) d > 0 \quad \forall d \neq 0 \wedge \nabla h_j(\bar{x})^T d = 0$$

Proof. Goal is to show that kernel is trivial.

Sidenote:

$$h'(x)^T = [\nabla h_1(x) \quad \dots \quad \nabla h_p(x)] \in \mathbb{R}^{n \times p}$$

Observe that

$$\Phi(x, \mu) = \begin{pmatrix} \nabla_{xx}^2 L(x, \mu) & h'(x)^T \\ h'(x) & 0 \end{pmatrix} \quad \forall (x, \mu) \in \mathbb{R}^n \times \mathbb{R}^p$$

Hence,

$$\begin{aligned} \Phi'(\bar{x}, \bar{\mu}) \begin{pmatrix} q \\ r \end{pmatrix} &= 0 \\ \iff \nabla_{xx}^2 L(\bar{x}, \bar{\mu}) q + h'(\bar{x})^T r &= 0 \\ \text{and} \quad h'(\bar{x})^T q &= 0 \end{aligned}$$

Note: $\Phi' \in \mathbb{R}^{(n+p) \times (n+p)}$

□

Nov 30 class

Lagrange-Newton Equation Update with $(x^{k+1}, \mu^{k+1}) := (x^k, \mu^k) + (\Delta x^k, \Delta \mu^k)$

$$\begin{aligned} \Phi'(x^k, \mu^k) \begin{pmatrix} \Delta x \\ \Delta \mu \end{pmatrix} &= -\Phi(x^k, \mu^k) \\ \iff \begin{bmatrix} \nabla_{xx}^2 L(x^k, \mu^k) & h'(x^k)^T \\ h'(x^k) & 0 \end{bmatrix} \begin{pmatrix} \Delta x \\ \Delta \mu \end{pmatrix} &= - \begin{bmatrix} \nabla f(x^k) - h'(x^k)^T \\ h'(x^k) \end{bmatrix} \end{aligned}$$

Ordinary Differential Equations

First order differential equations

Result 1.1.1 (Separable)

$$\frac{dy}{dx} = \frac{p(x)}{q(y)}$$

To solve, we do

$$\int p(x) dx = \int q(y) dy$$

Result 1.1.2 (Homogeneous)

$$\frac{dy}{dx} = g\left(\frac{y}{x}\right)$$

An easier way to check if $\frac{dy}{dx}$ satisfies this form is by letting $f(x, y) := \frac{dy}{dx}$ and verifying that $f(x, y) = f(kx, ky)$.

To solve, let $u := y/x$ and rewrite the DE in terms of u and x .

Result 1.1.3 (Linear)

$$\frac{dy}{dx} + p(x)y = q(x)$$

To solve, let $\ln u := \int p(x) dx$ and jump to

$$\frac{d}{dx}uy = u \cdot q(x)$$

(which is just the product rule of differentiation)

Result 1.1.4 (Bernoulli)

$$\frac{dy}{dx} + p(x)y = q(x) \cdot y^n$$

If $n \in \{0, 1\}$, we have the **linear case**.

Use $u := y^{1-n}$, eliminate all ys , and reduce to a linear DE in u :

$$\frac{du}{dx} + (1-n)p(x)u = (1-n)q(x)$$

Result 1.1.5 (Riccati)

$$\frac{dy}{dx} = p(x)y^2 + q(x)y + r(x)$$

To solve, first find a basic solution y_1 . (By inspection, hopefully. Usually a solution will be given as part of the homework problem)

Then let $y_2 := y_1 + \frac{1}{u}$, substitute it into the original DE, and reduce it to a linear DE in u :

$$-u' = (2py_1 + q)u + p$$

Compute. First we obtain $y'_2 = y'_1 - \frac{u'}{u^2}$. Then substitute into the original DE:

$$\begin{aligned} y'_2 &= py_2^2 + qy_2 + r \\ y'_1 - \frac{u'}{u^2} &= p \left(y_1 + \frac{1}{u} \right)^2 + q \left(y_1 + \frac{1}{u} \right) + r \\ &= p \left(y_1^2 + \frac{2y_1}{u} + \frac{1}{u^2} \right) + q \left(y_1 + \frac{1}{u} \right) + r \\ &= (py_1^2 + qy_1 + r) + p \left(\frac{2y_1}{u} + \frac{1}{u^2} \right) + q \left(\frac{1}{u} \right) \end{aligned}$$

Since y_1 is a solution to the original DE,

$$\begin{aligned} -\frac{u'}{u^2} &= p \left(\frac{2y_1}{u} + \frac{1}{u^2} \right) + q \left(\frac{1}{u} \right) \\ -u' &= (2py_1 + q)u + p \end{aligned}$$

□

□

Result 1.1.6 (Exact)

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0$$

Criteria: $M_y = N_x$.

The idea is to work towards a function F where $F_x = M$, and $F_y = N$. (Because then based on the original DE we'll have $F_x = 0$)

To solve, integrate to find $F(x, y) = \int M(x, y) dx = \int N(x, y) dy$.

Then we have $F(x, y) = C$ for some constant C .

Second order differential equations

- $ay'' + by' + cy = g(t)$: **Method of undetermined coefficients**
- $ax^2y'' + bxy' + cy = 0$: **Euler equations**
- $y'' + p(x)y' + q(x)y = r(x)$: Variation of parameters for either a **particular** or **complementary** solution.

Definition 2.1.1 (Wronskian) The Wronskian of two differentiable functions f and g is $W(f, g) := fg' - gf'$.

More generally, for n complex-valued functions f_1, \dots, f_n which are $n - 1$ times differentiable on an interval I , the Wronskian $W(f_1, \dots, f_n)$ is a function on $x \in I$ defined by

$$W(f_1, \dots, f_n)(x) := \det \begin{bmatrix} f_1(x) & f_2(x) & \dots & f_n(x) \\ f'_1(x) & f'_2(x) & \dots & f'_n(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)}(x) & f_2^{(n-1)}(x) & \dots & f_n^{(n-1)}(x) \end{bmatrix}$$

Result 2.1.2 (Method of undetermined coefficients)

$$ay'' + by' + cy = g(t)$$

Define $\Sigma_{n,t}(A) := A_0t^n + A_1t^{n-1} + \dots + A_n$ and $P_n(t) := a_0t^n + a_1t^{n-1} + \dots + a_n$.

$g(t)$	$y(t)$
$P_n(t)$	$\Sigma_{n,t}(A)$
$P_n(t)e^{\alpha t}$	$t^s e^{\alpha t} \cdot \Sigma_{n,t}(A)$
$P_n(t)e^{\alpha t} \begin{cases} \sin \beta t \\ \cos \beta t \end{cases}$	$t^s [\Sigma_{n,t}(A)e^{\alpha t} \cos \beta t + \Sigma_{n,t}(B)e^{\alpha t} \sin \beta t]$

Here, s is the smallest non-negative integer that ensures that no term in $y(t)$ is a solution of the corresponding homogeneous equation.

Result 2.1.3 (Variation of parameters (particular: y_p)) import **Wronskian** for W .

$$y'' + p(x)y' + q(x)y = r(x)$$

Suppose we already know that y_1 and y_2 satisfy the corresponding homogeneous equation. Then we can jump to

$$y_p = v_1y_1 + v_2y_2$$

where $v_1 := \frac{-y_2r}{W(y_1, y_2)}$ and $v_2 := \frac{y_1r}{W(y_1, y_2)}$.

Result 2.1.4 (Variation of parameters (complementary: y_c))

$$y'' + p(x)y' + q(x)y = r(x)$$

Suppose we already know that y_1 is a solution. Then let

$$y_2 := vy_1$$

Then substituting y_2 back into the original DE, we have

$$v''y_1 + v'(2y_1' + py_1) = 0$$

Compute.

$$y_2 := vy_1; \quad y_2' = v'y_1 + vy_1'; \quad y_2'' = v''y_1 + 2v'y_1' + vy_1''$$

Substituting this back into the original DE:

$$(v''y_1 + 2v'y_1' + vy_1'') + p(v'y_1 + vy_1') + q(vy_1) = r$$

But since y_1 is known to be a solution:

$$\begin{aligned} (v''y_1 + 2v'y_1') + p(v'y_1) &= 0 \\ v''y_1 + v'(2y_1' + py_1) &= 0 \end{aligned}$$

□

□

Which is a first-order linear equation in v' . Use $u := v'$ to solve for u and then substitute everything back to find y .

Result 2.1.5 (Euler equations)

$$ax^2y'' + bxy' + cy = 0$$

Try $y = x^r$ for some $r \in \mathbb{C}$ to be found.

If two distinct roots: $y := Ax^{r_1} + Bx^{r_2}$.

If one distinct root: $y := Ax^r + B \ln(x)x^r$.

If complex roots ($r = \alpha \pm \beta i$): $y := Ax^\alpha \cos(\beta \ln x) + Bx^\alpha \sin(\beta \ln x)$

Higher order differential equations

$$y^{(n)} + p_1y^{(n-1)} + p_2y^{(n-2)} + \dots + p_{n-1}y' + p_ny = q$$

where $p_1, \dots, p_n, q : \mathbb{R} \rightarrow \mathbb{R}$.

For these we split into a few cases:

- **Constant coefficients**
- **Euler equations**
- **Variation of parameters**

Result 3.1.1 (Constant coefficients)

$$y^{(n)} + a_1y^{(n-1)} + a_2y^{(n-2)} + \dots + a_{n-1}y' + a_ny = 0 \quad (*)$$

where $a_1, \dots, a_n \in \mathbb{R}$ fixed.

Then try $y = e^{rt}$. Substituting that into (*), we'll obtain

$$a_0r^n + a_1r^{n-1} + \dots + a_n = 0$$

Since a_1, \dots, a_n are given, we can solve for r .

If all the roots of r are real and no two are equal, then we have n distinct solutions $e^{r_1t}, \dots, e^{r_nt}$ of equation (*).

Moreover, if these solutions are linearly independent, then the general solution to (*) is

$$y_g = c_1e^{r_1t} + c_2e^{r_2t} + \dots + c_ne^{r_nt}$$

One way to establish the linear independence is to evaluate their **Wronskian**.

Result 3.1.2 (Euler equations)

$$a_0x^n y^{(n)} + a_1x^{n-1}y^{(n-1)} + \dots + a_ny = 0 \quad (*)$$

where $a_1, \dots, a_n \in \mathbb{R}$ fixed.

Then try $y = x^r$. Substituting that into (*), we'll obtain

$$\sum_{i=0}^n \left(\prod_{j=r-i+1}^r j \right) a_i x^r = 0$$

Which is then

$$\sum_{i=0}^n \left(\prod_{j=r-i+1}^r j \right) a_i = 0$$

Result 3.1.3 (Variation of parameters)

Result 3.1.4 (Radius of convergence of power series) Consider the power series

$$\sum_{k=0}^n (ax)^k = 1 + ax + a^2x^2 + a^3x^3 + \dots$$

As $n \rightarrow \infty$,

$$\sum_{k=0}^n (ax)^k \begin{cases} \text{is convergent on } \frac{1}{1-ax} & \text{if } |ax| < 1 \\ \text{is divergent} & \text{otherwise} \end{cases}$$

We use this as a benchmark to check for convergence of an arbitrary power series:

$$\sum_{k=0}^{\infty} b_k x^k$$

where $b_1, b_2, \dots \in \mathbb{R}$.

If for all $k \in \mathbb{N}_0$ we have $|b_k| < a^k$, and the geometric series converges for x , then the arbitrary series converges too.

This is the same as requiring $(b_k)^{1/k} < a$ and $|x| < \frac{1}{|a|}$, and hence we obtain the radius of convergence of the arbitrary power series:

$$R := \frac{1}{\lim_{k \rightarrow \infty} (b_k)^{1/k}}$$

The series converges if $|x| < R$, and diverges if $|x| > R$.

Proposition 3.1.5 (Power series at zero) Suppose we have an arbitrary power series that is zero

$$\sum_{k=0}^n a_k x^k = 0$$

Then we must have

$$a_k = 0 \quad \forall (k \in \mathbb{N}_0)$$

Definition 3.1.6 (Laplace transform) The Laplace transform of a function $f(t)$, defined for all $t \in \mathbb{R}$, $t > 0$, is the function $F(s)$, a unilateral transform defined by

$$F(s) := \int_0^{\infty} f(t) e^{-st} dt$$

Here's a few fundamental Laplace transforms:

$f(t)$	$F(s)$
1	$\frac{1}{s}$
t	$\frac{1}{s^2}$
t^n	$\frac{n!}{s^{n+1}}$
$e^{\alpha t}$	$\frac{1}{s-\alpha}$
$te^{\alpha t}$	$\frac{1}{(s-\alpha)^2}$
$\cos(\beta t)$	$\frac{s}{s^2+\beta^2}$
$\sin(\beta t)$	$\frac{\beta}{s^2+\beta^2}$
$e^{\alpha t} \cos(\beta t)$	$\frac{s-\alpha}{(s-\alpha)^2+\beta^2}$
$e^{\alpha t} \sin(\beta t)$	$\frac{\beta}{(s-\alpha)^2+\beta^2}$

Let $\mathcal{L}(f) := \mathcal{L}\{f\}(s)$. Then we have

$$\mathcal{L}(f') = -f(0) + s\mathcal{L}(f)$$

Compute. Let's work through a few examples together.

Let $f(t) := 1$. Then

$$\mathcal{L}\{f\}(s) = \int_0^\infty e^{-st} dt = \left[-\frac{1}{s} e^{-st} \right]_0^\infty = \frac{1}{s}$$

Next, let $f(t) := t$. Then

$$\begin{aligned} \mathcal{L}\{f\}(s) &= \int_0^\infty te^{-st} dt \\ &= \left[-\frac{1}{s} e^{-st} \cdot t \right]_0^\infty - \int_0^\infty -\frac{1}{s} e^{-st} dt \\ &= \frac{1}{s^2} \end{aligned}$$

□

Exercise 3.1.7 Solve the differential equation

$$y'' + 2ty' - 4y = 5 \quad y(0) = 9, y'(0) = 0$$

Compute. Let $u(s) := \mathcal{L}(y)$

Let's start by preprocessing some Laplace transforms:

$$\begin{aligned} \mathcal{L}(y') &= -f(0) + su &= 9 + su \\ \mathcal{L}(y'') &= -f'(0) - sf(0) + s^2u = -9s + s^2u \end{aligned}$$

And then,

$$\mathcal{L}(ty') = -\frac{d}{ds}\mathcal{L}(y') = -u - su'$$

So then applying the Laplace transform on both sides:

$$\begin{aligned} (-9s + s^2u) + 2(-u - su') - 4(u) &= 5 \\ (-2s)u' + (s^2 - 6)u &= 9s + 5 \\ u' + \frac{6 - s^2}{2s}u &= -\frac{9}{2} - \frac{5}{2s} \\ u' + \left(\frac{3}{s} - \frac{s}{2}\right)u &= -\frac{9}{2} - \frac{5}{2s} \end{aligned} \quad (*)$$

This is a first-order linear equation in u .

$$\text{Let } \ln v := \int \left(\frac{3}{s} - \frac{s}{2}\right) ds$$

$$\ln v = 3 \ln s - \frac{s^2}{4} = \ln s^3 - \ln e^{(s^2/4)} = \ln(s^3 e^{-(s^2/4)})$$

$\Rightarrow v = s^3 e^{-(s^2/4)}$. Substituting back to $(*)$, we have

$$\begin{aligned} uv &= \int \frac{9s + 5}{-2s} \cdot s^3 e^{-(s^2/4)} ds \\ &= \int \left(-\frac{9}{2}s^2 - \frac{5}{2}s\right) e^{-(s^2/4)} ds \end{aligned}$$

$$\begin{aligned} \int x e^{-(x^2/4)} dx &= -2e^{-(x^2/4)} \\ \int x^2 e^{-(x^2/4)} dx &= x(-2e^{-(x^2/4)}) - \int -2e^{-(x^2/4)} dx \\ &= x(-2e^{-(x^2/4)}) + \int 2e^{-(x^2/4)} dx \end{aligned}$$

□